



## Antipsychotic agents FDA Guidelines for the dose and frequency

Antipsychotic Agent*	Maximum Adult Dose	Autism	Bipolar Disorder	MDD	Parkinson's Psychosis	"Psychotic Disorders"	Schizophrenia	Tourette's Disorder	Notes
<b>Ability Maintena® (Aripiprazole)</b>	400 mg/mo		A				A		Bipolar I maintenance monotherapy.
<b>Abilify Asimtufii (Aripiprazole)</b>	960 mg/2 mo		A				A		Bipolar I maintenance monotherapy.
<b>Abilify MyCite® (Aripiprazole)</b>			A	A			A		Bipolar mania or mixed monotherapy; Bipolar mania adjunct to mood stabilizer; MDD adjunct to antidepressant
<b>Adasuve® (Loxapine) inhalation powder</b>	10 mg/d		A				A		Acute tx
<b>Aripiprazole tabs, oral solution (Abilify®) and ODT (Abilify Discmelt®)</b>	30 mg/d	C (6-17 yo)	A, C (10-17 yo)	A			A, C (13-17 yo)	C (6-18 yo)	Bipolar mania monotherapy; Bipolar mania adjunct to mood stabilizer; MDD adjunct to antidepressant; Agitation associated with BP/Schizophrenia in adults – IM injection
<b>Aristada® (Aripiprazole Lauroxil)</b>	882mg/mo						A		Establish tolerability with oral Aripiprazole first.
<b>Aristada Initio® (Aripiprazole Lauroxil)</b>	675 mg X 1						A		Establish tolerability with oral Aripiprazole first. Only for initiation in combination with an oral preparation and Aristada.
<b>Asanapine (Saphris®)</b>	20 mg/d		A, C (10-17 yo)				A		Bipolar I
<b>Caplyta® (Lumateperone)</b>	42 mg/d		A				A		Depressive episodes
<b>Chlorpromazine (Thorazine®)</b>	1000 mg		A			A	A		Also, in Severe behavioral problems in children 1 to 12.
<b>Clozapine (Clozaril®)</b>	900 mg						A tx-resistant		Suicidality in pts with schizophrenia or schizoaffective d/o.
<b>Clozapine ODT (Fazaclor®)</b>	900 mg/d						A		Tx-resistant schizophrenia. Suicidal behavior in schizophrenia or schizoaffective disorder
<b>Fanapt® (Iloperidone)</b>	24 mg/d						A		Discontinue if QTc is more than 500 msec.
<b>Fluphenazine (Prolixin®) tabs, sol, inj</b>	40 mg/d					A	A		

\*Please click here to see current Preferred Drug List (PDL) status of the listed antipsychotic medication.

Antipsychotic Agent*	Maximum Adult Dose	Autism	Bipolar Disorder	MDD	Parkinson's Psychosis	"Psychotic Disorders"	Schizophrenia	Tourette's Disorder	Notes
<b>Fluphenazine Decanoate inj (Prolixin® Decanoate)</b>	100 mg/3-6 weeks					A	A		
<b>Geodon® IM (Ziprasidone Mesylate)</b>	40 mg/d						A		Acute tx of agitation
<b>Haloperidol (Haldol®) tabs, inj</b>	100 mg/ d					A, C (3-18 yo)	A	A, C (3-18 yo)	Also, in Acute agitation: A and in Severe behavioral disorders and agitation: 3-18
<b>Haloperidol Decanoate inj (Haldol® Decanoate)</b>	450 mg/mo						A		
<b>Invega® (Paliperidone)</b>	12 mg/d					A	A, C (12-17 yo)		Schizoaffective disorder
<b>Invega Hafyera® (Paliperidone)</b>	1560 mg/6 mo						A		After 4 months of Sustenna or 1 cycle of Trinza.
<b>Invega Sustenna® (Paliperidone)</b>	234 mg/mo					A	A		Schizoaffective Disorder
<b>Invega Trinza® (Paliperidone)</b>	819 mg/3 mo						A		After 4 months of Sustenna
<b>Latuda® (Lurasidone)</b>	160 mg/d		A, C (10-17 yo)				A, C (13-17 yo)		Depressive episode in Bipolar I
<b>Loxapine (Loxitane®)</b>	250 mg/d						A		
<b>Molindone (Moban®)</b>	225 mg/d						A, C (12-17 yo)		
<b>Nuplazid® (Pimavanserin)</b>	34 mg/d				A				
<b>Olanzapine (Zyprexa®)</b>	20 mg/d		A, C (13-17 yo)	A		A, C (13-17 yo)			Bipolar I – acute manic or mixed, maintenance.
									In Symbyax and in tx-resistant MDD
<b>Olanzapine IM (Zyprexa® IM)</b>	20 mg/d		A				A		Acute agitation
<b>Olanzapine ODT (Zyrexia Zydis®)</b>	20 mg/d		A, C (13-17 yo)				A, C (13-17 yo)		Bipolar I – acute manic or mixed, maintenance
<b>Olanzapine/Fluoxetine (Symbyax®)</b>	12/50 mg/d		A, C (10-17 yo)	A					Bipolar I acute depressive episodes
									Tx-resistant depression
<b>Perphenazine (Trilafon®)</b>	64 mg/d						A		Also, in nausea/vomiting in A.
<b>Perphenazine/Amitriptyline (Triavil®)</b>	32/200 mg/d			A					MDD with severe anxiety and or/agitation; MDD with anxiety and chronic physical disease
<b>Perseris® (Risperidone)</b>	120 mg/mo						A		Establish tolerability with oral Risperidone first
<b>Pimozide (Orap®)</b>	10 mg/d							A, C (2-18 yo)	EKG at baseline and periodically thereafter

\*Please click here to see current Preferred Drug List (PDL) status of the listed antipsychotic medication.

Antipsychotic Agent*	Maximum Adult Dose	Autism	Bipolar Disorder	MDD	Parkinson's Psychosis	"Psychotic Disorders"	Schizophrenia	Tourette's Disorder	Notes
Quetiapine IR (Seroquel®)	800 mg/d		A, C (10-17 yo)			A, C (13-17 yo)			Bipolar mania: A, 10-17. Bipolar depression: A.
Quetiapine ER (Seroquel® XR)	800 mg/d		A, C (10-17 yo)	A		A, C (13-17 yo)			Bipolar mania: A, 10-17. Bipolar depression and Bipolar mixed: A.
									MDD adjunctive with antidepressants
Rexulti® (Brexpiprazole)	4 mg/d			A		A, C (13-17 yo)			MDD adjunct to antidepressant
Risperdal Consta® (Risperidone)	50 mg/ 2 wk		A			A			Bipolar I maintenance tx
Risperdal M-Tab ODT® (Risperidone)	16 mg	C (5-17 yo)	A, C, adolescents			A, adolescents			Bipolar mania
Risperidone (Risperdal®) tabs and oral solution	16 mg	C (5-17 yo)	A, C (10-17 yo)			A, adolescents			Bipolar mania
Secuado® (Asenapine) transdermal	7.6 mg/ d					A			
Thioridazine (Mellaril®)	800 mg/d					A, C, adolescents			Tx refractory schizophrenia
									EKG and K level at baseline and periodically thereafter
Thiothixene (Navane®)	60 mg/d					A, C (6-12 yo) adolescents			
Trifluoperazine (Stelazine®)	40 mg/ d					A	A, C (6-12 yo) adolescents		Also, in Anxiety: A
Uzedy (Risperidone)	125 mg/mo 250 mg/2 mo					A			Establish tolerability with oral Risperidone first
Versacloz® (Clozapine) solution	900 mg/ d					A tx-resistant			Suicidality in pts with schizophrenia or schizoaffective d/o.
Vraylar® (Cariprazine)	6 mg/d		A			A			Bipolar I
Ziprasidone (Geodon®)	160 mg/d		A			A			Bipolar I manic/mixed acute tx
									Bipolar I maintenance adjunct to mood stabilizer
Zyprexa Relprevv® (Olanzapine)	300 mg/2 wk or 405mg/4wk					A			Establish tolerability with oral Olanzapine first

A: adult C: Children yo: years old

\*Please click here to see current Preferred Drug List (PDL) status of the listed antipsychotic medication.