



## Expedited Partner Therapy in Maryland for Chlamydia, Gonorrhea and Trichomoniasis

### RECOMMENDED PARTNER TREATMENT REGIMENS\*

<b>Chlamydia Only</b>	<input type="checkbox"/> Doxycycline (Vibramycin) 100 mg orally 2 times/day for 7 days  <i>NOTE: Substitute azithromycin (Zithromax) 1 gram orally in a single dose for partners who may be pregnant, or unlikely to adhere to a 7-day regimen</i>
<b>Gonorrhea Only</b>	<input type="checkbox"/> Cefixime (Suprax) 800 mg orally in a single dose
<b>Gonorrhea/Chlamydia Coinfection or Chlamydia Not Ruled Out</b>	<input type="checkbox"/> Cefixime (Suprax) 800 mg orally in a single dose <b>PLUS</b> <input type="checkbox"/> Doxycycline (Vibramycin) 100 mg orally 2 times/day for 7 days  <i>NOTE: Substitute azithromycin (Zithromax) 1 gram orally in a single dose for doxycycline for partners who may be pregnant, or unlikely to adhere to a 7-day regimen</i>
<b>Trichomoniasis</b>	<input type="checkbox"/> Metronidazole (Flagyl) 500 mg orally 2 times/day for 7 days (Female partners)  <input type="checkbox"/> Metronidazole (Flagyl) 2 g orally in a single dose (Male partners)  <input type="checkbox"/> Tinidazole (Tindamax) 2 g orally in a single dose (Alternative regimen for all non-pregnant partners)

\*Based on CDC 2021 STI Treatment Guidelines.  
<https://www.cdc.gov/std/treatment-guidelines/default.htm>  
Last reviewed August, 2024.