A large blue cargo ship is docked at a pier. The ship's hull is a vibrant blue with some rust spots. Stacks of shipping containers are visible on the deck. Two workers wearing white hard hats and high-visibility yellow safety vests are walking on the pier in the foreground. The background shows more of the pier and some industrial structures under a clear sky.

# Maryland Occupational Health Surveillance Report, 2000-2013

# Acknowledgements

For more information  
on this indicator or  
occupational health in  
Maryland:

[Maryland Occupational  
Health and Safety  
website](#)

Environmental Health  
Helpline:

1-866-703-3266

Environmental Health  
email:

[dhmh.envhealth@maryland.gov](mailto:dhmh.envhealth@maryland.gov)

## Maryland Department of Health and Mental Hygiene

**Clifford S. Mitchell, MS, MD, MPH**

Director, Environmental Health Bureau

**Ann Liu, PhD**

Chief Epidemiologist, Environmental Health Bureau

**Elisabeth Dissen, MSPH**

Epidemiologist, Environmental Health Bureau

This publication was supported by Cooperative Agreement Number 1U60OH010912 from the Centers for Disease Control and Prevention (CDC). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of CDC.



# Table of Contents

For more information  
on this indicator or  
occupational health in  
Maryland:

[Maryland Occupational  
Health and Safety  
website](#)

Environmental Health  
Helpline:

1-866-703-3266

Environmental Health  
email:

[dhmh.envhealth@maryland.gov](mailto:dhmh.envhealth@maryland.gov)

	Page
Executive Summary	5
Employment demographics profile	6
Indicator 1: Non-fatal injuries and illnesses reported by employers	9
Indicator 2: Work-related hospitalizations	10
Indicator 3: Fatal work-related injuries	13
Indicator 4: Amputations reported by employers	15
Indicator 6: Hospitalizations for work-related burns	16
Indicator 7: Musculoskeletal disorders reported by employers	18
Indicator 9: Pneumoconiosis hospitalizations	19
Indicator 10: Pneumoconiosis mortality	21
Indicator 11: Acute work-related pesticide poisonings reported to poison control centers	22
Indicator 12: Incidence of malignant mesothelioma	23
Indicator 14: Workers employed in industries with high risk for occupational morbidity	24
Indicator 15: Workers employed in occupations with high risk for occupational morbidity	26
Indicator 16: Workers employed in industries and occupations with high risk for occupational mortality	27
Indicator 17: Occupational safety and health professionals	28
Indicator 18: Occupational safety and health administration (OSHA) enforcement activities	29
Indicator 19: Workers' compensation awards	30

# Table of Contents

For more information on this indicator or occupational health in Maryland:

[Maryland Occupational Health and Safety website](#)

Environmental Health Helpline:

1-866-703-3266

Environmental Health email:

[dhmh.envhealth@maryland.gov](mailto:dhmh.envhealth@maryland.gov)

	Page
Indicator 20: Work-related low back disorder hospitalizations	31
Indicator 21: Asthma among adults caused or made worse by work	34
Indicator 22: Work-related severe traumatic injury hospitalizations	35
Indicator 24: Occupational heat-related ED visit	38
State Specific Indicator #1: Work-Related Emergency Department Visits	39

## Indicators not available in Maryland:

Indicator 5: Amputations identified in state workers' compensation systems

Indicator 8: Carpal tunnel syndrome cases identified in state workers' compensation systems

Indicator 13: Elevated blood lead levels among adults

Indicator 23: Influenza vaccination coverage among healthcare personnel (reporting will begin with 2014 data)



# Executive Summary

For more information on this indicator or occupational health in Maryland:

[Maryland Occupational Health and Safety website](#)

Environmental Health Helpline:

1-866-703-3266

Environmental Health email:

[dhmh.envhealth@maryland.gov](mailto:dhmh.envhealth@maryland.gov)

About 3 million people work in Maryland, and in 2013 more than 3 of every 100 workers experienced a work-related injury or illness, including a total of 79 work-related fatalities.

The Council of State and Territorial Epidemiologists (CSTE), in association with the National Institute of Occupational Safety and Health (NIOSH), has recommended that states conduct surveillance for a set of 24 occupational health indicators. Occupational health indicators are measures of health (work-related disease or injury) or factors associated with health (workplace exposures, hazards, or interventions) that allow a state to compare its health or risk status with that of other states and evaluate trends over time. These data can help guide priorities for prevention and intervention efforts. The indicators that have been developed represent the consensus view of state and NIOSH representatives and are intended as advisory to the states.

The 24 occupational health indicators include:

- **16 Health effect indicators** (measures of injury or illness that indicate adverse effects from exposure to known or suspected occupational hazards),
- **1 Exposure indicator** (measures of markers in human tissue or fluid that identify the presence of a potentially harmful substance resulting from exposure in the workplace),
- **4 Hazard indicators** (measures of potential for worker exposure to health and safety hazards in the workplace),
- **2 Intervention indicators** (measures of intervention activities or intervention capacity to reduce workplace health and safety hazards), and
- **1 Socioeconomic impact indicator** (measure of the economic impact of work-related injuries and illnesses).

Maryland currently tracks 20 indicators. Tracking occupational health indicators allows Maryland to:

- Measure baseline health of worker populations;
- Identify trends and patterns of work-related injury, illness, and death;
- Identify problem areas that require attention;
- Reduce preventable workplace injuries; and
- Increase consistency and availability of occupational disease and injury surveillance data.


Maryland has also elected to conduct surveillance for one state-specific occupational health indicator (Work-related emergency department visits), and has expanded surveillance for some indicators to include sub-state data and breakdowns by race and ethnicity.



# Employment demographics profile - civilian workforce

## Maryland State Occupational Health Indicators

Work-related injuries and illnesses are preventable, and control of occupational hazards is the most effective means of prevention. Estimating the burden and tracking these injuries can help target prevention programs and activities. Information on reported cases can be used to identify contributory factors and to develop improved or new prevention strategies or regulations to protect workers.



Year	Total employed	Unemployed	Self-employed	Employed part-time*
2000	2,697,000	3.9%	5.7%	14.6%
2001	2,722,000	4.1%	5.6%	15.3%
2002	2,772,000	4.4%	6.0%	15.8%
2003	2,773,000	4.5%	6.2%	15.8%
2004	2,762,000	4.2%	5.8%	15.5%
2005	2,800,000	4.2%	5.7%	14.9%
2006	2,886,000	3.9%	5.4%	15.0%
2007	2,852,000	3.6%	5.4%	15.1%
2008	2,875,000	4.2%	5.5%	15.5%
2009	2,821,000	7.1%	5.6%	16.0%
2010	2,819,000	7.7%	5.1%	16.5%
2011	2,873,000	7.0%	5.4%	16.0%
2012	2,903,000	7.0%	5.1%	16.5%
2013	2,919,000	6.7%	4.8%	15.9%

\* "Employed part-time" are individuals who work 1 to 34 hours per week. Employees are considered full-time if they work at least 35 hours per week.

Data Source for this Indicator: Bureau of Labor Statistics' Current Population Survey Geographic Profiles of Employment and Unemployment.

### About this Indicator:

#### Why is this Indicator Important?

The workforce in the United States is more diverse than ever. Age, race, sex, ethnicity, and levels of employment in certain industries/occupations varies across states. These characteristics can impact rates of work-related injury and illness.

#### Limitation of Indicator:

Variations occur regarding employer reporting compliance, accuracy and completeness, and employer utilization of restricted or light duty work for injured workers as a means of decreasing the number of days an injured worker is away from work. Employers may not be aware of work-related conditions for which employees obtained care. The industries for which data are available vary among states. It is not recommended to compare numbers between state or national data.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.



# Employment demographics profile - hours, sex, race/ethnicity

## About this Indicator:

### Why is this Indicator Important?

The workforce in the United States is more diverse than ever. Age, race, sex, ethnicity, and levels of employment in certain industries/occupations varies across states. These characteristics can impact rates of work-related injury and illness.

### Limitation of Indicator:

Variations occur regarding employer reporting compliance, accuracy and completeness, and employer utilization of restricted or light duty work for injured workers as a means of decreasing the number of days an injured worker is away from work. Employers may not be aware of work-related conditions for which employees obtained care. The industries for which data are available vary among states. It is not recommended to compare numbers between state or national data.


For more information on this indicator or occupational health in Maryland, visit the DHMH website.

## Maryland State Occupational Health Indicators


### Percent of civilian employment by number of hours worked, Maryland and U.S.



### Percent of civilian employment by sex, Maryland and U.S.



### Percent of civilian employment by race/ethnicity, Maryland and U.S.



Year	Civilian hours worked per week			Civilian employment by sex		Civilian employment by age group			Civilian employment by race/ethnicity			
	<40 hours	40 hours	41+ hours	Male	Female	16 to 17	18 to 64	65+	White	Black	Other	Hispanic origin*
2000	31.1%	38.0%	30.9%	52.6%	47.4%	2.5%	94.4%	3.1%	67.4%	29.3%	3.3%	4.8%
2001	33.3%	38.4%	28.4%	51.4%	48.6%	2.1%	94.5%	3.5%	68.1%	27.8%	4.1%	5.2%
2002	32.6%	38.2%	29.3%	51.4%	48.6%	1.8%	94.4%	3.7%	66.8%	27.6%	5.6%	5.8%
2003	32.0%	41.2%	26.8%	51.3%	48.7%	1.8%	94.3%	3.9%	66.2%	26.5%	7.2%	7.0%
2004	32.0%	42.4%	25.7%	51.8%	48.2%	2.0%	94.4%	3.7%	67.7%	26.2%	6.0%	8.0%
2005	31.4%	41.7%	27.0%	51.8%	48.2%	1.6%	94.6%	3.8%	67.0%	26.9%	6.1%	6.9%
2006	31.5%	42.6%	25.9%	51.6%	48.4%	1.7%	94.3%	4.1%	67.0%	26.8%	6.2%	8.1%
2007	31.6%	43.0%	25.5%	51.4%	48.6%	1.5%	94.2%	4.3%	65.7%	27.9%	6.3%	7.6%
2008	31.3%	44.6%	24.1%	50.7%	49.2%	1.5%	94.2%	4.3%	65.8%	28.3%	5.9%	7.7%
2009	34.2%	43.2%	22.5%	50.5%	49.5%	1.3%	94.3%	4.4%	65.1%	27.9%	7.0%	8.4%
2010	35.0%	43.2%	21.7%	51.5%	48.5%	1.1%	94.0%	4.9%	65.0%	27.5%	7.5%	8.2%
2011	32.0%	45.1%	23.0%	51.4%	48.6%	1.0%	93.9%	5.2%	64.5%	27.5%	8.0%	8.1%
2012	32.0%	44.6%	23.3%	50.3%	49.7%	0.9%	93.4%	5.7%	63.0%	28.6%	8.4%	9.3%
2013	30.7%	46.0%	23.3%	50.7%	49.3%	0.7%	93.3%	6.0%	61.4%	28.2%	10.4%	10.1%

\* Person's Identified as Hispanic may be of any race (white, black, other)

Data Source for this Indicator: Bureau of Labor Statistics' Current Population Survey Geographic Profiles of Employment and Unemployment.



# Employment demographics profile - industry/occupation

## Maryland State Occupational Health Indicators

### About this Indicator:

#### Why is this Indicator Important?

The workforce in the United States is more diverse than ever. Age, race, sex, ethnicity, and levels of employment in certain industries/occupations varies across states. These characteristics can impact rates of work-related injury and illness.

#### Limitation of Indicator:

Variations occur regarding employer reporting compliance, accuracy and completeness, and employer utilization of restricted or light duty work for injured workers as a means of decreasing the number of days an injured worker is away from work. Employers may not be aware of work-related conditions for which employees obtained care. The industries for which data are available vary among states. It is not recommended to compare numbers between state or national data.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.

### Percentage of civilian employment by industry\*, Maryland

Industries	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Mining	<0.05	<0.05	0.1	<0.05	<0.05	<0.05	0.1	<0.05	<0.05	<0.05	<0.05
Construction	7.8	8.5	8.6	9.2	8.1	8.3	7.6	7.1	7.4	7.3	7.4
Manufacturing-Durable Goods	3.5	3.4	3.3	3.1	3.3	3.0	3.0	2.9	2.4	2.9	2.5
Manufacturing-Nondurable Goods	2.3	2.4	2.5	2.0	2.0	2.1	1.7	1.9	2.1	1.9	2.2
Trade-Wholesale and Retail	14.0	13.5	12.3	11.2	12.0	12.4	11.5	11.8	11.2	11.4	12.3
Transportation/ utilities	4.5	4.5	4.5	4.8	4.4	4.1	4.2	4.5	4.0	4.1	4.1
Information	3.2	2.7	2.7	2.7	2.2	2.4	2.4	2.2	2.4	2.3	2.2
Financial activities	7.1	6.7	7.0	7.6	7.3	7.2	6.9	7.1	6.6	6.3	6.1
Professional and business services	13.3	14.0	13.7	15.2	15.2	14.4	14.8	15.1	14.8	14.8	15.4
Education / health services	21.4	21.4	21.3	21.4	21.5	22.5	23.0	23.3	23.8	23.5	22.9
Leisure / hospitality	6.5	6.6	7.1	6.5	6.9	7.6	7.9	7.8	8.0	7.7	7.6
Other services	5.4	5.4	5.6	5.4	5.4	5.5	5.1	4.9	5.6	5.4	5.6
Public administration	10.5	10.6	10.7	10.3	10.9	9.9	11.2	10.5	10.7	11.6	11.1
Agriculture	0.6	0.4	0.6	0.6	0.6	0.7	0.5	0.8	1.0	0.8	0.5

\* Industry list was updated for use from 2003 on.

### Percentage of civilian employment by occupation,\* Maryland

Occupations	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Management/business/financial operations	18.0	16.7	17.2	17.1	18.9	18.5	18.3	17.3	18.5	19.1	18.8
Professional/related	24.9	25.4	25.3	25.3	25.9	26.4	27.5	27.2	28.0	27.7	27.8
Service	15.1	15.6	15.7	15.5	15.0	15.7	16.4	16.7	17.1	16.4	16.4
Sales/related	10.3	11.1	10.1	9.3	9.4	9.8	9.5	9.7	9.4	9.0	9.6
Office/administrative support	14.0	13.2	13.9	14.0	13.5	12.4	12.6	13.3	11.9	12.3	12.2
Farming/fishing/forestry	0.3	0.3	0.4	0.2	0.3	0.3	0.3	0.3	0.4	0.4	0.3
Construction/extraction	5.7	6.2	6.2	6.9	5.9	5.6	5.3	5.2	5.0	4.7	4.9
Installation/maintenance/repair	2.8	2.9	3.1	3.5	3.5	3.5	2.9	3.3	2.8	2.9	2.9
Production	3.8	3.7	3.6	3.3	3.4	3.3	2.9	2.9	2.4	3.2	2.7
Transportation/material moving	4.9	4.9	4.6	4.9	4.4	4.6	4.4	4.1	4.5	4.3	4.4

\* Occupations list was updated for use from 2003 on.


Data Source for this Indicator: Bureau of Labor Statistics' Current Population Survey Geographic Profiles of Employment and Unemployment.




# Indicator #1: Non-fatal Work-related Injuries and Illnesses Reported by Employers

## Maryland State Occupational Health Indicators

Work-related injuries usually result from a single event, such as a fall, while work-related illnesses, such as asthma or asbestosis occur as the result of a longer-term exposure to a hazardous chemical, physical hazard or repeated stress or strain at work. Work-related injuries and illnesses are preventable, and control of occupational hazards is the most effective means of prevention.



**Indicator #1.1:** Rate of non-fatal work-related injuries and illnesses reported by employers, Maryland and U.S.



**Indicator #1.2:** Rate of non-fatal work-related injuries and illnesses involving days away from work reported by employers, Maryland and U.S.

Year	1.1. All Cases of Work-Related Injury and Illness		1.2. Work-Related Injury and Illness Cases Involving Days Away from Work		1.3. Work-Related Injury and Illness Cases Involving more than 10 Days Away from Work
	Number	Rate*	Number	Rate*	Number
2000	78,400	4,600	30,300	1,800	21,120
2001	73,600	4,300	27,000	1,600	-
2002	72,500	4,300	28,900	1,700	-
2003	68,600	4,100	28,000	1,700	11,190
2004	69,700	4,200	27,600	1,700	11,190
2005	72,700	4,200	26,400	1,500	9,410
2006	66,400	3,800	25,600	1,500	9,580
2007	65,700	3,700	23,800	1,300	9,190
2008	58,600	3,300	21,100	1,200	8,690
2009	56,700	3,300	20,500	1,200	8,430
2010	58,900	3,600	23,300	1,400	9,430
2011	48,700	3,000	18,200	1,100	7,380
2012	51,900	3,100	19,400	1,200	8,770
2013	51,500	3,000	19,400	1,100	8,790

\* Rate per 100,000 full time workers

Data Source for this Indicator: Bureau of Labor Statistics' Annual Survey of Occupational Injuries and Illnesses.

### About this Indicator:

#### Why is this Indicator Important?

Estimating the burden and tracking these injuries can help target prevention programs and activities. This information can be used to identify contributory factors and to develop prevention strategies or regulations to protect workers.

#### Limitation of Indicator:

The Survey of Occupational Injuries and Illnesses is conducted by the Bureau of Labor Statistics using a probability sample and not a census of all employers. It is based on injury and illness data reported by employers and is subject to sampling error. Military, self-employed individuals, farms with fewer than 11 employees, and Federal agencies are excluded from the survey.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.




# Indicator #2: Work-Related Hospitalizations

## Maryland State Occupational Health Indicators

Individuals hospitalized for work-related injuries and illnesses have some of the most serious and costly adverse work-related health conditions. Tracking of these hospitalizations can be used to document the burden of occupational injuries and illnesses, to design, target, and evaluate the impact of prevention efforts over time, and to identify previously recognized settings in which workers may continue to be at high risk.

**Indicator # 2: Rate of work-related hospitalizations, Maryland and U.S.**



**Indicator # 2: Work-related hospitalizations, Maryland**

Year	Number	Rate*
2000	3,665	135.9
2001	3,844	141.2
2002	3,650	131.7
2003	3,952	142.5
2004	3,641	131.8
2005	3,547	126.7
2006	3,413	118.3
2007	3,354	116.8
2008	3,195	111.1
2009	3,102	110.0
2010	2,932	104.0
2011	2,629	91.5
2012	2,742	94.5
2013	2,336	80.0

\* Rate per 100,000 workers

**Data Source for this Indicator:** Maryland Hospital Discharge Data (number of work-related hospitalizations); Bureau of Labor Statistics Current Population Survey (total number of employed persons). A condition was considered work-related if workers' compensation was listed as primary payer in the hospital discharge data.

### About this Indicator:

#### Why is this Indicator Important?

Information on work-related injury and illness hospitalizations can be used to document the burden of occupational injuries and illnesses, to design, target, and evaluate the impact of prevention efforts over time, and identify settings in which workers may continue to be at high risk.

#### Limitation of Indicator:

Different factors may affect decisions by health care providers to hospitalize patients, to correctly diagnose work-related conditions, and/or to list the condition as a discharge diagnosis. All admissions are counted, including multiple admissions for a single individual. Because hospital discharge data is not available in all states, nationwide estimates are incomplete.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.



# Indicator #2: Work-Related Hospitalizations - Sub-State Data

## About this Indicator:

### Why is this Indicator Important?

Information on work-related injury and illness hospitalizations can be used to document the burden of occupational injuries and illnesses, to design, target, and evaluate the impact of prevention efforts over time, and identify settings in which workers may continue to be at high risk.


### Limitation of Indicator:

Different factors may affect decisions by health care providers to hospitalize patients, to correctly diagnose work-related conditions, and/or to list the condition as a discharge diagnosis. All admissions are counted, including multiple admissions for a single individual. Because hospital discharge data is not available in all states, nationwide estimates are incomplete.


For more information on this indicator or occupational health in Maryland, visit the DHMH website.

## Maryland State Occupational Health Indicators


**Indicator # 2: 2009-2013 Average annual crude rate of work-related hospitalizations by ZIP code per 100,000 employed persons**



**Indicator # 2: 2011-2013 Average annual crude rate of work-related hospitalizations by county per 100,000 employed persons**



**Indicator # 2: 2011-2013 Average annual crude rate of work-related hospitalizations by region per 100,000 employed persons**



**Data Source for this Indicator:** Maryland Hospital Discharge Data (number of work-related hospitalizations). County and region level denominator: Bureau of Labor Statistics Current Population Survey (total number of employed persons). ZIP code level denominator: The Nielson Company. A condition was considered work-related if workers' compensation was listed as primary payer in the hospital discharge data.



# Indicator #2: Work-Related Hospitalizations - Race and Ethnicity Data

## About this Indicator:

### Why is this Indicator Important?


Information on work-related injury and illness hospitalizations can be used to document the burden of occupational injuries and illnesses, to design, target, and evaluate the impact of prevention efforts over time, and identify settings in which workers may continue to be at high risk.

### Limitation of Indicator:


Different factors may affect decisions by health care providers to hospitalize patients, to correctly diagnose work-related conditions, and/or to list the condition as a discharge diagnosis. All admissions are counted, including multiple admissions for a single individual. Because hospital discharge data is not available in all states, nationwide estimates are incomplete.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.


## Maryland State Occupational Health Indicators



### Indicator # 2: 2009-2013 Average annual crude rate of work-related hospitalizations by county per 100,000 black employed persons



### Indicator # 2: 2009-2013 Average annual crude rate of work-related hospitalizations by county per 100,000 white employed persons




Data Source for this Indicator: Maryland Hospital Discharge Data (number of work-related hospitalizations); denominator: Bureau of Labor Statistics Current Population Survey (total number of employed persons). A condition was considered work-related if workers' compensation was listed as primary payer in the hospital discharge data.

# Indicator #3: Fatal Work-Related Injuries

## Maryland State Occupational Health Indicators

► A fatal work-related injury is an injury occurring at work that results in death. The Bureau of Labor Statistics conducts the Census of Fatal Occupational Injuries (CFOI), using multiple data sources to provide counts of all fatal work-related injuries in every state. Fatalities resulting from non-intentional injuries (i.e., falls, acute poisonings, and motor vehicle crashes that occurred during travel for work) and intentional injuries (i.e., homicides and suicides) that occurred at work are included in this measure. Fatalities that occur during a person's commute to or from work are not counted.

**Indicator # 3: Rate of fatal work-related injuries, Maryland and U.S.**



**Indicator # 3: Fatal work-related injuries, Maryland**

Year	Number	Rate*
2000	84	3.2
2001	64	2.5
2002	102	3.8
2003	92	3.4
2004	81	3.0
2005	95	3.5
2006	106	3.8
2007	82	3.0
2008	60	2.2
2009	65	2.5
2010	71	2.7
2011	71	2.6
2012	72	2.6
2013	79	2.8

\* Annual crude fatality rate per 100,000 full-time workers age 16 years or older

**Data Source for this Indicator:** Census of Fatal Occupational Injuries (numbers of fatalities); Bureau of Labor Statistics Current Population Survey Data (employment statistics used to calculate rates).

### About this Indicator:

#### Why is this Indicator Important?

Many factors contribute to work-related fatalities, including workplace/process design, organization, worker characteristics, economics and other social factors. Surveillance of work-related fatalities can identify hazards and clusters, leading to the development of interventions and new or revised regulations to protect workers.

#### Limitation of Indicator:

Fatalities of people younger than 16 may be included in the numerator but not in the denominator, since employment statistics are only available for those 16 years of age and older. Also, CFOI reports data on work-related fatalities by the state in which the fatal incident occurred, which is not always the state of residence.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.



# Indicator #3: Fatal Work-Related Injuries - Race and Ethnicity Data

## About this Indicator:

### Why is this Indicator Important?

Many factors contribute to work-related fatalities, including workplace/process design, organization, worker characteristics, economics and other social factors. Surveillance of work-related fatalities can identify hazards and clusters, leading to the development of interventions and new or revised regulations to protect workers.


### Limitation of Indicator:

Fatalities of people younger than 16 may be included in the numerator but not in the denominator, since employment statistics are only available for those 16 years of age and older. Also, CFI reports data on work-related fatalities by the state in which the fatal incident occurred, which is not always the state of residence.


For more information on this indicator or occupational health in Maryland, visit the DHMH website.

## Maryland State Occupational Health Indicators

**Indicator #3: 2013 Crude rate of fatal work-related injuries by race and ethnicity, Maryland**



**Indicator #3: 2003-2013 Crude rate of fatal work-related injuries by race and ethnicity, Maryland**



**Indicator # 3: Fatal work-related injuries by race and ethnicity, Maryland**

Year	White (non-Hispanic)		Black (non-Hispanic)		Asian (non-Hispanic)		Hispanic (any race)	
	Number FTEs	Crude Rate*	Number FTEs	Crude Rate*	Number FTEs	Crude Rate*	Number FTEs	Crude Rate*
2003	1,600,351	3.1	698,359	3.7	150,823	3.3	189,914	5.8
2004	1,596,394	2.7	684,568	2.3	124,121	4.0	219,332	7.8
2005	1,623,865	3.4	726,358	2.9	135,327	6.7	192,411	4.2
2006	1,639,537	3.4	732,869	3.1	134,897	2.2	225,450	9.8
2007	1,601,044	2.7	764,974	3.4	141,425	4.2	209,666	3.3
2008	1,591,098	2.3	769,050	1.7	136,156	0.0	214,873	4.7
2009	1,505,511	3.1	733,803	1.8	146,300	2.1	218,296	1.4
2010	1,494,999	2.9	704,317	1.8	145,960	0.0	211,870	5.7
2011	1,549,143	2.1	721,838	3.7	178,120	1.7	211,766	3.8
2012	1,541,911	2.4	729,653	2.5	179,172	0.0	251,759	6.0
2013	1,517,149	2.8	743,481	2.4	208,185	1.4	278,665	5.4

\* Annual Crude Fatality Rate per 100,000 FTEs, Age 16 Years or Older

**Data Source for this Indicator:** Census of Fatal Occupational Injuries (numbers of fatalities); Bureau of Labor Statistics Current Population Survey Data (employment statistics used to calculate rates).




# Indicator #4: Amputations Reported by Employers

## Maryland State Occupational Health Indicators

► An amputation is defined as full or partial loss of a protruding body part - an arm, hand, finger, leg, foot, toe, ear, or nose. An amputation may greatly reduce a worker's job skills and earning potential as well as significantly affect general quality of life.

**Indicator # 4: Rate of work-related amputations involving days away from work reported by private sector employers, Maryland and U.S.**



**Indicator # 4: Work-related amputations with days away from work reported by employers, Maryland**

Year	Number	Rate**
2000	73	4
2001	58	3
2002	97	6
2003	230	14
2004	170	10
2005	60	3
2006	100	5
2007	110	6
2008	30	2
2009	70	4
2010	30	2
2011*	90	5
2012	210	13
2013	80	5

\* Revised version of the Occupational Injuries and Illnesses Classification System (OIICS)

\*\* Incidence Rate per 100,000 full-time workers

### About this Indicator:

#### Why is this Indicator Important?

Work-related amputations are a preventable serious injury, and control of occupational hazards is the most effective means of prevention. Information on reported cases can be used to identify contributory factors and to develop improved or new prevention strategies or regulations to protect workers.

#### Limitation of Indicator:

The Survey of Occupational Injuries and Illnesses is conducted by the Bureau of Labor Statistics using a probability sample and not a census of all employers. It is based on injury and illness data reported by employers and is subject to sampling error. Military, self-employed individuals, farms with fewer than 11 employees, and Federal agencies are excluded from the survey.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.




Data Source for this Indicator: Bureau of Labor Statistics' Annual Survey of Occupational Injuries and Illnesses.

# Indicator #6: Hospitalizations for Work-Related Burns

## Maryland State Occupational Health Indicators

► Burns include injuries to tissues caused by contact with dry heat (fire), moist heat (steam), chemicals, electricity, friction, or radiation. Burns are among the most expensive work-related injuries to treat and can result in significant disability. Thermal and chemical burns are the most frequent types of work-related burn injury. A substantial proportion of burns occur in the service industry, especially in food service, often disproportionately affecting working adolescents.

**Indicator # 6: Rate of hospitalizations for work-related burns, Maryland and U.S.**



**Indicator # 6: Hospitalizations for work-related burns, Maryland**

Year	Number	Rate*
2000	17	0.6
2001	17	0.6
2002	40	1.4
2003	57	2.1
2004	52	1.9
2005	48	1.7
2006	74	2.6
2007	65	2.3
2008	55	1.9
2009	55	2.0
2010	44	1.6
2011	45	1.6
2012	43	1.5
2013	34	1.2

\* Rate per 100,000 workers

**Data Source for this Indicator:** Maryland Hospital Discharge Data (number of work-related hospitalizations); Bureau of Labor Statistics Current Population Survey (total number of employed persons).

### About this Indicator:

#### Why is this Indicator Important?

Although hospitalized burns are unusual events, they are painful, disabling, and expensive to treat. Many result in significant disfigurement.

#### Limitation of Indicator:

These data include inpatient hospitalizations of individuals who were hospitalized in acute care hospitals. These data are based only on primary discharge diagnosis codes, and do not include individuals who were seen by an Emergency Department, but not admitted to the hospital. Hospital Discharge records are only available for non-federal, acute care hospitals.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.



# Indicator #6: Hospitalizations for Work-Related Burns - Sub-State Data

## Maryland State Occupational Health Indicators

### About this Indicator:

#### Why is this Indicator Important?


Although hospitalized burns are unusual events, they are painful, disabling, and expensive to treat. Many result in significant disfigurement.

#### Limitation of Indicator:


These data include inpatient hospitalizations of individuals who were hospitalized in acute care hospitals. These data are based only on primary discharge diagnosis codes, and do not include individuals who were seen by an Emergency Department, but not admitted to the hospital. Hospital Discharge records are only available for non-federal, acute care hospitals.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.

**Indicator # 6: 2009-2013 Average annual crude rate of hospitalizations for work-related burns by county per 100,000 workers**



**Indicator # 6: 2009-2013 Average annual crude rate of hospitalizations for work-related burns by region per 100,000 workers**




Data Source for this Indicator: Maryland Hospital Discharge Data (number of work-related hospitalizations); Bureau of Labor Statistics Current Population Survey (total number of employed persons).



# Indicator #7: Musculoskeletal Disorders Reported by Employers

## Maryland State Occupational Health Indicators

Work-related musculoskeletal disorders (MSDs) are injuries or disorders of muscles, tendons, nerves, ligaments, joints, or spinal discs that are caused or aggravated by work activities. Workplace risk factors for MSDs include repetitive forceful motions, awkward postures, use of vibrating tools or equipment, and manual handling of heavy, awkward loads. These disorders also can be caused by single, traumatic events such as falls.



**Indicator # 7.1:** Rates of work-related musculoskeletal disorders involving days away from work reported by private sector employers, Maryland and U.S.

**Indicator # 7.2:** Rates of work-related disorders of the neck, shoulder, and upper extremities involving days away from work reported by private sector employers, Maryland and U.S.

**Indicator # 7.3:** Rates of work-related Carpal Tunnel Syndrome cases involving days away from work reported by private sector employers, Maryland and U.S.

**Indicator # 7.4:** Rates of work-related disorders of the back involving days away from work reported by private sector employers, Maryland and U.S.

**Indicator # 7: Work-related musculoskeletal disorders with days away from work reported by employers, Maryland**

Year	1.All Musculoskeletal Disorders		2. Disorders of the Neck, Shoulder, and Upper Extremities		3.Carpal Tunnel Syndrome Cases		4. Disorders of the Back	
	Number	Rate*	Number	Rate*	Number	Rate*	Number	Rate*
2000	9,075	530	1,962	115	160	9	4,509	263
2001	9,053	558	2,201	130	199	12	4,699	276
2002	7,625	456	1,933	115	108	7	3,747	224
2003	7,740	462	1,750	104	160	10	3,760	225
2004	8,300	506	2,280	139	420	25	3,630	221
2005	7,020	406	1,520	88	150	9	3,590	208
2006	6,840	390	1,830	104	110	7	3,040	173
2007	6,240	351	1,440	82	100	6	3,050	172
2008	5,690	322	1,320	74	50	3	7,890	163
2009	5,370	313	1,210	70	50	3	2,830	165
2010	5,950	362	1,490	90	110	7	3,000	182
2011	5,650	347	1,250	77	30	2	2,720	167
2012	5,880	349	1,440	86	60	4	2,870	170
2013	5,430	319	1,480	87	30	2	2,490	146

\* Rate per 100,000 full-time workers

**Data Source for this Indicator:** Bureau of Labor Statistics' Annual Survey of Occupational Injuries and Illnesses.

### About this Indicator:

#### Why is this Indicator Important?

Work-related musculoskeletal disorders are preventable and control of occupational hazards is the most effective means of prevention. Estimating the burden and tracking these injuries helps target prevention programs and activities.

#### Limitation of Indicator:

The Survey of Occupational Injuries and Illnesses is conducted by the Bureau of Labor Statistics using a probability sample and not a census of all employers. It is based on injury and illness data reported by employers and is subject to sampling error. Military, self-employed individuals, farms with fewer than 11 employees, and Federal agencies are excluded from the survey.


For more information on this indicator or occupational health in Maryland, visit the DHMH website.



# Indicator #9: Pneumoconiosis Hospitalizations

## Maryland State Occupational Health Indicators

► Pneumoconiosis is a term for a class of non-malignant lung diseases caused by the inhalation of mineral dust, nearly always in occupational settings. Most cases of pneumoconiosis develop only after many years of cumulative exposure; thus they are usually diagnosed in older individuals, often long after the onset of exposure. Pneumoconiosis includes: silicosis, asbestosis, coal workers' pneumoconiosis (CWP), and, less commonly, pneumoconiosis due to a variety of other mineral dusts, including talc, aluminum, bauxite, and graphite.



**Indicator #9.1:** Age-standardized rate of hospitalizations from or with Pneumoconiosis, Maryland and U.S.

**Indicator #9.2:** Age-standardized rate of hospitalizations from or with Coal Workers' Pneumoconiosis, Maryland and U.S.

**Indicator #9.3:** Age-standardized rate of hospitalizations from or with Asbestosis, Maryland and U.S.

**Indicator #9.4:** Age-standardized rate of hospitalizations from or with Silicosis, Maryland and U.S.

**Indicator #9.5:** Age-standardized rate of hospitalizations from or with other or unspecified Pneumoconiosis, Maryland and U.S.

Year	1. Total Pneumoconiosis			2. Coal Workers' Pneumoconiosis			3. Asbestosis			4. Silicosis			5. Other and Unspecified Pneumoconiosis		
	Number	Age-Adjusted Rate*	Adjusted Rate*	Number	Age-Adjusted Rate*	Adjusted Rate*	Number	Age-Adjusted Rate*	Adjusted Rate*	Number	Age-Adjusted Rate*	Adjusted Rate*	Number	Age-Adjusted Rate*	Adjusted Rate*
2000	1,114	266.9	291.6	101	24.2	26.8	988	236.7	258.2	17	4.1	4.6	10	2.4	2.6
2001	1,228	290.1	315.3	76	18.0	19.5	1,125	265.8	288.9	24	5.7	6.0	9	2.1	2.4
2002	1,413	329.1	355.2	92	21.4	23.1	1,292	300.9	325.5	25	5.8	6.0	9	2.1	2.0
2003	1,320	303.6	328.1	68	15.6	16.7	1,218	280.1	303.1	25	5.7	6.1	11	2.5	2.7
2004	1,307	297.3	319.1	51	11.6	12.4	1,233	280.5	301.3	18	4.1	4.3	9	2.0	2.1
2005	1,434	322.6	346.4	59	13.3	14.0	1,339	301.2	324.1	26	5.8	6.0	11	2.5	2.5
2006	1,295	288.8	308.3	43	9.6	10.0	1,228	273.8	292.8	18	4.0	4.1	9	2.0	2.1
2007	1,195	264.7	281.5	63	14.0	14.4	1,097	243.0	259.0	23	5.1	5.3	15	3.3	3.4
2008	1,415	311.4	326.3	59	13.0	13.1	1,334	293.5	308.2	19	4.2	4.4	10	2.2	2.3
2009	1,129	246.3	255.0	43	9.4	9.3	1,057	230.6	239.8	18	3.9	3.6	14	3.1	3.0
2010	1,016	217.3	221.2	47	10.1	10.3	946	202.4	206.3	15	3.2	3.1	13	2.8	3.6
2011	991	193.1	213.9	33	7.0	6.9	928	196.7	201.3	19	4.0	3.6	12	2.5	2.5
2012	908	190.2	190.2	34	7.1	7.1	852	178.5	178.9	15	3.1	2.9	**	**	**
2013	785	163.1	160.3	40	8.3	8.2	727	151.0	148.5	**	**	**	**	**	**

\* Rate of hospitalizations per million residents

\*\* HSCRC data counts <11 suppressed

**Data Source for this Indicator:** Maryland Hospital Discharge Data (number of work-related hospitalizations); U.S. Census Bureau (population statistics to calculate rates). 19

### About this Indicator:

#### Why is this Indicator Important?

Tracking of pneumoconiosis is essential for measuring progress towards elimination of the disease, as well as for targeting prevention and disease management programs.

#### Limitation of Indicator:

These data are based on primary discharge diagnosis codes for patients admitted to a hospital, and do not include individuals who were seen by an Emergency Department, but not admitted. Hospital Discharge records are only available for non-federal, acute care hospitals.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.



# Indicator #9: Pneumoconiosis Hospitalizations - Race Data

## Maryland State Occupational Health Indicators

### About this Indicator:

#### Why is this Indicator Important?


Tracking of pneumoconiosis is essential for measuring progress towards elimination of the disease, as well as for targeting prevention and disease management programs.

#### Limitation of Indicator:

These data are based on primary discharge diagnosis codes for patients admitted to a hospital, and do not include individuals who were seen by an Emergency Department, but not admitted. Hospital Discharge records are only available for non-federal, acute care hospitals.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.

**Indicator # 9: 2013 number of hospitalizations from or with Pneumoconiosis by race, Maryland**



**Indicator # 9: 2013 hospitalizations from or with Pneumoconiosis by race, Maryland**

Year	Total Pneumoconiosis				Asbestosis			
	White		Black		White		Black	
	Number	Crude Rate*	Number	Crude Rate*	Number	Crude Rate*	Number	Crude Rate*
2003	570	186.9	120	81.7	529	173.5	111	75.6

\* Rate per million residents




Data Source for this Indicator: Maryland Hospital Discharge Data (number of work-related hospitalizations); U.S. Census Bureau (population statistics to calculate rates).


# Indicator #10: Pneumoconiosis Mortality

## Maryland State Occupational Health Indicators

► Pneumoconiosis is a term for a class of non-malignant lung diseases caused by the inhalation of mineral dust, nearly always in occupational settings. Most cases of pneumoconiosis develop only after many years of cumulative exposure; thus they are often diagnosed in older individuals, long after the onset of exposure. These diseases, which include silicosis and asbestosis, are incurable and may ultimately result in death.



**Indicator #10.1:** Age-standardized mortality rate from or with pneumoconiosis, Maryland and U.S.



**Indicator #10.3:** Age-standardized mortality rate from or with asbestosis, Maryland and U.S.

**Indicator #10: Mortality from or with Pneumoconiosis, Maryland**

Year	1. Total pneumoconiosis deaths			2. Coal workers' pneumoconiosis deaths			3. Asbestosis deaths			4. Silicosis deaths			5. Other and unspecified pneumoconiosis deaths		
	Number	Crude Rate*	Age-Adjusted Rate*	Number	Crude Rate*	Age-Adjusted Rate*	Number	Crude Rate*	Age-Adjusted Rate*	Number	Crude Rate*	Age-Adjusted Rate*	Number	Crude Rate*	Age-Adjusted Rate*
2000	46	11.0	12.4	<5	-	-	42	10.1	11.3	<5	-	-	<5	-	-
2001	46	10.9	12.3	<5	-	-	43	10.2	11.5	0	0.0	0.0	0	0.0	0.0
2002	56	13.0	14.5	<5	-	-	45	10.5	11.7	6	1.4	1.5	<5	-	-
2003	46	10.6	11.7	<5	-	-	37	8.5	9.4	<5	-	-	5	1.1	1.2
2004	59	13.4	14.7	<5	-	-	53	12.1	13.1	<5	-	-	<5	-	-
2005	36	8.1	8.9	0	0.0	0.0	35	7.9	8.7	<5	-	-	0	0.0	0.0
2006	55	12.3	13.4	<5	-	-	52	11.6	12.7	0	0.0	0.0	<5	-	-
2007	49	10.9	11.8	<5	-	-	45	10.0	10.9	0	0.0	0.0	<5	-	-
2008	49	10.8	11.5	<5	-	-	44	9.7	10.4	<5	-	-	<5	-	-
2009	45	9.8	9.8	<5	-	-	40	8.7	8.8	0	0.0	0.0	<5	-	-
2010	29	6.2	6.4	<5	-	-	26	5.6	5.7	0	0.0	0.0	<5	-	-
2011	33	7.0	7.4	0	0.0	0.0	32	6.8	7.2	0	0.0	0.0	<5	-	-
2012	31	6.5	6.5	<5	-	-	25	5.2	5.3	<5	-	-	<5	-	-
2013	27	5.6	5.6	<5	-	-	24	5.0	5.0	<5	-	-	0	0.0	0.0

\* Deaths per million residents, rates not calculated for types with less than five deaths

Data Source for this Indicator: Maryland Vital Statistics Records (number of deaths); U.S. Census Bureau (population statistics to calculate rates).



**About this Indicator:**  
**Why is this Indicator Important?**  
 Tracking of pneumoconiosis is essential for tracking progress towards elimination of the disease, as well as for targeting prevention and disease management programs.

**Limitation of Indicator:**  
 Because pneumoconioses are typically chronic diseases with a long latency, current incidence is not necessarily indicative of current exposures, and it may be several years before reductions in exposures affect mortality. In addition, people may not die in the state where they were exposed.


For more information on this indicator or occupational health in Maryland, visit the DHMH website.

# Indicator #11: Acute Work-Related Pesticide Poisonings Reported to Poison Control Centers

## Maryland State Occupational Health Indicators

▶ A pesticide is a substance or mixture of substances used to prevent or control undesired insects, plants, animals, or fungi. Although the value of pesticides in protecting the food supply and controlling disease vectors is well recognized, it is also recognized that pesticides can cause harm to people and the environment. Adverse health effects from exposure vary depending on the amount and route of exposure and the type of chemical used. Agricultural workers and pesticide applicators are at greatest risk for the more severe pesticide poisonings.

**Indicator # 11: Rate of work-related pesticide associated poisonings, Maryland and U.S.**



**Indicator # 11: Acute work-related pesticide associated illness and injury reported to Poison Control Centers, Maryland**

Year	Number	Rate*
2000	39	1.5
2001	40	1.5
2002	32	1.2
2003	40	1.4
2004	56	2.0
2005	34	1.2
2006	48	1.7
2007	37	1.3
2008	33	1.2
2009	42	1.5
2010	57	2.0
2011	67	2.3
2012	43	1.5
2013	83	2.8

\* Annual incidence rate per 100,000 employed persons age 16 years or older

**Data Source for this Indicator:** American Association of Poison Control Centers (Numbers of pesticide-associated illness and injury); Bureau of Labor Statistics Current Population Survey (total number of employed persons).

### About this Indicator:

#### Why is this Indicator Important?

Tracking pesticide poisoning data can be useful in targeting prevention efforts.

#### Limitation of Indicator:

Poison Control Centers (PCC) capture only a small proportion of acute occupational pesticide-related illness cases, an estimated 10%. PCCs do not systematically collect information on industry and occupation.


For more information on this indicator or occupational health in Maryland, visit the DHMH website.



# Indicator #12: Acute Work-Related Pesticide Poisonings Reported to Poison Control Centers

## Maryland State Occupational Health Indicators

➤ Malignant mesothelioma is a rare but highly fatal cancer of the thin membranes surrounding the chest cavity (pleura) or abdominal cavity (peritoneum). Much less frequently, this tumor affects other sites (e.g., pericardium). The only well-established risk factor for mesothelioma is exposure to asbestos fibers. Prior asbestos exposure, primarily from exposure in the workplace, has been reported in 62 to 85 percent of all mesothelioma cases.



**Indicator # 12: Cases of Malignant Mesothelioma, Maryland**

Year	Number	Incidence Rate*	Age-standardized Incidence Rate*
2000	47	11.3	9.0
2001	51	12.0	10.0
2002	64	14.9	13.0
2003	57	13.1	11.0
2004	48	10.9	9.0
2005	46	10.3	9.0
2006	44	9.8	8.0
2007	57	12.6	10.0
2008	50	11.0	9.0
2009	42	10.1	9.0
2010	70	15.0	15.0
2011	65	13.8	10.9
2012	46	9.6	7.7
2013	45	9.3	7.2

\* Cases per million residents

Data Source for this Indicator: SEER Stat Static data (Numbers of mesothelioma cases); U.S. Census Bureau (population statistics to calculate rates).

### About this Indicator:

#### Why is this Indicator Important?

Malignant mesothelioma, is a fatal cancer largely attributable to workplace exposure to asbestos. Tracking of malignant mesothelioma can be used to document the burden of the disease, design, target, and evaluate the impact of prevention efforts over time, and to identify previously unrecognized settings in which workers may continue to be at risk of asbestos exposure.

#### Limitation of Indicator:

Not all cases of mesothelioma are caused by occupational exposures. Because cancer is a disease of long latency, current incidence is not indicative of current exposures and it may be many years before reductions in exposures affect incidence.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.




# Indicator #14: Workers Employed in Industries with High Risk for Occupational Morbidity

## Maryland State Occupational Health Indicators

Workers in certain industries sustain non-fatal injuries and illnesses at much higher rates than the overall workforce. The proportion of the workforce that is employed in these high-risk industries varies by state. This variation can help explain differences in injury and illness rates among states.

**Indicator #14: Percentage of workers in industries with high risk for occupational morbidity, Maryland and U.S.**



\* List of High Risk Industries Updated in Guidelines

**Indicator #14: Workers in industries at high risk for occupational morbidity, Maryland**

Year	Number	Percentage
2000	77,907	3.8
2001	79,641	3.8
2002	78,244	3.8
2003*	111,995	5.4
2004	114,887	5.3
2005	112,306	5.2
2006	115,668	5.2
2007	114,358	5.1
2008*	144,551	6.5
2009	141,903	6.7
2010	139,937	6.7
2011	139,245	6.6
2012	138,622	6.4
2013*	105,641	4.8

\* List of High Risk Industries Updated in Guidelines

Data Source for this Indicator: Census Bureau County Business Patterns.

### About this Indicator:

#### Why is this Indicator Important?

Work-related injuries and illnesses are preventable, and control of occupational hazards is the most effective means of prevention. Concentrating on high-risk industries for non-fatal injuries and illnesses helps prioritize limited resources.

#### Limitation of Indicator:

It is possible that some new employers are not counted in the County Business Patterns mid-March survey. In addition, differences in regional industrial practices may cause the ranking of high-risk industries within a specific State to differ from those identified from national data.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.



# Indicator #14: Workers Employed in Industries with High Risk for Occupational Morbidity - Sub-State Data

## Maryland State Occupational Health Indicators

### About this Indicator:

#### Why is this Indicator Important?


Work-related injuries and illnesses are preventable, and control of occupational hazards is the most effective means of prevention. Concentrating on high-risk industries for non-fatal injuries and illnesses helps prioritize limited resources.

#### Limitation of Indicator:


It is possible that some new employers are not counted in the County Business Patterns mid-March survey. In addition, differences in regional industrial practices may cause the ranking of high-risk industries within a specific State to differ from those identified from national data.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.

Indicator # 14: 2013 Percentage of workers in industries with high risk for occupational morbidity by county



Indicator # 14: 2013 Percentage of workers in industries with high risk for occupational morbidity by region



# Indicator #15: Workers Employed in Occupations with High Risk for Occupational Morbidity

**About this Indicator:**

**Why is this Indicator Important?**

Work-related injuries and illnesses are preventable, and control of occupational hazards is the most effective means of prevention. Concentrating on high-risk occupations for non-fatal injuries and illnesses helps prioritize limited resources.

**Limitation of Indicator:**


Differences in regional industrial practices may cause the ranking of high-risk occupations within a specific state or industry to differ from those identified from national data.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.

## Maryland State Occupational Health Indicators

Workers in certain occupations sustain non-fatal injuries and illnesses at much higher rates than the overall workforce. The proportion of the workforce that is employed in these high-risk occupations varies by state. This variation can help explain differences in injury and illness rates among states.

**Indicator # 15: Percentage of workers in occupations with high risk of occupational morbidity, Maryland and U.S.**



\* List of High Risk Industries Updated in Guidelines

**Indicator # 15: Percentage of workers in occupations at high risk for occupational morbidity, Maryland**

Year	Number	Percentage
2000	228,906	7.5
2001	204,523	6.1
2002	145,725	5.3
2003*	252,956	9.1
2004	246,527	8.9
2005	244,934	8.7
2006	287,293	10.0
2007	270,002	9.4
2008*	292,328	14.9
2009	268,627	14.2
2010	292,295	14.9
2011	291,925	14.9
2012	293,357	14.7
2013	308,205	15.4


\* List of High Risk Occupations Updated in Guidelines




# Indicator #16: Workers Employed in Industries and Occupations with High Risk for Occupational Mortality

## Maryland State Occupational Health Indicators

Multiple factors and risks contribute to work-related fatalities, including workplace and process design, work organization, worker characteristics, economics and other social factors. Workers in certain industries and occupations sustain fatal injuries at much higher rates than the overall workforce. The proportion of the workforce that is employed in these high-risk industries and occupations varies by state. This variation can help explain differences in injury mortality rates among states.



**Indicator #16.1:** Percentage of workers employed in industries with high risk for occupational mortality, Maryland and U.S.



**Indicator #16.2:** Percentage of workers employed in occupations with high risk for occupational mortality, Maryland and U.S.

\* List of High Risk Industries Updated in Guidelines

**Indicator # 16: Percentage of workers employed in industries and occupations at high risk for occupational mortality, Maryland**

Year	1. Industries		2. Occupations	
	Number	Percentage	Number	Percentage
2000	332,573	10.9	165,104	5.4
2001	429,229	12.7	135,357	4.0
2002	342,263	12.4	127,263	4.6
2003*	332,181	12.0	253,335	9.2
2004	361,682	13.1	252,201	9.1
2005	362,092	12.9	252,580	9.0
2006	384,467	13.3	274,269	9.5
2007	350,495	12.2	248,518	8.7
2008*	331,222	14.8	257,145	11.5
2009	306,088	14.2	227,360	10.5
2010	314,269	14.2	235,024	10.6
2011	322,878	14.5	251,296	11.3
2012	318,659	14.2	226,828	10.1
2013	318,802	14.1	238,299	10.5

\* Lists of High Risk Industries and Occupations Updated in Guidelines

Data Source for this Indicator: Bureau of Labor Statistics Current Population Survey.

### About this Indicator:

#### Why is this Indicator Important?

Surveillance of work-related fatalities can identify new hazards and case clusters, leading to the development of new interventions and development of new or revised regulations to protect workers. Concentrating on high-risk occupations and industries for fatalities helps prioritize limited resources.

#### Limitation of Indicator:

Differences in regional industrial practices may cause the ranking of high-risk occupations and industries within a specific State to differ from those identified from national data.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.



# Indicator #17: Occupational Safety and Health Professionals

## Maryland State Occupational Health Indicators

► This occupational health indicator provides information about occupational safety and health professionals who are board-certified occupational medicine physicians, members of the American College of Occupational and Environmental Medicine (ACOEM), board-certified occupational health nurses, members of the American Association of Occupational Health Nurses (AAOHN), board-certified industrial hygienists, members of the American Industrial Hygiene Association (AIHA), board-certified safety professionals, and members of the American Society of Safety Engineers (ASSE).

**Indicator # 17: Occupational safety and health professionals, Maryland**

Year	American Board of Preventive Medicine (ABPM)		American College of Occupational and Environmental Medicine (ACOEM)		American Board of Occupational Health Nurses (ABOHN)		American Association of Occupational Health Nurses (AAOHN)	
	Number	Rate*	Number	Rate*	Number	Rate*	Number	Rate*
2000**	108	4.0	156	5.8	143	5.3	143	5.3
2001**	108	4.0	156	5.7	143	5.3	143	5.3
2002**	108	3.9	156	5.6	143	5.2	143	5.2
2003	108	3.9	156	5.6	143	5.2	143	5.2
2004	113	4.1	142	5.1	141	5.1	198	7.2
2005	113	4.0	148	5.3	135	4.8	176	6.3
2006	112	3.9	135	4.7	171	5.9	169	5.9
2007	115	4.0	130	4.6	153	5.4	173	6.1
2008	113	3.9	126	4.4	150	5.2	149	5.2
2009	113	4.0	132	4.7	150	5.3	138	4.9
2010	122	4.3	137	4.9	145	5.1	N/A	N/A
2011	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
2012	131	4.5	128	4.4	N/A	N/A	116	4.0
2013	N/A	N/A	125	4.3	131	4.5	N/A	N/A

**Indicator # 17: Occupational safety and health professionals, Maryland, Continued**

Year	American Board of Industrial Hygiene (ABIH)		American Industrial Hygiene Association (AIHA)		Board of Certified Safety Professionals (BCSP)		American Society of Safety Engineers (ASSE)	
	Number	Rate*	Number	Rate*	Number	Rate*	Number	Rate*
2000**	221	8.2	359	13.3	191	7.1	588	21.8
2001**	221	8.1	359	13.2	191	7.0	588	21.6
2002**	221	8.0	359	13.0	191	6.9	588	21.2
2003	221	8.0	359	12.9	191	6.9	588	21.2
2004	231	8.4	341	12.3	198	7.2	630	22.8
2005	238	8.5	323	11.5	204	7.3	685	24.5
2006	239	8.3	312	10.8	201	7.0	566	19.6
2007	247	8.7	281	9.9	206	7.2	562	19.7
2008	246	8.6	266	9.3	220	7.7	544	18.9
2009	244	8.7	272	9.6	238	8.4	502	17.8
2010	241	8.6	259	9.2	242	8.6	666	23.6
2011	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
2012	177	6.1	306	10.5	271	9.3	546	18.8
2013	210	7.2	198	6.8	285	9.8	556	19.0

\* Rate per 100,000 employed persons age 16 years or older

\*\* Records of past membership totals do not exist for most organizations. For this reason, for years 2000-2002, membership counts for 2003 were used in the numerator of this indicator, while the number employed for each respective year was used in the denominator. For 2003 onward, numerator and denominator data come from the same year.

**Data Source for this Indicator:** Current membership rosters of cited organizations (Numbers of health and safety professionals); Bureau of Labor Statistics Current Population Survey (Employment estimates used to calculate rates).

### About this Indicator:

#### Why is this Indicator Important?

Work-related injuries and illnesses are preventable. It is important to determine if there are sufficient trained personnel to implement occupational health preventive services.

#### Limitation of Indicator:

Other important occupational health specialties such as fire prevention, health physicists, and ergonomists are not included.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.



# Indicator #18: Occupational Safety and Health Administration (OSHA) Enforcement Activities

## About this Indicator:

### Why is this Indicator Important?

The measures of frequency for this indicator may approximate the added health and safety benefits and protections felt by workers as a result of their worksites being inspected.


### Limitation of Indicator:

This indicator only measures enforcement activity and no other measures of OSHA activity such as education and compliance assistance. OSHA may conduct multiple inspections of one establishment during a calendar year, therefore the percent of establishments inspected may be overestimated.


For more information on this indicator or occupational health in Maryland, visit the DHMH website.

## Maryland State Occupational Health Indicators

The Occupational Safety and Health Act of 1970 was passed by Congress to assure safe and healthy working conditions for every worker in the nation. Under the Act, the U.S. Department of Labor's Occupational Safety and Health Administration (OSHA) is authorized to conduct worksite inspections to determine whether employers are complying with health and safety standards. OSHA may issue citations and impose fines on employers if violations are found. This indicator provides a measure of the workers and worksites potentially benefiting directly from Federal/State OSHA inspection activity.



**Indicator #18.1:** Percentage of establishments under OSHA jurisdiction inspected by OSHA, Maryland and U.S.



**Indicator #18.2:** Percentage of workers in establishments under OSHA jurisdiction whose work areas were inspected by OSHA, Maryland and U.S.

**Indicator #18: OSHA enforcement activities, Maryland**

Year	1. Number of Establishments Inspected by OSHA		2. Number of Establishments Eligible for Inspection	3. Number of Employees at Inspected Sites	4. Number of Covered Employees in State	
	Average Number	Percentage			Average Number	Percentage
2000	1,814	1.2	146,148	184,584	2,400,659	2.5
2001	1,703	1.2	146,748	173,884	2,417,010	1.8
2002	1,504	1.0	149,176	136,432	2,422,473	1.5
2003	1,440	1.0	149,999	162,965	2,429,457	1.7
2004	1,074	0.7	154,973	49,955	2,454,451	1.0
2005	1,156	0.7	159,636	67,007	2,492,242	1.2
2006	1,149	0.7	162,536	87,834	2,524,798	1.5
2007	1,454	0.9	166,617	87,815	2,542,297	1.5
2008	1,487	0.9	165,484	95,642	2,532,654	3.1
2009	1,303	0.8	163,736	63,271	2,456,465	2.6
2010	1,206	0.7	163,239	31,841	2,448,540	1.3
2011	1,367	0.8	164,221	57,078	2,473,732	2.3
2012	1,747	1.0	167,221	54,778	2,506,593	2.2
2013	1,465	0.9	168,565	30,228	2,527,159	1.20

Data Source for this Indicator: Bureau of Labor Statistics' Covered Employers and Wages (Number of establishments and workforce estimates).




# Indicator #19: Workers' Compensation Awards

## Maryland State Occupational Health Indicators

Workers' compensation is a state-based social insurance program, developed to provide compensation for workers with work-related injuries or illnesses while limiting the liability exposure of employers. Workers' compensation provides benefits to partially replace lost wages and pay for medical expenses associated with a work-related injury or illness. Workers' compensation awards are reviewed to establish whether the reported medical condition is work-related. Accepted awards represent known work-related injuries and illnesses, and often more severe cases. The total and average amounts of benefits paid estimate the burden of these events.

**Indicator #19: Average Workers' Compensation benefit paid per covered worker, Maryland and U.S.**



**Indicator # 19: Workers compensation awards, Maryland**

Year	Total Amount of workers' compensation benefits paid	Average amount of workers' compensation paid per covered worker
2000	\$1,194,629,000	\$524.65
2001	\$787,442,000	\$343.11
2002	\$783,686,000	\$340.88
2003	\$628,510,000	\$272.55
2004	\$767,576,000	\$329.15
2005	\$769,563,000	\$324.44
2006	\$815,351,000	\$339.02
2007	\$830,927,000	\$343.07
2008	\$935,948,000	\$388.84
2009	\$895,905,000	\$385.17
2010	\$953,533,000	\$412.78
2011	\$1,006,998,000	\$432.19
2012	\$970,734,000	\$410.81
2013	\$944,612,000	\$396.23

Data Source for this Indicator: National Academy of Social Insurance (total amount and average benefits paid).

**About this Indicator:**  
**Why is this Indicator Important?**  
 Accepted awards represent known work-related injuries and illnesses, and often more severe cases. The total and average amounts of benefits paid estimate the burden of these events, which can help justify prevention programs and activities.

**Limitation of Indicator:**  
 Workers' compensation data is not complete, as many individuals with work-related illnesses do not file for workers' compensation. Differences among states in benefits paid could be due to a variety of factors other than injury and illness incidence. For this reason, this occupational health indicator should be used to monitor trends within states over time rather than to compare states.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.



# Indicator #20: Work-related low back disorder hospitalizations

## About this Indicator:

### Why is this Indicator Important?

Hospitalizations for work-related back disorders have serious and costly effects, including: high direct medical costs, significant functional impairment and disability, high absenteeism, reduced work performance, and lost productivity. Well-recognized prevention efforts can be implemented for high risk job activities and reduce the burden of work-related low back disorders.

### Limitation of Indicator:


Hospital discharge records are only available for non-federal, acute care hospitals. Many individuals with work-related injuries do not file for workers' compensation or fail to recognize work as the cause of their injury. Self-employed individuals are not covered by state workers' compensation systems.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.




## Maryland State Occupational Health Indicators

Each year 15-20% of Americans report back pain, resulting in over 100 million workdays lost and more than 10 million physician visits. National Health Interview survey data estimates 2/3 of all low back pain cases are attributable to occupational activities. The cost of back pain is also disproportionate, as it represents about 20% of workers' compensation claims but nearly 40% of the costs. In 2003, 3.2% of the total US workforce experienced a loss in productive time due to back pain. The total cost of this productive time lost to back pain is estimated to be in excess of \$19.8 billion dollars.



**Indicator #20.1:** Crude Rate of Work-Related Surgical Low Back Disorder Hospitalizations by State and U.S.



**Indicator #20.2:** Crude Rate of Work-Related Low Back Disorder Hospitalizations by State and U.S.

**Indicator # 20: Hospitalizations for work-related low back disorders**

Year	20.1. Surgical Low Back Disorder Hospitalizations		20.2. Low Back Disorder Hospitalizations	
	Number	Rate	Number	Rate
2008	531	18.5	670	23.8
2009	498	17.7	613	21.7
2010	433	15.4	545	19.3
2011	380	13.2	460	16.0
2012	328	11.3	420	14.5
2013	284	9.7	356	12.2

\* per 100,000 employed persons age 16 years or older

Data Source for this Indicator: Hospital discharge data (numerator); BLS Current Population Survey Data (denominator).

**About this Indicator:**

**Why is this Indicator Important?**

Hospitalizations for work-related back disorders have serious and costly effects, including: high direct medical costs, significant functional impairment and disability, high absenteeism, reduced work performance, and lost productivity. Well-recognized prevention efforts can be implemented for high risk job activities and reduce the burden of work-related low back disorders.


**Limitation of Indicator:**

Hospital discharge records are only available for non-federal, acute care hospitals. Many individuals with work-related injuries do not file for workers' compensation or fail to recognize work as the cause of their injury. Self-employed individuals are not covered by state workers' compensation systems.


For more information on this indicator or occupational health in Maryland, visit the DHMH website.

## Maryland State Occupational Health Indicators

**Indicator # 20.2: 2011-2013 crude rate of work-related low back disorder hospitalizations by county**



**Indicator # 20.2: 2011-2013 crude rate of work-related low back disorder hospitalizations by region**



Data Source for this Indicator: Hospital discharge data (numerator); BLS Current Population Survey Data (denominator).



# Indicator #20: Work-Related Low Back Disorder Hospitalizations - Race and Ethnicity Data

## About this Indicator:

### Why is this Indicator Important?

Hospitalizations for work-related back disorders have serious and costly effects, including: high direct medical costs, significant functional impairment and disability, high absenteeism, reduced work performance, and lost productivity. Well-recognized prevention efforts can be implemented for high risk job activities and reduce the burden of work-related low back disorders.


### Limitation of Indicator:

Hospital discharge records are only available for non-federal, acute care hospitals. Many individuals with work-related injuries do not file for workers' compensation or fail to recognize work as the cause of their injury. Self-employed individuals are not covered by state workers' compensation systems.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.

## Maryland State Occupational Health Indicators

**Indicator #20: 2013 rate of work-related low back disorder hospitalizations by race and ethnicity, Maryland**



\* Persons whose ethnicity is Hispanic may be of any race and, therefore, are classified by ethnicity as well as by race

**Indicator #20: 2013 work-related low back disorder hospitalizations by race, Maryland**

Year	White		Black		Hispanic*	
	Number	Crude Rate**	Number	Crude Rate**	Number	Crude Rate**
2013	225	12.6	82	10.0	16	5.4

\* Persons whose ethnicity is Hispanic may be of any race and, therefore, are classified by ethnicity as well as by race

\*\* Rate per 100,000 employed persons 16 years or older



# Indicator #21: Asthma Among Adults Caused or Made Worse by Work

## About this Indicator:

### Why is this Indicator Important?

Asthma is a chronic inflammatory disease of the airways that is life threatening and can be managed but not cured. Adult asthma can be caused by or made worse by workplace exposures.


### Limitation of Indicator:

This population-based estimate is subject to measurement, non-response and sampling errors. The indicator does not distinguish between new-onset and work-aggravated asthma. The Asthma Call Back Survey began new weighting methods in 2011 and the wording and order of questions changed in 2012. Any trend analysis would need to be restricted to 2012 forward. States using landline only versus landline and cellphone methodology do not have comparable estimates.


For more information on this indicator or occupational health in Maryland, visit the DHMH website.

## Maryland State Occupational Health Indicators

Work-related asthma is preventable but often goes undiagnosed by physicians. Research has shown that work-related asthma can have adverse effects on the worker, including increased morbidity, adverse socioeconomic impacts and difficulty getting and sustaining work. Estimating the burden of asthma caused or made worse by work can help target prevention programs and activities. If diagnosed early, work-related asthma may be partially or completely reversible if exposures can be identified and properly stopped or controlled. The Asthma Call-Back Survey (ACBS) contains multiple questions related to the work-relatedness of a respondent's asthma and these questions are administered to adults 18 years or older.



**Indicator #21.2:** Estimated proportion of ever-employed adults with current asthma who report that their asthma was caused or made worse by exposures at work by state and U.S. (Landline only survey methodology)



**Indicator #21.2:** Estimated proportion of ever-employed adults with current asthma who report that their asthma was caused or made worse by exposures at work by state and U.S. (Landline & Cellphone survey methodology)

### Indicator # 21: Asthma Among Adults Caused or Made Worse By Work

Year	Landline Only Survey Methodology		Landline & Cellphone Survey Methodology	
	21.1. Weighted Frequency	21.2. Estimated Proportion	21.1. Weighted Frequency	21.2. Estimated Proportion
2011	194,859	46.3	N/A	N/A
2012	N/A	N/A	N/A	N/A
2013	202,747	42.3	229,988	51.5

Data Source for this Indicator: Asthma Call-Back Survey (numerator & denominator).



# Indicator #22: Work-Related Severe Traumatic Injury Hospitalizations

## Maryland State Occupational Health Indicators

### About this Indicator:

### Why is this Indicator Important?

Acute work-related trauma is a leading cause of death and disability for U.S. workers. Severe traumatic injury can lead to long-term pain and disability and is very costly for workers' compensation systems and society as a whole.


### Limitation of Indicator:

Hospital discharge records are only available for non-federal, acute care hospitals. Many individuals with work-related injuries do not file for workers' compensation or fail to recognize work as the cause of their injury. Self-employed individuals are not covered by state workers' compensation systems.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.

► Acute work-related trauma is a leading cause of death and disability among U.S. workers. Changes in hospitalization practices and workers' compensation coverage/reporting may increasingly reduce capture of minor injuries but have little effect on severe injuries. Use of a severity threshold can decrease the impact of changing utilization and service delivery patterns on observed injury trends. When hospitalization data are used to calculate occupational injury trends in the absence of severity restriction, observed trends are biased downward. Accurate characterization of injury trends is critical to understanding how we are doing as a nation with regard to occupational injury prevention.

**Indicator # 22: Work-related severe traumatic injury hospitalizations, Maryland**



**Indicator # 22: Work-related severe traumatic injury hospitalizations, Maryland**

Year	Number	Rate*
2008	292	10.2
2009	276	9.8
2010	273	9.7
2011	236	8.2
2012	253	8.7
2013	239	8.2

\* Rate per 100,000 employed persons

Data Source for this Indicator: Hospital discharge data (numerator); BLS Current Population Survey Data (denominator).



# Indicator #22: Work-Related Severe Traumatic Injury Hospitalizations - Sub-State Data

About this Indicator:

Why is this Indicator Important?

Acute work-related trauma is a leading cause of death and disability for U.S. workers. Severe traumatic injury can lead to long-term pain and disability and is very costly for workers' compensation systems and society as a whole.


Limitation of Indicator:

Hospital discharge records are only available for non-federal, acute care hospitals. Many individuals with work-related injuries do not file for workers' compensation or fail to recognize work as the cause of their injury. Self-employed individuals are not covered by state workers' compensation systems.


For more information on this indicator or occupational health in Maryland, visit the DHMH website.

## Maryland State Occupational Health Indicators

Indicator # 22: 2009-2013 crude rate of work-related severe traumatic injury hospitalizations by county



Indicator # 22: 2009-2013 crude rate of work-related severe traumatic injury hospitalizations by region



Data Source for this Indicator: Hospital discharge data (numerator); BLS Current Population Survey Data (denominator).



# Indicator #22: Work-Related Severe Traumatic Injury Hospitalizations - Race and Ethnicity Data

## About this Indicator:

## Why is this Indicator Important?

Acute work-related trauma is a leading cause of death and disability for U.S. workers. Severe traumatic injury can lead to long-term pain and disability and is very costly for workers' compensation systems and society as a whole.


## Limitation of Indicator:

Hospital discharge records are only available for non-federal, acute care hospitals. Many individuals with work-related injuries do not file for workers' compensation or fail to recognize work as the cause of their injury. Self-employed individuals are not covered by state workers' compensation systems.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.

## Maryland State Occupational Health Indicators

**Indicator #22: 2013 rate of work-related severe traumatic injury hospitalizations by race and ethnicity, Maryland**



\* Persons whose ethnicity is Hispanic may be of any race and, therefore, are classified by ethnicity as well as by race

**Indicator #22: 2013 work-related severe traumatic injury hospitalizations by race, Maryland**

Year	White		Black		Hispanic*	
	Number	Crude Rate**	Number	Crude Rate**	Number	Crude Rate**
2013	130	7.3	50	6.1	26	8.8

\* Persons whose ethnicity is Hispanic may be of any race and, therefore, are classified by ethnicity as well as by race

\*\* Rate per 100,000 employed persons 16 years or older




# Indicator #24: Occupational Heat-Related Emergency Department Visits

## Maryland State Occupational Health Indicators

Minimal epidemiological information about occupational heat-related morbidity is available. Tracking occupational heat-related illness using emergency department data will establish a baseline for occupational epidemiologist to understand the magnitude of the disease burden in the population and support implementation and evaluation of prevention measures.

**Indicator # 24: Occupational Heat-Related Emergency Department Visits, Maryland**



**Indicator # 24: Occupational Heat-Related Emergency Department Visits**

Year	Number	Rate*
2012	58	2.0
2013	43	1.5

\* Rate per 100,000 employed persons

### About this Indicator:

#### Why is this Indicator Important?

Exposure to environmental heat is a hazard for many workers who are not able to maintain thermal equilibrium due to their work environment, required clothing, and usage of protective equipment.

#### Limitation of Indicator:

The number of diagnostic fields in ED records and utilization of EDs varies by state. The majority of individuals with work-related illnesses and injuries do not file for workers' compensation. Self-employed individuals and out-of-state workers are not captured. Attribution of payer in may not be accurate. The effectiveness of identifying work-relatedness through E-codes will vary within each facility.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.




Data Source for this Indicator: Emergency department visits data (numerator); BLS Current Population Survey Data (denominator). A condition was considered work-related if workers' compensation was listed as primary payer in the hospital discharge data.

# State Specific Indicator #1: Work-Related Emergency Department Visits

## Maryland State Occupational Health Indicators

Collecting data on emergency department visits for occupational illness and injury allows for tracking of emergency department utilization for work-related diseases and helps document the burden of occupational injuries and illnesses, including less severe cases that do not require inpatient hospitalization. These data can be used to design, target, and evaluate the impact of prevention efforts over time, and identify settings in which workers may continue to be at high risk.

**State Specific Indicator # 1: Work-Related Emergency Department Visits, Maryland**



**State Specific Indicator # 1: Work-Related Emergency Department Visits**

Year	Number	Rate*
2012	24,839	855.6
2013	24,510	839.0

\* Rate per 100,000 employed persons

**Data Source for this Indicator:** Emergency department visits data (numerator); BLS Current Population Survey Data (denominator). A condition was considered work-related if workers' compensation was listed as primary payer in the hospital discharge data.

### About this Indicator:

#### Why is this Indicator Important?

Information on work-related emergency department visits can be used to document the burden of occupational injuries and illnesses. Work-related visits to emergency departments may differ compared to inpatient hospitalizations over time and between sub groups of Maryland residents based on differences in utilization patterns and severity of injury.

#### Limitation of Indicator:

The majority of individuals with work-related illnesses and injuries do not file for workers' compensation. Self-employed individuals and out-of-state workers are not captured. Attribution of payer may not be accurate.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.



# State Specific Indicator #1: Work-Related Emergency Department Visits - Sub-State Data

**About this Indicator:**  
**Why is this Indicator Important?**


Information on work-related emergency department visits can be used to document the burden of occupational injuries and illnesses. Work-related visits to emergency departments may differ compared to inpatient hospitalizations over time and between sub groups of Maryland residents based on differences in utilization patterns and severity of injury.

**Limitation of Indicator:**


The majority of individuals with work-related illnesses and injuries do not file for workers' compensation. Self-employed individuals and out-of-state workers are not captured. Attribution of payer may not be accurate.

For more information on this indicator or occupational health in Maryland, visit the DHMH website.


## Maryland State Occupational Health Indicators



**State Specific Indicator # 1: 2013 crude rate of work-related emergency department visits by ZIP code per 100,000 workers**



**State Specific Indicator # 1: 2013 crude rate of work-related emergency department visits by county per 100,000 workers**



**State Specific Indicator # 1: 2013 crude rate of work-related emergency department visits by region per 100,000 workers**

**Data Source for this Indicator:** Maryland emergency department visits data (numerator). County and region level denominator: Bureau of Labor Statistics Current Population Survey (total number of employed persons). ZIP code level denominator: The Nielson Company. A condition was considered work-related if workers' compensation was listed as primary payer in the hospital discharge data.



# State Specific Indicator #1: Work-Related Emergency Department Visits - Race and Ethnicity Data

## About this Indicator:

### Why is this Indicator Important?

Information on work-related emergency department visits can be used to document the burden of occupational injuries and illnesses. Work-related visits to emergency departments may differ compared to inpatient hospitalizations over time and between sub groups of Maryland residents based on differences in utilization patterns and severity of injury.


### Limitation of Indicator:

The majority of individuals with work-related illnesses and injuries do not file for workers' compensation. Self-employed individuals and out-of-state workers are not captured. Attribution of payer may not be accurate.


For more information on this indicator or occupational health in Maryland, visit the DHMH website.

## Maryland State Occupational Health Indicators


### State Specific Indicator # 1: Work-Related Emergency Department Visits, Maryland



### State Specific Indicator # 1: 2013 crude rate of work-related emergency department visits by county per 100,000 black employed persons



### State Specific Indicator # 1: 2013 crude rate of work-related emergency department visits by county per 100,000 white employed persons



Data Source for this Indicator: Maryland emergency department visits data (numerator). County and region level denominator: Bureau of Labor Statistics Current Population Survey (total number of employed persons). A condition was considered work-related if workers' compensation was listed as primary payer in the hospital discharge data.

