

Studies that show **Cell Tower** Health Effects

Table of Contents

78 Studies Showing Health Effects from Cell Tower Radio Frequency Radiation

Health Effects	Pages
Cancer	Pages 2-4
Neurological Effects	Pages 5-7
Effects on Reproductive Organs and Fertility Effects	Pages 7-9
Effects on Wellbeing	Pages 9-14
Effects on Sleep	Pages 14-15
Effects on Cells	Pages 16
Effects on Eyes	Pages 17-18
Oxidative Stress	Pages 18-20
Effects on Blood	Pages 20-21
Death	Page 22-23
Effects on Hormones	Pages 23-24
Genetic Effects	Pages 24-29
Effects on Glands	Pages 29
Effects on Animals and Environment	Page 29-32
Effects on Skin	Page 32
Effects on Protein	Pages 33
Effects on Immune Function	Page 33-34
Electromagnetic Hyper-Sensitivity	Pages 34
RF Levels in Cities	Pages 34-36
Effects on Water	Page 36
RF Levels From Cell Towers	Pages 36-37
Danger Perception and Symptoms	Pages 37-38
Reason Why Not A Lot of Studies on Cell Towers	Page 38-39
Miscellaneous Effects	Pages 39

Studies that show **Cell Tower** Health Effects

Cancer

Wolf R, Wolf D. Increased incidence of cancer near a cell-phone transmitter station. Inter J Cancer Prev 1(2):123-128, 2004.

Significant concern has been raised about possible health effects from exposure to radiofrequency (RF) electromagnetic fields, especially after the rapid introduction of mobile telecommunication systems. Parents are especially concerned with the possibility that children might develop cancer after exposure to the RF emissions from mobile telephone base stations erected in or near schools. The few epidemiologic studies that did report on cancer incidence in relation to RF radiation have generally presented negative or inconsistent results, and thus emphasized the need for more studies that should investigate cohorts with high RF exposure for changes in cancer incidence. The aim of this study is to investigate whether there is an increased cancer incidence in populations, living in a small area, and exposed to RF radiation from a cell-phone transmitter station.

This is an epidemiologic assessment, to determine whether the incidence of cancer cases among individuals exposed to a cell-phone transmitter station is different from that expected in Israel, in Netanya, or as compared to people who lived in a nearby area. Participants are people (n=622) living in the area near a cell-phone transmitter station for 3-7 years who were patients of one health clinic (of DW). The exposure began 1 year before the start of the study when the station first came into service. A second cohort of individuals (n=1222) who get their medical services in a clinic located nearby with very closely matched, environment, workplace and occupational characteristics was used for comparison.

In the area of exposure (area) eight cases of different kinds of cancer were diagnosed in a period of only one year. This rate of cancers was compared both with the rate of 31 cases per 10,000 per year in the general population and the 2/1222 rate recorded in the nearby clinic (area B). Relative cancer rates for female were 10.5 for area A. 0.6 for area B and 1 for the whole town of Netanya. Cancer incidence of women in area A was thus significantly higher ($p < 0.0001$) compared with that of area B and the whole city. A comparison of the relative risk revealed that there were 4.15 times more cases in area than in the entire population. The study indicates an association between increased incidence of cancer and living in proximity to a cell-phone transmitter station.

Yakymenko I, Sidorik E, Krylenko S, Chekhun V. Long-term exposure to microwave radiation provokes cancer growth: evidences from radars and mobile communication systems. Exp Oncol. 33(2):62-70, 2011.

In this review we discuss alarming epidemiological and experimental data on possible carcinogenic effects of long term exposure to low intensity microwave (MW) radiation. Recently, a number of reports revealed that under certain conditions the irradiation by low intensity MW can substantially induce cancer progression in humans and in animal models. The carcinogenic effect of MW irradiation is typically manifested after long term (up to 10 years and more) exposure. Nevertheless, even a year of operation of a

Studies that show **Cell Tower** Health Effects

powerful base transmitting station for mobile communication reportedly resulted in a dramatic increase of cancer incidence among population living nearby. In addition, model studies in rodents unveiled a significant increase in carcinogenesis after 17-24 months of MW exposure both in tumor-prone and intact animals. To that, such metabolic changes, as overproduction of reactive oxygen species, 8-hydroxi-2-deoxyguanosine formation, or ornithine decarboxylase activation under exposure to low intensity MW confirm a stress impact of this factor on living cells. We also address the issue of standards for assessment of biological effects of irradiation. It is now becoming increasingly evident that assessment of biological effects of non-ionizing radiation based on physical (thermal) approach used in recommendations of current regulatory bodies, including the International Commission on Non-Ionizing Radiation Protection (ICNIRP) Guidelines, requires urgent reevaluation. We conclude that recent data strongly point to the need for re-elaboration of the current safety limits for non-ionizing radiation using recently obtained knowledge. We also emphasize that the everyday exposure of both occupational and general public to MW radiation should be regulated based on a precautionary principles which imply maximum restriction of excessive exposure.

Michelozzi P, Ancona C, Fusco D, Forastiere F, Perucci CA, Risk of leukemia and residence near a radio transmitter in Italy. Epidemiology 9 (Suppl) 354p, 1998.

We conducted a small area study to investigate a cluster of leukemia near a high power radio-transmitter in a peripheral area of Rome. The leukemia mortality within 3.5 km (5,863 inhabitants) was higher than expected (SMR=2.5, 95% confident interval 1.07-4.83); the excess was due to a significant higher mortality among men (7 cases observed, SMR=3.5). The results of the Stone's test, after adjusting for socio-economic confounding, showed a significant decline in risk with distance from the transmitter only among men (p=0.005), whereas the p-value for both sexes was p=0.07.

Lourencini da Silva R , Albano F, Lopes dos Santos LR , Tavares AD, Felzenszwalb I, The effect of electromagnetic field exposure on the formation of DNA lesions. Redox Rep 5(5):299-301, 2000.

In an attempt to determine whether electromagnetic field (EMF) exposure might lead to DNA damage, we exposed SnCl₂-treated pBR322 plasmids to EMF and analysed the resulting conformational changes using agarose gel electrophoresis. An EMF-dependent potentiation of DNA scission (i.e. the appearance of relaxed plasmids) was observed. In confirmation of this, plasmids pre-exposed to EMF also were less capable of transforming Escherichia coli. The results indicate that EMF, in the presence of a transition metal, is capable of causing DNA damage. These observations support the idea that EMF, probably through secondary generation of reactive oxygen species, can be clastogenic and provide a possible explanation for the observed correlation between EMF exposure and the frequency of certain types of cancers in humans.

Li CY, Liu CC, Chang YH, Chou LP, Ko MC. A population-based case-control study of radiofrequency exposure in relation to childhood neoplasm. Sci Total Environ. 435-436:472-478, 2012.

Studies that show **Cell Tower** Health Effects

This population-based case-control study in Taiwan considered incident cases aged 15 years or less and admitted in 2003 to 2007 for all neoplasm (ICD-9-CM: 140-239) (n=2606), including 939 leukemia and 394 brain neoplasm cases. Controls were randomly selected, with a case/control ratio of 1:30 and matched on year of birth, from all non-neoplasm children insured in the same year when the index case was admitted. Annual summarized power (ASP, watt-year) was calculated for each of the 71,185 mobile phone base stations (MPBS) in service between 1998 and 2007. Then, the **annual power density (APD, watt-year/km(2))** of each township (n=367) was computed as a ratio of the total ASP of all MPBS in a township to the area of that particular township. Exposure of each study subject to radio frequency (RF) was indicated by the averaged APD within 5 years prior to the neoplasm diagnosis (cases) or July 1st of the year when the index case was admitted (controls) in the township where the subject lived. Unconditional logistic regression model with generalized estimation equation was employed to calculate the covariate-**adjusted odds ratio [AOR]** of childhood neoplasm in relation to RF exposure. A higher than median averaged APD (approximately 168 WYs/km(2)) was significantly associated with an increased AOR for all neoplasms (1.13; 1.01 to 1.28), but not for leukemia (1.23; 0.99 to 1.52) or brain neoplasm (1.14, 0.83 to 1.55). This study noted a significantly increased risk of all neoplasms in children with higher-than-median RF exposure to MPBS. The slightly elevated risk was seen for leukemia and brain neoplasm, but was not statistically significant. These results may occur due to several methodological limitations.

Johnson EH, Chima SC, Muirhead DE, A cerebral primitive neuroectodermal tumor in a squirrel monkey (*Saimiri sciureus*). J Med Primatol 28(2):91-96, 1999.

An adult squirrel monkey with a history of long-term exposure to microwave radiation was found at necropsy to have a malignant tumor of the right cerebral cortex. Gross examination revealed a mass with expanding borders in the right frontoparietal cortex with compression of the adjacent lateral ventricle. Microscopy revealed a tumor composed of sheets of moderate-sized cells, resembling an oligodendroglioma, with clear cytoplasm and central nuclei interrupted by delicate vasculature. Malignant features were present in the form of marked nuclear pleomorphism, frequent mitotic figures, and focal necrosis. A neuronal cell origin for this tumor was supported by immunohistochemical analysis, which revealed immunopositivity for neurofilament proteins and neuron-specific enolase. Staining for vimentin and glial fibrillary acid protein was negative, except in reactive astrocytes at the tumor margins and adjacent to intra-tumoral blood vessels. Antibody activity against Ki-67 antigen, a marker of rapidly proliferating tumor cells, and p53 oncoprotein was strongly positive, indicative of the aggressive and malignant nature of this tumor. The tumor was diagnosed as a cerebral primitive neuroectodermal tumor.

Neurological Effects

Khurana VG, Hardell L, Everaert J, Bortkiewicz A, Carlberg M, Ahonen M.

Studies that show **Cell Tower** Health Effects

Epidemiological evidence for a health risk from mobile phone base stations. *Int J Occup Environ Health.* 16(3):263-267, 2010.

Human populations are increasingly exposed to microwave/radiofrequency (RF) emissions from wireless communication technology, including mobile phones and their base stations. By searching PubMed, we identified a total of 10 epidemiological studies that assessed for putative health effects of mobile phone base stations. Seven of these studies explored the association between base station proximity and neurobehavioral effects and three investigated cancer. We found that eight of the 10 studies reported increased prevalence of adverse neurobehavioral symptoms or cancer in populations living at distances < 500 meters from base stations. None of the studies reported exposure above accepted international guidelines, suggesting that current guidelines may be inadequate in protecting the health of human populations. We believe that comprehensive epidemiological studies of long-term mobile phone base station exposure are urgently required to more definitively understand its health impact.

Hocking B, Westerman R. Neurological abnormalities associated with CDMA exposure. *Occup Med (Lond)* 51(6):410-413, 2001.

Dysaesthesiae of the scalp and neurological abnormality after mobile phone use have been reported previously, but the roles of the phone per se or the radiations in causing these findings have been questioned. We report finding a neurological abnormality in a patient after accidental exposure of the left side of the face to mobile phone radiation [code division multiple access (CDMA)] from a down-powered mobile phone base station antenna. He had headaches, unilateral left blurred vision and pupil constriction, unilateral altered sensation on the forehead, and abnormalities of current perception thresholds on testing the left trigeminal ophthalmic nerve. His nerve function recovered during 6 months follow-up. His exposure was 0.015-0.06 mW/cm(2) over 1-2 h. The implications regarding health effects of radiofrequency radiation are discussed.

Abdel-Rassoul G, El-Fateh OA, Salem MA, Michael A, Farahat F, El-Batanouny M, Salem E. Neurobehavioral effects among inhabitants around mobile phone base stations. *Neurotoxicology.* 28(2):434-40, 2007.

BACKGROUND: There is a general concern on the possible hazardous health effects of exposure to radiofrequency electromagnetic radiations (RFR) emitted from mobile phone base station antennas on the human nervous system. AIM: To identify the possible neurobehavioral deficits among inhabitants living nearby mobile phone base stations. METHODS: A cross-sectional study was conducted on (85) inhabitants living nearby the first mobile phone station antenna in Menoufiya governorate, Egypt, 37 are living in a building under the station antenna while 48 opposite the station. A control group (80) participants were matched with the exposed for age, sex, occupation and educational level. All participants completed a structured questionnaire containing: personal, educational and medical histories; general and neurological examinations; neurobehavioral test battery (NBTB) [involving tests for visuomotor speed, problem solving, attention and memory]; in addition to Eysenck personality questionnaire (EPQ). RESULTS: The prevalence of neuropsychiatric complaints as headache (23.5%), memory

Studies that show **Cell Tower** Health Effects

changes (28.2%), dizziness (18.8%), tremors (9.4%), depressive symptoms (21.7%), and sleep disturbance (23.5%) were significantly higher among exposed inhabitants than controls: (10%), (5%), (5%), (0%), (8.8%) and (10%), respectively (P<0.05). The NBTB indicated that the exposed inhabitants exhibited a significantly lower performance than controls in one of the tests of attention and short-term auditory memory [Paced Auditory Serial Addition Test (PASAT)]. Also, the inhabitants opposite the station exhibited a lower performance in the problem solving test (block design) than those under the station. All inhabitants exhibited a better performance in the two tests of visuomotor speed (Digit symbol and Trailmaking B) and one test of attention (Trailmaking A) than controls. The last available measures of RFR emitted from the first mobile phone base station antennas in Menoufiya governorate were less than the allowable standard level. CONCLUSIONS AND RECOMMENDATIONS: Inhabitants living nearby mobile phone base stations are at risk for developing neuropsychiatric problems and some changes in the performance of neurobehavioral functions either by facilitation or inhibition. So, revision of standard guidelines for public exposure to RER from mobile phone base station antennas and using of NBTB for regular assessment and early detection of biological effects among inhabitants around the stations are recommended.

Akbari A, Jelodar G, Nazifi S. Vitamin C protects rat cerebellum and encephalon from oxidative stress following exposure to radiofrequency wave generated by a BTS antenna model. Toxicol Mech Methods. 24(5):347-352, 2014.

Radio frequency wave (RFW) generated by base transceiver station has been reported to produce deleterious effects on the central nervous system function, possibly through oxidative stress. This study was conducted to evaluate the effect of RFW-induced oxidative stress in the cerebellum and encephalon and the prophylactic effect of vitamin C on these tissues by measuring the antioxidant enzymes activity, including: glutathione peroxidase, superoxide dismutase, catalase, and malondialdehyde (MDA). Thirty-two adult male Sprague-Dawley rats were randomly divided into four equal groups. The control group; the control-vitamin C group received L-ascorbic acid (200 mg/kg of body weight/day by gavage) for 45 days. The RFW group was exposed to RFW and the RFW+ vitamin C group was exposed to RFW and received vitamin C. At the end of the experiment, all groups were killed and encephalon and cerebellum of all rats were removed and stored at -70 °C for measurement of antioxidant enzymes activity and MDA. The results indicate that exposure to RFW in the test group decreased antioxidant enzymes activity and increased MDA compared with the control groups (p < 0.05). The protective role of vitamin C in the treated group improved antioxidant enzymes activity and reduced MDA compared with the test group (p < 0.05). It can be concluded that RFW causes oxidative stress in the brain and vitamin C improves the antioxidant enzymes activity and decreases MDA.

Bak M, Dudarewicz A, Zmyślony M, Sliwinska-Kowalska M. Effects of GSM signals during exposure to event related potentials (ERPs). Int J Occup Med Environ Health. 23(2):191-199, 2010.

Studies that show **Cell Tower** Health Effects

Objectives: The primary aim of this work was to assess the effect of electromagnetic field (EMF) from the GSM mobile phone system on human brain function. The assessment was based on the assay of event related potentials (ERPs). **Material and Methods:** The study group consisted of 15 volunteers, including 7 men and 8 women. The test protocol comprised determination of P300 wave in each volunteer during exposure to the EMF. To eliminate possible effects of the applied test procedure on the final result, the test was repeated without EMF exposure. P300 latency, amplitude, and latency of the N1, N2, P2 waves were analysed. **Results:** The statistical analysis revealed an effect of EMF on P300 amplitude. In the experiment with EMF exposure, lower P300 amplitudes were observed only at the time in which the volunteers were exposed to EMF; when the exposure was discontinued, the values of the amplitude were the same as those observed before EMF application. No such change was observed when the experiment was repeated with sham exposure, which may be considered as an indirect proof that lower P300 amplitude values were due to EMF exposure. No statistically significant changes were noted in the latencies of the N1, N2, P2 waves that precede the P300 wave, nor in the latency of the P300 itself. **Conclusions:** The results suggest that exposure to GSM EMF exerts some effects on CNS, including effects on long latency ERPs.

Hinrichs H, Heinze HJ. Effects of GSM electromagnetic field on the MEG during an encoding-retrieval task. *Neuroreport*. 15(7):1191-1194, 2004.

Potential effects of GSM 1800 electromagnetic fields (EMF) on verbal memory encoding were investigated by recording event-related magnetic fields (ERMF) from the brain during subsequent memory retrieval. Twelve normal subjects participated in the study. After encoding words from a study list presented in the first phase they had to discriminate old from new words mixed together in a test list presented during the second phase. All subjects performed two experimental sessions, one with exposure to EMF during the study phase, and one without. Exposure to EMF changed an early (350-400 ms) task-specific component of the ERMF indicating an interference of EMF and item encoding. Behavioural measures were not significantly affected. Adverse health effects cannot be derived from these data.

Effects on Reproductive Organs and Fertility Effects

Magras, IN, Xenos, TD, RF radiation-induced changes in the prenatal development of mice. *Bioelectromagnetics* 18(6):455-461, 1997.

The possible effects of radiofrequency (RF) radiation on prenatal development has been investigated in mice. This study consisted of RF level measurements and in vivo experiments at several places around an "antenna park." At these locations RF power densities between 168 nW/cm² and 1053 nW/cm² were measured. Twelve pairs of mice, divided in two groups, were placed in locations of different power densities and were repeatedly mated five times. One hundred eighteen newborns were collected. They were measured, weighed, and examined macro- and microscopically. A progressive decrease in the number of newborns

Studies that show **Cell Tower** Health Effects

per dam was observed, which ended in irreversible infertility. The prenatal development of the newborns, however, evaluated by the crown-rump length, the body weight, and the number of the lumbar, sacral, and coccygeal vertebrae, was improved.

Jelodar G, Nazifi S, Akbari A. The prophylactic effect of vitamin C on induced oxidative stress in rat testis following exposure to 900 MHz radio frequency wave generated by a BTS antenna model. Electromagn Biol Med. 2013 Jan 16. [Epub ahead of print]
Radio frequency wave (RFW) generated by base transceiver station (BTS) has been reported to make deleterious effects on reproduction, possibly through oxidative stress. This study was conducted to evaluate the effect of RFW generated by BTS on oxidative stress in testis and the prophylactic effect of vitamin C by measuring the antioxidant enzymes activity, including glutathione peroxidase, superoxide dismutase (SOD) and catalase, and malondialdehyde (MDA). Thirty-two adult male Sprague-Dawley rats were randomly divided into four experimental groups and treated daily for 45 days as follows: sham, sham+vitamin C (l-ascorbic acid 200 mg/kg of body weight/day by gavage), RFW (exposed to 900 MHz RFW) 'sham' and 'RFW' animals were given the vehicle, i.e., distilled water and the RFW+vitamin C group (received vitamin C in addition to exposure to RFW). At the end of the experiment, all the rats were sacrificed and their testes were removed and used for measurement of antioxidant enzymes and MDA activity. The results indicate that exposure to RFW in the test group decreased antioxidant enzymes activity and increased MDA compared with the control groups ($p < 0.05$). In the treated group, vitamin C improved antioxidant enzymes activity and reduced MDA compared with the test group ($p < 0.05$). It can be concluded that RFW causes oxidative stress in testis and vitamin C improves the antioxidant enzymes activity and decreases MDA.

Lukac N, Massanyi P, Roychoudhury S, Capcarova M, Tvrda E, Knazicka Z, Kolesarova A, Danko J. In vitro effects of radiofrequency electromagnetic waves on bovine spermatozoa motility. J Environ Sci Health A Tox Hazard Subst Environ Eng. 46(12):1417-1423, 2011.

In this study the effects of 1800 MHz GSM-like radiofrequency electromagnetic waves (RF-EMW) exposure on bovine semen was monitored. The experimental samples were analyzed in vitro in four time periods (0, 30, 120 and 420 min) and compared with unexposed samples (control). Spermatozoa motility was determined by computer assisted semen analyzer (CASA). Evaluation of the percentage of motile spermatozoa showed significant ($P < 0.001$) decrease in experimental groups after 120 and 420 min of culture when exposed to microwaves, in comparison to control. Similar spermatozoa motility inhibition was detected for the percentage of progressively motile spermatozoa, too. Average path distance decreased significantly ($p < 0.001$) in experimental groups after 30 and 420 min of culture. Path velocity increased in the experimental groups exposed to RF-EMW after 30 minutes of culture, but subsequently decreased after 420 min of culture, in comparison to control. This indicates a possible initial stimulation and subsequent velocity inhibition of bovine spermatozoa under RF-EMW exposure. Changes in spermatozoa motility were also detected for some fine parameters, too. A significant decrease ($P < 0.001$) was noted for amplitude of lateral

Studies that show **Cell Tower** Health Effects

head displacement in the experimental group after 420 minutes of culture. Detailed in vitro motility analysis of bovine spermatozoa exposed to microwave radiation suggested that the parameters of path and velocity at the beginning of the culture significantly increase, but after longer culture (420 minutes) a significant decrease occur in the experimental group as compared to control. In general, results of this experiment indicate a negative time-dependent effect of 1800 MHz RF-EMW radiation on bovine spermatozoa motility.

Effects on Wellbeing

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. Study of the health of people living in the vicinity of mobile phone base stations: I. Influence of distance and sex. Pathol Biol (Paris) 50(6):369-373, 2002. [Article in French]

A survey study using questionnaire was conducted in 530 people (270 men, 260 women) living or not in vicinity of cellular phone base stations, on 18 Non Specific Health Symptoms. Comparisons of complaints frequencies (CHI-SQUARE test with Yates correction) in relation with distance from base station and sex, show significant ($p < 0.05$) increase as compared to people living > 300 m or not exposed to base station, till 300 m for tiredness, 200 m for headache, sleep disturbance, discomfort, etc. 100 m for irritability, depression, loss of memory, dizziness, libido decrease, etc. Women significantly more often than men ($p < 0.05$) complained of headache, nausea, loss of appetite, sleep disturbance, depression, discomfort and visual perturbations. This first study on symptoms experienced by people living in vicinity of base stations shows that, in view of radioprotection, minimal distance of people from cellular phone base stations should not be < 300 m.

Santini R, Santini P, Le Ruz P, Danze JM, Seigne M, Survey study of people living in the vicinity of cellular phone base stations. Electromag Biol Med 22:41-49, 2003.

A survey study was conducted, using a questionnaire, on 530 people (270 men, 260 women) living or not in proximity to cellular phone base stations. Eighteen different symptoms (Non Specific Health Symptoms-NSHS), described as radiofrequency sickness, were studied by means of the chi-square test with Yates correction. The results that were obtained underline that certain complaints are experienced only in the immediate vicinity of base stations (up to 10 m for nausea, loss of appetite, visual disturbances), and other at greater distances from base stations (up to 100 m for irritability, depressive tendencies, lowering of libido, and up to 200 m for headaches, sleep disturbance, feeling of discomfort). In the 200 m to 300 m zone, only the complaint of fatigue is experienced significantly more often when compared with subjects residing at more than 300 m or not exposed (reference group). For seven of the studied symptoms and for the distance up to 300 m, the frequency of reported complaints is significantly higher ($P < 0.05$) for women in comparison to men. Significant differences are also observed in relation to the ages of subjects, and for the location of subjects in relation to the antennas and to other electromagnetic factors.

Studies that show **Cell Tower** Health Effects

Navarro EA, Sequera J, Portoles M, Gomez-Perretta de Mateo C. The Microwave Syndrome: A Preliminary Study in Spain. *Electromag Biol Med* 22:161-169, 2003.

A health survey was carried out in Murcia, Spain, in the vicinity of a Cellular Phone Base Station working in DCS-1800 MHz. This survey contained health items related to "microwave sickness" or "RF syndrome." The microwave power density was measured at the respondents' homes. Statistical analysis showed significant correlation between the declared severity of the symptoms and the measured power density. The separation of respondents into two different exposure groups also showed an increase of the declared severity in the group with the higher exposure.

Lerchl A, Krüger H, Niehaus M, Streckert JR, Bitz AK, Hansen V. Effects of mobile phone electromagnetic fields at nonthermal SAR values on melatonin and body weight of Djungarian hamsters (*Phodopus sungorus*). *J Pineal Res.* 44(3):267-272, 2008.

In three experiments, adult male Djungarian hamsters (*Phodopus sungorus*) were exposed 24 hr/day for 60 days to radio frequency electromagnetic fields (RF-EMF) at 383, 900, and 1800 MHz, modulated according to the TETRA (383 MHz) and GSM standards (900 and 1800 MHz), respectively. A radial waveguide system ensured a well defined and uniform exposure at whole-body averaged specific absorption rates of 80 mW/kg, which is equal to the upper limit of whole-body exposure of the general population in Germany and other countries. For each experiment, using two identical waveguides, hamsters were exposed (n = 120) and sham-exposed (n = 120) in a blind fashion. In all experiments, pineal and serum melatonin levels as well as the weights of testes, brain, kidneys, and liver were not affected. At 383 MHz, exposure resulted in a significant transient increase in body weight up to 4%, while at 900 MHz this body weight increase was more pronounced (up to 6%) and not transient. At 1800 MHz, no effect on body weight was seen. The results corroborate earlier findings which have shown no effects of RF-EMF on melatonin levels in vivo and in vitro. The data are in accordance with the hypothesis that absorbed RF energy may result in metabolic changes which eventually cause body weight increases in exposed animals. The data support the notion that metabolic effects of RF-EMFs need to be investigated in more detail in future studies.

Kato Y, Johansson O. Reported functional impairments of electrosensitive Japanese: A questionnaire survey. *Pathophysiology*.19(2) 95-100, 2012.

An increasing number of people worldwide complain that they have become electromagnetic hypersensitive (EHS). We conducted a questionnaire survey of EHS persons in Japan. The aim was to identify electromagnetic fields (EMF) and plausible EMF sources that caused their symptoms. Postal questionnaires were distributed via a self-help group, and 75 participants (95% women) responded. Reported major complaints were "fatigue/tiredness" (85%), "headache", "concentration, memory, and thinking" difficulty (81%, respectively). Seventy-two per cent used some form of complementary/alternative therapy. The most plausible trigger of EHS onset was a mobile phone base station or personal handy-phone system (37%). Sixty-five percent experienced health problems to be due to the radiation from other passengers' mobile

Studies that show **Cell Tower** Health Effects

phones in trains or buses, and 12% reported that they could not use public transportation at all. Fifty-three percent had a job before the onset, but most had lost their work and/or experienced a decrease in income. Moreover, 85.3% had to take measures to protect themselves from EMF, such as moving to low EMF areas, or buying low EMF electric appliances. EHS persons were suffering not only from their symptoms, but also from economical and social problems.

Hutter HP, Moshammer H, Wallner P, Kundi M. Subjective symptoms, sleeping problems, and cognitive performance in subjects living near mobile phone base stations. *Occup Environ Med.* 63(5):307-313, 2006.

BACKGROUND: The erection of mobile telephone base stations in inhabited areas has raised concerns about possible health effects caused by emitted microwaves.

METHODS: In a cross-sectional study of randomly selected inhabitants living in urban and rural areas for more than one year near to 10 selected base stations, 365 subjects were investigated. Several cognitive tests were performed, and wellbeing and sleep quality were assessed. Field strength of high-frequency electromagnetic fields (HF-EMF) was measured in the bedrooms of 336 households. RESULTS: Total HF-EMF and exposure related to mobile telecommunication were far below recommended levels (max. 4.1 mW/m²). Distance from antennae was 24-600 m in the rural area and 20-250 m in the urban area. Average power density was slightly higher in the rural area (0.05 mW/m²) than in the urban area (0.02 mW/m²). Despite the influence of confounding variables, including fear of adverse effects from exposure to HF-EMF from the base station, there was a significant relation of some symptoms to measured power density; this was highest for headaches. Perceptual speed increased, while accuracy decreased insignificantly with increasing exposure levels. There was no significant effect on sleep quality. CONCLUSION: Despite very low exposure to HF-EMF, effects on wellbeing and performance cannot be ruled out, as shown by recently obtained experimental results; however, mechanisms of action at these low levels are unknown.

Bortkiewicz A, Zmyslony M, Szyjkowska A, Gadzicka E. [Subjective symptoms reported by people living in the vicinity of cellular phone base stations: a review of the studies] *Med Pr.* 55(4):345-351, 2004. [Article in Polish]

The problem of health effects of electromagnetic fields (EMF) emitted by cellular phone base stations evokes much interest in view of the fact that people living in their vicinity are fated to continuous exposure to EMF. None of the studies carried out throughout the world have revealed excessive values of standards adopted by the International Commission on Non-Ionizing Radiation Protection (ICNIRP). A questionnaire was used as a study tool. The results of the questionnaire survey reveal that people living in the vicinity of base stations report various complaints mostly of the circulatory system, but also of sleep disturbances, irritability, depression, blurred vision, concentration difficulties, nausea, lack of appetite, headache and vertigo. The performed studies showed the relationship between the incidence of individual symptoms, the level of exposure, and the distance between a residential area and a base station. This association was observed in both groups of persons, those who linked their complaints

Studies that show **Cell Tower** Health Effects

with the presence of the base station and those who did not notice such a relation. Further studies, clinical and those based on questionnaires, are needed to explain the background of reported complaints.

Bortkiewicz A, Gadzicka E, Szykowska A, Politański P, Mamrot P, Szymczak W, Zmyślony M. Subjective complaints of people living near mobile phone base stations in Poland. *Int J Occup Med Environ Health*. 25(1):31-40, 2012.

OBJECTIVES: The aim of our study was to assess the health conditions and subjective symptoms of the inhabitants living in the base stations vicinity and to analyse the relationship between the complaints and level of exposure to electromagnetic fields (EMF). **MATERIALS AND METHODS:** Our study was performed in housing estates located in five regions of Łódź. The electric field measurements were performed in the buildings located closest to the azimuth of the antennas. Respondents were selected by trained interviewers using an uniform procedure. The number of the households to be examined was set at a minimum of 420. The questionnaire contained: demographic data, occupational and environmental exposure to EMF, health condition, subjective complaints. Results were adjusted for confounders (age, gender, EMF at the workplace and EMF emitted by household equipment) using multiple regression model. **RESULTS:** 181 men and 319 women from 500 households were examined. Electric field above 0.8 V/m was recorded in 12% of flats. There was no significant correlation between electric field strength and the distance of examined flats from the base stations. To make possible comparison with relevant literature, we analysed also the frequency of the reported symptoms vs. the distance. Headache was declared by 57% people, most frequently (36.4%) living 100-150 m away from the base station compared to people living at longer distances ($p = 0.013$). 24.4% subjects, mostly living at a distance above 150 m, declared impaired memory. Difference was statistically significant in comparison with people living at other distances ($p = 0.004$). **CONCLUSIONS:** The explanation why we did not find any correlation between the electric field strength and frequency of subjective symptoms but found a correlation between subjective symptoms and distance from base station needs further studies. Maybe new metrics of exposure assessment should be adopted for this purpose.

Augner C, Florian M, Pauser G, Oberfeld G, Hacker GW. GSM base stations: Short-term effects on well-being. *Bioelectromagnetics*. 30:73-80, 2009.

The purpose of this study was to examine the effects of short-term GSM (Global System for Mobile Communications) cellular phone base station RF-EMF (radiofrequency electromagnetic fields) exposure on psychological symptoms (good mood, alertness, calmness) as measured by a standardized well-being questionnaire. Fifty-seven participants were selected and randomly assigned to one of three different exposure scenarios. Each of those scenarios subjected participants to five 50-min exposure sessions, with only the first four relevant for the study of psychological symptoms. Three exposure levels were created by shielding devices in a field laboratory, which could be installed or removed during the breaks between sessions such that double-blinded conditions prevailed. The overall median power flux densities were 5.2 microW/m(2) during "low," 153.6 microW/m(2) during "medium," and 2126.8 microW/m(2) during

Studies that show **Cell Tower** Health Effects

"high" exposure sessions. For scenario HM and MH, the first and third sessions were "low" exposure. The second session was "high" and the fourth was "medium" in scenario HM; and vice versa for scenario MH. Scenario LL had four successive "low" exposure sessions constituting the reference condition. Participants in scenarios HM and MH (high and medium exposure) were significantly calmer during those sessions than participants in scenario LL (low exposure throughout) ($P = 0.042$). However, no significant differences between exposure scenarios in the "good mood" or "alertness" factors were obtained. We conclude that short-term exposure to GSM base station signals may have an impact on well-being by reducing psychological arousal.

Augner C, Hacker GW. Are people living next to mobile phone base stations more strained? Relationship of health concerns, self-estimated distance to base station, and psychological parameters. *Indian J Occup Environ Med.* 13(3):141-145, 2009.

BACKGROUND AND AIMS: Coeval with the expansion of mobile phone technology and the associated obvious presence of mobile phone base stations, some people living close to these masts reported symptoms they attributed to electromagnetic fields (EMF). Public and scientific discussions arose with regard to whether these symptoms were due to EMF or were nocebo effects. The aim of this study was to find out if people who believe that they live close to base stations show psychological or psychobiological differences that would indicate more strain or stress. Furthermore, we wanted to detect the relevant connections linking self-estimated distance between home and the next mobile phone base station (DBS), daily use of mobile phone (MPU), EMF-health concerns, electromagnetic hypersensitivity, and psychological strain parameters. DESIGN, MATERIALS AND METHODS: Fifty-seven participants completed standardized and non-standardized questionnaires that focused on the relevant parameters. In addition, saliva samples were used as an indication to determine the psychobiological strain by concentration of alpha-amylase, cortisol, immunoglobulin A (IgA), and substance P. RESULTS: Self-declared base station neighbors (DBS \leq 100 meters) had significantly higher concentrations of alpha-amylase in their saliva, higher rates in symptom checklist subscales (SCL) somatization, obsessive-compulsive, anxiety, phobic anxiety, and global strain index PST (Positive Symptom Total). There were no differences in EMF-related health concern scales. CONCLUSIONS: We conclude that self-declared base station neighbors are more strained than others. EMF-related health concerns cannot explain these findings. Further research should identify if actual EMF exposure or other factors are responsible for these results.

Blettner M, Schlehofer B, Breckenkamp J, Kowall B, Schmiedel S, Reis U, Potthoff P, Schuez J, Berg-Beckhoff G. Mobile phone base stations and adverse health effects: Phase 1: A population-based cross-sectional study in Germany. *Occup Environ Med.* 66(2):118-123. 2009.

Abstract OBJECTIVE: The aim of this first phase of a cross-sectional study from Germany was to investigate whether proximity of residence to mobile phone base stations as well as risk perception is associated with health complaints. METHODS: We conducted a population-based multi-phase cross-sectional study within the context of a large panel

Studies that show **Cell Tower** Health Effects

survey regularly carried out by a private research institute in Germany. In the initial phase, which we will report on in this paper, 30,047 persons from a total of 51,444 who took part in the nationwide survey also answered questions on how mobile phone base stations affect their health. A list of 38 health complaints was used. A multiple linear regression model was used to identify predictors of health complaints including proximity of residence to mobile phone base stations and risk perception. RESULTS: Of the 30,047 participants (response rate 58.6%), 18.7% of participants were concerned about adverse health effects of mobile phone base stations, while an additional 10.3% attributed their personal adverse health effects to the exposure from them. Participants who are concerned about or attribute adverse health effects to mobile phone base stations and those living in the vicinity of a mobile phone base station (500 m) reported slightly more health complaints than others. CONCLUSIONS: A substantial proportion of the German population is concerned about adverse health effects caused by exposure from mobile phone base stations. The observed slightly higher prevalence of health complaints near base stations can however not be fully explained by attributions or concerns.

Effects on Sleep

Mohammed HS, Fahmy HM, Radwah NM, Elsayed AA. Non-thermal continuous and modulated electromagnetic radiation fields effects on sleep EEG of rats. J Adv Res 4(2) 181-187, 2013.

In the present study, the alteration in the sleep EEG in rats due to chronic exposure to low-level non-thermal electromagnetic radiation was investigated. Two types of radiation fields were used; 900 MHz unmodulated wave and 900 MHz modulated at 8 and 16 Hz waves. Animals has exposed to radiation fields for 1 month (1 h/day). EEG power spectral analyses of exposed and control animals during slow wave sleep (SWS) and rapid eye movement sleep (REM sleep) revealed that the REM sleep is more susceptible to modulated radiofrequency radiation fields (RFR) than the SWS. The latency of REM sleep increased due to radiation exposure indicating a change in the ultradian rhythm of normal sleep cycles. The cumulative and irreversible effect of radiation exposure was proposed and the interaction of the extremely low frequency radiation with the similar EEG frequencies was suggested.

Liu H, Chen G, Pan Y, Chen Z, Jin W, Sun C, Chen C, Dong X, Chen K, Xu Z, Zhang S, Yu Y. (2014) Occupational Electromagnetic Field Exposures Associated with Sleep Quality: A Cross-Sectional Study. PLoS ONE 9(10): e110825. doi:10.1371/journal.pone.0110825.

BACKGROUND: Exposure to electromagnetic field (EMF) emitted by mobile phone and other machineries concerns half the world's population and raises the problem of their impact on human health. The present study aims to explore the effects of electromagnetic field exposures on sleep quality and sleep duration among workers from electric power plant. METHODS: A cross-sectional study was conducted in an

Studies that show **Cell Tower** Health Effects

electric power plant of Zhejiang Province, China. A total of 854 participants were included in the final analysis. The detailed information of participants was obtained by trained investigators using a structured questionnaire, which including socio-demographic characteristics, lifestyle variables, sleep variables and electromagnetic exposures. Physical examination and venous blood collection were also carried out for every study subject. **RESULTS:** After grouping daily occupational electromagnetic exposure into three categories, subjects with long daily exposure time had a significantly higher risk of poor sleep quality in comparison to those with short daily exposure time. The adjusted odds ratios were 1.68 (95%CI: 1.18, 2.39) and 1.57 (95%CI: 1.10, 2.24) across tertiles. Additionally, among the subjects with long-term occupational exposure, the longer daily occupational time apparently increased the risk of poor sleep quality (OR (95%CI): 2.12 (1.23~3.66) in the second tertile; 1.83 (1.07~3.15) in the third tertile). There was no significant association of long-term occupational exposure duration, monthly electric fee or years of mobile-phone use with sleep quality or sleep duration. **CONCLUSIONS:** The findings showed that daily occupational EMF exposure was positively associated with poor sleep quality. It implies EMF exposure may damage human sleep quality rather than sleep duration.

Hung CS, Anderson C, Horne JA, McEvoy P. Mobile phone 'talk-mode' signal delays EEG-determined sleep onset. *Neurosci Lett.* 421: 82-86, 2007.

Mobile phones signals are pulse-modulated microwaves, and EEG studies suggest that the extremely low-frequency (ELF) pulse modulation has sleep effects. However, 'talk', 'listen' and 'standby' modes differ in the ELF (2, 8, and 217Hz) spectral components and specific absorption rates, but no sleep study has differentiated these modes. We used a GSM900 mobile phone controlled by a base-station simulator and a test SIM card to simulate these three specific modes, transmitted at 12.5% (23dBm) of maximum power. At weekly intervals, 10 healthy young adults, sleep restricted to 6h, were randomly and single-blind exposed to one of: talk, listen, standby and sham (nil signal) modes, for 30min, at 13:30h, whilst lying in a sound-proof, lit bedroom, with a thermally insulated silent phone beside the right ear. Bipolar EEGs were recorded continuously, and subjective ratings of sleepiness obtained every 3min (before, during and after exposure). After exposure the phone and base-station were switched off, the bedroom darkened, and a 90min sleep opportunity followed. We report on sleep onset using: (i) visually scored latency to onset of stage 2 sleep, (ii) EEG power spectral analysis. There was no condition effect for subjective sleepiness. Post-exposure, sleep latency after talk mode was markedly and significantly delayed beyond listen and sham modes. This condition effect over time was also quite evident in 1-4Hz EEG frontal power, which is a frequency range particularly sensitive to sleep onset. It is possible that 2, 8, 217Hz modulation may differentially affect sleep onset.

Studies that show **Cell Tower** Health Effects

Effects on Cells

Neshev NN, Kirilova EI, Environmental-health aspects of pulse-modulated microwaves. Rev Environ Health 11(1-2):85-88, 1996.

Our theoretical model describes the potential influence of irradiation with pulse-modulated microwaves on the conformational oscillations of enzymes in living organisms. Certain values of pulse-repetition time, determined by the period of conformational oscillations of the corresponding type of enzyme, can produce the effect at extremely low power levels. Synchronized oscillations in identical enzyme molecules produce in turn large-scale oscillations within living cells. Thus, short periods of exposure to pulse-modulated microwaves could be beneficial to cellular function, whereas maintaining the amplitude of such oscillations at a maximum for long periods may have a stressful effect on biochemical processes. The model discloses the possible environmental-health risks of long-term exposure in ambient fields that are created by radar, navigation, and communication systems.

Kwee S, Raskmark P, Changes in cell proliferation due to environmental non-ionizing radiation 2. Microwave radiation. Bioelectrochem Bioenerg 44(2) 251-255, 1998.

Due to the use of mobile telephones, there is an increased exposure of the environment to weak radiofrequency (RF) electromagnetic fields, emitted by these devices. This study was undertaken to investigate if the microwave radiation from these fields will have a similar effect on cell proliferation as weak electromagnetic (ELF) fields. The field was generated by signal simulation of the Global System for Mobile communications (GSM) of 960 MHz. Cell cultures, growing in microtiter plates, were exposed in a specially constructed chamber, a Transverse Electromagnetic (TEM) cell. The Specific Absorption Rate (SAR) values for each cell well were calculated for this exposure system. Experiments were performed on cell cultures of transformed human epithelial amnion cells (AMA), which were exposed to 960 MHz microwave fields at three different power levels and three different exposure times, respectively. It was found that cell growth in the exposed cells was decreased in comparison to that in the control and sham exposed cells. Cell proliferation during the period following exposure varied not only with the various SAR levels, but also with the length of exposure time. On the other hand, repeated periods of exposure did not seem to change the effects. There was a general linear correlation between power level and growth change. However, the exposure time required to obtain the maximum effect was not the same for the various power levels. It turned out that at low power level, a maximum effect was first reached after a longer exposure time than at higher power level. A similar phenomenon was registered in the studies on ELF electromagnetic fields. Here, it was found that there was a linear correlation between the length of exposure time to obtain maximum effect and field strength.

Studies that show **Cell Tower** Health Effects

Effects on Eyes

Lu L, Xu H, Wang X, Guo G. Increased nitric oxide synthase activity is essential for electromagnetic-pulse-induced blood-retinal barrier breakdown in vivo. *Brain Res.* 1264:104-10, 2009.

PURPOSE: To examine whether electromagnetic pulses (EMPs) affected the permeability of the **blood-retinal barrier (BRB)**, gene expression of occludin and activity of nitric oxide synthase (NOS). **METHODS:** Sprague-Dawley (SD) rats were used and randomized into EMP and control groups. Retinas were removed immediately, and 2 h or 24 h after EMP radiation. BRB permeability was analyzed by transmission electron microscopy and Evans Blue staining. Retinal NOS activity and concentrations of nitrite and nitrate were measured. Occludin mRNA and protein levels were detected by RT-PCR and Western blotting. **RESULTS:** Exposure of SD rats to EMP resulted in increased BRB permeability, with the greatest decrease in occludin at 24 h. Moreover, this permeability defect was also correlated with significant increases in the formation of NO and induction of NOS activity in SD rats. Furthermore, we found that treatment with NOS inhibitor N-nitro-L-arginine methyl ester (L-NAME) blocked BRB breakdown and prevented the increase in NO formation and induction of NOS activity, as well as the decrease in occluding expression. **CONCLUSION:** Taken together, these results support the view that NOS-dependent NO production is an important factor that contributes to EMP-induced BRB dysfunction, and suggests that NOS induction may play an important role in BRB breakdown.

Hässig M, Jud F, Naegeli H, Kupper J, Spiess B. Prevalence of nuclear cataract in Swiss veal calves and its possible association with mobile telephone antenna base stations. *Schweiz Arch Tierheilkd.* 151(10):471-478, 2009.

The purpose of this study was to evaluate the prevalence of nuclear cataract in veal calves and to elucidate a possible impact by mobile phone base stations (MPBS). For this experiment a cohort study was conducted. A follow-up of the geographical location of each dam and its calf from conception through the fetal period up to slaughter was performed. The first trimester of gestation (organogenesis) was particularly emphasized. The activities of selected protective antioxidants (superoxide dismutase, catalase, glutathione peroxidase [GPx]) were assessed in aqueous humor of the eye to evaluate the redox status. Of 253 calves, 79 (32 %) had various degrees of nuclear cataract, but only 9 (3.6 %) calves had severe nuclear cataract. Results demonstrate a relation between the location of veal calves with nuclear cataracts in the first trimester of gestation and the strength of antennas. The number of antennas within 100 to 199 meters was associated with oxidative stress and there was an association between oxidative stress and the distance to the nearest MPBS. Oxidative stress was increased in eyes with cataract (OR per kilometer: 0.80, confidence interval 95 % 0.62, 0.93). It has not been shown that the antennas actually affected stress. Hosmer-Lemeshow statistics showed an accuracy of 100 % in negative cases with low radiation, and only 11.11 % accuracy in positive cases with high radiation. This reflects, that there are a lot of other

Studies that show **Cell Tower** Health Effects

possibilities for nuclear cataract beside MPBS. Further studies on the influence of electromagnetic fields during embryonic development animal or person at risk are indicated.

Hässig M, Jud F, Spiess B. [Increased occurrence of nuclear cataract in the calf after erection of a mobile phone base station]. *Schweiz Arch Tierheilkd.* 154(2):82-86, 2012.[Article in German]

We examined and monitored a dairy farm in which a large number of calves were born with nuclear cataracts after a mobile phone base station had been erected in the vicinity of the barn. Calves showed a 3.5 times higher risk for heavy cataract if born there compared to Swiss average. All usual causes such as infection or poisoning, common in Switzerland, could be excluded. The real cause of the increased incidence of cataracts remains unknown.

Jelodar G, Akbari A, Nazifi S. The prophylactic Effect of Vitamin C on Oxidative Stress Indexes in Rat Eyes Following Exposure to Radiofrequency Wave Generated by a BTS Antenna Model. *Int J Radiat Biol.* 89(2):128-131, 2013.

Purpose: This study was conducted to evaluate the effect of radiofrequency wave (RFW)-induced oxidative stress in the eye and the prophylactic effect of vitamin C on this organ by measuring the antioxidant enzymes activity including: glutathione peroxidase (GPx), superoxide dismutase (SOD) and catalase (CAT), and malondialdehyde (MDA). Materials and methods: Thirty-two adult male Sprague-Dawley rats were randomly divided into four experimental groups and treated daily for 45 days as follows: control, vitamin C (L-ascorbic acid 200 mg/kg of body weight/day by gavage), test (exposed to 900 MHz RFW) and the treated group (received vitamin C in addition to exposure to RFW). At the end of the experiment all animals were killed, their eyes were removed and were used for measurement of antioxidant enzymes and MDA activity. Results: The results indicate that exposure to RFW in the test group decreased antioxidant enzymes activity and increased MDA compared with the control groups (P<0.05). In the treated group vitamin C improved antioxidant enzymes activity and reduced MDA compared to the test group (P<0.05). Conclusions: It can be concluded that RFW causes oxidative stress in the eyes and vitamin C improves the antioxidant enzymes activity and decreases MDA.

Oxidative Stress

Achudume A, Onibere B, Aina F, Tchokossa P. Induction of oxidative stress in male rats subchronically exposed to electromagnetic fields at non-thermal intensities. *J Electromagnetic Analysis and Applications* 2(8), 482-487, 2010.

To investigate the oxidative stress-inducing potential of non-thermal electromagnetic fields in rats. Male Wister rats were exposed to electrical field intensity of $2.3 \pm 0.82 \mu\text{V/m}$. Exposure was in three forms: continuous waves, or modulated at 900 MHz or modulated GSM-nonDTX. The radio frequency radiation (RFR) was 1800 MHz, specific

Studies that show **Cell Tower** Health Effects

absorption radiation (SAR) (0.95-3.9 W/kg) for 40 and/or 60 days continuously. Control animals were located > 300 m from base station, while sham control animals were located in a similar environmental conditions, but in the vicinity of a non-functional base station. The rats were assessed for thiobarbituric and reactive species (TBARS), reduced glutathione (GSH) content, catalase activity, glutathione reductase (GR) and glucose residue after 40 and 60 days of exposure. At 40 days, electromagnetic radiation failed to induce any significant alterations. However, at 60 days of exposure various attributes evaluated decreased. The respective decreases in both nicotinamide adenine dinucleotide phosphate (NADPH) and Ascorbate- linked lipid peroxidation (LPO) with concomitant diminution in enzymatic antioxidative defense systems resulted in decreased glucose residue. The present studies showed some biochemical changes that may be associated with a prolong exposure to electromagnetic fields and its relationship to the activity of antioxidant system in rat Regular assessment and early detection of antioxidative defense system among people working around the base stations are recommended.

Augner C, Hacker GW, Oberfeld G, Florian M, Hitzl W, Hutter J, Pauser G. Effects of Exposure to GSM mobile phone base station signals on salivary cortisol, alpha-amylase, and Immunoglobulin A. *Biomed Environ Sci.* 23(3):199-207, 2010.

OBJECTIVE: The present study aimed to test whether exposure to radiofrequency electromagnetic fields (RF-EMF) emitted by mobile phone base stations may have effects on salivary alpha-amylase, immunoglobulin A (IgA), and cortisol levels. METHODS: Fifty seven participants were randomly allocated to one of three different experimental scenarios (22 participants to scenario 1, 26 to scenario 2, and 9 to scenario 3). Each participant went through five 50-minute exposure sessions. The main RF-EMF source was a GSM-900-MHz antenna located at the outer wall of the building. In scenarios 1 and 2, the first, third, and fifth sessions were "low" (median power flux density 5.2 $\mu\text{W}/\text{m}^2$) exposure. The second session was "high" (2126.8 $\mu\text{W}/\text{m}^2$), and the fourth session was "medium" (153.6 $\mu\text{W}/\text{m}^2$) in scenario 1, and vice versa in scenario 2. Scenario 3 had four "low" exposure conditions, followed by a "high" exposure condition. Biomedical parameters were collected by saliva samples three times a session. Exposure levels were created by shielding curtains. RESULTS: In scenario 3 from session 4 to session 5 (from "low" to "high" exposure), an increase of cortisol was detected, while in scenarios 1 and 2, a higher concentration of alpha-amylase related to the baseline was identified as compared to that in scenario 3. IgA concentration was not significantly related to the exposure. CONCLUSIONS: RF-EMF in considerably lower field densities than ICNIRP-guidelines may influence certain psychobiological stress markers.

Marzook EA, Abd El Moneim AE, Elhadary AA. Prootective role of seame oil against mobile phone base station-induced oxidative stress. *J Rad Res Appl Sci* 7(1):1-6, 2014.

The present study was undertaken to shed the light on the environmental threats associated with the wireless revolution and the health hazards associated with exposure to mobile base station (MBS). Besides, studying the possible protective role of sesame oil (SO) as an antioxidant against oxidative stress. Therefore, the present work was

Studies that show **Cell Tower** Health Effects

designed to study the effect of chronic exposure to electromagnetic radiations (EMR), produced by a cellular tower for mobile phone and the possible protective role of sesame oil on glutathione reductase (GSH-Rx), superoxide dismutase (SOD), catalase (CAT), total testosterone and lipid profile (total cholesterol (Tch), triglycerides (TG), low density lipoprotein cholesterol (LDL-c) and high density lipoprotein cholesterol (HDL-c) in male albino rats. Rats were arranged into four groups: the control unexposed, the exposed untreated and the exposed treated groups (1.5 and 3 ml oil). Exposed groups were subjected to electromagnetic field at frequency of 900 MHz, for 24 h/day for 8 weeks, at the same time both treated groups were supplied with oral injection of sesame oil three times per week. At the end of the experiment, blood samples were obtained for determination of the above mentioned variables in serum. The results obtained revealed that TG and testosterone were raised significantly over control in all groups and the significant increase in oil groups occurred in dose dependent manner. SOD and CAT activities were reduced significantly in exposed rats than control and increased significantly in sesame oil groups as the dose of oil increased. Total cholesterol only showed remarkable reduction in the group treated with 3 ml sesame oil. Also, in this latter group, significant elevation of GSH-Rx was recorded. Changes in serum HDL-c and LDL-c followed an opposite trend in exposed and sesame oil groups reflecting their affection by EMR or sesame oil. In conclusion, all results of the current study proved that sesame oil can be used as an edible oil to attenuate the oxidative stress which could be yielded as a result of chronic exposure to EMR.

Effects on Blood

Kismali G, Ozgur E, Guler G, Akcay A, Sel T, Seyhan N. The influence of 1800 MHz GSM-like signals on blood chemistry and oxidative stress in non-pregnant and pregnant rabbits. *Int J Radiat Biol.* 88(5):414-419, 2012.

PURPOSE: Environmental electromagnetic fields originate from man-made sources, such as mobile phones and base stations, and have led to increasing public concern about their possible adverse health effects. We aimed to investigate the possible effects of radiofrequency radiation (RFR) generated from these devices on oversensitive animals, such as pregnant rabbits. **MATERIALS AND METHODS:** In the present study, the effects of whole body 1800 MHz Global System for Mobile Communications (GSM)-like RFR exposure for 15 min/day for seven days on blood chemistry and lipid peroxidation levels in both non-pregnant and pregnant New Zealand White rabbits were investigated. Thirteen-month-old rabbits were studied in the following four groups: Non-pregnant control, non-pregnant RFR-exposed, pregnant control and pregnant RFR-exposed. **RESULTS:** Lipid peroxidation, namely malondialdehyde (MDA) levels, did not change after RFR exposure. However, blood chemistry parameters, such as cholesterol (CHO), total protein (TP), albumin (ALB), uric acid, creatinin and **creatine kinase (CK)** and **creatine kinase-myocardial band isoenzyme (CK-MB)** changed due to both pregnancy and RFR exposure. **CONCLUSION:** Our investigations have been shown that no indication for oxidative stress was detected in the blood of pregnant rabbits upon RF exposure at specific conditions employed in the present study. Minor

Studies that show **Cell Tower** Health Effects

changes in some blood chemistry parameters were detected but CK-MB and CK increases were found remarkable. Studies on RFR exposure during pregnancy will help establish international standards for the protection of pregnant women from environmental RFR.

Yurekli AI, Ozkan M, Kalkan T, Saybasili H, Tuncel H, Atukeren P, Gumustas K, Seker S. GSM Base Station Electromagnetic Radiation and Oxidative Stress in Rats. *Electromagn Biol Med.* 2006;25(3):177-188, 2006.

The ever increasing use of cellular phones and the increasing number of associated base stations are becoming a widespread source of nonionizing electromagnetic radiation. Some biological effects are likely to occur even at low-level EM fields. In this study, a gigahertz transverse electromagnetic (GTEM) cell was used as an exposure environment for plane wave conditions of far-field free space EM field propagation at the GSM base transceiver station (BTS) frequency of 945 MHz, and effects on oxidative stress in rats were investigated. When EM fields at a power density of 3.67 W/m² (specific absorption rate = 11.3 mW/kg), which is well below current exposure limits, were applied, MDA (malondialdehyde) level was found to increase and GSH (reduced glutathione) concentration was found to decrease significantly ($p < 0.0001$). Additionally, there was a less significant ($p = 0.0190$) increase in SOD (superoxide dismutase) activity under EM exposure.

Jin YB, Lee HJ, Seon Lee J, Pack JK, Kim N, Lee YS. One-year, simultaneous combined exposure of CDMA and WCDMA radiofrequency electromagnetic fields to rats. *Int J Radiat Biol.* 87(4):416-423, 2011.

PURPOSE: We investigated whether one-year, long-term, simultaneous exposure to code division multiple access (CDMA; 849 MHz) and wideband code division multiple access (WCDMA; 1.95 GHz) radiofrequencies (RF) would induce chronic illness in Sprague-Dawley (SD) rats. **MATERIALS AND METHODS:** Two groups of 40 SD rats (50% males and females in sham and exposed groups) were exposed to CDMA and WCDMA RF simultaneously at 2.0 W/kg for 45 min/day (total 4.0 W/kg), 5 days per week for a total of one year. Body and organ weight measurements, urinalysis, haematological and blood biochemical analysis, and histopathological evaluations were performed.

RESULTS: The mortality patterns in male and female rats exposed to RF were compared with those found in gender-matched sham control animals. No significant alteration in body weight was observed with the simultaneous combined RF exposure. Most RF-exposed rats showed no significant alteration, based on urinalysis, haematology, blood biochemistry, or histopathology. However, some altered parameters of the complete blood count and serum chemistry were seen in RF-exposed rats. The total tumour incidence was not different between sham-exposed and RF-exposed animals.

CONCLUSIONS: Our results suggest that one-year chronic exposure to CDMA (849 MHz) and WCDMA (1.95 GHz) RF simultaneously at 2.0 W/kg for 45-min RF exposure periods (total, 4 W/kg) did not increase chronic illness in rats, although there were some altered parameters in the complete blood count and serum chemistry.

Studies that show **Cell Tower** Health Effects

Death

Dode AC, Leão MM, Tejo Fde A, Gomes AC, Dode DC, Dode MC, Moreira CW, Condessa VA, Albinatti C, Caiaffa WT. Mortality by neoplasia and cellular telephone base stations in the Belo Horizonte municipality, Minas Gerais state, Brazil. Sci Total Environ. 409(19):3649-3665, 2011.

Pollution caused by the electromagnetic fields (EMFs) of radio frequencies (RF) generated by the telecommunication system is one of the greatest environmental problems of the twentieth century. The purpose of this research was to verify the existence of a spatial correlation between **base station (BS)** clusters and cases of deaths by neoplasia in the Belo Horizonte municipality, Minas Gerais state, Brazil, from 1996 to 2006 and to measure the human exposure levels to EMF where there is a major concentration of cellular telephone transmitter antennas. A descriptive spatial analysis of the BSs and the cases of death by neoplasia identified in the municipality was performed through an ecological-epidemiological approach, using georeferencing. The database employed in the survey was composed of three data banks: 1. death by neoplasia documented by the Health Municipal Department; 2. BSs documented in ANATEL ("Agência Nacional de Telecomunicações": 'Telecommunications National Agency'); and 3. census and demographic city population data obtained from official archives provided by IBGE ("Instituto Brasileiro de Geografia e Estatística": 'Brazilian Institute of Geography and Statistics'). The results show that approximately 856 BSs were installed through December 2006. Most (39.60%) of the BSs were located in the "Centro-Sul" ('Central-Southern') region of the municipality. Between 1996 and 2006, 7191 deaths by neoplasia occurred and within an area of 500 m from the BS, the mortality rate was 34.76 per 10,000 inhabitants. Outside of this area, a decrease in the number of deaths by neoplasia occurred. The greatest accumulated incidence was 5.83 per 1000 in the Central-Southern region and the lowest incidence was 2.05 per 1000 in the Barreiro region. During the environmental monitoring, the largest accumulated electric field measured was 12.4 V/m and the smallest was 0.4 V/m. The largest density power was 40.78 $\mu\text{W}/\text{cm}^2$, and the smallest was 0.04 $\mu\text{W}/\text{cm}^2$.

Adang D, Remacle C, Vorst AV Results of a long-term low-level microwave exposure of rats. IEEE Trans Microwave Theor Tech 57: 2488-2497, 2009.

This paper summarizes the results of experimental research on biological effects induced by electromagnetic exposure to low-level microwaves. We exposed four-month-old Wistar albino rats during 21 months to two different microwave frequencies and exposure modes, 2 h a day, seven days a week. In order to assess possible biological effects of microwaves, we selected among others the following parameters: leucocytes, erythrocytes, monocytes, neutrophils, lymphocytes, hemoglobin, mean corpuscular hemoglobin concentration, and mortality rate. After three and eight months of exposure, we found a statistically significant difference of about 20% between the 970-MHz continuous wave group and sham-exposed group regarding the monocytes in both considered periods. After 14 and 18 months of exposure, we observed a significant

Studies that show **Cell Tower** Health Effects

increase in white blood cells and neutrophils of about 15% and 25%, respectively. Lymphocytes fell down after 18 months of exposure with about 15% compared to the sham-exposed group. No other statistically significant differences were found, except for minor changes with little biological significance. The most obvious effect we detected is the increase in mortality rate of the exposed groups with respect to the sham-exposed group after 21 months of exposure at the age of 25 months. This increase even increases when observing rats until the age of 28 months: mortality in exposed groups then reaches almost twice the value observed in the sham-exposed group.

Effects on Hormones

Eskander EF, Estefan SF, Abd-Rabou AA. How does long term exposure to base stations and mobile phones affect human hormone profiles? Clin Biochem. 45(1-2):157-161, 2012

OBJECTIVES: This study is concerned with assessing the role of exposure to radio frequency radiation (RFR) emitted either from mobiles or base stations and its relations with human's hormone profiles. **DESIGN AND METHODS:** All volunteers' samples were collected for hormonal analysis. **RESULTS:** This study showed significant decrease in volunteers' ACTH, cortisol, thyroid hormones, prolactin for young females, and testosterone levels. **CONCLUSION:** The present study revealed that high RFR effects on pituitary-adrenal axis.

Eşmekaya MA, Seyhan N, Omeroğlu S. Pulse modulated 900 MHz radiation induces hypothyroidism and apoptosis in thyroid cells: A light, electron microscopy and immunohistochemical study. Int J Radiat Biol. 86(12):1106-1116, 2010.

Purpose: In the present study we investigated the possible histopathological effects of pulse modulated Radiofrequency (RF) fields on the thyroid gland using light microscopy, electron microscopy and immunohistochemical methods. Materials and methods: Two months old male Wistar rats were exposed to a 900 MHz pulse-modulated RF radiation at a specific absorption rate (SAR) of 1.35 Watt/kg for 20 min/day for three weeks. The RF signals were pulse modulated by rectangular pulses with a repetition frequency of 217 Hz and a duty cycle of 1:8 (pulse width 0.576 ms). To assess thyroid endocrine disruption and estimate the degree of the pathology of the gland, we analysed structural alterations in follicular and colloidal diameters and areas, colloid content of the follicles, and height of the follicular epithelium. Apoptosis was confirmed by Transmission Electron Microscopy and assessing the activities of an initiator (caspase-9) and an effector (caspase-3) caspases that are important markers of cells undergoing apoptosis. Results: Morphological analyses revealed hypothyrophy of the gland in the 900 MHz RF exposure group. The results indicated that thyroid hormone secretion was inhibited by the RF radiation. In addition, we also observed formation of apoptotic bodies and increased caspase-3 and caspase-9 activities in thyroid cells of the rats that were exposed to modulated RF fields. Conclusion: The overall findings indicated that whole body exposure to pulse-modulated RF radiation that is similar to that emitted by

Studies that show **Cell Tower** Health Effects

global system for mobile communications (GSM) mobile phones can cause pathological changes in the thyroid gland by altering the gland structure and enhancing caspase-dependent pathways of apoptosis.

Genetic Effects

Gandhi G, Kaur G, Nisar U. A cross-sectional case control study on genetic damage in individuals residing in the vicinity of a mobile phone base station. Electromagn Biol Med. 2014 Jul 9:1-11. [Epub ahead of print]

Mobile phone base stations facilitate good communication, but the continuously emitting radiations from these stations have raised health concerns. Hence in this study, genetic damage using the single cell gel electrophoresis (comet) assay was assessed in peripheral blood leukocytes of individuals residing in the vicinity of a mobile phone base station and comparing it to that in healthy controls. The power density in the area within 300 m from the base station exceeded the permissive limits and was significantly ($p = 0.000$) higher compared to the area from where control samples were collected. The study participants comprised 63 persons with residences near a mobile phone tower, and 28 healthy controls matched for gender, age, alcohol drinking and occupational sub-groups. Genetic damage parameters of DNA migration length, damage frequency (DF) and damage index were significantly ($p = 0.000$) elevated in the sample group compared to respective values in healthy controls. The female residents ($n = 25$) of the sample group had significantly ($p = 0.004$) elevated DF than the male residents ($n = 38$). The linear regression analysis further revealed daily mobile phone usage, location of residence and power density as significant predictors of genetic damage. The genetic damage evident in the participants of this study needs to be addressed against future disease-risk, which in addition to neurodegenerative disorders, may lead to cancer.

Fucic A, Garaj-Vrhovac V, Skara M, Dimitrovic B, X-rays, microwaves and vinyl chloride monomer: their clastogenic and aneugenic activity, using the micronucleus assay on human lymphocytes. Mutat Res 282(4):265-271, 1992.

Chromosome aberration assays, sister-chromatid exchange techniques and micronucleus assays are commonly used methods for biomonitoring genetic material damaged by chemical or physical agents. On the other hand, their aneugenic activity, which can lead to hypoploidy and may also be associated with carcinogenesis, has not been thoroughly investigated. In our study we chose the micronucleus assay with a new mathematical approach to separate clastogenic from aneugenic activity of three well-known mutagens (vinyl chloride monomer, X-rays and microwaves) on the genome of human somatic cells. The comparison of frequencies of size distribution of micronuclei in the lymphocytes of humans exposed to each of these three mutagens showed that X-rays and microwaves were preferentially clastogens while vinyl chloride monomer showed aneugenic activity as well. Microwaves possess some mutagenic characteristics typical of chemical mutagens.

Studies that show **Cell Tower** Health Effects

Esmekaya MA, Aytekin E, Ozgur E, Güler G, Ergun MA, Omeroğlu S, Seyhan N. Mutagenic and morphologic impacts of 1.8GHz radiofrequency radiation on human peripheral blood lymphocytes (hPBLs) and possible protective role of pre-treatment with Ginkgo biloba (EGb 761). *Sci Total Environ.* 410-411:59-64, 2011.

The mutagenic and morphologic effects of 1.8GHz Global System for Mobile Communications (GSM) modulated RF (radiofrequency) radiation alone and in combination with Ginkgo biloba (EGb 761) pre-treatment in human peripheral blood lymphocytes (hPBLs) were investigated in this study using Sister Chromatid Exchange (SCE) and electron microscopy. Cell viability was assessed with 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide (MTT) reduction assay. The lymphocyte cultures were exposed to GSM modulated RF radiation at 1.8GHz for 6, 8, 24 and 48h with and without EGb 761. We observed morphological changes in pulse-modulated RF radiated lymphocytes. Longer exposure periods led to destruction of organelle and nucleus structures. Chromatin change and the loss of mitochondrial crista occurred in cells exposed to RF for 8h and 24h and were more pronounced in cells exposed for 48h. Cytoplasmic lysis and destruction of membrane integrity of cells and nuclei were also seen in 48h RF exposed cells. There was a significant increase ($p < 0.05$) in SCE frequency in RF exposed lymphocytes compared to sham controls. EGb 761 pre-treatment significantly decreased SCE from RF radiation. RF radiation also inhibited cell viability in a time dependent manner. The inhibitory effects of RF radiation on the growth of lymphocytes were marked in longer exposure periods. EGb 761 pre-treatment significantly increased cell viability in RF+EGb 761 treated groups at 8 and 24h when compared to RF exposed groups alone. The results of our study showed that RF radiation affects cell morphology, increases SCE and inhibits cell proliferation. However, EGb 761 has a protective role against RF induced mutagenity. We concluded that RF radiation induces chromosomal damage in hPBLs but this damage may be reduced by EGb 761 pre-treatment.

Kim JY, Hong SY, Lee YM, Yu SA, Koh WS, Hong JR, Son T, Chang SK, Lee M. In vitro assessment of clastogenicity of mobile-phone radiation (835 MHz) using the alkaline comet assay and chromosomal aberration test. *Environ Toxicol.* 23(3):319-327, 2008.

Recently we demonstrated that 835-MHz radiofrequency radiation electromagnetic fields (RF-EMF) neither affected the reverse mutation frequency nor accelerated DNA degradation in vitro. Here, two kinds of cytogenetic endpoints were further investigated on mammalian cells exposed to 835-MHz RF-EMF (the most widely used communication frequency band in Korean CDMA mobile phone networks) alone and in combination with model clastogens: in vitro alkaline comet assay and in vitro chromosome aberration (CA) test. No direct cytogenetic effect of 835-MHz RF-EMF was found in the in vitro CA test. The combined exposure of the cells to RF-EMF in the presence of ethylmethanesulfonate (EMS) revealed a weak and insignificant cytogenetic effect when compared to cells exposed to EMS alone in CA test. Also, the comet assay results to evaluate the ability of RF-EMF alone to damage DNA were nearly negative, although showing a small increase in tail moment. However, the applied RF-EMF had potentiation

Studies that show **Cell Tower** Health Effects

effect in comet assay when administered in combination with model clastogens (cyclophosphamide or 4-nitroquinoline 1-oxide). Thus, our results imply that we cannot confidently exclude any possibility of an increased risk of genetic damage, with important implications for the possible health effects of exposure to 835-MHz electromagnetic fields.

Nikolova T, Czyz J, Rolletschek A, Blyszczuk P, Fuchs J, Jovtchev G, Schuderer J, Kuster N, Wobus AM. Electromagnetic fields affect transcript levels of apoptosis-related genes in embryonic stem cell-derived neural progenitor cells. ASEB J. 19(12):1686-1688, 2005.

Mouse embryonic stem (ES) cells were used as an experimental model to study the effects of electromagnetic fields (EMF). ES-derived nestin-positive neural progenitor cells were exposed to extremely low frequency EMF simulating power line magnetic fields at 50 Hz (ELF-EMF) and to radiofrequency EMF simulating the Global System for Mobile Communication (GSM) signals at 1.71 GHz (RF-EMF). Following EMF exposure, cells were analyzed for transcript levels of cell cycle regulatory, apoptosis-related, and neural-specific genes and proteins; changes in proliferation; apoptosis; and cytogenetic effects. Quantitative RT-PCR analysis revealed that ELF-EMF exposure to ES-derived neural cells significantly affected transcript levels of the apoptosis-related bcl-2, bax, and cell cycle regulatory "growth arrest DNA damage inducible" GADD45 genes, whereas mRNA levels of neural-specific genes were not affected. RF-EMF exposure of neural progenitor cells resulted in down-regulation of neural-specific Nurr1 and in up-regulation of bax and GADD45 mRNA levels. Short-term RF-EMF exposure for 6 h, but not for 48 h, resulted in a low and transient increase of DNA double-strand breaks. No effects of ELF- and RF-EMF on mitochondrial function, nuclear apoptosis, cell proliferation, and chromosomal alterations were observed. We may conclude that EMF exposure of ES-derived neural progenitor cells transiently affects the transcript level of genes related to apoptosis and cell cycle control. However, these responses are not associated with detectable changes of cell physiology, suggesting compensatory mechanisms at the translational and posttranslational level.

Maes A, Collier M, Slaets D, Verschaeve L, 954 MHz microwaves enhance the mutagenic properties of mitomycin C. Environ Mol Mutagen 28(1):26-30, 1996.

This paper focuses on the combined effects of microwaves from mobile communication frequencies and a chemical DNA damaging agent mitomycin C (MMC). The investigation was performed in vitro by exposing whole blood samples to a 954 MHz emitting antenna from a GSM (Global System for Mobile Communication) base station, followed by lymphocyte cultivation in the presence of MMC. A highly reproducible synergistic effect was observed as based on the frequencies of sister chromatid exchanges in metaphase figures.

Fritze K, Wiessner C, Kuster N, Sommer C, Gass P, Hermann DM, Kiessling M, Hossmann KA, Effect of global system for mobile communication microwave exposure on the genomic response of the rat brain. Neuroscience 81(3):627-639, 1997.

Studies that show **Cell Tower** Health Effects

The acute effect of global system for mobile communication (GSM) microwave exposure on the genomic response of the central nervous system was studied in rats by measuring changes in the messenger RNAs of hsp70, the transcription factor genes c-fos and c-jun and the glial structural gene GFAP using in situ hybridization histochemistry. Protein products of transcription factors, stress proteins and marker proteins of astroglial and microglial activation were assessed by immunocytochemistry. Cell proliferation was evaluated by bromodeoxyuridine incorporation. A special GSM radiofrequency test set, connected to a commercial cellular phone operating in the discontinuous transmission mode, was used to simulate GSM exposure. The study was conducted at time averaged and brain averaged specific absorption rates of 0.3 W/kg (GSM exposure), 1.5 W/kg (GSM exposure) and 7.5 W/kg (continuous wave exposure), respectively. Immediately after exposure, in situ hybridization revealed slight induction of hsp70 messenger RNA in the cerebellum and hippocampus after 7.5 W/kg exposure, but not at lower intensities. A slightly increased expression of c-fos messenger RNA was observed in the cerebellum, neocortex and piriform cortex of all groups subjected to immobilization, but no differences were found amongst different exposure conditions. C-jun and GFAP messenger RNAs did not increase in any of the experimental groups. 24 h after exposure, immunocytochemical analysis of FOS and JUN proteins (c-FOS, FOS B, c-JUN JUN B, JUN D), of HSP70 or of KROX-20 and -24 did not reveal any alterations. Seven days after exposure, neither increased cell proliferation nor altered expression of astroglial and microglial marker proteins were observed. In conclusion, acute high intensity microwave exposure of immobilized rats may induce some minor stress response but does not result in lasting adaptive or reactive changes of the brain.

Baohong Wang, Jiliang H, Lifan J, Deqiang L, Wei Z, Jianlin L, Hongping D. Studying the synergistic damage effects induced by 1.8GHz radiofrequency field radiation (RFR) with four chemical mutagens on human lymphocyte DNA using comet assay in vitro. *Mutat Res.* 578(1-2):149-157, 2005.

The aim of this investigation was to study the synergistic DNA damage effects in human lymphocytes induced by 1.8GHz radiofrequency field radiation (RFR, SAR of 3W/kg) with four chemical mutagens, i.e. mitomycin C (MMC, DNA crosslinker), bleomycin (BLM, radiomimetic agent), methyl methanesulfonate (MMS, alkylating agent), and 4-nitroquinoline-1-oxide (4NQO, UV-mimetic agent). The DNA damage of lymphocytes exposed to RFR and/or with chemical mutagens was detected at two incubation time (0 or 21h) after treatment with comet assay in vitro. Three combinative exposure ways were used. Cells were exposed to RFR and chemical mutagens for 2 and 3h, respectively. Tail length (TL) and tail moment (TM) were utilized as DNA damage indexes. The results showed no difference of DNA damage indexes between RFR group and control group at 0 and 21h incubation after exposure ($P>0.05$). There were significant difference of DNA damage indexes between MMC group and RFR+MMC co-exposure group at 0 and 21h incubation after treatment ($P<0.01$). Also the significant difference of DNA damage indexes between 4NQO group and RFR+4NQO co-exposure group at 0 and 21h incubation after treatment was observed ($P<0.05$ or $P<0.01$). The DNA damage in RFR+BLM co-exposure groups and RFR+MMS co-exposure groups was not significantly

Studies that show **Cell Tower** Health Effects

increased, as compared with corresponding BLM and MMS groups ($P>0.05$). The experimental results indicated 1.8GHz RFR (SAR, 3W/kg) for 2h did not induce the human lymphocyte DNA damage effects in vitro, but could enhance the human lymphocyte DNA damage effects induced by MMC and 4NQO. The synergistic DNA damage effects of 1.8GHz RFR with BLM or MMS were not obvious.

Baohong W, Lifen J, Lanjuan L, Jianlin L, Deqiang L, Wei Z, Jiliang H. Evaluating the combinative effects on human lymphocyte DNA damage induced by ultraviolet ray C plus 1.8GHz microwaves using comet assay in vitro. *Toxicology*. 232(3):311-316, 2007.

The objective of this study was to observe whether 1.8GHz microwaves (MW) (SAR, 3 W/kg) exposure can influence human lymphocyte DNA damage induced by ultraviolet ray C (UVC). The lymphocytes, which were from three young healthy donors, were exposed to 254 nm UVC at the doses of 0.25, 0.5, 0.75, 1.0, 1.5 and 2.0 J m⁻², respectively. The lymphocytes were irradiated by 1.8GHz MW (SAR, 3 W/kg) for 0, 1.5 and 4 h. The combinative exposure of UVC plus MW was conducted. The treated cells were incubated for 0, 1.5 and 4 h. Finally, comet assay was used to measure DNA damage of above treated lymphocytes. The results indicated that the difference of DNA damage induced between MW group and control group was not significant ($P>0.05$). The MTLs induced by UVC were 1.71 \pm 0.09, 2.02 \pm 0.08, 2.27 \pm 0.17, 2.27 \pm 0.06, 2.25 \pm 0.12, 2.24 \pm 0.11 microm, respectively, which were significantly higher than that (0.96 \pm 0.05 microm) of control ($P<0.01$). MTLs of some sub-groups in combinative exposure groups at 1.5-h incubation were significantly lower than those of corresponding UVC sub-groups ($P<0.01$ or $P<0.05$). However, MTLs of some sub-groups in combinative exposure groups at 4-h incubation were significantly higher than those of corresponding UVC sub-groups ($P<0.01$ or $P<0.05$). In this experiment it was found that 1.8GHz (SAR, 3 W/kg) MW exposure for 1.5 and 4 h did not enhance significantly human lymphocyte DNA damage, but could reduce and increase DNA damage of human lymphocytes induced by UVC at 1.5-h and 4-h incubation, respectively.

Canseven AG, Esmekaya MA, Kayhan H, Tuysuz MZ, Seyhan N. Effects of microwave exposure and Gemcitabine treatment on apoptotic activity in Burkitt's lymphoma (Raji) cells. *Electromagn Biol Med*. 2014 Jun 5:1-5. [Epub ahead of print]

We investigated the effects of 1.8 MHz Global System for Mobile Communications (GSM)-modulated microwave (MW) radiation on apoptotic level and cell viability of Burkitt's lymphoma (Raji) cells with or without Gemcitabine, which exhibits cell phase specificity, primarily killing cells undergoing DNA synthesis (S-phase). Raji cells were exposed to 1.8 GHz GSM-modulated MW radiation at a specific absorption rate (SAR) of 0.350 W/kg in a CO₂ incubator. The duration of the exposure was 24 h. The amount of apoptotic cells was analyzed using Annexin V-FITC and propidium iodide (PI) staining with flow cytometer. The apoptotic activity of MW exposed Raji cells was increased significantly. In addition, cell viability of exposed samples was significantly decreased. Combined exposure of MW and Gemcitabine increased the amount of apoptotic cells than MW radiation alone. Moreover, viability of MW + Gemcitabine exposed cells was lower than that of cells exposed only to MW. These results demonstrated that MW

Studies that show **Cell Tower** Health Effects

radiation exposure and Gemcitabine treatment have a synergistic effect on apoptotic activity of Raji cells.

Effects on Glands

Aydogan F, Unlu I, Aydin E, Yumusak N, Devrim E, Samim EE, Ozgur E, Unsal V, Tomruk A, Ozturk GG, Seyhan N. The effect of 2100 MHz radiofrequency radiation of a 3G mobile phone on the parotid gland of rats. Am J Otolaryngol. 2014 Oct 5. pii: S0196-0709(14)00207-5. doi: 10.1016/j.amjoto.2014.10.001. [Epub ahead of print]

PURPOSE: We aimed to evaluate the effect of 2100 MHz radiofrequency radiation on the parotid gland of rats in short and relatively long terms. **MATERIAL AND METHODS:** Thirty Wistar albino rats were divided into four groups. Groups A and B served as the control groups (for 10 days and 40 days, respectively), and each group included six rats. Groups C and D were composed of nine rats each, and they were the exposure groups. The rats were exposed to 2100 MHz radiofrequency radiation emitted by a generator, simulating a third generation mobile phone for 6 hours/day, 5 days/week, for 10 or 40 days. Following exposure, the rats were sacrificed and parotid glands were removed. Histopathological and biochemical examinations were performed. **RESULTS:** Although there were no histopathological changes in the control groups except for two animals in group A and three animals in group B, the exposure groups C (10 days) and D (40 days) showed numerous histopathological changes regarding salivary gland damage including acinar epithelial cells, interstitial space, ductal system, vascular system, nucleus, amount of cytoplasm and variations in cell size. The histopathological changes were more prominent in group D compared to group C. There was statistically significant different parameter regarding variation in cell size between the groups B and D ($p=0.036$). **CONCLUSION:** The parotid gland of rats showed numerous histopathological changes after exposure to 2100 MHz radiofrequency radiation, both in the short and relatively long terms. Increased exposure duration led to an increase in the histopathological changes.

Effects on Animals and Environment

Panagopoulos DJ, Chavdoula ED, Margaritis LH. Bioeffects of mobile telephony radiation in relation to its intensity or distance from the antenna. Int J Radiat Biol. 86(5):345-357, 2010.

PURPOSE: To examine the bioactivity of GSM 900 and 1800 (Global System for Mobile Telecommunications) radiations, in relation to the distance from the antenna or to the radiation-field intensities. **MATERIALS AND METHODS:** *Drosophila melanogaster* adult insects were exposed to the radiation of a GSM 900/1800 mobile phone antenna at different distances ranging from 0 to 100 cm, and the effect on their reproductive capacity and cell death induction in the gonads by the use of TUNEL (Terminal

Studies that show **Cell Tower** Health Effects

deoxynucleotide transferase dUTP Nick End Labeling) assay, was studied. RESULTS: These radiations/fields decreased the reproductive capacity by cell death induction, at all the different distances tested. The effect diminished with the distance/decreasing intensities. An increased bioactivity 'window' was revealed at distances of 20-30 cm from the mobile phone antenna, (radiation intensity around 10 microW/cm(2)) where the effect became highest, in relation to smaller or longer distances. The effect diminished considerably for distances longer than 40-50 cm and became not evident for distances longer than 1 m or radiation intensities smaller than 1 microW/cm(2). CONCLUSIONS: GSM bioactivity is highest for intensities down to less than 10 microW/cm(2) and still evident until 1 microW/cm(2) exhibiting 'window' effects

Loscher W, Kas G, Extraordinary behavior disorders in cows in proximity to transmission stations. Der Praktische Tierarz 79:437-444, 1998. (Article in German) In addition to reduction of milk yield and increased health problems, behavioral abnormalities were observed over a period of two years in a herd of dairy cows maintained in close proximity to a TV and cell phone transmitting antenna. Evaluation of possible factors which could explain the abnormalities in the live stock did not disclose any factors other than the high-frequency electromagnetic fields. An experiment in which a cow with abnormal behavior was brought to a stable 20 km away from the antenna resulted in a complete normalization of the cow within five days, whereas symptoms returned when the cow was brought back to the stable nearby the antenna. In view of the previous described effects of electromagnetic fields, it might be possible that the observed abnormalities in cows are related to electromagnetic field exposure. (power densities measured 0.02-7 mW/m2).

Koldayev VM, Shchepin YV, Effects of electromagnetic radiation on embryos of sea-urchins. Bioelectrochem Bioenerg 43:161-164, 1997. Electromagnetic radiation (EMR) causes a decrease in the number of fertilized eggs and an increase in the number of zygotes with abnormal fertilization envelopes in sea-urchins. The microstructural impairments of the cellular surface, the increase of lipid peroxidation and the changes of amino acid metabolism show that the impairments of the development of embryos exposed to EMR are caused by the damages of the membrane structures.

Cammaerts MC, De Doncker P, Patris X, Bellens F, Rachidi Z, Cammaerts D. GSM 900 MHz radiation inhibits ants' association between food sites and encountered cues. Electromagn Biol Med. 31(2):151-165, 2012.

The kinetics of the acquisition and loss of the use of olfactory and visual cues were previously obtained in six experimental colonies of the ant *Myrmica sabuleti* meinert 1861, under normal conditions. In the present work, the same experiments were conducted on six other naive identical colonies of *M. sabuleti*, under electromagnetic radiation similar to those surrounding GSM and communication masts. In this situation, no association between food and either olfactory or visual cues occurred. After a recovery period, the ants were able to make such an association but never reached the

Studies that show **Cell Tower** Health Effects

expected score. Such ants having acquired a weaker olfactory or visual score and still undergoing olfactory or visual training were again submitted to electromagnetic waves. Not only did they lose all that they had memorized, but also they lost it in a few hours instead of in a few days (as under normal conditions when no longer trained). They kept no visual memory at all (instead of keeping 10% of it as they normally do). The impact of GSM 900 MHz radiation was greater on the visual memory than on the olfactory one. These communication waves may have such a disastrous impact on a wide range of insects using olfactory and/or visual memory, i.e., on bees.

Senavirathna MD, Asaeda T, Thilakarathne BL, Kadono H. Nanometer-scale elongation rate fluctuations in the *Myriophyllum aquaticum* (Parrot feather) stem were altered by radio-frequency electromagnetic radiation. *Plant Signal Behav.* 2014 Mar 26;9(3). pii: e28590. [Epub ahead of print]

The emission of radio-frequency electromagnetic radiation (EMR) by various wireless communication base stations has increased in recent years. While there is wide concern about the effects of EMR on humans and animals, the influence of EMR on plants is not well understood. In this study, we investigated the effect of EMR on the growth dynamics of *Myriophyllum aquaticum* (Parrot feather) by measuring the nanometric elongation rate fluctuation (NERF) using a statistical interferometry technique. Plants were exposed to 2 GHz EMR at a maximum of 1.42 Wm⁻² for 1 h. After continuous exposure to EMR, *M. aquaticum* plants exhibited a statistically significant 51 ± 16% reduction in NERF standard deviation. Temperature observations revealed that EMR exposure did not cause dielectric heating of the plants. Therefore, the reduced NERF was due to a non-thermal effect caused by EMR exposure. The alteration in NERF continued for at least 2.5 h after EMR exposure and no significant recovery was found in post-EMR NERF during the experimental period.

Balmori A. Mobile Phone Mast Effects on Common Frog (*Rana temporaria*) Tadpoles: The City Turned into a Laboratory. *Electromagn Biol Med.* 29(1-2):31-35, 2010.

An experiment has been made exposing eggs and tadpoles of the common frog (*Rana temporaria*) to electromagnetic radiation from several mobile (cell) phone antennae located at a distance of 140 meters. The experiment lasted two months, from the egg phase until an advanced phase of tadpole prior to metamorphosis. Measurements of electric field intensity (radiofrequencies and microwaves) in V/m obtained with three different devices were 1.8 to 3.5 V/m. In the exposed group (n = 70), low coordination of movements, an asynchronous growth, resulting in both big and small tadpoles, and a high mortality (90%) was observed. Regarding the control group (n = 70) under the same conditions but inside a Faraday cage, the coordination of movements was normal, the development was synchronous, and a mortality of 4.2% was obtained. These results indicate that radiation emitted by phone masts in a real situation may affect the development and may cause an increase in mortality of exposed tadpoles. This research may have huge implications for the natural world, which is now exposed to high microwave radiation levels from a multitude of phone masts.

Studies that show **Cell Tower** Health Effects

Balode, Z, Assessment of radio-frequency electromagnetic radiation by the micronucleus test in bovine peripheral erythrocytes. Sci Total Environ 180(1):81-85, 1996.

Previous bioindicative studies in the Skrunda Radio Location Station area have focused on the somatic influence of electromagnetic radiation on plants, but it is also important to study genetic effects. We have chosen cows as test animals for cytogenetical evaluation because they live in the same general exposure area as humans, are confined to specific locations and are chronically exposed to radiation. Blood samples were obtained from female Latvian Brown cows from a farm close to and in front of the Skrunda Radar and from cows in a control area. A simplified alternative to the Schiff method of DNA staining for identification of micronuclei in peripheral erythrocytes was applied. Microscopically, micronuclei in peripheral blood erythrocytes were round in shape and exhibited a strong red colour. They are easily detectable as the only coloured bodies in the uncoloured erythrocytes. From each individual animal 2000 erythrocytes were examined at a magnification of x 1000 for the presence of micronuclei. The counting of micronuclei in peripheral erythrocytes gave low average incidences, 0.6 per 1000 in the exposed group and 0.1 per 1000 in the control, but statistically significant ($P < 0.01$) differences were found in the frequency distribution between the control and exposed groups.

Effects on Skin

Cam ST, Seyhan N, Kavaklı C, Celikbıçak O. Effects of 900 MHz Radiofrequency Radiation on Skin Hydroxyproline Contents. Cell Biochem Biophys. 2014 Apr 24. [Epub ahead of print]

The present study aimed to investigate the possible effect of pulse-modulated radiofrequency radiation (RFR) on rat skin hydroxyproline content, since skin is the first target of external electromagnetic fields. Skin hydroxyproline content was measured using liquid chromatography mass spectrometer method. Two months old male wistar rats were exposed to a 900 MHz pulse-modulated RFR at an average whole body specific absorption rate (SAR) of 1.35 W/kg for 20 min/day for 3 weeks. The radiofrequency (RF) signals were pulse modulated by rectangular pulses with a repetition frequency of 217 Hz and a duty cycle of 1:8 (pulse width 0.576 ms). A skin biopsy was taken at the upper part of the abdominal costa after the exposure. The data indicated that whole body exposure to a pulse-modulated RF radiation that is similar to that emitted by the global system for mobile communications (GSM) mobile phones caused a statistically significant increase in the skin hydroxyproline level ($p = 0.049$, Mann-Whitney U test). Under our experimental conditions, at a SAR less than the International Commission on Non-Ionizing Radiation Protection safety limit recommendation, there was evidence that GSM signals could alter hydroxyproline concentration in the rat skin.

Studies that show **Cell Tower** Health Effects

Effects on Protein

Hässig M, Wullschleger M, Naegeli HP, Kupper J, Spiess B, Kuster N, Capstick M, Murbach M. Influence of non ionizing radiation of base stations on the activity of redox proteins in bovines. *BMC Vet Res.* 2014 Jun 19;10(1):136. [Epub ahead of print]

BACKGROUND: The influence of electromagnetic fields on the health of humans and animals is still an intensively discussed and scientifically investigated issue (Prakt Tierarzt 11:15-20, 2003; Umwelt Medizin Gesellschaft 17:326-332, 2004; J Toxicol Environment Health, Part B 12:572-597, 2009). We are surrounded by numerous electromagnetic fields of variable strength, coming from electronic equipment and its power cords, from high-voltage power lines and from antennas for radio, television and mobile communication. Particularly the latter cause's controversy, as everyone likes to have good mobile reception at anytime and anywhere, whereas nobody wants to have such a base station antenna in their proximity. **RESULTS:** In this experiment, the **non-ionizing radiation (NIR)** has resulted in changes in the enzyme activities. Certain enzymes were disabled, others enabled by NIR. Furthermore, individual behavior patterns were observed. While certain cows reacted to NIR, others did not react at all, or even inversely. **CONCLUSION:** The present results coincide with the information from the literature, according to which NIR leads to changes in redox proteins, and that there are individuals who are sensitive to radiation and others that are not. However, the latter could not be distinctly attributed - there are cows that react clearly with one enzyme while they do not react with another enzyme at all, or even the inverse. The study approach of testing ten cows each ten times during three phases has proven to be appropriate. Future studies should however set the post-exposure phase later on.

Effects on Immune Function

Li CY, Liao MH, Lin CW, Tsai WS, Huang CC, Tang TK. Inhibitory Effects of Microwave Radiation on LPS-Induced NFκB Expression in THP-1 Monocytes. *Chin J Physiol.* 55(6):421-427, 2012.

Microwave radiations can be encountered regularly in daily lives. When WHO announced that microwave radiations were a kind of environmental energy which interfere with the physiological functions of the human body, great concerns have been raised over the damages microwave frequencies can do to human physiology. The immunological performance and the activities of the cellular inflammatory factor NFκB have been closely related in monocyte. Due to the effect of phorbol 12-myristate 13-acetate (PMA) on THP-1 monocytes, THP-1 monocytes would differentiate into macrophages and would then react with lipopolysaccharides (LPS), and the amount of NFκB increased in the THP-1 monocytes. Expression of cytokine is affected when cells are exposed to a frequency of 2450 MHz and at 900 W. Thus, in our experiments, an observation was made when THP-1 monocytes were stimulated with PMA and LPS to differentiate into macrophage, the amount of NFκB in cells increased exponentially, and

Studies that show **Cell Tower** Health Effects

the levels of NFκB expression were decreased by the exposure of microwave radiation. In conclusion, microwave radiations were found to inhibit the activity functions of THP-1 monocytes stimulated with PMA and LPS.

Electro Hypersensitivity

Nordin S, Neely G, Olsson D, Sandström M. Odor and Noise Intolerance in Persons with Self-Reported Electromagnetic Hypersensitivity. Int J Environ Res Public Health. 11(9):8794-8805, 2014.

Lack of confirmation of symptoms attributed to electromagnetic fields (EMF) and triggered by EMF exposure has highlighted the role of individual factors. Prior observations indicate intolerance to other types of environmental exposures among persons with electromagnetic hypersensitivity (EHS). This study assessed differences in odor and noise intolerance between persons with EHS and healthy controls by use of subscales and global measures of the Chemical Sensitivity Scale (CSS) and the Noise Sensitivity Scale (NSS). The EHS group scored significantly higher than the controls on all CSS and NSS scales. Correlation coefficients between CSS and NSS scores ranged from 0.60 to 0.65 across measures. The findings suggest an association between EHS and odor and noise intolerance, encouraging further investigation of individual factors for understanding EMF-related symptoms.

RF Levels in Cities

Urbiniello D, Huss A, Beekhuizen J, Vermeulen R, Rösli M. Use of portable exposure meters for comparing mobile phone base station radiation in different types of areas in the cities of Basel and Amsterdam. Sci Total Environ. 468-469:1028-1033, 2014.

BACKGROUND: Radiofrequency electromagnetic fields (RF-EMF) are highly variable and differ considerably within as well as between areas. Exposure assessment studies characterizing spatial and temporal variation are limited so far. Our objective was to evaluate sources of data variability and the repeatability of daily measurements using portable exposure meters (PEMs). METHODS: Data were collected at 12 days between November 2010 and January 2011 with PEMs in four different types of urban areas in the cities of Basel (BSL) and Amsterdam (AMS). RESULTS: Exposure from mobile phone base stations ranged from 0.30 to 0.53 V/m in downtown and business areas and in residential areas from 0.09 to 0.41 V/m. Analysis of variance (ANOVA) demonstrated that measurements from various days were highly reproducible (measurement duration of approximately 30 min) with only 0.6% of the variance of all measurements from mobile phone base station radiation being explained by the measurement day and only 0.2% by the measurement time (morning, noon, afternoon), whereas type of area (30%) and city (50%) explained most of the data variability. CONCLUSIONS: We conclude that mobile monitoring of exposure from mobile phone base station radiation with PEMs is

Studies that show **Cell Tower** Health Effects

useful due to the high repeatability of mobile phone base station exposure levels, despite the high spatial variation.

Urbiniello D, Joseph W, Verloock L, Martens L, Rösli M. Temporal trends of radio-frequency electromagnetic field (RF-EMF) exposure in everyday environments across European cities. Environ Res. 2014 Aug 12;134C:134-142. doi: 10.1016/j.envres.2014.07.003. [Epub ahead of print]

BACKGROUND: The rapid development and increased use of wireless telecommunication technologies led to a substantial change of radio-frequency electromagnetic field (RF-EMF) exposure in the general population but little is known about temporal trends of RF-EMF in our everyday environment. **OBJECTIVES:** The objective of our study is to evaluate temporal trends of RF-EMF exposure levels in different microenvironments of three European cities using a common measurement protocol. **METHODS:** We performed measurements in the cities of Basel (Switzerland), Ghent and Brussels (Belgium) during one year, between April 2011 and March 2012. RF-EMF exposure in 11 different frequency bands ranging from FM (Frequency Modulation, 88MHz) to WLAN (Wireless Local Area Network, 2.5GHz) was quantified with portable measurement devices (exposimeters) in various microenvironments: outdoor areas (residential areas, downtown and suburb), public transports (train, bus and tram or metro rides) and indoor places (airport, railway station and shopping centers). Measurements were collected every 4s during 10-50min per environment and measurement day. Linear temporal trends were analyzed by mixed linear regression models. **RESULTS:** Highest total RF-EMF exposure levels occurred in public transports (all public transports combined) with arithmetic mean values of 0.84V/m in Brussels, 0.72V/m in Ghent, and 0.59V/m in Basel. In all outdoor areas combined, mean exposure levels were 0.41V/m in Brussels, 0.31V/m in Ghent and 0.26V/m in Basel. Within one year, total RF-EMF exposure levels in all outdoor areas in combination increased by 57.1% (p<0.001) in Basel by 20.1% in Ghent (p=0.053) and by 38.2% (p=0.012) in Brussels. Exposure increase was most consistently observed in outdoor areas due to emissions from mobile phone base stations. In public transports RF-EMF levels tended also to increase but mostly without statistical significance. **DISCUSSION:** An increase of RF-EMF exposure levels has been observed between April 2011 and March 2012 in various microenvironments of three European cities. Nevertheless, exposure levels were still far below regulatory limits of each country. A continuous monitoring is needed to identify high exposure areas and to anticipate critical development of RF-EMF exposure at public places.

Estenberg J, Augustsson T. Extensive frequency selective measurements of radiofrequency fields in outdoor environments performed with a novel mobile monitoring system. Bioelectromagnetics. 2013 Dec 27. doi: 10.1002/bem.21830. [Epub ahead of print]

A novel, car based, measuring system for estimation of general public outdoor exposure to radiofrequency fields (RF) has been developed. The system enables fast, large area, isotropic spectral measurements with a bandwidth covering the frequency range of

Studies that show **Cell Tower** Health Effects

30 MHz to 3 GHz. Measurements have shown that complete mapping of a town with 15000 inhabitants and a path length of 115 km is possible to perform within 1 day. The measured areas were chosen to represent typical rural, urban and city areas of Sweden. The data sets consist of more than 70000 measurements. All measurements were performed during the daytime. The median power density was $16 \mu\text{W}/\text{m}^2$ in rural areas, $270 \mu\text{W}/\text{m}^2$ in urban areas, and $2400 \mu\text{W}/\text{m}^2$ in city areas. In urban and city areas, base stations for mobile phones were clearly the dominating sources of exposure.

Effects on Water

Hinrikus H, Lass J, Karai D, Pilt K, Bachmann M. Microwave effect on diffusion: a possible mechanism for non-thermal effect. Electromagn Biol Med. 23:1-7, 2014.

In this study, we assume that microwave radiation affects hydrogen bonding between dipolar water molecules and through that diffusion in water at constant temperature. The experimental study was performed on the setup of two identical reservoirs filled with pure water and 0.9% NaCl solution and connected by a thin tube. Alterations of NaCl concentration in the reservoir initially filled with pure water were measured using the resistance of the solution as an indicator. The applied 450 MHz continuous-wave microwave field had the maximal specific absorption rate of 0.4 W/kg on the connecting tube. The standard deviation of water temperature in the setup was 0.02 °C during an experiment. Our experimental data demonstrated that microwave exposure makes faster the process of diffusion in water. The time required for reduction of initial resistance of the solution by 10% was 1.7 times shorter with microwave. This result is consistent with the proposed mechanism of low-level microwave effect: microwave radiation, rotating dipolar water molecules, causes high-frequency alterations of hydrogen bonds between water molecules, thereby affects its viscosity and makes faster diffusion.

RF Levels From Cell Towers

Martinez-Burdalo M, Martin A, Anguiano M, Villar R. On the safety assessment of human exposure in the proximity of cellular communications base-station antennas at 900, 1800 and 2170 MHz. Phys Med Biol. 50(17):4125-4137, 2005.

In this work, the procedures for safety assessment in the close proximity of cellular communications base-station antennas at three different frequencies (900, 1800 and 2170 MHz) are analysed. For each operating frequency, we have obtained and compared the distances to the antenna from the exposure places where electromagnetic fields are below reference levels and the distances where the specific absorption rate (SAR) values in an exposed person are below the basic restrictions, according to the European safety guidelines. A high-resolution human body model has been located, in front of each base-station antenna as a worst case, at different distances, to compute whole body averaged SAR and maximum 10 g averaged SAR

Studies that show **Cell Tower** Health Effects

inside the exposed body. The finite-difference time-domain method has been used for both electromagnetic fields and SAR calculations. This paper shows that, for antenna-body distances in the near zone of the antenna, the fact that averaged field values be below the reference levels could, at certain frequencies, not guarantee guidelines compliance based on basic restrictions.

Hu J, Lu Y, Zhang H, Xie H, Yang X. [Level of microwave radiation from mobile phone base stations built in residential districts] Wei Sheng Yan Jiu. 38(6):712-716, 2009.
[Article in Chinese]

OBJECTIVE: To investigate the condition of microwave radiation pollution from mobile phone base station built in populated area. **METHODS:** Random selected 18 residential districts where had base station and 10 residential districts where had no base stations. A TES-92 electromagnetic radiation monitor were used to measure the intensity of microwave radiation in external and internal living environment. **RESULTS:** The intensities of microwave radiation in the exposure residential districts were more higher than those of the control residential districts ($p < 0.05$). There was a intensity peak at about 10 m from the station, it would gradually weaken with the increase of the distance. The level of microwave radiation in antenna main lobe region is not certainly more higher than the side lobe direction, and the side lobe direction also is not more lower. At the same district, where there were two base stations, the electromagnetic field nestification would take place in someplace. The intensities of microwave radiation outside the exposure windows in the resident room not only changed with distance but also with the height of the floor. The intensities of microwave radiation inside the aluminum alloys security net were more lower than those of outside the aluminum alloys security net ($p < 0.05$), but the inside or outside of glass-window appears almost no change ($p > 0.05$). **CONCLUSIONS:** Although all the measure dates on the ground around the base station could be below the primary standard in "environment electromagnetic wave hygienic standard" (GB9175-88), there were still a minorities of windows which exposed to the base station were higher, and the outside or inside of a few window was even higher beyond the primary safe level defined standard. The aluminum alloys security net can partly shield the microwave radiation from the mobile phone base station.

Danger Perception and Symptoms

Hutter HP, Moshammer H, Wallner P, Kundi M. Public perception of risk concerning cell towers and mobile phones. Soz Praventivmed. 49(1):62-66, 2004.

OBJECTIVE: The controversy about health risks of electromagnetic fields (EMF) has contributed in raising fears concerning emissions from celltowers. The study was to examine whether or not neighbours of celltowers are particularly concerned about adverse health effects of mobile phones and their base stations. **METHODS:** Prior to information delivered by medical doctors of the Institute of Environmental Health at public hearings a questionnaire was handed out to participants asking for their personal

Studies that show **Cell Tower** Health Effects

rating of several environmental health risks including those of mobile telecommunication (n = 123, response rate approx. 48%). Medical students (n = 366) served as a contrast group. RESULTS: Participants rated health risk for both, mobile phones and celltowers higher as students. A trend for higher ratings was also seen with older subjects and female sex. The risk ratings of both exposures correlated well with each other. The magnitude of the perceived risks, however, resembled that of other ubiquitous exposures like traffic noise and air pollution. CONCLUSION: Contrary to the claims of the telecommunication industry, opponents of celltowers generally do not express unusual fears concerning electromagnetic field exposure. The outcome of our study indicates that the risk rating is comparable with other perceived common hazards of the civilised world. It is hypothesised that offering information and participation to the concerned population will be efficient in reducing exaggerated fears.

Reason Why Not A Lot Of Studies on Cell Towers

Kundi M, Hutter HP. Mobile phone base stations-Effects on wellbeing and health. Pathophysiology. 16(2-3):123-135, 2009.

Studying effects of mobile phone base station signals on health have been discouraged by authoritative bodies like WHO International EMF Project and COST 281. WHO recommended studies around base stations in 2003 but again stated in 2006 that studies on cancer in relation to base station exposure are of low priority. As a result only few investigations of effects of base station exposure on health and wellbeing exist. Cross-sectional investigations of subjective health as a function of distance or measured field strength, despite differences in methods and robustness of study design, found indications for an effect of exposure that is likely independent of concerns and attributions. Experimental studies applying short-term exposure to base station signals gave various results, but there is weak evidence that UMTS and to a lesser degree GSM signals reduce wellbeing in persons that report to be sensitive to such exposures. Two ecological studies of cancer in the vicinity of base stations report both a strong increase of incidence within a radius of 350 and 400m respectively. Due to the limitations inherent in this design no firm conclusions can be drawn, but the results underline the urgent need for a comprehensive investigation of this issue. Animal and in vitro studies are inconclusive to date. An increased incidence of DMBA induced mammary tumors in rats at a SAR of 1.4W/kg in one experiment could not be replicated in a second trial. Indications of oxidative stress after low-level in vivo exposure of rats could not be supported by in vitro studies of human fibroblasts and glioblastoma cells. From available evidence it is impossible to delineate a threshold below which no effect occurs, however, given the fact that studies reporting low exposure were invariably negative it is suggested that power densities around 0.5-1mW/m² must be exceeded in order to observe an effect. The meager data base must be extended in the coming years. The difficulties of investigating long-term effects of base station exposure have been exaggerated, considering that base station and handset exposure have almost nothing

Studies that show **Cell Tower** Health Effects

in common both needs to be studied independently. It cannot be accepted that studying base stations is postponed until there is firm evidence for mobile phones.

Miscellaneous Effects

Panagopoulos, D. J., Johansson O. & Carlo G.L. Polarization: A Key Difference between Man-made and Natural Electromagnetic Fields, in regard to Biological Activity. *Sci. Rep.* 5, 14914; doi: 10.1038/srep14914 (2015). Published online Oct 12, 2015.

In the present study we analyze the role of polarization in the biological activity of Electromagnetic Fields (EMFs)/Electromagnetic Radiation (EMR). All types of man-made EMFs/EMR - in contrast to natural EMFs/EMR - are polarized. Polarized EMFs/EMR can have increased biological activity, due to: 1) Ability to produce constructive interference effects and amplify their intensities at many locations. 2) Ability to force all charged/polar molecules and especially free ions within and around all living cells to oscillate on parallel planes and in phase with the applied polarized field. Such ionic forced-oscillations exert additive electrostatic forces on the sensors of cell membrane electro-sensitive ion channels, resulting in their irregular gating and consequent disruption of the cell's electrochemical balance. These features render man-made EMFs/EMR more bioactive than natural non-ionizing EMFs/EMR. This explains the increasing number of biological effects discovered during the past few decades to be induced by man-made EMFs, in contrast to natural EMFs in the terrestrial environment which have always been present throughout evolution, although human exposure to the latter ones is normally of significantly higher intensities/energy and longer durations. Thus, polarization seems to be a trigger that significantly increases the probability for the initiation of biological/health effects.

Dhami AK. Study of electromagnetic radiation pollution in an Indian city. *Environ Monit Assess.*184(11):6507-6512, 2012.

Abstract. Electromagnetic radiation emitted by cell phone towers is a form of environmental pollution and is a new health hazard, especially to children and patients. The present studies were taken to estimate the microwave/RF pollution by measuring radiation power densities near schools and hospitals of Chandigarh city in India. The cell phone radiations were measured using a handheld portable power density meter TES 593 and specific absorption rates were estimated from the measured values. These values of electromagnetic radiation in the environment were compared with the levels at which biological system of humans and animals starts getting affected. The values were also compared with the international exposure limits set by the International Commission on Non-Ionizing Radiation Protection (ICNIRP). The highest measured power density was 11.48 mW/m(2) which is 1,148% of the biological limit. The results indicated that the exposure levels in the city were below the ICNIRP limit, but much above the biological limit.

Report of Partial Findings from the National Toxicology Program Carcinogenesis Studies of Cell Phone Radiofrequency Radiation in Hsd: Sprague Dawley® SD rats (Whole Body Exposures)

Draft 5-19-2016

Table of Contents

Abstract.....	2
Summary.....	4
Study Rationale	5
Description of the NTP Cell Phone RFR Program	6
Study Design	7
Results	8
Brain.....	9
Heart.....	10
Discussion.....	13
Conclusions	15
Next Steps	15
Appendix A – Contributors.....	17
Appendix B – Statistical Analysis.....	18
Appendix C - Pathology.....	21
Appendix D – Historical Controls	26
Appendix E – Time on Study to Appearance of Tumors	27
Appendix F – Reviewer’s Comments	29
Appendix G – NIH Reviewer’s Comments	32
Appendix G1: Reviewer’s Comments	33
Appendix G2: NTP’s Responses to NIH Reviewer’s Comments	65

Abstract

The US National Toxicology Program (NTP) has carried out extensive rodent toxicology and carcinogenesis studies of radiofrequency radiation (RFR) at frequencies and modulations used in the US telecommunications industry. This report presents partial findings from these studies. The occurrences of two tumor types in male Harlan Sprague Dawley rats exposed to RFR, malignant gliomas in the brain and schwannomas of the heart, were considered of particular interest, and are the subject of this report. The findings in this report were reviewed by expert peer reviewers selected by the NTP and National Institutes of Health (NIH). These reviews and responses to comments are included as appendices to this report, and revisions to the current document have incorporated and addressed these comments. Supplemental information in the form of 4 additional manuscripts has or will soon be submitted for publication. These manuscripts describe in detail the designs and performance of the RFR exposure system, the dosimetry of RFR exposures in rats and mice, the results to a series of pilot studies establishing the ability of the animals to thermoregulate during RFR exposures, and studies of DNA damage.

Capstick M, Kuster N, Kühn S, Berdinas-Torres V, Wilson P, Ladbury J, Koepke G, McCormick D, Gauger J, Melnick R. A radio frequency radiation reverberation chamber exposure system for rodents

Yijian G, Capstick M, McCormick D, Gauger J, Horn T, Wilson P, Melnick RL and Kuster N. Life time dosimetric assessment for mice and rats exposed to cell phone radiation

- 1 Wyde ME, Horn TL, Capstick M, Ladbury J, Koepke G, Wilson P, Stout MD, Kuster N,
- 2 Melnick R, Bucher JR, and McCormick D. Pilot studies of the National Toxicology Program's
- 3 cell phone radiofrequency radiation reverberation chamber exposure system
- 4
- 5 Smith-Roe SL, Wyde ME, Stout MD, Winters J, Hobbs CA, Shepard KG, Green A, Kissling
- 6 GE, Tice RR, Bucher JR, Witt KL. Evaluation of the genotoxicity of cell phone radiofrequency
- 7 radiation in male and female rats and mice following subchronic exposure

Report of Partial Findings from the National Toxicology Program Carcinogenesis Studies of Cell Phone Radiofrequency Radiation in Hsd: Sprague Dawley® SD rats (Whole Body Exposures)

Draft 5-19-2016

SUMMARY

The purpose of this communication is to report partial findings from a series of radiofrequency radiation (RFR) cancer studies in rats performed under the auspices of the U.S. National Toxicology Program (NTP).¹ This report contains peer-reviewed, neoplastic and hyperplastic findings only in the brain and heart of Hsd:Sprague Dawley® SD® (HSD) rats exposed to RFR starting *in utero* and continuing throughout their lifetimes. These studies found low incidences of malignant gliomas in the brain and schwannomas in the heart of male rats exposed to RFR of the two types [Code Division Multiple Access (CDMA) and Global System for Mobile Communications (GSM)] currently used in U.S. wireless networks. Potentially preneoplastic lesions were also observed in the brain and heart of male rats exposed to RFR.

The review of partial study data in this report has been prompted by several factors. Given the widespread global usage of mobile communications among users of all ages, even a very small increase in the incidence of disease resulting from exposure to RFR could have broad implications for public health. There is a high level of public and media interest regarding the safety of cell phone RFR and the specific results of these NTP studies.

¹ NTP is a federal, interagency program, headquartered at the National Institute of Environmental Health Sciences, part of the National Institutes of Health, whose goal is to safeguard the public by identifying substances in the environment that may affect human health. For more information about NTP and its programs, visit <http://ntp.niehs.nih.gov>

Lastly, the tumors in the brain and heart observed at low incidence in male rats exposed to GSM- and CDMA-modulated cell phone RFR in this study are of a type similar to tumors observed in some epidemiology studies of cell phone use. These findings appear to support the International Agency for Research on Cancer (IARC) conclusions regarding the possible carcinogenic potential of RFR.²

It is important to note that this document reviews only the findings from the brain and heart and is not a complete report of all findings from the NTP's studies. Additional data from these studies in Hsd:Sprague Dawley[®] SD[®] (Harlan) rats and similar studies conducted in B6C3F₁/N mice are currently under evaluation and will be reported together with the current findings in two forthcoming NTP Technical Reports.

STUDY RATIONALE

Cell phones and other commonly used wireless communication devices transmit information via non-ionizing radiofrequency radiation (RFR). In 2013, IARC classified RFR as a *possible human carcinogen* based on “limited evidence” of an association between exposure to RFR from heavy wireless phone use and glioma and acoustic neuroma (vestibular schwannoma) in human epidemiology studies, and “limited evidence” for the carcinogenicity of RFR in experimental animals. While ionizing radiation is a well-accepted human carcinogen, theoretical arguments have been raised against the possibility that non-ionizing radiation could induce tumors (discussed in IARC, 2013). Given the extremely large number of people who use wireless

² IARC (International Agency for Research on Cancer). 2013. Non-Ionizing Radiation, Part 2: Radiofrequency Electromagnetic Fields. IARC Monogr Eval Carcinog Risk Hum 102. Available: <http://monographs.iarc.fr/ENG/Monographs/vol102/mono102.pdf> [accessed 26 May 2016].

communication devices, even a very small increase in the incidence of disease resulting from exposure to the RFR generated by those devices could have broad implications for public health.

DESCRIPTION OF THE NTP CELL PHONE RFR PROGRAM

RFR emitted by wireless communication devices, especially cell phones, was nominated to the NTP for toxicology and carcinogenicity testing by the U.S. Food and Drug Administration (FDA). After careful and extensive evaluation of the published literature and experimental efforts already underway at that time, the NTP concluded that additional studies were warranted to more clearly define any potential health hazard to the U.S. population. Due to the technical complexity of such studies, NTP staff worked closely with RFR experts from the National Institute of Standards and Technology (NIST). With support from NTP, engineers at NIST evaluated various types of RFR exposure systems and demonstrated the feasibility of using a specially designed exposure system (reverberation chambers), which resolved the inherent limitations identified in existing systems.

In general, NTP chronic toxicity/carcinogenicity studies expose laboratory rodents to a test article for up to 2 years and are designed to determine the potential for the agent tested to be hazardous and/or carcinogenic to humans.³ For cell phone RFR, a program of study was designed to evaluate potential, long-term health effects of whole-body exposures. These studies were conducted in three phases: (1) a series of pilot studies to establish field strengths that do not raise body temperature, (2) 28-day toxicology studies in rodents exposed to various low-level field strengths, and (3) chronic toxicology and carcinogenicity studies. The studies were carried out under contract at IIT Research Institute (IITRI) in Chicago, IL following Good Laboratory

³ Specifications for the Conduct of NTP Studies, http://ntp.niehs.nih.gov/ntp/test_info/finalntp_toxcarspecsjan2011.pdf

Practices (GLP). These studies were conducted in rats and mice using a reverberation chamber exposure system with two signal modulations [Code Division Multiple Access (CDMA) and Global System for Mobile Communications (GSM)] at two frequencies (900 MHz for rats and 1900 MHz for mice), the modulations and frequency bands that are primarily used in the United States.

STUDY DESIGN

Hsd:Sprague Dawley[®] SD[®] (Harlan) rats were housed in custom-designed reverberation chambers and exposed to cell phone RFR. Experimentally generated 900 MHz RF fields with either GSM or CDMA modulation were continuously monitored in real-time during all exposure periods via RF sensors located in each exposure chamber that recorded RF field strength (V/m). Animal exposure levels are reported as whole-body specific absorption rate (SAR), a biological measure of exposure based on the deposition of RF energy into an absorbing organism or tissue. SAR is defined as the energy (watts) absorbed per mass of tissue (kilograms). Rats were exposed to GSM- or CDMA-modulated RFR at 900 MHz with whole-body SAR exposures of 0, 1.5, 3, or 6 W/kg. RFR field strengths were frequently adjusted based on changes in body weight to maintain desired SAR levels.

Exposures to RFR were initiated *in utero* beginning with the exposure of pregnant dams (approximately 11-14 weeks of age) on Gestation Day (GD) 5 and continuing throughout gestation. After birth, dams and pups were exposed in the same cage through weaning on postnatal day (PND) 21, at which point the dams were removed and exposure of 90 pups per sex per group was continued for up to 106 weeks. Pups remained group-housed from PND 21 until they were individually housed on PND 35. Control and treatment groups were populated with no

more than 3 pups per sex per litter. All RF exposures were conducted over a period of approximately 18 hours using a continuous cycle of 10 minutes on (exposed) and 10 minutes off (not exposed), for a total daily exposure time of approximately 9 hours a day, 7 days/week. A single, common group of unexposed animals of each sex served as controls for both RFR modulations. These control rats were housed in identical reverberation chambers with no RF signal generation. Each chamber was maintained on a 12-hour light/dark cycle, within a temperature range of $72 \pm 3^{\circ}\text{F}$, a humidity range of $50 \pm 15\%$, and with at least 10 air changes per hour. Throughout the studies, all animals were provided *ad libitum* access to feed and water.

RESULTS

In pregnant rats exposed to 900 MHz GSM- or CDMA-modulated RFR, no exposure-related effects were observed on the percent of dams littering, litter size, or sex distribution of pups. Small, exposure-level-dependent reductions (up to 7%) in body weights compared to controls were observed throughout gestation and lactation in dams exposed to GSM- or CDMA-modulated RFR. In the offspring, litter weights tended to be lower (up to 9%) in GSM and CDMA RFR-exposed groups compared to controls. Early in the lactation phase, body weights of male and female pups were lower in the GSM-modulated (8%) and CDMA-modulated (15%) RFR groups at 6 W/kg compared to controls. These weight differences in the offspring for both GSM and CDMA exposures tended to lessen (6% and 10%, respectively) as lactation progressed. Throughout the remainder of the chronic study, no RFR exposure-related effects on body weights were observed in male and female rats exposed to RFR, regardless of modulation.

At the end of the 2-year study, survival was lower in the control group of males than in all groups of male rats exposed to GSM-modulated RFR. Survival was also slightly lower in control females than in females exposed to 1.5 or 6 W/kg GSM-modulated RFR. In rats exposed to CDMA-modulated RFR, survival was higher in all groups of exposed males and in the 6 W/kg females compared to controls.

Brain

A low incidence of malignant gliomas and glial cell hyperplasia was observed in all groups of male rats exposed to GSM-modulated RFR (Table 1). In males exposed to CDMA-modulated RFR, a low incidence of malignant gliomas occurred in rats exposed to 6 W/kg (Table 1). Glial cell hyperplasia was also observed in the 1.5 W/kg and 6 W/kg CDMA-modulated exposure groups. No malignant gliomas or glial cell hyperplasias were observed in controls. There was not a statistically significant difference between the incidences of lesions in exposed male rats compared to control males for any of the GSM- or CDMA-modulated RFR groups. However, there was a statistically significant positive trend in the incidence of malignant glioma ($p < 0.05$) for CDMA-modulated RFR exposures.

Table 1. Incidence of brain lesions in male Hsd:Sprague Dawley[®] SD[®] (Harlan) rats exposed to GSM- or CDMA-modulated RFR[§]

	Control	GSM			CDMA		
	0 W/kg	1.5 W/kg	3 W/kg	6 W/kg	1.5 W/kg	3 W/kg	6 W/kg
Number examined	90	90	90	90	90	90	90
Malignant glioma ^{†‡}	0*	3 (3.3%)	3 (3.3%)	2 (2.2%)	0	0	3 (3.3%)
Glial cell hyperplasia	0	2 (2.2%)	3 (3.3%)	1 (1.1%)	2 (2.2%)	0	2 (2.2%)

[§] Data presented as number of animals per group with lesions (percentage of animals per group with lesions).

* Significant SAR-dependent trend for CDMA exposures by poly-6 ($p < 0.05$). See appendix B

[†] Poly-6 survival adjusted rates for malignant gliomas were 0/53.48 in controls; GSM: 3/67.96 (4.4%), 3/72.10 (4.2%), and 2/72.65 (2.8%) in the 1.5, 3, and 6 W/kg groups, respectively; CDMA: 0/65.94, 0/73.08, and 3/57.49 (5.2%) for the 1.5, 3, and 6 W/kg groups, respectively.

[‡] Historical control incidence in NTP studies: 11/550 (2.0%), range 0-8%

In females exposed to GSM-modulated RFR, a malignant glioma was observed in a single rat exposed to 6 W/kg, and glial cell hyperplasia was observed in a single rat exposed to 3 W/kg (Table 2). In females exposed to CDMA-modulated RFR, malignant gliomas were observed in two rats exposed to 1.5 W/kg. Glial cell hyperplasia was observed in one female in each of the CDMA-modulation exposure groups (1.5, 3, and 6 W/kg). There was no glial cell hyperplasia or malignant glioma observed in any of the control females. Detailed descriptions of the malignant gliomas and glial cell hyperplasias are presented in Appendix C.

Table 2. Incidence of brain lesions in female Hsd:Sprague Dawley[®] SD[®] (Harlan) rats exposed to GSM- or CDMA-modulated RFR[§]

	Control	GSM			CDMA		
	0 W/kg	1.5 W/kg	3 W/kg	6 W/kg	1.5 W/kg	3 W/kg	6 W/kg
Number examined	90	90	90	90	90	90	90
Malignant glioma [‡]	0	0	0	1 (1.1%)	2 (2.2%)	0	0
Glial cell hyperplasia	0	0	1 (1.1%)	0	1 (1.1%)	1 (1.1%)	1 (1.1%)

[§] Data presented as number of animals per group with lesions (percentage of animals per group with lesions).

[‡] Historical control incidence in NTP studies: 1/540 (0.18%), range 0-2%

Heart

Cardiac schwannomas were observed in male rats in all exposed groups of both GSM- and CDMA-modulated RFR, while none were observed in controls (Table 3). For both modulations (GSM and CDMA), there was a significant positive trend in the incidence of schwannomas of the heart with respect to exposure SAR. Additionally, the incidence of schwannomas in the 6 W/kg males was significantly higher in CDMA-modulated RFR-exposed males compared to controls. The incidence of schwannomas in the 6 W/kg GSM-modulated RFR-exposed males was higher, but not statistically significant ($p = 0.052$) compared to controls. Schwann cell

hyperplasia of the heart was also observed in three males exposed to 6 W/kg CDMA-modulated RFR. In the GSM-modulation exposure groups, a single incidence of Schwann cell hyperplasia was observed in a 1.5 W/kg male.

Table 3. Incidence of heart lesions in male Hsd:Sprague Dawley[®] SD[®] (Harlan) rats exposed to GSM- or CDMA-modulated cell phone RFR[§]

	Control	GSM			CDMA		
	0	1.5	3	6	1.5	3	6
	W/kg	W/kg	W/kg	W/kg	W/kg	W/kg	W/kg
Number examined	90	90	90	90	90	90	90
Schwannoma ^{† ‡}	0*	2 (2.2%)	1 (1.1%)	5 (5.5%)	2 (2.2%)	3 (3.3%)	6 (6.6%)**
Schwann cell hyperplasia	0	1 (1.1%)	0	2 (2.2%)	0	0	3 (3.3%)

[§] Data presented as number of animals per group with lesions (percentage of animals per group with lesions).

* Significant SAR level-dependent trend for GSM and CDMA by poly-3 ($p < 0.05$). See appendix B

** Significantly higher than controls by poly-3 ($p < 0.05$)

[†] Poly-3 survival adjusted rates for schwannomas were 0/65.47 in controls; GSM: 2/74.87 (2.7%), 1/77.89 (1.3%), and 5/78.48 (6.4%) in the 1.5, 3, and 6 W/kg groups, respectively; CDMA: 2/74.05 (2.7%), 3/78.67 (3.8%), and 6/67.94 (8.8%) for the 1.5, 3, and 6 W/kg groups, respectively.

[‡] Historical control incidence in NTP studies: 9/699 (1.3%) range 0-6%

In females, schwannomas of the heart were also observed at 3 W/kg GSM-modulated RFR and 1.5 and 6 W/kg CDMA-modulated RFR. Schwann cell hyperplasia was observed in one female in each of the CDMA-modulation exposure groups (1.5, 3, and 6 W/kg).

Table 4. Incidence of heart lesions in female Hsd:Sprague Dawley[®] SD[®] (Harlan) rats exposed to GSM- or CDMA-modulated cell phone RFR[§]

	Control	GSM			CDMA		
	0	1.5	3	6	1.5	3	6
	W/kg	W/kg	W/kg	W/kg	W/kg	W/kg	W/kg
Number examined	90	90	90	90	90	90	90
Schwannoma [‡]	0	0	2 (2.2%)	0	2 (2.2%)	0	2 (2.2%)
Schwann cell hyperplasia	0	0	0	0	1 (1.1%)	1 (1.1%)	1 (1.1%)

[§] Data presented as number of animals per group with tumors (percentage of animals per group with tumors).

[‡] Historical control incidence in NTP studies: 4/699 (0.6%), range 0-4%

Schwann cells are present in the peripheral nervous system and are distributed throughout the whole body, not just in the heart. Therefore, organs other than the heart were examined for schwannomas and Schwann cell hyperplasia. Several occurrences of schwannomas were observed in the head, neck, and other sites throughout the body of control and GSM and CDMA RFR-exposed male rats. In contrast to the significant increase in the incidence of schwannomas in the heart of exposed males, the incidence of schwannomas observed in other tissue sites of exposed males (GSM and CDMA modulations) was not significantly different than in controls (Table 5). Additionally, Schwann cell hyperplasia was not observed in any tissues other than the heart. The combined incidence of schwannomas from all sites was generally higher in GSM- and CDMA-modulated RFR exposed males, but not significantly different than in controls. The Schwann cell response to RFR appears to be specific to the heart of male rats.

Table 5. Incidence of schwannomas in male Hsd:Sprague Dawley[®] SD[®] (Harlan) rats exposed to GSM- or CDMA-modulated RFR[§]

	Control	GSM				CDMA	
	0 W/kg	1.5 W/kg	3 W/kg	6 W/kg	1.5 W/kg	3 W/kg	6 W/kg
Number examined	90	90	90	90	90	90	90
Heart [‡]	0*	2 (2.2%)	1 (1.1%)	5 (5.5%)	2 (2.2%)	3 (3.3%)	6 (6.6%)**
Other sites [†]	3 (3.3%)	1 (1.1%)	4 (4.4%)	2 (2.2%)	2 (2.2%)	1 (1.1%)	1 (1.1%)
All sites (total)	3 (3.3%)	3 (3.3%)	5 (5.5%)	7 (7.7%)	4 (4.4%)	4 (4.4%)	7 (7.7%)

[§] Data presented as number of animals per group with tumors (percentage of animals per group with tumors).

* Significant SAR level-dependent trend for GSM and CDMA, poly 3 test ($p < 0.05$)

** Significantly higher than controls, poly-3 test ($p < 0.05$)

[‡] Historical control incidence in NTP studies: 9/699 (1.3%), range 0-6%

[†] Mediastinum, thymus, and fat

In female rats, there was no statistically significant or apparent exposure-related effect on the incidence of schwannomas in the heart or the combined incidence in the heart or other sites (Table 6).

Table 6. Incidence of schwannomas in female Hsd:Sprague Dawley[®] SD[®] (Harlan) rats exposed to GSM- or CDMA-modulated RFR[§]

Schwannoma site	Control	GSM				CDMA		
	0 W/kg	1.5 W/kg	3 W/kg	6 W/kg		1.5 W/kg	3 W/kg	6 W/kg
Number examined	90	90	90	90		90	90	90
Heart [‡]	0	0	2 (2.2%)	0		2 (2.2%)	0	2 (2.2%)
Other sites [†]	4 (4.4%)	1 (1.1%)	3 (3.3%)	1 (1.1%)		0	2 (2.2%)	2 (2.2%)
All sites (total)	4 (4.4%)	1 (1.1%)	5 (5.5%)	2 (2.2%)		2 (2.2%)	2 (2.2%)	4 (4.4%)

[§] Data presented as number of animals per group with tumors (percentage of animals per group with tumors).

[‡] Historical control incidence in NTP studies: 4/699 (0.6%), range 0-4%

[†] Ovary, uterus, vagina, thymus, abdomen, and clitoral gland

DISCUSSION

The two tumor types, which are the focus of this report, are malignant gliomas of the brain and schwannomas of the heart. Glial cells are a collection of specialized, non-neuronal, support cells whose functions include maintenance of homeostasis, formation of myelin, and providing support and protection for neurons of the peripheral nervous system (PNS) and the central nervous system (CNS). In the CNS, glial cells include astrocytes, oligodendroglia, microglial cells, and ependymal cells. Schwann cells are classified as glial cells of the PNS. In the PNS, Schwann cells produce myelin and are analogous to oligodendrocytes of the CNS. Generally, glial neoplasms in the rat are aggressive, poorly differentiated, and usually classified as malignant.

In the heart, exposure to GSM or CDMA modulations of RFR in male rats resulted in a statistically significant, positive trend in the incidence of schwannomas. There was also a statistically significant, pairwise increase at the highest CDMA exposure level tested compared to controls. Schwann cell hyperplasias also occurred at the highest exposure level of CDMA-

modulated RFR. Schwann cell hyperplasia in the heart may progress to cardiac schwannomas. No Schwann cell hyperplasias or schwannomas of the heart were observed in the single, common control group of male rats. The historical control rate of schwannomas of the heart in male Harlan Sprague Dawley rats is 1.30% (7/539) and ranges from 0-6% for individual NTP studies (Table D2, Appendix D). The 5.5-6.6% observed in the 6 W/kg GSM- and CDMA-modulated RFR groups exceeds the historical incidence, and approaches or exceeds the highest rate observed in a single study (6%). The increase in the incidence of schwannomas in the heart of male rats in this study is likely the result of whole-body exposures to GSM- or CDMA-modulated RFR.

In the brain, there was a significant, positive trend in the incidences of malignant gliomas in males exposed to CDMA-modulated RFR, and a low incidence was observed in males at all exposure levels of GSM-modulated RFR that was not statistically different than in control males. Glial cell hyperplasia, a preneoplastic lesion distinctly different from gliosis, was also observed at low incidences in rats exposed to either GSM or CDMA modulation. Glial cell hyperplasia may progress to malignant glioma. Neither of these lesions was observed in the control group of male rats. Although not observed in the current control group, malignant gliomas have been observed in control male Harlan Sprague Dawley rats from other completed NTP studies. Currently in males, the historical control rate of malignant glioma for those studies is 2.0% (11/550) and ranges from 0-8% for individual studies (Table D1, Appendix D). The 2.2-3.3% observed in all of the GSM-modulation groups and in the 6 W/kg CDMA-modulated group only slightly exceeds the mean historical control rate and falls within the observed range.

The survival of the control group of male rats in the current study (28%) was relatively low compared to other recent NTP studies in Hsd:Sprague Dawley[®] SD[®] (Harlan) rats (average 47%, range 24-72%). If malignant gliomas or schwannomas are late-developing tumors, the absence of these lesions in control males in the current study could conceivably be related to the shorter longevity of control rats in this study. Appendix E lists the time on study for each animal with a malignant glioma or heart schwannoma. Most of the gliomas were observed in animals that died late in the study, or at the terminal sacrifice. However, a relatively high number of the heart schwannomas in exposed groups were observed by 90 weeks into the study, a time when approximately 60 of the 90 control male rats remained alive and at risk for developing a tumor.

CONCLUSIONS

Under the conditions of these 2-year studies, the hyperplastic lesions and glial cell neoplasms of the heart and brain observed in male rats are considered likely the result of whole-body exposures to GSM- or CDMA-modulated RFR. There is higher confidence in the association between RFR exposure and the neoplastic lesions in the heart than in the brain. No biologically significant effects were observed in the brain or heart of female rats regardless of modulation.

NEXT STEPS

The results reported here are limited to select findings of concern in the brain and heart and do not represent a complete reporting of all findings from these studies of cell phone RFR. The complete results for all NTP studies on the toxicity and carcinogenicity of GSM and CDMA-modulated RFR are currently being reviewed and evaluated according to the established NTP process and will be reported together with the current findings in two forthcoming NTP

1 Technical Reports. Given the large scale and scope of these studies, completion of this process is
2 anticipated by fall 2017, and the draft NTP Technical Reports are expected to be available for
3 peer review and public comment by the end of 2017. We anticipate that the results from a series
4 of initial studies investigating the tolerance to various power levels of RFR, including
5 measurements of body temperatures in both sexes of young and old rats and mice and in
6 pregnant female rats, will be published in the peer-reviewed literature later in 2016.

APPENDIX A – CONTRIBUTORS

NTP CONTRIBUTORS

Participated in the evaluation and interpretation of results and the reporting of findings.

M.E. Wyde, Ph.D. (NTP study scientist)

M.F. Cesta, D.V.M., Ph.D. (NTP pathologist)

C.R. Blystone, Ph.D.

J.R. Bucher, Ph.D.

S.A. Elmore, D.V.M., M.S.

P.M. Foster, Ph.D.

M.J. Hooth, Ph.D.

G.E. Kissling, Ph.D.

D.E. Malarkey, D.V.M., Ph.D.

R.C. Sills, D.V.M., Ph.D.

M.D. Stout, Ph.D.

N.J. Walker, Ph.D.

K.L. Witt, M.S.

M.S. Wolfe, Ph.D.

APPENDIX B – STATISTICAL ANALYSIS

The Poly-k test (Bailer and Portier, 1988; Portier and Bailer, 1989; Piegorsch and Bailer, 1997) was used to assess neoplasm prevalence. This test is a survival-adjusted quantal-response procedure that modifies the Cochran-Armitage linear trend test to take survival differences into account. More specifically, this method modifies the denominator in the quantal estimate of lesion incidence to approximate more closely the total number of animal years at risk. For analysis of lesion incidence at a given site, each animal is assigned a risk weight. This value is one if the animal had a lesion at that site or if it survived until terminal sacrifice; if the animal died prior to terminal sacrifice and did not have a lesion at that site, its risk weight is the fraction of the entire study time that it survived, raised to the kth power. This method yields a lesion prevalence rate that depends only upon the choice of a shape parameter, k, for a Weibull hazard function describing cumulative lesion incidence over time (Bailer and Portier, 1988). A further advantage of the Poly-k method is that it does not require lesion lethality assumptions.

Unless otherwise specified, the NTP uses a value of $k=3$ in the analysis of site-specific lesions (Portier et al., 1986). Bailer and Portier (1988) showed that the Poly-3 test gives valid results if the true value of k is anywhere in the range from 1 to 5. In addition, Portier et al. (1986) modeled a collection of relatively common tumors observed in control animals from two-year NTP rodent carcinogenicity studies, showing that the Weibull distribution with values of k ranging between 1 and 5 was a reasonable fit to tumor incidence in most cases. In cases of early tumor onset or late tumor onset, however, $k=3$ may not be the optimal choice. Tumors with early onset would require a value of k much less than 3, while tumors with late onset would require a value of k much greater than 3. In the current studies, malignant brain gliomas occurred only in animals surviving more than 88% of the length of the study. For these brain tumors, a Weibull distribution with $k=6$ is a better fit to survival time than with $k=3$ (Portier, 1986). Malignant schwannomas of the heart occurred in animals surviving at least 65% of the length of the study; a Weibull distribution with $k=3$ adequately fits these heart tumor incidences. Therefore, poly-6 tests were used for analyses of brain tumors and poly-3 tests were used for schwannomas.

Variation introduced by the use of risk weights, which reflect differential mortality, was accommodated by adjusting the variance of the Poly-k statistic as recommended by Bieler and Williams (1993) and a continuity correction modified from Thomas et al. (1977) was applied.

Tests of significance for tumors and nonneoplastic lesions included pairwise comparisons of each dosed group with controls and a test for an overall dose-related trend. Continuity-corrected Poly-k tests were used in the analysis of lesion incidence, and reported P values are one sided.

Body weights and litter weights were compared to the control group using analysis of variance and Dunnett's test (1955). The probability of survival was estimated by the product-limit procedure of Kaplan and Meier (1958). Statistical analyses for possible exposure-related effects on survival used Cox's (1972) method for testing two groups for equality and Tarone's (1975) life table test to identify exposure-related trends. Survival analysis p-values are two-sided.

REFERENCES

- Bailer, A.J., and Portier, C.J. (1988). Effects of treatment-induced mortality and tumor-induced mortality on tests for carcinogenicity in small samples. *Biometrics* 44, 417-431.
- Bieler, G.S., and Williams, R.L. (1993). Ratio estimates, the delta method, and quantal response tests for increased carcinogenicity. *Biometrics* 49, 793-801.
- Cox, D.R. (1972). Regression models and life-tables. *Journal of the Royal Statistical Society B* 34, 187-220.
- Dunnett, C. W. (1955). A multiple comparison procedure for comparing several treatments with a control. *Journal of the American Statistical Association* 50, 1096-1121.
- Kaplan, E.L. and Meier, P. (1958). Nonparametric estimation from incomplete observations. *Journal of the American Statistical Association* 53, 457-481.
- Piegorsch, W.W., and Bailer, A.J. (1997). *Statistics for Environmental Biology and Toxicology*, Section 6.3.2. Chapman and Hall, London.
- Portier, C.J. (1986) Estimating the tumour onset distribution in animal carcinogenesis experiments. *Biometrika* 72, 371-378.

- 1 Portier, C.J., Hedges, J.C., Hoel, D.G. (1986) Age-specific models of mortality and tumor onset
2 for historical control animals in the National Toxicology Program's carcinogenicity experiments.
3 Cancer Research 46, 4372-4378.
4
- 5 Portier, C.J., and Bailer, A.J. (1989). Testing for increased carcinogenicity using a
6 survivaladjusted quantal response test. Fundam. Appl. Toxicol. 12, 731-737.
7
- 8 Thomas, D.G., Breslow, N., Gart, J.J. (1977). Trend and homogeneity analyses of proportions
9 and life table data. Computers and Biomedical Research 10, 373-381.

APPENDIX C – PATHOLOGY

Pathology data presented in this report on cell phone RFR were subjected to a rigorous peer review process. The primary goal of the NTP peer-review process is to reach consensus agreement on treatment-related findings, confirm the diagnosis of all neoplasms, and confirm any unusual lesions. At study termination, a complete necropsy and histopathology evaluation was conducted on every animal. The initial pathology examination was performed by a veterinary pathologist, who recorded all neoplastic and nonneoplastic lesions. This examination identified several potential treatment-related lesions in target organs of concern (brain and heart), which were chosen for immediate review.¹ The initial findings of glial cell tumors and hyperplasias in the brain and schwannomas, Schwann cell hyperplasia, and schwannomas from all sites were subjected to an expedited, multilevel NTP pathology peer-review process. The data were locked² prior to receipt of the finalized, study-laboratory reports to ensure that the raw data did not change during the review.

The pathology peer review consisted of a quality assessment (QA) review of all slides with tissues from the central nervous system (7 sections of brain and 3 sections of spinal cord), trigeminal nerve and ganglion, and heart. Additionally, the schwannomas of the head and neck region were reviewed. The QA review of the central nervous system and head and neck schwannomas was performed by Dr. Margarita Gruebbl of Experimental Pathology Laboratories, Inc. (EPL), and the QA review of the hearts and trigeminal nerves and ganglia was performed by Dr. Cynthia Shackelford, EPL.

The QA review pathologists then met with Dr. Mark Cesta, NTP pathologist for these studies, and Dr. David Malarkey, head of the NTP Pathology Group, to review lesions and select slides for the Pathology Working Group (PWG) reviews. All PWG reviews were conducted blinded with respect to treatment group and only identified the test articles as “test agent A” or “test

¹ Pathology peer review of remaining lesions from the cell phone RFR studies continues and is not addressed in this report.

² Locking data refers to restricting access to the computer database so the data for a particular study cannot be changed.

agent B”. Due to the large number of slides for review, the PWG was held in three separate sessions:

- January 29, 2016, for review of glial lesions in the brain and Schwann cell lesions in the heart
- February 11, 2016, for review of schwannomas of the head and neck
- February 12, 2016, for review of granular cell lesions of the brain

The reviewing PWG pathologists largely agreed on the diagnostic criteria for the lesions and on the diagnoses of schwannomas in the head and neck, and granular cell lesions in the brain.

However, there was much discussion on the criteria for differentiating glial cell hyperplasia from malignant glioma and Schwann cell hyperplasia from schwannoma. The lack of PWG agreement on definitive criteria for the glial cell and Schwann cell lesions, and the requirement for a high level of confidence in the diagnoses prompted NTP to convene two additional PWGs (organized and conducted by the NTP pathologist, Dr. Mark Cesta) with selected experts in the organ under review. These second level PWG reviews were also conducted as noted above and held in two separate sessions:

- February 25, 2016, for review of glial lesions in the brain
- March 3, 2016, for review of cardiac schwannomas, schwannomas in other organs (except the head and neck), and right ventricular degeneration

In both PWGs, the participants came to consensus on the diagnoses of the lesions and the criteria used for those diagnoses. Participants of the individual PWGs are listed below.

Table C-1. NTP Pathology Working Group (PWG) Attendees

PWG member	Affiliation
<i>January 29, 2016 - Evaluated glial lesions in the brain and Schwann cell lesions in the heart</i>	
A.E. Brix, D.V.M., Ph.D.	Experimental Pathology Laboratories, Inc. RTP, NC
M.F. Cesta, D.V.M., Ph.D.	National Institute of Environmental Health Sciences (NTP study pathologist)
S.A. Elmore, D.V.M., MS	National Institute of Environmental Health Sciences
G.P. Flake, M.D.	National Institute of Environmental Health Sciences
R.H. Garman, D.V.M.	Consultants in Veterinary Pathology, Inc. Monroeville, PA
M.M. Gruebbel, D.V.M., Ph.D.	Experimental Pathology Laboratories, Inc. RTP, NC (observer)
R.A. Herbert, D.V.M., Ph.D.	National Institute of Environmental Health Sciences
J.S. Hoane, D.V.M.	Charles River Laboratories, Inc. Durham, NC (contract study pathologist)
K.S. Janardhan, BVSc, MVSc, Ph.D.	Integrated Laboratory System
R. Kovi, BVSc, MVSc, Ph.D.	Experimental Pathology Laboratories, Inc. RTP, NC (observer)
D.E. Malarkey, D.V.M., Ph.D.	National Institute of Environmental Health Sciences
R.A. Miller, D.V.M., Ph.D.	Experimental Pathology Laboratories, Inc. RTP, NC
J.P. Morrison, D.V.M.	Charles River Laboratories, Inc. Durham, NC

PWG member	Affiliation
A.R. Pandiri, BVSc & AH, Ph.D.	National Institute of Environmental Health Sciences
C.C. Shackelford, D.V.M., Ph.D.	Experimental Pathology Laboratories, Inc. RTP, NC (observer)
J.A. Swenberg, D.V.M., Ph.D.	University of North Carolina – Chapel Hill, NC
G. Willson, BVMS, Dip RC Path, FRC Path, MRCVS	Experimental Pathology Laboratories, Inc. RTP, NC (PWG coordinator)
<i>February 11, 2016 - Evaluated schwannomas of the head and neck</i>	
A.E. Brix, D.V.M., Ph.D.	Experimental Pathology Laboratories, Inc. RTP, NC
M.F. Cesta, D.V.M., Ph.D.	National Institute of Environmental Health Sciences (NTP study pathologist)
S.A. Elmore, D.V.M., MS	National Institute of Environmental Health Sciences
G.P. Flake, M.D.	National Institute of Environmental Health Sciences
M.M. Gruebel, D.V.M., Ph.D.,	Experimental Pathology Laboratories, Inc. RTP, NC (PWG coordinator)
K.S. Janardhan, BVSc, MVSc, Ph.D.	Integrated Laboratory System RTP, NC
D.E. Malarkey, D.V.M., Ph.D.	National Institute of Environmental Health Sciences
A.R. Pandiri, BVSc & AH, Ph.D.	National Institute of Environmental Health Sciences
R.R. Maronpot, D.V.M.	Experimental Pathology Laboratories, Inc. RTP, NC
<i>February 12, 2016 - Evaluated granular cell lesions of the brain</i>	
A.E. Brix, D.V.M., Ph.D.	Experimental Pathology Laboratories, Inc. RTP, NC
M.F. Cesta, D.V.M., Ph.D.	National Institute of Environmental Health Sciences (NTP study pathologist)
S.A. Elmore, D.V.M., MS	National Institute of Environmental Health Sciences
M.M. Gruebel, D.V.M., Ph.D.,	Experimental Pathology Laboratories, Inc. RTP, NC (PWG coordinator)
J.S. Hoane, D.V.M.	Charles River Laboratories, Inc. Durham, NC (contract study pathologist)
K.S. Janardhan, BVSc, MVSc, Ph.D.	Integrated Laboratory System RTP, NC
A.R. Pandiri, BVSc. & AH, Ph.D.	National Institute of Environmental Health Sciences
R.R. Moore, D.V.M.	Integrated Laboratory System RTP, NC
<i>February 25, 2016 - Evaluated glial lesions in the brain</i>	
D. Bigner, M.D., Ph.D.	Duke University Durham, NC
B. Bolon, D.V.M., MS, Ph.D.	GEMpath, Inc. Longmont, CO
V. Chen, D.V.M., Ph.D.	National Institute of Environmental Health Sciences (observer)
M.F. Cesta, D.V.M., Ph.D.	National Institute of Environmental Health Sciences (PWG coordinator, NTP study pathologist)
S.A. Elmore, D.V.M., MS	National Institute of Environmental Health Sciences (observer)
G.P. Flake, M.D.	National Institute of Environmental Health Sciences (observer)
J.S. Hardisty, D.V.M.	Experimental Pathology Laboratories, Inc. RTP, NC
R.A. Herbert, D.V.M., Ph.D.,	National Institute of Environmental Health Sciences (observer)
R. Kovi, BVSc, MVSc, Ph.D.	Experimental Pathology Laboratories, Inc. (observer)
P.B. Little, D.V.M.	Experimental Pathology Laboratories, Inc.
D.E. Malarkey, D.V.M., Ph.D.	National Institute of Environmental Health Sciences
J.P. Morrison, D.V.M., Ph.D.	Charles River Laboratories, Inc.
A. Sharma, BVSc, MVSc, MS, Ph.D.	Covance
<i>March 3, 2016 - Evaluated heart lesions, and schwannomas in other organs (except head and neck)</i>	
B. Berridge, D.V.M., Ph.D.	GlaxoSmithKline RTP, NC
M.C. Boyle, D.V.M., Ph.D.	Amgen Thousand Oaks, CA
V. Chen, D.V.M., Ph.D.	National Institute of Environmental Health Sciences (observer)
M.F. Cesta, D.V.M., Ph.D.	National Institute of Environmental Health Sciences (PWG coordinator, NTP study pathologist)
S.A. Elmore, D.V.M., MS	National Institute of Environmental Health Sciences (observer)
M. Elwell, D.V.M., Ph.D.	Covance Chantilly, VA

PWG member	Affiliation
J.R. Hailey, D.V.M.	Covance Chantilly, VA
M. Novilla, D.V.M., MS, Ph.D.	SNBL Everett, WA

LESION DESCRIPTIONS

Brain

Malignant gliomas were infiltrative lesions, usually of modest size, with indistinct tumor margins. The neoplastic cells were typically very densely packed with more cells than neuropil. The cells were typically small and had round to oval, hyperchromatic nuclei. Mitoses were infrequent. In some of the neoplasms, invasion of the meninges, areas of necrosis surrounded by palisading neoplastic cells, cuffing of blood vessels, and neuronal satellitosis were observed. The malignant gliomas did not appear to arise from any specific anatomic subsite of the brain.

Glial cell hyperplasia consisted of small, proliferative, and poorly demarcated foci of poorly differentiated glial cells that accumulated and invaded into the surrounding parenchyma. In some cases, there was a small amount of perivascular cuffing. The hyperplastic cells appeared morphologically identical to those in the gliomas but were typically less dense with more neuropil than glial cells. There were no necrotic or degenerative elements present, so there was no evidence that the increased number of glial cells was a reaction to brain injury.

Heart

The intracardiac schwannomas were either endocardial or myocardial (intramural). The endocardial schwannomas lined the ventricles and atria and invaded into the myocardium. Two morphologic cell types were observed, but indistinct cell margins and eosinophilic cytoplasm were common to both types. Groups of cells with widely spaced small, round nuclei and moderate amounts of cytoplasm were interspersed among bands or sheets of parallel, elongated cells with thin, spindle-shaped, hyperchromatic nuclei. The myocardial schwannomas were typically less densely cellular and infiltrated amid, sometimes replacing, the cardiomyocytes. The cell types described for the endocardial neoplasms were both present, but in fewer numbers. In both subtypes of schwannomas, there was a minimal amount of cellular pleomorphism. In some larger neoplasms, Antoni type A and B patterns were present.

- 1 The Schwann cell hyperplasias were similar in appearance to the schwannomas, but were smaller
- 2 and had less pleomorphism of the cells. In the case of the endocardial Schwann cell hyperplasia,
- 3 there was no invasion of the myocardium.

APPENDIX D – HISTORICAL CONTROLS

Table D1. Incidence of astrocytoma, glioma, and/or oligodendroglioma in brains of male Harlan Sprague Dawley rats in NTP studies

Chemical	First dose	N	Control incidence
Dibutylphthalate	8/30/2010	49	4%
2-Hydroxy-4-methoxybenzophenone	11/8/2010	50	0%
p-Chloro-a,a,a-trifluorotoluene	1/17/2011	50	4%
Di-(2-ethylhexyl)phthalate	2/17/2011	50	8%
Di-(2-ethylhexyl)phthalate (perinatal)	6/27/2011	50	0%
Tris (chloroisopropyl) phosphate	12/12/2011	50	0%
Sodium tungstate	12/23/2011	50	4%
Resveratrol	5/7/2012	50	0%
Black cohosh	7/2/2012	50	2%
Radiofrequency radiation (GSM/CDMA)	9/16/2012	90	0%

Historical control rate: 11/550 (2.0%)

Table D2. Incidence of schwannoma in the heart of male Harlan Sprague Dawley rats in NTP studies

Chemical	First dose	N	Control incidence
Indole-3-carbinol	3/14/2007	50	2%
Perfluorooctanoic acid	6/19/2009	50	0%
Dietary zinc	9/3/2009	50	0%
Dibutylphthalate	8/30/2010	49	4%
2-Hydroxy-4-methoxybenzophenone	11/8/2010	50	2%
p-Chloro-a,a,a-trifluorotoluene	1/17/2011	50	0%
Di-(2-ethylhexyl)phthalate	2/17/2011	50	6%
Di-(2-ethylhexyl)phthalate (perinatal)	6/27/2011	50	4%
Tris (chloroisopropyl) phosphate	12/12/2011	50	0%
Sodium tungstate	12/23/2011	50	0%
Resveratrol	5/7/2012	50	0%
Black Cohosh	7/2/2012	50	0%
Radiofrequency radiation (GSM/CDMA)	9/16/2012	90	0%

Historical control rate: 9/699 (1.30%)

APPENDIX E – TIME ON STUDY TO APPEARANCE OF TUMORS

Malignant Glioma

SAR (W/kg)	Animal ID number	Time on study (weeks)
GSM-modulated exposed males		
1.5	717	105
	735	102
	786	104
3.0	924	101
	943	105
	1014	93
6.0	1135	104
	1137	102
CDMA-modulated exposed males		
6.0	1795	105
	1799	104
	1852	105
GSM-modulated exposed females		
6.0	1246	96
CDMA-modulated exposed females		
1.5	1463	105
	1474	105

Time to Malignant Schwannoma in Heart

SAR (W/kg)	Animal ID number	Length of survival (weeks)
GSM-modulated exposed males		
1.5	758	104
	801	105
3.0	931	105
6.0	1149	83
	1155	105
	1187	104
	1206	104
	1230	91
CDMA-modulated exposed males		
1.5	1364	105
	1352	105
3.0	1559	92
	1617	105
	1622	104
6.0	1801	76
	1821	70
	1829	104
	1833	89
	1849	104
	1860	105
GSM-modulated exposed females		
3.0	1037	105
	1077	83
CDMA-modulated exposed females		
1.5	1461	106
	1480	93
6.0	1888	105
	1965	106

APPENDIX F – REVIEWER’S COMMENTS

National Toxicology Program

Peer Review Charge and Summary Comments

Purpose: To provide independent peer review of an initial draft of this partial report. The peer reviewers were blind to the test agents under study. Introductory materials on RFR and details of the methods dealing with the field generation and animal housing were redacted from the version sent to the reviewers. The reviewers were provided a study data package, also blinded to test agents, containing basic in life study information such as body weight and survival curves and information concerning the generation of pups from the *in utero* exposures.

Report Title: Draft Report of Partial Findings from the National Toxicology Program Carcinogenesis Studies of Test Articles A and B (and associated Study Data Package)

Reviewers’ Names:

David Dorman, D.V.M., Ph.D., North Carolina State University
Russell Cattley, D.V.M., Ph.D., Auburn University
Michael Pino, D.V.M., Ph.D., Pathology consultant

Charge: To peer review the draft report and comment on whether the scientific evidence supports NTP’s conclusion(s) for the study findings.

1. Scientific criticisms:

- a. Please comment on whether the information presented in the draft report, including presentation of data in any tables, is clearly and objectively presented. Please suggest any improvements.

All three reviewers found the results to be clearly and objectively presented, although there were suggestions to provide historical control information for brain and heart lesions for female Harlan Sprague Dawley rats, clarify statements about the specific statistical tests used and the presence or lack of statistical significance of the brain

gliomas in the Results, and expand the conclusions statements to clarify the basis for the conclusions.

- b. Please comment on whether NTP’s scientific interpretations of the data are objective and reasonable. Please explain why or why not.

The reviewers stated that the NTP had performed an adequate and objective peer review of the pathology data, and the statistical approaches used were consistent with other NTP studies. The methods were described as objective and reasonable. The interpretations of the data, including the limitations, were also reasonable and objective. One reviewer found the data on schwannomas of the heart to be more compelling with respect to an association with treatment than the brain gliomas. This reviewer summarized the findings as:

“In the heart the evidence for a carcinogenic effect can be based on 1) the presence of the tumors in all six of the test article groups versus none in the controls 2) the statistically significant trend for schwannomas with both compounds and the statistically significant increase in incidence in the 4X (top) dose for test article B; 3) the fact that the incidence of the tumors in both 4X dose groups approaches or exceeds the high end of the historical control range; and 4) the tumors in the 4X group of test article B are accompanied by a higher incidence of Schwann cell hyperplasia. Using the NTP’s guide for levels of evidence for carcinogenic activity, I would consider the heart schwannomas as ‘Some Evidence’ of carcinogenic activity.

The proliferative lesions in the brain are more difficult to interpret because 1) their low incidence that was well within the historical control range, 2) lack of clear dose response; and 3) lack of statistical significance (except for the significant exposure-dependent trend for test article B. . . . However, the presence of malignant gliomas and/or foci of glial cell hyperplasia in 5 of 6 test article groups for both sexes vs none in controls of either sex is suggestive of a test

article effect. . . I would consider the malignant gliomas as ‘Equivocal Evidence’ of carcinogenic activity.”

2. Please identify any Information that should be added or deleted:

One reviewer suggested that more information be given on the time when tumors were observed (e.g., at terminal necropsy, or early in the study) to help assess the possible impact of the decreased survival times in the control animals on tumor incidence. This reviewer also suggested a discussion of how the survival of control male rats in this study compared to the historical control data. There was also concern that the diagnostic criteria developed by the PWG and used in the current study would impact the historical control incidence rates reported in Table D.

3. The scientific evidence supports NTP’s conclusion(s) for the study findings:

The NTP’s overall draft conclusion was as follows: “Under the conditions of these studies, the observed hyperplastic lesions and neoplasms outlined in this partial report are considered likely the result of exposures to test article A and test article B. The findings in the heart were statistically stronger than the findings in the brain.”

The reviewers had the option of agreeing, agreeing in principle, or disagreeing with the draft conclusions. All three reviewers agreed in principle, reiterating issues discussed above.

APPENDIX G – NIH REVIEWER’S COMMENTS

National Institutes of Health

Peer Review Charge and Reviewer’s Comments

Purpose: To provide independent peer review of the pathology diagnoses and statistical evaluation of the partial findings from NTP’s studies. Background materials included the draft NTP report, introductory materials on RFR, and details on the methods dealing with the field generation and statistical analyses references and guidance. The reviewers were provided a study data package, containing basic in life study information such as body weight and survival curves, information concerning the generation of pups from the *in utero* exposures, and raw pathology data.

Report Title: Draft Report of Partial Findings from the National Toxicology Program Carcinogenesis Studies of Test Articles A and B (and associated Study Data Package)

Reviewers’ Names:

Diana C. Haines, D.V.M., Frederick National Laboratory
Michael S. Lauer, M.D., Office of Extramural Research, NIH
Maxwell P. Lee, Ph.D., Laboratory of Cancer Biology and Genetics, NCI,
Aleksandra M. Michalowski, M.Sc., Ph.D., Laboratory of Cancer Biology and Genetics, NCI
R. Mark Simpson, D.V.M., Ph.D., Laboratory of Cancer Biology and Genetics, NCI
[Sixth reviewer's name and comments are withheld.]

Charge: To peer review the draft report, statistical analyses, and pathology data and comment on whether the scientific evidence supports NTP’s conclusion(s) for the study findings.

Reviewer’s comments and NTP responses to the comments are provided.

- Appendix G1: Reviewer’s comments
- Appendix G2: NTP’s responses to NIH reviewer’s comments

Appendix G1: Reviewer's comments

Reviewer: Diana C. Haines, D.V.M., Frederick National Laboratory

April 5, 2016

Dr. Tabak,

I've always relied on experts, not myself, for statistical analysis, and so do not feel qualified to address the statistical methods used. My training and experience has been in veterinary pathology, including QA review of NTP studies, and serving on PWGs, so will give my opinion on the pathology interpretation (biological significance rather than statistical significance).

Having perused the 3 RFR Draft Report and the raw data, all appears to be in order, including QA of the histopathology (technique) as well as PWG review (diagnosis). Looking at the data, I agree with the report's conclusion: *Under the conditions of these studies, the hyperplastic lesions and neoplasms observed in male rats are considered likely the result of exposures to GSM- an CDMA-modulated RFR. The findings in the heart were statistically stronger than the findings in the brain.* But note, it is "considered likely" not "definitely is".

There may be also several caveats relating to "under the conditions of these studies", including how well the conditions recapitulate actual human exposure: whole body exposure from in utero to old age; 18.5 hours/day (10 min on/10 min off, for total of 9hr actual exposure); and dose^A. I'm not physicist, so have to presume experts analyzed and accepted concept of the reverberation chamber, including "doses"^A as being relevant to human exposure.

^A Dosimetric Assessment paper: "As could be expected in a study following NTP protocols, the exposure levels for the rodents in this project exceed the limits for the wbSAR and psSAR defined in the IEEE Std C95.1-2005 safety standard for human exposure to mobile phone radiation. In the low dose exposure group the exposure level in the organs exceeds or is close to the localized SAR limit for the general public, except for a few low-water content tissues. More specifically, the psSAR over 1 g in the human head, is limited by the safety standards to <2W/kg, whereas, in the low dose rodents the SAR averaged over the whole brain is >2.4 W/kg for mice, and >1.3 W/kg for rats, hence similar to the limit. Furthermore, the psSAR and oSAR have larger uncertainty compared to the wbSAR. Deviations of the exposure level from the target dose, especially during the early exposure period, should be carefully evaluated in the interpretation of the final biological studies.

Results from the companion mouse study will hopefully add some insight.

Diana Copeland Haines, DVM

Diplomate, American College of Veterinary Pathologists
 Senior Staff Pathologist, Pathology Section
 Pathology/Histotechnology Laboratory
 Leidos Biomedical Research, Inc.
 Frederick National Laboratory for Cancer Research
 P.O. Box B, Frederick, MD 21702
 Phone: 301-846-5921 Fax: 301-846-1953
Diana.Haines@fnlcr.nih.gov
<http://ncifrederick.cancer.gov/rtp/lasp/phl/>

Appendix G1: Reviewer's comments

Reviewer: Michael S. Lauer, M.D., Office of Extramural Research, NIH

Michael S Lauer, MD (OER)

Review of NTP paper: "Report of Partial Findings from the National Toxicology Program Carcinogenesis Studies of Cell Phone Radiofrequency Radiation (Whole Body Exposures)"
March 20, 2016

Summary of findings:

This is a partial report, a report which is presumably part of a larger set of studies involving 2 species (mice and rats), 2 sexes (male, female), and multiple tissue types, all based on 90-week studies of two different types (GSM and CDMA) of cell phone radiofrequency radiation (RFR). In this partial report, we are given findings regarding brain gliomas and heart schwannomas in male and female Harlan Sprague Dawley rats which were exposed to control or 3 different levels (1.5, 3.0, 6.0) of two types (GSM and CDMA) of RFR. There were 90 rats in each group. Using the poly-3 test with the Bieler-Williams variance adjustment, the authors found a statistically significant increase in the rate of brain gliomas in males exposed to CDMA RFR. Using the poly-6 test, the authors found a statistically significant increase in the rates of heart schwannomas in males exposed to GSM and CDMA. There were no statistically significant differences in rates of gliomas or schwannomas in females; also there was no statistically significant increase in rates of gliomas in males exposed to GSM RFR.

Comments:

- 1) Why aren't we being told, at least at a high level, of the results of other experiments (i.e., male and female mice, tissues other than heart and brain, tumors other than glioma and schwannoma)? Given the multiple comparisons inherent in this kind of work (see pages 27-30 and Table 13 of the FDA guidance document), there is a high risk of false positive discoveries. In the absence of knowing other findings, we must worry about selective reporting bias.
- 2) I was able to reproduce the authors' positive P-value findings (see Appendix 1, R code) using the [MCPAN R package](#). However, I'm getting slightly different values for adjusted denominators (also in Appendix 1).
- 3) I was able to reproduce the authors' findings of longer survival with RFR (see Appendix 1, R code).
- 4) I have a number of questions about the study design:
 - a. Were control rats selected in utero like the exposed rats were?
 - b. Were pregnant dams assigned to different groups by formal randomization? If not, why not?
 - c. Why were pups in the same litter included? Did the authors take any steps in their analyses to account for the resulting absence of i.i.d?
 - d. The authors state that at most 3 pups were chosen per litter. How were the 3 pups chosen (and the others presumably not used for this experiment)? Were the 3 pups that were chosen selected by formal randomization? If not, why not?

- e. Were all analyses based on the intent-to-treat principle? Were there any crossovers? Were all rats accounted for by the end of the experiment and were all rats who started in the experiment included in the final analyses?
 - f. Blinding: The authors state that “All PWG reviewer were conducted blinded with respect to treatment group,” but in the very next phrase write “only identifying the test articles as ‘test agent A’ or ‘test agent B.’” Why was this information (test agent A or B) given? The blinding was not complete.
- 5) Sample size:
- a. Did the authors perform a prospective (that is before initiation of the work) sample size calculation? If so, what were the prior assumptions? In other words, why did the authors choose to study 90 rats in each group and why did they set the maximum duration to 90 weeks (instead of 104 weeks)?
 - b. I used a [publicly available](#) simulation package¹ to calculate the study power for male rats based on the following (see Appendix 2, power calculation simulation studies):
 - i. Control tumor rate of ~1.5%.
 - ii. Risk ratio 2.5 in the group receiving the highest dose
 - iii. 2-sided Alpha = 0.005 (based on Table 13 of the FDA guidance document). Note this low alpha of 0.005 for poly-k trend tests is recommended to minimize the risk of false positive discoveries.
 - iv. Sample size of 90 for each group with one planned sacrifice.
 - v. Low lethality with lethality parameters set according to study duration and Weibull shape parameter (see Table 3 of Moon et al¹). When I re-ran the simulations using intermediate lethality, results were not materially changed.
 - vi. Study duration 90 weeks
 - vii. 5000 simulations
 - viii. Note – I used dose levels of 0,1,2, and 4 because I was unable to adjust these on the web site (despite trying 3 different browsers).
 - c. Based on these inputs, the recommendations in Table 13 of the FDA guidance document, and a sample size of 90 rats in each group, I find very low power (<5%, see Appendix 2). Even allowing for a risk ratio of 5.0 (a level that is clinically unlikely), the power for 2-sided alpha=0.005, k=3 and low lethality is only ~14% (see Appendix 2).
 - d. The low power implies that there is a high risk of false positive findings², especially since the epidemiological literature questions the purported association between cell phone exposure and cancer.³
- 6) Summary: I am unable to accept the authors’ conclusions:
- a. We need to know all other findings of these experiments (mice, other tumor types) given the risk of false positive findings and reporting bias. It would be helpful to have a copy of the authors’ statistical code.
 - b. We need to know whether randomization was employed to assign dams to specific groups (control and intervention).

- c. We need to know whether randomization was employed to determine which pups from each litter were chosen for continued participation in the experiment.
- d. We need to know whether there was a formal power/sample size calculation performed prior to initiation of the experiment. If not, why not? If yes, we need to see the details. In particular, we need to know whether the authors followed the recommendations of the FDA guidance document (in particular Table 13).
- e. I suspect that this experiment is substantially underpowered and that the few positive results found reflect false positive findings.² The higher survival with RFR, along with the prior epidemiological literature, leaves me even more skeptical of the authors' claims.

References:

1. Moon H, Lee JJ, Ahn H, Nikolova RG. A Web-based Simulator for Sample Size and Power Estimation in Animal Carcinogenicity Studies. *J Stat Software*; Vol 1, Issue 13 . 2002. doi:10.18637/jss.v007.i13.
2. Ioannidis JPA. Why most published research findings are false. Jantsch W, Schaffler F, eds. *PLoS Med*. 2005;2(8):e124. doi:10.1371/journal.pmed.0020124.
3. Frei P, Poulsen AH, Johansen C, Olsen JH, Steding-Jessen M, Schüz J. Use of mobile phones and risk of brain tumours: update of Danish cohort study. *BMJ*. 2011;343.

Appendix 1: Attempted replication of positive findings

Review of NTP paper on cell phone RFR and certain cancers

Attempt to reproduce the positive findings

Data from Larry Tabak

Code by Mike Lauer

```
setwd("~/Desktop/Files to save")
```

```
library(MCPAN)
```

```
library(rms)
```

```
library(Hmisc)
```

Read in CDMA NTP data

```
CDMA <- read.csv("~/Desktop/Files to save/NTP CDMA Raw Tumor Data.csv")
```

Survival and treatment group, adjusting for sex, by Cox proportional hazards

```
CDMA$status<-1
```

```
CDMA$S<-Surv(CDMA$Removal.Day, CDMA$status)
```

```
f<-cph(S~Treatment+Sex, data=CDMA)
```

```
f
```

Survival greater (better) for 3.0W, P=0.0157, for 6.0W, P=0.0260

Table 1 -- Poly-3 test for malignant glioma in males CDMA

```
males_CDMA<-subset(CDMA, Sex=='M')
```

```
poly3test(time=males_CDMA$Removal.Day, status=males_CDMA$Brain.Glioma.Malignant,
           f=males_CDMA$Dose, k=3, type='Williams', method='BW', alternative='greater')
```

P=0.039

```
poly3ci(time=males_CDMA$Removal.Day, status=males_CDMA$Brain.Glioma.Malignant,
         f=males_CDMA$Dose, k=3, type='Williams', method='BW', alternative='greater')
```

Call result:

Sample estimates, using poly- 3 -adjustment

	0	1.5	3	6
x	0.0000	0.0000	0.0000	3.0000
n	90.0000	90.0000	90.0000	90.0000
adjusted n	63.8258	72.3688	76.6821	64.8154
adjusted estimate	0.0000	0.0000	0.0000	0.0463

Table 3 -- Poly-6 test for malignant Schwannoma in males CDMA

```
poly3test(time=males_CDMA$Removal.Day,
           status=males_CDMA$Heart.Schwannoma.Malignant, f=males_CDMA$Dose, k=6,
           type='Williams', method='BW', alternative='greater')
```

P=0.0005

```
poly3ci(time=males_CDMA$Removal.Day,
         status=males_CDMA$Heart.Schwannoma.Malignant, f=males_CDMA$Dose,
         k=3, type='Williams', method='BW')
```

Call result:

Sample estimates, using poly- 3 -adjustment

	0	1.5	3	6
x	0.0000	2.0000	3.0000	6.0000
n	90.0000	90.0000	90.0000	90.0000
adjusted n	63.8258	72.3971	77.0575	66.5582
adjusted estimate	0.0000	0.0276	0.0389	0.0901

Read in GSM NTP data

```
GSM <- read.csv("~/Desktop/Files to save/NTP GSM Raw Tumor data.csv")
```

Survival and treatment group, adjusting for sex, by Cox proportional hazards

```
GSM$status<-1
GSM$S<-Surv(GSM$Removal.Day, GSM$status)
f<-cph(S~Treatment+Sex, data=GSM)
f
```

Survival greater (better) for 6.0W, P=0.0048

```
males_GSM<-subset(GSM, Sex=='M')
```

Table 3 -- Poly-6 test for malignant Schwannomas in males GSM

```
poly3test(time=males_GSM$Removal.Day, status=males_GSM$Heart.Schwannoma.Malignant,
          f=males_CDMA$Dose, k=6, type='Williams', method='BW', alternative='greater')
```

```
# P=0.004
```

```
poly3ci(time=males_GSM$Removal.Day, status=males_GSM$Heart.Schwannoma.Malignant,
         f=males_CDMA$Dose, k=3, type='Williams', method='BW', alternative='greater')
```

Call result:

Sample estimates, using poly- 3 -adjustment

	0	1.5	3	6
x	0.0000	2.0000	1.0000	5.0000
n	90.0000	90.0000	90.0000	90.0000
adjusted n	63.8258	73.1547	76.1127	77.0723
adjusted estimate	0.0000	0.0273	0.0131	0.0649

Appendix 2: Simulations for power calculations

Power Simulations for NTP Cell Phone RFR paper (from
<https://biostatistics.mdanderson.org/acss/Login.aspx> and
<https://www.jstatsoft.org/article/view/v007i13>)¹

Michael Lauer, MD (OER)
March 19, 2016

- 1) For malignant gliomas (Table 1), $P = 0.005$, $HR = 2.5$, $k=3$

The University of Texas M. D. Anderson Cancer Center
Sample Size and Power Estimation for Animal Carcinogenicity Studies

Reference: "A Web-based Simulator for Sample Size and Power
Estimation in Animal Carcinogenicity Studies."
Hojin Moon, J. Jack Lee, Hongshik Ahn and Rumiana G. Nikolova,
Journal of Statistical Software. (2002)¹

*** Input Parameters ***

Selected Seed = 3000
Number of Groups = 4
Dose metric of each group:
0.00 1.00 2.00 4.00
Number of animals in each group
90 90 90 90
Number of sacrifices including a terminal sacrifice = 1
Sacrifice time points in weeks:

Study duration = 90 weeks
Number of INTERIM sacrificed animals in each interval:
Background tumor onset probability at the end of the study = 0.01
Tumor onset distribution assumed: Weibull with a shape parameter 3.00
Hazard ratio(s) of dose vs. control group
1.50 2.00 2.50
Competing Risks Survival Rate (CRSR) for each group:
0.70 0.70 0.70 0.70
Tumor lethality parameter entered = 23.00
Level of the test = 0.01
One-sided or two-sided test = 2 sided test
Number of simulation runs = 5000

*** Simulation Results ***

dose group 0:

average tumor rate = 0.0149

average competing risks survival rate = 0.6990

average lethality = 0.0816

sacrifice time	d	a1	b1	a2	b2
45	0.0000	0.0000	0.0060	0.0000	0.0000
67	0.0002	0.0002	0.0334	0.0000	0.0000
78	0.0003	0.0005	0.0729	0.0000	0.0000
90	0.0005	0.0023	0.1855	0.0094	0.6887

dose group 1:

average tumor rate = 0.0225

average competing risks survival rate = 0.7000

average lethality = 0.0784

sacrifice time	d	a1	b1	a2	b2
45	0.0001	0.0000	0.0059	0.0000	0.0000
67	0.0003	0.0002	0.0325	0.0000	0.0000
78	0.0004	0.0008	0.0720	0.0000	0.0000
90	0.0007	0.0034	0.1851	0.0145	0.6842

dose group 2:

average tumor rate = 0.0297

average competing risks survival rate = 0.6997

average lethality = 0.0772

sacrifice time	d	a1	b1	a2	b2
45	0.0001	0.0000	0.0059	0.0000	0.0000
67	0.0004	0.0003	0.0331	0.0000	0.0000
78	0.0005	0.0012	0.0721	0.0000	0.0000
90	0.0010	0.0045	0.1829	0.0191	0.6790

dose group 3:

average tumor rate = 0.0366

average competing risks survival rate = 0.7007

average lethality = 0.0772

sacrifice time	d	a1	b1	a2	b2
45	0.0001	0.0000	0.0059	0.0000	0.0000
67	0.0005	0.0003	0.0330	0.0000	0.0000

78	0.0006	0.0013	0.0716	0.0000	0.0000
90	0.0012	0.0054	0.1812	0.0238	0.6749

Positive Trend (Power): 0.0238

2) For malignant Schwannomas (Table 3), $P = 0.005$, $HR = 2.5$, $k=6$

The University of Texas M. D. Anderson Cancer Center
Sample Size and Power Estimation for Animal Carcinogenicity Studies

Reference: "A Web-based Simulator for Sample Size and Power
Estimation in Animal Carcinogenicity Studies."
Hojin Moon, J. Jack Lee, Hongshik Ahn and Rumiana G. Nikolova,
Journal of Statistical Software. (2002)¹

*** Input Parameters ***

Selected Seed = 3000
Number of Groups = 4
Dose metric of each group:
0.00 1.00 2.00 4.00
Number of animals in each group
90 90 90 90
Number of sacrifices including a terminal sacrifice = 1
Sacrifice time points in weeks:

Study duration = 90 weeks
Number of INTERIM sacrificed animals in each interval:
Background tumor onset probability at the end of the study = 0.01
Tumor onset distribution assumed: Weibull with a shape parameter 6.00
Hazard ratio(s) of dose vs. control group
1.50 2.00 2.50
Competing Risks Survival Rate (CRSR) for each group:
0.70 0.70 0.70 0.70
Tumor lethality parameter entered = 45.00
Level of the test = 0.01
One-sided or two-sided test = 2 sided test
Number of simulation runs = 5000

*** Simulation Results ***

dose group 0:

average tumor rate = 0.0149

average competing risks survival rate = 0.6990

average lethality = 0.0631

sacrifice time	d	a1	b1	a2	b2
45	0.0000	0.0000	0.0060	0.0000	0.0000
67	0.0001	0.0001	0.0335	0.0000	0.0000
78	0.0002	0.0003	0.0732	0.0000	0.0000
90	0.0005	0.0019	0.1859	0.0096	0.6887

dose group 1:

average tumor rate = 0.0225

average competing risks survival rate = 0.7000

average lethality = 0.0602

sacrifice time	d	a1	b1	a2	b2
45	0.0000	0.0000	0.0059	0.0000	0.0000
67	0.0001	0.0001	0.0326	0.0000	0.0000
78	0.0003	0.0005	0.0723	0.0000	0.0000
90	0.0006	0.0029	0.1856	0.0148	0.6842

dose group 2:

average tumor rate = 0.0297

average competing risks survival rate = 0.6997

average lethality = 0.0582

sacrifice time	d	a1	b1	a2	b2
45	0.0000	0.0000	0.0059	0.0000	0.0000
67	0.0002	0.0001	0.0333	0.0000	0.0000
78	0.0004	0.0007	0.0726	0.0000	0.0000
90	0.0009	0.0038	0.1837	0.0195	0.6790

dose group 3:

average tumor rate = 0.0366

average competing risks survival rate = 0.7007

average lethality = 0.0588

sacrifice time	d	a1	b1	a2	b2
45	0.0000	0.0000	0.0059	0.0000	0.0000
67	0.0003	0.0001	0.0332	0.0000	0.0000
78	0.0005	0.0007	0.0722	0.0000	0.0000
90	0.0011	0.0046	0.1821	0.0243	0.6749

Positive Trend (Power): 0.0230

3) For further consideration, $P = 0.005$, $HR = 5$, $k=3$

The University of Texas M. D. Anderson Cancer Center
Sample Size and Power Estimation for Animal Carcinogenicity Studies

Reference: "A Web-based Simulator for Sample Size and Power
Estimation in Animal Carcinogenicity Studies."
Hojin Moon, J. Jack Lee, Hongshik Ahn and Rumiana G. Nikolova,
Journal of Statistical Software. (2002) In Press.

*** Input Parameters ***

Selected Seed = 3000
Number of Groups = 4
Dose metric of each group:
0.00 1.00 2.00 4.00
Number of animals in each group
90 90 90 90
Number of sacrifices including a terminal sacrifice = 1
Sacrifice time points in weeks:

Study duration = 90 weeks
Number of INTERIM sacrificed animals in each interval:
Background tumor onset probability at the end of the study = 0.01
Tumor onset distribution assumed: Weibull with a shape parameter 3.00
Hazard ratio(s) of dose vs. control group
2.00 3.50 5.00
Competing Risks Survival Rate (CRSR) for each group:
0.70 0.70 0.70 0.70
Tumor lethality parameter entered = 23.00
Level of the test = 0.01
One-sided or two-sided test = 2 sided test
Number of simulation runs = 5000

*** Simulation Results ***

dose group 0:
average tumor rate = 0.0149
average competing risks survival rate = 0.6990

average lethality = 0.0816

sacrifice time	d	a1	b1	a2	b2
45	0.0000	0.0000	0.0060	0.0000	0.0000
67	0.0002	0.0002	0.0334	0.0000	0.0000
78	0.0003	0.0005	0.0729	0.0000	0.0000
90	0.0005	0.0023	0.1855	0.0094	0.6887

dose group 1:

average tumor rate = 0.0301

average competing risks survival rate = 0.7000

average lethality = 0.0743

sacrifice time	d	a1	b1	a2	b2
45	0.0001	0.0000	0.0059	0.0000	0.0000
67	0.0004	0.0003	0.0324	0.0000	0.0000
78	0.0005	0.0011	0.0717	0.0000	0.0000
90	0.0009	0.0045	0.1839	0.0194	0.6789

dose group 2:

average tumor rate = 0.0515

average competing risks survival rate = 0.6997

average lethality = 0.0774

sacrifice time	d	a1	b1	a2	b2
45	0.0002	0.0000	0.0058	0.0000	0.0000
67	0.0007	0.0006	0.0328	0.0000	0.0000
78	0.0009	0.0020	0.0713	0.0000	0.0000
90	0.0017	0.0076	0.1795	0.0331	0.6638

dose group 3:

average tumor rate = 0.0727

average competing risks survival rate = 0.7007

average lethality = 0.0804

sacrifice time	d	a1	b1	a2	b2
45	0.0003	0.0000	0.0059	0.0000	0.0000
67	0.0010	0.0006	0.0327	0.0000	0.0000
78	0.0013	0.0028	0.0701	0.0000	0.0000
90	0.0025	0.0107	0.1755	0.0470	0.6496

Positive Trend (Power): 0.1420

4) For further consideration, same as in baseline (1) but with intermediate lethality

*** Input Parameters ***

Selected Seed = 3000

Number of Groups = 4

Dose metric of each group:

0.00 1.00 2.00 4.00

Number of animals in each group

90 90 90 90

Number of sacrifices including a terminal sacrifice = 1

Sacrifice time points in weeks:

Study duration = 90 weeks

Number of INTERIM sacrificed animals in each interval:

Background tumor onset probability at the end of the study = 0.01

Tumor onset distribution assumed: Weibull with a shape parameter 3.00

Hazard ratio(s) of dose vs. control group

1.50 2.00 2.50

Competing Risks Survival Rate (CRSR) for each group:

0.70 0.70 0.70 0.70

Tumor lethality parameter entered = 225.00

Level of the test = 0.01

One-sided or two-sided test = 2 sided test

Number of simulation runs = 5000

*** Simulation Results ***

dose group 0:

average tumor rate = 0.0149

average competing risks survival rate = 0.6990

average lethality = 0.3936

sacrifice time	d	a1	b1	a2	b2
45	0.0004	0.0000	0.0060	0.0000	0.0000
67	0.0014	0.0001	0.0334	0.0000	0.0000
78	0.0014	0.0004	0.0729	0.0000	0.0000
90	0.0019	0.0015	0.1855	0.0063	0.6887

dose group 1:

average tumor rate = 0.0225

average competing risks survival rate = 0.7000

average lethality = 0.3852

sacrifice time	d	a1	b1	a2	b2
45	0.0006	0.0000	0.0059	0.0000	0.0000
67	0.0022	0.0001	0.0325	0.0000	0.0000
78	0.0020	0.0006	0.0720	0.0000	0.0000
90	0.0029	0.0023	0.1851	0.0097	0.6842

dose group 2:

average tumor rate = 0.0297

average competing risks survival rate = 0.6997

average lethality = 0.3839

sacrifice time	d	a1	b1	a2	b2
45	0.0008	0.0000	0.0059	0.0000	0.0000
67	0.0029	0.0003	0.0331	0.0000	0.0000
78	0.0027	0.0008	0.0721	0.0000	0.0000
90	0.0039	0.0031	0.1829	0.0127	0.6790

dose group 3:

average tumor rate = 0.0366

average competing risks survival rate = 0.7007

average lethality = 0.3897

sacrifice time	d	a1	b1	a2	b2
45	0.0009	0.0000	0.0059	0.0000	0.0000
67	0.0037	0.0003	0.0330	0.0000	0.0000
78	0.0033	0.0009	0.0716	0.0000	0.0000
90	0.0048	0.0037	0.1812	0.0157	0.6749

Positive Trend (Power): 0.0219

References:

1. Moon H, Lee JJ, Ahn H, Nikolova RG. A Web-based Simulator for Sample Size and Power Estimation in Animal Carcinogenicity Studies. *J Stat Software*; Vol 1, Issue 13 . 2002. doi:10.18637/jss.v007.i13.
2. Ioannidis JPA. Why most published research findings are false. Jantsch W, Schaffler F, eds. *PLoS Med*. 2005;2(8):e124. doi:10.1371/journal.pmed.0020124.
3. Frei P, Poulsen AH, Johansen C, Olsen JH, Steding-Jessen M, Schüz J. Use of mobile phones and risk of brain tumours: update of Danish cohort study. *BMJ*. 2011;343.

Appendix G1: Reviewer's comments

Reviewer: Maxwell P. Lee, Ph.D., Laboratory of Cancer Biology and Genetics, NCI

I think the study was well designed and the analyses and results were clearly presented.

My main concern is the control data. Since the main finding was the increased incidence rates of heart schwannomas and brain gliomas in male Harlan Sprague Dawley rats exposed to GSM- or CDMA-modulated cell phone RFR, my analyses and evaluation below were focused on the male rats.

My concern regarding the control data came from the following two considerations. First, we need to consider sample variation. The incidence rates of the current controls for brain gliomas and heart schwannomas were 0. However, the historical controls were 1.67% for gliomas (range 0-8%) and 1.30% for schwannomas (0-6%). Given that there were substantial variations among the historical controls and the concurrent control is at the lowest end of the range, it is important to evaluate how different estimates of control incidence rates may impact the results of analyses. Supplementary Table S1 shows that for gliomas with 1.7% incidence rate we have 40%, 37%, 17%, and 6% of chance to observe 0 tumor, 1 tumor, 2 tumors, and greater than 2 tumors, respectively; heart schwannomas has similar distribution. Given the low incidence rate and moderate sample size of the control, even after observing 0 tumor in the current study, the 'true' incidence rate may be higher than 0. If we were repeating the experiment, we may see some control studies have 1 or more tumors. Second, it is puzzling why the control had short survival rate. Given that most of the gliomas and heart schwannomas are late-developing tumors, it is possible that if the controls were living longer some tumors might develop. Although the use of poly-3 (or poly-6) test intended to adjust the number of rats used in the study, it is still important to re-evaluate the analysis by considering the incidence rate in controls not being 0.

Therefore I have performed the analyses using the original data as well as the data modified by adding 1 tumor to the control. I implemented the poly-3 (or poly-6) trend test in R using the formula described in the file, Poly3 correction factor[1].docx.

The results are summarized in Table 1 for brain gliomas

Table 1. Incidence of brain gliomas in male rats exposed to GSM- or CDMA-modulated RFR, comparing control data with 0 vs. 1 tumor.

RFR	W/kg				pvalue
	0	1.5	3	6	
GSM	0	3	3	2	0.9771
GSM	1	3	3	2	0.8668
CDMA	0	0	0	3	0.0233
CDMA	1	0	0	3	0.1077

Poly-6 adjusted rates were used in the chi-square trend test. The 1st and 3rd rows correspond to the original data with 0 tumor observed in the control group (The numbers in Table 1 here are identical to those in Table 1 in the original report). The test is significant for CDMA exposures (pvalue = 0.0233). However, it is not significant after adding 1 tumor to the control group (pvalue = 0.1077, the 4th row).

Similar analysis was performed for heart schwannomas. The results are summarized in Table 2.

Table 2. Incidence of heart schwannomas in male rats exposed to GSM- or CDMA-modulated RFR, comparing control data with 0 vs. 1 tumor.

RFR	W/kg				pvalue
	0	1.5	3	6	
GSM	0	2	1	5	0.0431
GSM	1	2	1	5	0.1079
CDMA	0	2	3	6	0.0144
CDMA	1	2	3	6	0.0365

Poly-3 adjusted rates were used in the chi-square trend test. The 1st and 3rd rows correspond to the original data with 0 tumor observed in the control group (The numbers in Table 2 here are identical to those in Table 3 in the original report). The tests are significant for both GSM (pvalue = 0.0431) and CDMA (pvalue = 0.0144) exposures. However, only CDMA exposure remains significant after adding 1 tumor to the control group (pvalue = 0.0365, the 4th row).

Since the incidence of heart schwannomas in the 6 W/kg males was significantly higher in CDMA exposed males than the control group in the original report, I also analyzed the impact of adding 1 tumor to the control group

Table 3. Incidence of heart schwannomas in male rats exposed to 6 W/kg CDMA-modulated RFR, comparing control data with 0 vs. 1 tumor.

RFR	W/kg		pvalue
	0	6	
CDMA	0	6	0.0381
CDMA	1	6	0.0986

Poly-3 adjusted rates were used in the chi-square trend test. The 1st row corresponds to the original data with 0 tumor observed in the control group. The test was significant for CDMA exposures (pvalue = 0.0381). However, it was not significant after adding 1 tumor to the control group (pvalue = 0.0986, the 2nd row).

Conclusions

Increased incidence of heart schwannomas in male rats exposed to GSM- or CDMA-modulated RFR is statistically significant by the chi-square trend test. The evidence is better for CDMA exposure than GSM exposure. I think additional experiments are needed to assess if the incidence of brain gliomas in male rats exposed to GSM- or CDMA-modulated RFR is significantly higher than the control group or not.

My additional comments are summarized below.

1. I compared poly-3 adjusted number from Table 3 in the original report versus the poly-3 adjusted number that I calculated using the raw data from the excel files. Supplementary Figure S1 shows that these two sets of numbers agree with each other in general. This is in contrast to the comparison for poly-6 adjusted number from Table 1 in the original report versus the poly-6 adjusted number that I calculated using the raw data from the excel files (Supplementary Figure S2). In fact, the adjusted rat numbers from Table 1 and Table 3 of the original report look quite similar (Supplementary Figure S3). This suggests that the poly-3 adjusted number was used in the footnotes in both Table 1 and Table 3 in the original report.
2. I noted that in Table S2 the adjusted numbers in from.original.report and poly3 are identical at Dose 0 and 1.5 for both CDMA and GSM as well as at Dose 3 for GSM but differ slightly in the other treatment doses for heart schwannomas. One possible cause of the difference is that the version of the raw data in the excel files differs from that used to generate the original report. The second possibility is typ in the footnote in Table 3. I also generated Table S3 that has the poly-6 adjusted numbers for brain gliomas. The two sets of the poly-6 adjusted numbers are ver different.
3. There are a couple of errors in the footnote of Table in the original report. 2/74.05 (5%) should be 2/74.05 (2.7%). 3/78.67 (4%) should be 3/78.67 (3.8%).

Supplementary Information

Table S1. Expected percentage of observing different numbers of tumors in the controls based on binomial distribution.

	0 tumor	1 tumor	2 tumors	>2 tumors
control for glioma	40%	37%	17%	6%
control for heart schwannoma	43%	37%	15%	5%

The percentage was calculated with 1.7% historical control rate for male rats (gliomas) and with poly-6 adjusted animal number, 53. Similarly, the percentage was calculated with 1.3% historical control rate for male (heart schwannoma) and with poly-3 adjusted animal number, 65.

Table S2. The poly-3 adjusted rat numbers in Table in the original report and those calculated from the raw data.

RFR	Dose	from.original.report	poly3
CDMA	0	65.47	65.47
CDMA	1.5	74.05	74.05
CDMA	3	78.67	78.35
CDMA	6	67.94	66.24
GSM	0	65.47	65.47
GSM	1.5	74.87	74.87
GSM	3	77.89	77.89
GSM	6	78.48	77.66


The numbers in from.original.report refers to the poly-3 adjusted rat number from Table 3 in the original report. The numbers in poly3 refers to the poly-3 adjusted rat numbers that I calculated from the raw data for heart schwannoma.

Table S3. The poly-6 adjusted rat numbers in Table in the original report and those calculated from the raw data.

RFR	Dose	from.original.report	poly6
CDMA	0	65.47	53.48
CDMA	1.5	74.05	65.94
CDMA	3	78.35	73.08
CDMA	6	66.24	57.5
GSM	0	65.47	53.48
GSM	1.5	74.93	67.84
GSM	3	78.27	71.43
GSM	6	77.1	72.55


The numbers in from.original.report refers to the poly-6 adjusted rat number from Table 1 in the original report. The numbers in poly6 refers to the poly-6 adjusted rat numbers that I calculated from the raw data for brain gliomas.

Figure S1. Comparison of poly-3 adjusted rat numbers between those from the original report versus those calculated from the raw data.




The poly-3 adjusted rat number from Table 3 of the original report is compare with the poly-3 adjusted rat number that I calculated from the raw data for heart schwannomas experiment

Figure S2. Comparison of poly-6 adjusted rat numbers between those from the original report versus those calculated from the raw data.



The poly-6 adjusted rat number from Table 1 of the original report is compared with the poly-6 adjusted rat number that I calculated from the raw data for brain gliomas experiment

Figure S3. Comparison of poly-6 adjusted rat numbers between those from the original report versus those calculated from the raw data.



The adjusted rat numbers from Table 1 and Table 3 of the original report are compared with each other.

Appendix G1: Reviewer's comments

Reviewer: Aleksandra M. Michalowski, M.Sc., Ph.D., Laboratory of Cancer Biology and Genetics, NCI

REVIEWER COMMENTS

Reviewer's Name:

Aleksandra M. Michalowski, Ph.D., M.Sc., National Cancer Institute/LCBG

Report Title:

Report of Partial Findings from the National Toxicology Program Carcinogenesis Studies of Cell Phone Radiofrequency Radiation (Whole Body Exposures); Draft 3-16-2016

Charge: To peer review the draft report and comment on whether the scientific evidence supports NTP's conclusion(s) for the study findings.

1. Scientific criticisms:

- a. *Please comment on whether the information presented in the draft report, including presentation of data in any tables, is clearly and objectively presented. Please suggest any improvements.*

Overall, the information included in the report is presented in a comprehensive and accurate manner. Specifically, the experimental design and conditions are sufficiently documented and the choice of statistical approaches is explained; the results are well organized and necessary details are provided.

Nevertheless, a few additions could be suggested:

(1) Appendix tables for all poly-k tests performed could be added. I believe this would enhance the presentation of the adjusted rates and the strength of the statistical evidence. As a possible example I prepared the below table using the R package *MCPAN* and its *poly3test()* function.

poly-3	Heart Schwannoma Malignant, Male				Heart Schwannoma Malignant, Female			
CDMA exposure	0	1.5	3	6	0	1.5	3	6
X	0	2	3	6	0	2	0	2
N	90	90	90	90	90	90	90	90
adjusted n	63.8	72.4	77.1	66.6	67.9	71.8	70.3	78.0
Dunnett contrast	—	1.5 - 0	3 - 0	6 - 0	—	1.5 - 0	3 - 0	6 - 0
Estimate	0	0.03	0.04	0.09	0	0.03	0	0.03
Statistic	—	1.24	1.58	2.45	—	1.26	0	1.24
p-value	—	0.2704	0.1542	0.0209	—	0.2466	0.7992	0.2562
Williams contrast	—	(6,3,1.5) - 0	(6,3) - 0	6 - 0	—	(6,3,1.5) - 0	(6,3) - 0	6 - 0
Estimate	0	0.05	0.06	0.09	0	0.02	0.01	0.03
Statistic	—	2.78	2.75	2.45	—	1.27	0.88	1.24
p-value	—	0.0056	0.0060	0.0138	—	0.1661	0.2871	0.1744

(2) In the portion of the text describing poly-k test results, p-values are given for significant pairwise comparisons; I would also give the p-values estimated for the significant trends (maximum test).

(3) Information could be included regarding the software or programming environment used for the computations.

(4) In the portion of the text describing differences in survival at the end of the study between control and RFR-exposed animals (page 5§2) the compared characteristic is not named (median survival, TSAC?) and also no numerical values of the estimates or the range of differences are given. I would add numbers in the text or an Appendix table showing the group survival estimates described in this paragraph.


Median survival			TSAC percentage		
CDMA	Female	Male	GSM	Female	Male
0	737	662.5	0	737	662.5
1.5	734	719	1.50	738	729
3	737	731	3	737	730
6	738.5	717	6	738	731

- b. *Please comment on whether NTP's scientific interpretations of the data are objective and reasonable. Please explain why or why not.*

Appropriate statistical design and methods were applied in accord with the FDA/NTP guidelines for conducting long-term rodent carcinogenicity studies and analyses. The results and limiting issues were objectively discussed. The critical issue of shorter survival in the male control group was addressed with regard to the percentage of animals surviving to terminal sacrifice in historical control data (avg. 47%, range 24% to 72%) and the possible impact of the observed age of tumor occurrence on the statistical inference.

I believe detailed information about animal selection and randomization procedures should be given so that the potential for allocation bias could be judged. As shown in the figure below, the lower survival rate to terminal sacrifice (28%) in the male control is accompanied by the higher rate of moribund sacrifice (49%); in the male group exposed to CDMA with 6 W/kg, a higher rate of natural death was observed (46%).

It has been reported that insufficient randomization can lead to differences in survival rates. As an example, in a carcinogenicity study on aspartame it was suggested that lack of randomization to different rooms may have possibly been the cause of low survival rates (27%) in the control female group due to a high background infection rate (EFSA, 2006; Magnuson, B., Williams, G.M., 2008).



2. Please identify any information that should be added or deleted:

A statement of the required statistical significance level should be added. FDA guidance suggests the use of significance levels of 0.025 and 0.005 for tests for positive trends in incidence rates of rare tumors and common tumors, respectively; for testing pairwise differences in tumor incidence the use of significance levels of 0.05 and 0.01 is recommended for rare and common tumors, respectively. If power calculations to determine the required sample size were performed, the results should also be included.

3. The scientific evidence supports NTP's conclusion(s) for the study findings:

The NTP's overall draft conclusion was as follows: "Under the conditions of these studies, the observed hyperplastic lesions and neoplasms outlined in this partial report are considered likely the result of exposures to test article A and test article B. The findings in the heart were statistically stronger than the findings in the brain."

In my view, the results support the conclusion of likely carcinogenic effect of the RFR-exposure on Schwannoma heart lesions in male Harlan Sprague Dawley rats.

Possible carcinogenic effects in the brain are marginal and are not sufficiently supported by statistical evidence in the male Harlan Sprague Dawley rats.

In the female Harlan Sprague Dawley rats very few lesions were observed in either site and statistical significance was not reached at all.

Appendix G1: Reviewer's comments

Reviewer: R. Mark Simpson, D.V.M., Ph.D., Laboratory of Cancer Biology and Genetics, NCI

Analysis of National Toxicology Program (NTP) study evaluating risk in rat lifetime exposure to GSM or CDMA RFR.

Notes:

The NTP study document acknowledges several study limitations [page 10, discussion section]. Potential limitations should prominently factor into considerations regarding the context of the findings, as well as their interpretation and application.

Working list of limitations potentially impacting NTP study interpretations

- Difficulty in achieving diagnostic consensus in lesions classifications of rare, unusual, and incompletely understood lesion association
- Document appears to indicate that the second Pathology Working Group (PWG) empaneled to review and obtain lesion classification consensus, following the inability of the initial PWG to do so, may have reviewed different lesions sets
- No record of clinical disease manifestations due to lesions involving heart and brain [note lesions in heart and brain are mutually exclusive; affected rats have either one or the other and do not appear to have the involvement of both organs together (appendix E)]
- Lesions, including malignancies, do not appear to materially shorten lifespan, except for a subgroup of rats (less than 1/3 of affected rats) with malignant Schwannomas in heart
- Lack of shortened lifespan as a consequence of malignancy for the majority of affected rats contrasts with shortened lifespan of male control rats for which there is absence of attributable cause of death. The survival of the control group of male rats in the current study (28%) was relatively low compared to other recent NTP studies (avg 47%, range 24 to 72%).
Creates greater reliance on statistical controlling for survival disparities and reliance on historical controls
- Reliance on historical controls made up of rats of different genetic strain background, held under different environmental conditions
- Absence of data on incidence of more frequently expected tumor occurrences in rats (background lesions)

Documenting the nature of the brain and cardiac lesions observed in RFR exposed rats and placing them into test article exposure-related context, in contrast to potential for their occurring spontaneously, are important and challenging goals. The NTP study limitations make the interpretation of reasonable risk more complicated. NTP acknowledgements of study limitations appear factored into one of NTP's reviewer's study conclusion, i.e., findings represent "some evidence" for a test article effect in statistically significant trend for Schwannomas; an opinion which is coupled with a conclusion for "equivocal evidence" of an effect in relation to malignant gliomas of the brain [NTP Appendix F, Reviewer Comments].

The summation from Appendix F reviewers regarding existence of test article effect is less than conclusive. The NTP study documents a series of cytoproliferative changes

in heart and brain. The nature of some of the changes is challenging diagnostically and appears to be incompletely understood. These findings are presented in the absence of complete analysis of the entire consequences of the study effects. For example, no potential significance for test article effect context is given to any of granular cell proliferative lesions of the brain, a finding mentioned only as a contrast to what was less well understood pathologically (NTP Appendix C, Pathology). It is noteworthy that the lesion types analyzed in the NTP RFR study under review are uncommon historically in rats, in the organs discussed. Furthermore, the malignancies of neuroglia appear to be paired with the occurrence of poorly understood changes involving neuroglial cell hyperplasias in the central and peripheral nervous systems. Little information can be gleaned from the literature about the nature and significance of these latter proliferative changes, interpreted by NTP as nonneoplastic and non-inflammation-reactive neuroglial cell in nature. Although unclear in the NTP study document, it is plausible that the particular lesion constellation, along with the relative novelty of some lesions, contributed to the lack of consensus regarding the nature of the lesions on the part of the initial PWG study pathologists. Concern raised by one of the reviewers (Appendix F, Reviewer Comments) regarding how this difficulty in ability to classify lesions might impact comparisons to historical control lesion incidence data (NTP Table D) is certainly principled.

The extraordinary PWG process, presumably posed by the difficult diagnostic interpretations, has the potential to influence the reliance on historical controls. In this regard, study limitations concerning determination of whether or not there is a test article effect include the substantially poor survival of male rats in the control group. The survival of the control group of male rats in the study under review (28%) was relatively low compared to other recent NTP studies (avg 47%, range 24 to 72%). This apparently led to greater statistical construction to account for the impact of study matched controls, and created increased reliance upon historical data of rare tumor incidences in control animals taken from other chronic carcinogenicity studies. NTP acknowledges a limitation in using the historical incident data and a small study match control group due to poor survivability. There are potential sources of variability when using historical controls of different rat strains and fluctuating study conditions (environment, vehicle, route of exposure, etc.), as is the case here. It seems less than clear what appropriate background lesion incidence is, as NTP indicates some data involve other strains of rats. The range of lesion incidence in historical controls could mean that the true incidence of some lesions varies considerably and might be considered rare or more common depending upon the incidence rate.

The guidance manual on Statistical Aspects of the Design, Analysis and Interpretation of Chronic Rodent Carcinogenicity Studies of Pharmaceuticals by the FDA provided for this review discusses applying comparisons using historical control lesion incidences at some length [beginning page 27, line 996]. Considering lesions as being rare or more common appears to influence selection of the level of statistical significance for comparisons. It appears that analysis for significant differences in tumor incidence between the control and the dose groups for these NTP studies has been established at the 0.05 level (NTP Tables 1,3,5). Interpretations of trend tests may be influenced by the choice of decision rule applied. Such choices can result in

about twice as large overall false positive error as that associated with control-high pairwise comparison tests [page 28, line 1012-1026]. The FDA guidance manual [page 31, line 1136] highlights concern regarding reliance upon historical control incidence data, stating that using historical control data in the interpretation of statistical test results is not very satisfactory because the range of historical control rates is usually too wide. This is especially true in situations in which the historical tumor rates of most studies used are clustered together, but a few other studies give rates far away from the cluster. When the range of historical control data is simply calculated as the difference between the maximum and the minimum of the historical control rates, the range does not consider the shape of the distribution of the rates. These circumstances may impose some limitations on optimal risk assessment designs.

Somewhat paradoxically then, NTP study limitations including that imposed due to reliance upon less than optimal historical control lesion incidence data for much of the comparisons between treated and untreated rats, is confronted by existence of a difficult to classify and incompletely understood lesion constellation interpreted to include neuroglial cell hyperplasia. Notwithstanding, this confounding proliferative lesion occurring in the context along with malignancies of apparently similar histogeneses, sustains a level of concern for a rare injury mechanism related to test article effect. Additional information about the study together with an assessment of the statistical analyses may enhance the value of this analysis.

R. Mark Simpson, D.V.M., Ph.D.

Appendix G2: NTP's responses to NIH reviewer's comments

Appendix G2: NTP's Responses to NIH Reviewer's Comments

NTP Responses to Pathology Reviewer' Comments

April 12, 2016

Reviewers: R. Mark Simpson, D.V.M., Ph.D. and Diana Copeland Haines, D.V.M.

Responses Relating to the Pathology Review Process

Drafts of the PWG reports are provided. As described in the PWG report, the specific task of the first PWG (January 29th 2016) was to: 1) confirm the presence of glial cell hyperplasia and malignant gliomas in the brain and Schwann cell hyperplasia and schwannomas in the heart; 2) develop specific diagnostic criteria in the brain for distinguishing glial cell hyperplasia from malignant glioma and gliosis, and in the heart for distinguishing between Schwann cell hyperplasia and schwannoma. The PWG participants confirmed the malignant gliomas and schwannomas, but the criteria for distinguishing between hyperplasia and neoplasia differed between the participants.

In order to clearly establish specific diagnostic criteria for the differentiation between hyperplastic and neoplastic lesions in the brain and heart, two additional PWGs were convened. The participants for the second (February 25, 2016) and third (March 3, 2016) PWGs were selected based on their distinguished expertise in the fields of neuropathology and cardiovascular pathology, respectively. Some of the participants were leaders in the International Harmonization of Nomenclature and Diagnostic Criteria initiative. The neuropathology experts of the second PWG confirmed the malignant gliomas in the brain, established diagnostic criteria for glial cell hyperplasia, and agreed that the hyperplastic lesions are within a continuum leading to malignant glioma. The cardiovascular pathology experts of the third PWG established specific diagnostic criteria for Schwann cell hyperplasia and schwannoma in the endocardium and myocardium, and reviewed and confirmed all cases of Schwann cell hyperplasia and schwannoma observed in these studies. The outcome of the PWG provided a very high degree of confidence in the diagnoses.

The participants of the first PWG (January 29th 2016) only reviewed a subset of the glial lesions that were observed in the studies. The review for the second PWG (February 25, 2016) included all glial lesions in the studies including the subset that was reviewed in the first PWG.

Responses Relating to Considerations of Historical Control Data

For NTP toxicology and carcinogenicity studies, the concurrent controls are always the primary comparison group. However, historical control information is useful particularly in instances when there is differential survival between controls and exposed groups, as was observed in the RFR studies. Rates for glial cell neoplasms and heart schwannomas from control groups of male Harlan Sprague Dawley rats from other recently completed NTP studies are presented in Appendix D of the 3-16-2016 draft report. While Harlan Sprague Dawley rats are an outbred strain, they are considered a single genetic strain in the same sense as other outbred strains, such as the Long-Evans or Wistar rat. Therefore, these historical control tumor rates are applicable to this study. However, it's important to note that the studies listed in Appendix D were carried out at laboratories other than the RFR studies, and under different housing and environmental conditions. At the time of the 3-16-2016 draft report, not all of these studies had undergone a complete pathology peer review. In the past several weeks NTP pathologists have reviewed brain and heart slides from these male rat control groups, and have confirmed, with few exceptions, the low rates of hyperplastic and neoplastic lesions reported in Appendix D, applying the diagnostic criteria established during the PWGs outlined in Appendix C.

NTP Comments on Statistical Issues Raised by the Reviewers

April 12, 2016

Given the multiple comparisons inherent in this kind of work, there is a high risk of false positive discoveries (Michael S. Lauer).

Although the NTP conducts statistical tests on multiple cancer endpoints in any given study, numerous authors have shown that the study-wide false positive rate does not greatly exceed 0.05 (Fears et al., 1977; Haseman, 1983; Office of Science and Technology Policy, 1985; Haseman, 1990; Haseman and Elwell, 1996; Lin and Rahman, 1998; Rahman and Lin, 2008; Kissling et al., 2014). One reason for this is that NTP's carcinogenicity decisions are not based solely on statistics and in many instances statistically significant findings are not concluded to be due to the test agent. Many factors go into this determination including whether there were pre-neoplastic lesions, whether there was a dose-response relationship, biological plausibility, background rates and variability of the tumor, etc. Additionally, with rare tumors especially, the actual false positive rate of each individual test is well below 0.05, due to the discrete nature of the data, so the cumulative false positive rate from many such tests is less than one person would expect by multiplying 0.05 by the number of tests conducted (Fears et al., 1977; Haseman, 1983; Kissling et al., 2015).

I'm getting slightly different values for poly-k adjusted denominators (Michael S. Lauer).

I compared poly---3 adjusted number from Table 3 in the original report versus the poly---3 adjusted number that I calculated using the raw data from the excel files. Supplementary Figure S1 shows that these two sets of numbers agree with each other in general. This is in contrast to the comparison for poly---6 adjusted number from Table 1 in the original report versus the poly---6 adjusted number that I calculated using the raw data from the excel files (Supplementary Figure S2). In fact, the adjusted rat numbers from Table 1 and Table 3 of the original report look quite similar (Supplementary Figure S3). This suggests that the poly---3 adjusted number was used in the footnotes in both Table 1 and Table 3 in the original report. (Max Lee)

I noted that in Table S2 the adjusted numbers in from.original.report and poly3 are identical at Dose 0 and 1.5 for both CDMA and GSM as well as at Dose 3 for GSM but differ slightly in the other treatment doses for heart schwannomas. One possible cause of the difference is that the version of the raw data in the excel files differs from that used to generate the original report. The second possibility is typo in the footnote in Table 3. I also generated Table S3 that has the poly---6 adjusted numbers for brain gliomas. The two sets of the poly---6 adjusted numbers are very different. (Max Lee)

Information could be included regarding the software or programming environment used for the computations. (Aleksandra M. Michalowski)

The adjusted denominators in Table 1 of the original report were labeled as poly-6 denominators, but were actually poly-3 denominators. This error was noted and brought to Dr Tabak's attention by Dr. Bucher in a March 22 email.

The p-values and adjusted denominators calculated by NTP are correct, except as noted for Table 1, and were calculated using validated poly-k software. This software is coded in Java and is embedded within NTP's TDMSE (Toxicology Data Management System Enterprise) system. Poly-k

calculations conducted by the reviewers in R may vary slightly from the NTP's calculation due to selection of study length and the NTP's use of the Bieler-Williams variance adjustment and a continuity correction. In his calculations, Dr. Lauer used 90 weeks as the study length, whereas the actual study length was 10 weeks. It is not apparent from the R documentation that the Bieler-Williams adjustment or the continuity correction is incorporated into the poly-3 calculations in R. In his calculations, Dr. Lee used two-sided p-values. In NTP statistical tests for carcinogenicity, the expectation is that if the test article is carcinogenic, tumor rates should increase with increasing exposure; thus, the NTP employs one-sided tests and p-values are one-sided. Using one-sided p-values in Dr. Lee's Table 1, the GSM trend if there were brain glioma in the control group remains nonsignificant, but the CDMA trend approaches 0.05 ($p = 0.054$) if there were brain glioma in the control group. In Dr. Lee's Table 2, the one-sided p-value for the GSM trend if there were 1 heart schwannoma in the control group approaches 0.05 ($p = 0.054$) and the one-sided p-value for the CDMA trend in heart schwannomas remains significant at $p = 0.018$ if there were 1 heart schwannoma in the control group. In Dr. Lee's Table 3, the one-sided p-value for the CDMA pairwise comparison is significant at $p = 0.049$ if there were 1 heart schwannoma in the control group.

statement of the required statistical significance level should be added. FDA guidance suggests the use of significance levels of 0.025 and 0.005 for tests for positive trends in incidence rates of rare tumors and common tumors, respectively; for testing pairwise differences in tumor incidence the use of significance levels of 0.05 and 0.01 is recommended for rare and common tumors, respectively. (Aleksandra M. Michalowski)

Although the FDA guidance suggests lowering the significance level for most tests of trend and pairwise differences, this guidance is based on a misunderstanding of findings reported by Haseman (1983). In this paper, Haseman discusses several rules proposed by others for setting the significance level lower than 0.05. *If* these rules are rigidly followed, Haseman showed that study conclusions will be consistent with the NTP's more complex decision-making process, for which 0.05 is the nominal significance level and p-values are taken into consideration along with other factors (outlined above in response to comment 1) in determining whether the tumor increase is biologically significant. The NTP does not strictly adhere to a specific statistical significance level in determining whether a carcinogenic effect is present.

Appendix tables for all poly-k tests performed could be added. (Aleksandra M. Michalowski)

Dr. Michalowski proposed a sample table. The rows corresponding to X, N, adjusted n are already included in the tables or appear the footnotes in the tables. The rows corresponding to "Dunnett contrast" and "Williams contrast" are not appropriate for dichotomous tumor data. Both Dunnett's test and Williams' test assume that the data are continuous and normally distributed.

In the portion of the text describing poly-k test results, p-values are given for significant pairwise comparisons; I would also give the p-values estimated for the significant trends. (Aleksandra M. Michalowski)

Indicators of significant trends are given in the tables in the form of asterisks next to control group tumor counts.

There are a couple of errors in the footnote of Table 3 in the original report. 2/74.05 (5%) should be 2/74.05 (2.7%). 3/78.67 (4%) should be 3/78.67 (3.8%). (Max Lee)

Thank you for pointing this out. The percentages will be corrected in our final report.

Were control rats selected in utero like the exposed rats were? Were pregnant dams assigned to different groups by formal randomization? How were the pups per litter chosen? (Michael S. Lauer).

believe detailed information about animal selection and randomization procedures should be given so that the potential for allocation bias could be judged. (Aleksandra M. Michalowski)

Pregnant dams were assigned to groups, including the control group, using formal randomization that sought to also equalize mean body weights across groups. The three pups per sex per litter were selected using formal randomization, as well. Tumors in the heart and brain were not observed in littermates, indicating that there was no litter-based bias in the results.

Were all analyses based on the intent-to-treat principle? Were there any crossovers? Were all rats accounted for by the end of the experiment and were all rats who started in the experiment included in the final analyses? (Michael S. Lauer)

The intent-to-treat principle is not relevant to this animal experiment, in which all animals that were assigned to treatment group received the full and equal treatment of that group. There were no crossovers. All animals that started the experiment were accounted for by the end of the experiment and included in the final analyses.

The PWG review blinding was not complete. (Michael S. Lauer)

PWG reviewers were blinded to the identity of the test article and the level of exposure but were not blinded to the fact that there were two different, yet related, test articles (modulations of cell phone RFR), to emphasize the fact that there was a common control group.

Did the authors perform a prospective sample size calculation? (Michael S. Lauer)

If power calculations to determine the required sample size were performed, the results should also be included. (Aleksandra M. Michalowski)

Sample size calculations were conducted for this study. However, for detecting carcinogenesis, sample size and power will depend on the baseline (control) tumor rate and the expected magnitude of the increase in tumors. For example, at 80% power, sample size requirements will be quite different for detecting a 2-fold increase in a rare tumor having a spontaneous occurrence of 0.5% compared to 2-fold increase in a more common tumor having a spontaneous occurrence of 10%. Because many different tumor types having wide range of spontaneous occurrence are involved in these studies, there is no “one-size-fits-all” sample size; rather, the sample size is a

compromise among several factors, including obtaining reasonable power to detect moderate to large increases for most tumor types, while staying within budgets of time, space, and funding. A sample of 90 animals per sex per group was selected as providing as much statistical power as possible across the spectrum of tumors, under the constraints imposed by the exposure system.

The NTP's carcinogenicity studies are similar in structure to the OECD's 45 Guideline for carcinogenicity studies and the FDA's guidance for rodent carcinogenicity studies of pharmaceuticals. These guidelines recommend at least 50 animals of each sex per group, but also mention that an increase in group size provides relatively little increase in statistical power. In the NTP's RFR studies, the group sizes were 90 animals of each sex per group, nearly twice as many as the minimum recommendation. Increasing the group sizes further provides diminishing returns, for which additional animals do not substantially increase power.

The low power implies that there is high risk of false positive findings (citing Ioannidis, 2005). ... suspect that this experiment is substantially underpowered and that the few positive results found reflect false positive findings (citing Ioannidis, 2005). (Michael S. Lauer)

It is true that the power is low for detecting moderate increases above a low background tumor rate of approximately 1%, as was seen in the brain and heart tumors. However, this low power does not correspond to high risk of false positive findings. The paper by Ioannidis that was cited correctly states that when studies are small or effect sizes are small (i.e., statistical power is low), "the less likely the research findings are to be true." Research findings can be "not true" if the result is a false positive or a false negative. With low statistical power, false negatives are much more likely than false positives. Therefore, the vast majority of false research findings in a low power situation will result from the failure to detect an effect when it exists. The false positive rate on any properly constructed statistical test will not exceed its significance level, alpha. By definition, the significance level of a statistical test is its false positive rate, and it is typically selected by the researcher, often at a low fixed value such as 0.05 or 5%.

If we were repeating the experiment, we may see some control studies have 1 or more tumors. (Max Lee) (Dr. Lee also presented analyses of the male rat data, inserting hypothetical data on one tumor-bearing animal in the control group.)

In light of the historical control data, Dr. Lee demonstrated that several associations became less or not significant with the insertion of a tumor data point in the control group. While we appreciate that some other studies had one or more tumors, the NTP considers the concurrent control group as the most important comparator to the treated groups. We took the historical control tumor rates into account in a more subjective manner in our interpretation of the findings. In 2010, we asked to adopt a more formal method of incorporating historical control data in our statistical testing, but our Board of Scientific Counselors voted against adopting the method.

It is puzzling why the control had short survival rate. Given that most of the gliomas and heart schwannomas are late-developing tumors, it is possible that if the controls were living longer some tumors might develop. Although the use of poly-3 (or poly-6) test intended to adjust the number of rats

used in the study, it is still important to re-evaluate the analysis by considering the incidence rate in controls not being 0. (Max Lee)

We do not know why the male rat control group had a low survival rate. We generally do observe lower survival rates in studies such as the RFR studies in which animals are singly- rather than group housed. While some tumors might possibly have arisen in controls if they lived longer, it was notable that no glial cell or Schwann cell hyperplasias were found in these animals as well.

The poly-k (e.g., poly-3 or poly-6) test was developed to adjust for the fact that not all animals survive to the end of a two-year study, and survival rates may differ among groups. The test is essentially a Cochran-Armitage trend test in which the denominator of the tumor rate in each group is adjusted downward to better reflect the number of animal-years at risk during the study. Each animal that develops the tumor or survives to the end of the study is counted as one animal. Each animal that does not develop the tumor and dies (or is moribund sacrificed) before the end of the study is counted as a fractional animal. The fraction is calculated as the proportion of the study that it survived, raised to the k-th power; $k = 3$ or $k = 6$ in this study. The survival-adjusted tumor rate in each group is then the number of animals having the tumor of interest divided by the total count of animals at risk of developing the tumor in the group. These survival-adjusted rates are used in the Cochran-Armitage formula to provide the poly-k test for dose-related trends and pairwise comparisons with the control group.

The poly-k test has been shown to yield valid inferences about tumor rates in NTP two-year rat and mouse carcinogenicity studies (Bailer and Portier, 1988; Portier and Bailer, 1989; Portier et al., 1986). Its theoretical basis is that tumor incidence, while not directly observed unless the tumor is immediately lethal, follows a Weibull distribution with a shape parameter, k . Verification using NTP studies has shown that if k is between 1 and 5, setting $k = 3$ yields a valid statistical test (Portier and Bailer, 1989; Portier et al., 1986). Thus, most of the time, the NTP uses the poly-3 test. If tumor type is late-occurring, as we observed with the brain gliomas, $k = 6$ is a better fit to the data and the poly-6 test has more validity.

In the portion of the text describing differences in survival at the end of the study between control and RFR-exposed animals the compared characteristic is not named and also no numerical values of the estimates or the range of differences are given. I would add numbers in the text of a Appendix table showing the group survival estimates described in this paragraph. (Aleksandra M. Michalowski)

The Statistical Methods section describes the method for comparing survival distributions between the control and RFR-exposed groups, namely, Tarone's (1975) life table test to identify exposure-related trends in survival and Cox's (1972) method for testing two groups for equality of survival distributions.

References

- Bailer, A.J., and Portier, C.J. (1988). Effects of treatment-induced mortality and tumor-induced mortality on tests for carcinogenicity in small samples. *Biometrics* 44, 417-431.
- Bieler, G.S., and Williams, R.L. (1993). Ratio estimates, the delta method, and quantal response tests for increased carcinogenicity. *Biometrics* 49, 793-801.
- Cox, D.R. (1972). Regression models and life-tables. *Journal of the Royal Statistical Society B* 34, 187-220.
- Dunnett, C. W. (1955). A multiple comparison procedure for comparing several treatments with control. *Journal of the American Statistical Association* 50, 1096-1121.
- Fears, T. R., R. E. Tarone, et al. (1977). "False-positive and false-negative rates for carcinogenicity screens." *Cancer Research* 37(7 Pt 1): 1941-1945.
- Haseman, J. K. (1983). A reexamination of false-positive rates for carcinogenesis studies. *Fundamental and Applied Toxicology* 3(4): 334-339.
- Haseman, J. K. (1990). Use of statistical decision rules for evaluating laboratory animal carcinogenicity studies. *Fundamental and Applied Toxicology* 14(4): 637-648.
- Haseman, J. K. and M. R. Elwell (1996). Evaluation of false positive and false negative outcomes in NTP long-term rodent carcinogenicity studies. *Risk Analysis* 16(6): 813-820.
- Ioannidis, J.P.A. (2005). Why most published research findings are false. *PLoS Medicine* 2(8):e124.
- Kissling, G.E., Haseman, J.K., Zeiger, E. (2015). Proper interpretation of chronic toxicity studies and their statistics: A critique of "Which level of evidence does the US National Toxicology Program provide? Statistical considerations using the Technical Report 57 on *Ginkgo biloba* as an example." *Toxicology Letters* 237: 161-164.
- Lin, K. K. and M. A. Rahman (1998). Overall false positive rates in tests for linear trend in tumor incidence in animal carcinogenicity studies of new drugs. *Journal of Biopharmaceutical Statistics* 8(1): 1-15; discussion 17-22.
- Office of Science and Technology Policy (1985). Chemical carcinogens: A review of the science and its associated principles. *Federal Register* 10:10371-10442.
- Piegorsch, W.W., and Bailer, A.J. (1997). *Statistics for Environmental Biology and Toxicology*, Section 6.3.2. Chapman and Hall, London.
- Portier, C.J. (1986) Estimating the tumour onset distribution in animal carcinogenesis experiments. *Biometrika* 72, 371-378.
- Portier, C.J., Hedges, J.C., Hoel, D.G. (1986) Age-specific models of mortality and tumor onset for historical control animals in the National Toxicology Program's carcinogenicity experiments. *Cancer Research* 46, 4372-4378.

Portier, C.J., and Bailer, A.J. (1989). Testing for increased carcinogenicity using a survivaladjusted quantal response test. *Fundamental and Applied Toxicology* 12, 731-737.

Rahman, M. A. and K. K. Lin (2008). comparison of false positive rates of Peto and poly-3 methods for long-term carcinogenicity data analysis using multiple comparison adjustment method suggested by Lin and Rahman. *Journal of Biopharmaceutical Statistics* 18(5): 949-958.

Tarone, R.E. (1975). Tests for trend in life table analysis. *Biometrika* 62:679-682.

Thomas, D.G., Breslow, N., Gart, J.J. (1977). Trend and homogeneity analyses of proportions and life table data. *Computers and Biomedical Research* 10, 373-381.

ADDITIONAL RESPONSE:

Dear All,

Thanks again for all your helpful comments on the NTP RFR studies. I did want to follow up on one remaining point of disagreement that Mike Lauer alluded to in his comments about low powered studies. Although we agree that our study design had low power to detect statistically significant neoplastic effects in the brain and heart, which occurred with both RFR modulations in male rats, we disagree over the assertion that low power in and of itself, creates false positive results. We cited a handful of publications outlining the statistical arguments against this with specific respect to the NTP rodent cancer study design in our response to comments document sent earlier. Although Mike referred to the example of positive findings in underpowered epidemiology studies that could not be replicated in larger follow up studies, there is a growing literature alluding to this problem with respect to experimental animal studies as well. An example is a relatively recent article by one of our collaborators in CAMARADES, Malcolm MacLeod.

<http://www.nature.com/news/2011/110928/full/477511a.html>

It's important to distinguish between low power to detect effects, and the constellation of other factors that often accompany low powered experimental animal studies in contributing to this problem. We've addressed this issue in a recent editorial, and these factors are captured in our published systematic review process for evaluating study quality in environmental health sciences (Rooney et al., 2014).

<http://ehp.niehs.nih.gov/wp-content/uploads/122/7/ehp.1408671.pdf>

<http://ehp.niehs.nih.gov/wp-content/uploads/122/7/ehp.1307972.pdf>

Table 1 in the Rooney et al. report outlines risk of bias considerations that commonly plague studies carried out by academic researchers that are accounted for in NTP studies.

I provide these examples to assure you that we are completely cognizant of these issues and take them very seriously. Again, we appreciate the help you've provided in assuring that we appropriately interpret and communicate our findings.

Best
John Bucher

SPIN vs FACT: *National Toxicology Program* report on cancer risk from cellphone radiation

The National Toxicology Program (NTP) of the National Institutes of Health [reported](#) partial findings from their \$25 million study of the cancer risk from cellphone radiofrequency radiation (RFR). Controlled studies of rats showed that RFR caused two types of tumors, glioma and schwannoma. The results "...could have broad implications for public health." Below are some biased statements, or "Spin," about the study that tend to create doubt about data quality and implications, as well as "Facts" from decades of previous research.

SPIN	FACT
Conclusions are faulty. Dr. Michael Lauer, deputy director for extramural research at the National Institutes of Health, "I am unable to accept the authors' conclusions."	The NTP is world-renowned for toxicology research. This is "by far ... the most carefully done cell phone" toxicology study of RFR carcinogenic effects. Criticisms by Dr. Lauer and other scientists who reviewed the study were rebutted in the study report .
Study reports a "low incidence" of tumors in the brain and heart in rats exposed to RFR.	The study found that one in twelve (8.5%) of the 540 male rats exposed to cellphone radiation developed cancer or pre-cancerous cells as compared to none of the 90 rats in the control condition.
Relevance of animal studies to humans is questionable.	The cells that developed tumors are the same cells that display elevated tumor risk in studies of long-term, heavy cellphone users . Rats are the preferred animal model for carcinogenicity studies.
International Agency for Research on Cancer (IARC) , rated cellphone radiation a "possible" human carcinogen (Group 2B), the same rating given to coffee, pickled vegetables, and talc.	The report provides strong evidence that RFR exposure causes cancer. Major studies published since the 2011 IARC meeting consistently find that long-term, heavy cellphone users have increased risk of brain tumors. Group 2B carcinogens also include DDT, lead, and diesel fumes.
Prior research contradicts NTP study results (e.g., Danish Cohort Study , British Million Women Study).	The Danish study has been criticized by many scientists for excluding heavy cellphone users. The British Study has also been criticized; but, it found evidence for acoustic neuroma (a form of schwannoma).
Epidemiological studies fail to show an increase in brain tumor incidence since 1992 even though cellphone use has mushroomed.	The incidence of nonmalignant tumors has significantly increased in the U.S. since cellphones. Moreover, the incidence of glioblastoma multiforme , the most serious type of brain cancer, has increased in parts of the brain proximal to where cellphones are held . Brain cancer can take decades to develop, so it is premature to see overall increases in malignant tumors in the general population.
There is no mechanism to explain how cellphones could cause cancer. Unlike ionizing radiation, <i>non-ionizing</i> radiation from cellphones cannot damage DNA.	A review paper reported that in 93 of 100 studies RFR produced a cellular stress response which can lead to DNA damage and cancer. The NTP study also found evidence of DNA damage . Several published papers present evidence for different mechanisms by which RFR may cause cancer.
The research has not been peer-reviewed.	The NTP report has been peer-reviewed by experts. Some reviews appear in the report along with the authors' responses.
Findings are preliminary, it is premature to conclude we should take precautions or change policy.	These are not preliminary findings. According to NTP, the effects of RFR on these two tumors, glioma and schwannoma, are final. The federal government released this partial report because the results "could have broad implications" for the public due to widespread cellphone use . The NTP posted on its website a link to the FDA's recommendations on how to reduce cellphone radiation exposure.

Apple says:

“Read all safety information below and operating instructions before using iPad to **avoid injury.**”

iPad Operating Manual

“iPad contains radio transmitters and receivers. When on, iPad sends and receives radio frequency (RF) energy through its antenna. The Wi-Fi and Bluetooth® antennas are located behind the screen to the left of the Home button, and behind the Apple logo... A cellular antenna is located at the top edge of iPad Wi-Fi+3G, when oriented with the Home button at the bottom.”

“... to be sure that human exposure to RF energy does not exceed the FCC, IC, and European Union guidelines, always follow these instructions and precautions: Orient the device in portrait mode with the Home button at the bottom of the display, or in landscape mode with the cellular antenna (located under the black edge at the top of the device) away from your body or other objects...”

“you can further limit your exposure by limiting the amount of time using iPad Wi-Fi+3G in wireless mode, since time is a factor in how much exposure a person receives, and by placing more distance between your body and iPad Wi-Fi + 3G, since exposure level drops off dramatically with distance.”

Seizures, Blackouts, and Eyestrain

A small percentage of people may be susceptible to blackouts or seizures (even if they have never had one before) when exposed to flashing lights or light patterns such as when playing games or watching video. If you have experienced seizures or blackouts or have a family history of such occurrences, you should consult a physician before playing games or watching videos on your iPad. Discontinue use of iPad and consult a physician if you experience headaches, blackouts, seizures, convulsion, eye or muscle twitching, loss of awareness, involuntary movement, or disorientation. To reduce risk of headaches, blackouts, seizures, and eyestrain, avoid prolonged use, hold iPad some distance from your eyes, use iPad in a well-lit room, and take frequent breaks.

Read more:

<http://support.apple.com/manuals/#ipad>

iPad Operating Manual

Do you know where the WiFi/Bluetooth Antennas are on your child's iPad?



WHAT YOU DON'T SEE



Parents for Safe Schools

Parents for Safe Schools

More Information:

<https://www.facebook.com/pages/Parents-for-Safe-Schools/428808610553840>

www.citizensforsafetechnology.org

www.safeinschool.org

Email: Parentsforsafeschools@gmail.com



Did you know?

- When ifixit.com dissected a Wi-Fi+3G iPad, they discovered **five antennas**, including the frame of the LCD screen that is one giant antenna.
- When an iPad's WiFi is turned on, it emits a burst of microwave radiation approximately every four seconds. That makes 900 blasts per hour in the child's hands, on their lap, or at their face and does not include any additional data signals resulting from uploading and downloading activities. The beacon signal from the iPad occurs every 4 seconds even when the user is not accessing the Internet. In other words, even if a child is only drawing or playing games, they will still receive 900 bursts of microwave radiation an hour as long as the WiFi antenna is left 'On.'
- The user manual recommends: "you can further limit your exposure by limiting the amount of time using iPad Wi-Fi+3G in wireless mode, since time is a factor in how much exposure a person receives, and by placing more distance between your body and iPad Wi-Fi + 3G, since exposure level drops off dramatically with distance."
- In the user manuals for the 2nd generation and newer models of iPad, Apple misinforms users that the WiFi and Bluetooth antennas are at the back of the iPad behind the Apple logo. That is inaccurate. The WiFi and Bluetooth antennas are actually at the bottom of the iPad next to the HOME button. With this fact being undisclosed, users (children) will be resting the iPad WiFi-Bluetooth antennas directly on their laps and bodies without being warned of the risks.
- iPad WiFi Radiation with Wireless Router On - Video
<http://bit.ly/1j6jA6y>



What can you do to reduce the RF radiation exposure from your child's iPad?

- 1** Never let your child hold an iPad in portrait mode on his/her lap or near his chest, where microwave radiation is penetrating through internal organs.
- 2** Never let your child hold an iPad in landscape mode with his/her hands covering the constantly transmitting antennas and absorbing microwave radiation directly into their bodies.
- 3** Never put an iPad under your child's pillow or close to his body while he is sleeping.
- 4** Never give an iPad to a child unless the iPad's airplane mode is turned on AND the Wi-Fi and Bluetooth antennas are turned off.



Note

Putting an iPad on 'airplane mode' will initially turn off all antennas; however, WiFi and Bluetooth antennas can be **re-activated** respectively without switching off the airplane mode. Therefore, an iPad showing 'airplane mode' is not a guarantee that all the antennas are off. It is important to check the antennas separately to ensure they are ALL turned 'Off'. In addition, when the WiFi function on an iPad shows "Not Connected," it does not mean the antenna is off, it only means the iPad is not logged onto any available network. The WiFi antenna is still sending out microwave radiation.

Dec 8 (4 days ago)

to me

----- Forwarded message -----

From: **Amelia McGhee** <acmcghee@verizon.net>

Date: Sat, Sep 10, 2016 at 10:45 PM

Subject: Please recommend a prohibition of cell towers on school grounds

To: ematsui@jhmi.edu, [rachel.hess-](mailto:rachel.hess-mutinda@maryland.gov)

mutinda@maryland.gov, emily@marylandpirg.org, guy.guzzone@senate.state.md.us, amezu@msde.state.md.us, allison@mdehn.org, Angela.Angel@house.state.md.us, benoythomas@gmail.com, brandi.stocksdales@maryland.gov, [Varney-](mailto:Varney-Alvarado@dhcd.state.md.us)

Alvarado@dhcd.state.md.us, christina.church@maryland.gov, dbishai@jhsph.edu, dianna.abney@maryland.gov, gdielte@jhmi.edu, lornegarr@msn.com, bgitterman@aol.com, jicarella@msn.com, julianlevy@comcast.net, megan.latschaw@aphl.org, cliff.mitchell@maryland.gov, nancy.servatius@maryland.gov, nobot@cehn.org, rob.hofstetter@maryland.gov

Dear Maryland Children's Environmental Health and Protection Advisory Council Members,

Schools should be safe environments for children. Currently schools across the state are placing cell towers on school grounds unaware of the many health and safety risks they pose. Cell towers emit radio-frequency radiation and children will be exposed long term to this radiation when cell towers are installed on school grounds. In addition, the tower installation includes diesel tanks and lead acid batteries, and thus is considered a HAZMAT area.

Please make a strong recommendation to the State Department of Education that schools PROHIBIT cell towers on school grounds.

Please make a strong recommendation to the General Assembly that schools PROHIBIT cell towers on school grounds.

Please make a strong recommendation to the General Assembly that the Department of Health be allocated funds to do research on electromagnetic radiation and children's health.

[See 25 Reasons Cell Towers Have No Business On or Near School Property by The Maryland Coalition To Halt Cell Towers at Schools](#) for a detailed list of health and safety concerns. <http://www.safeschoolspg.org/joint-statement-of-the-maryland-coalition-to-halt-cell-towers-at-schools.html> and this recent Washington Post article which highlights the need for tougher standards nationwide. <http://www.washingtonpost.com/wp-dyn/articles/A35421-2004Dec29.html>

Many thanks,

Amelia McGhee

**Protect
the ones
you love.**♥



American Academy of Pediatrics Recommends:

CELL PHONE SAFETY TIPS FOR FAMILIES

- Make only short or essential calls on cell phones.
- Use cell phones in speaker mode or with the use of hands-free kits.
- Avoid carrying your phone against the body like in a pocket, sock, or bra. Cell phone manufacturers can't guarantee that the amount of radiation you're absorbing will be at a safe level.
- If you plan to watch a movie on your device, download it first, then switch to airplane mode while you watch in order to avoid unnecessary radiation exposure.
- Keep an eye on your signal strength (i.e. how many bars you have). The weaker your signal, the harder your phone has to work and the more radiation it gives off.
- Avoid making calls in cars, elevators, trains, and buses. The cell phone works harder (and emits more radiation) to get a signal through metal.
- Remember that cell phones are not toys or teething items.

GOOD TO KNOW:

Wireless devices (such as **tablets, laptops, baby monitors, video game consoles and Wi-Fi routers**) emit the same type of radiation as cell phones.

Airplane mode = Wireless Off.

AT HOME:

Easily reduce wireless radiation at home. Get a **corded landline phone** and forward cell phones to it.

Use corded plug-in (non-wireless) connections for printers, mouse, computers, laptops, tablets, e-readers, routers, keyboards, headphones, speakers, and other accessories—with all wireless functions off.

MORE AT **EHTrust.org**
#PracticeSafeTech
@saferphones





AAP Headquarters

141 Northwest Point Blvd
Elk Grove Village, IL 60007-1019
Phone: 847/434-4000
Fax: 847/434-8000
E-mail: kidsdocs@aap.org
www.aap.org

Reply to

Department of Federal Affairs

Homer Building, Suite 400 N
601 13th St NW
Washington, DC 20005
Phone: 202/347-8600
Fax: 202/393-6137
E-mail: kids1st@aap.org

Executive Committee

President

Thomas K. McInerney, MD, FAAP

President-Elect

James M. Perrin, MD, FAAP

Immediate Past President

Robert W. Block, MD, FAAP

Executive Director/CEO

Errol R. Alden, MD, FAAP

Board of Directors

District I

Carole E. Allen, MD, FAAP
Arlington, MA

District II

Danielle Laraque, MD, FAAP
Brooklyn, NY

District III

David I. Bromberg, MD, FAAP
Frederick, MD

District IV

Francis E. Rushton, Jr, MD, FAAP
Beaufort, SC

District V

Marilyn J. Bull, MD, FAAP
Indianapolis, IN

District VI

Pamela K. Shaw, MD, FAAP
Kansas City, KS

District VII

Kenneth E. Matthews, MD, FAAP
College Station, TX

District VIII

Kyle Yasuda, MD, FAAP
Seattle, WA

District IX

Stuart A. Cohen, MD, MPH, FAAP
San Diego, CA

District X

Sara H. Goza, MD, FAAP
Fayetteville, GA

August 29, 2013

The Honorable Mignon L. Clyburn
Acting Commissioner
Federal Communications Commission
445 12th Street SW
Washington, DC 20054

The Honorable Dr. Margaret A. Hamburg
Commissioner
U.S. Food and Drug Administration
10903 New Hampshire Avenue
Silver Spring, MD 20993

Dear Acting Chairwoman Clyburn and Commissioner Hamburg:

The American Academy of Pediatrics (AAP), a non-profit professional organization of 60,000 primary care pediatricians, pediatric medical sub-specialists, and pediatric surgical specialists dedicated to the health, safety and well-being of infants, children, adolescents, and young adults appreciates this opportunity to comment on the Proposed Rule "Reassessment of Exposure to Radiofrequency Electromagnetic Fields Limits and Policies" published in the Federal Register on June 4, 2013.

In the past few years, a number of American and international health and scientific bodies have contributed to the debate over cell phone radiation and its possible link to cancer. The International Agency for Research on Cancer (IARC), part of the United Nations' World Health Organization, said in June 2011 that a family of frequencies that includes mobile-phone emissions is "possibly carcinogenic to humans." The National Cancer Institute has stated that although studies have not demonstrated that RF energy from cell phones definitively causes cancer, more research is needed because cell phone technology and cell phone use are changing rapidly. These studies and others clearly demonstrate the need for further research into this area and highlight the importance of reassessing current policy to determine if it is adequately protective of human health.

As radiation standards are reassessed, the AAP urges the FCC to adopt radiation standards that:

- **Protect children's health and well-being.** Children are not little adults and are disproportionately impacted by all environmental exposures, including cell phone radiation. Current FCC standards do not account for the unique vulnerability and use patterns specific to pregnant women and children. It is essential that any new standard for cell phones or other wireless devices be based on

protecting the youngest and most vulnerable populations to ensure they are safeguarded throughout their lifetimes.

- **Reflect current use patterns.** The FCC has not assessed the standard for cell phone radiation since 1996. Approximately 44 million people had mobile phones when the standard was set; today, there are more than 300 million mobile phones in use in the United States. While the prevalence of wireless phones and other devices has skyrocketed, the behaviors around cell phone uses have changed as well. The number of mobile phone calls per day, the length of each call, and the amount of time people use mobile phones has increased, while cell phone and wireless technology has undergone substantial changes. Many children, adolescents and young adults, now use cell phones as their only phone line and they begin using wireless phones at much younger ages. Pregnant women may carry their phones for many hours per day in a pocket that keeps the phone close to their uterus. Children born today will experience a longer period of exposure to radio-frequency fields from cellular phone use than will adults, because they start using cellular phones at earlier ages and will have longer lifetime exposures. FCC regulations should reflect how people are using their phones today.
- **Provide meaningful consumer disclosure.** The FCC has noted that it does not provide consumers with sufficient information about the RF exposure profile of individual phones to allow consumers to make informed purchasing decisions. The current metric of RF exposure available to consumers, the Specific Absorption Rate, is not an accurate predictor of actual exposure. AAP is supportive of FCC developing standards that provide consumers with the information they need to make informed choices in selecting mobile phone purchases, and to help parents to better understand any potential risks for their children. To that end, we support the use of metrics that are specific to the exposure children will experience.

The AAP supports the reassessment of radiation standards for cell phones and other wireless products and the adoption of standards that are protective of children and reflect current use patterns. If you have questions, please contact Clara Filice in the AAP's Washington Office at 202/347-8600.

Sincerely,

A handwritten signature in cursive script that reads "Thomas K. McInerney".

Thomas K. McInerney, MD FAAP
President

TKM/cf

**AAP Headquarters**

141 Northwest Point Blvd
Elk Grove Village, IL 60007-1019
Phone: 847/434-4000
Fax: 847/434-8000
E-mail: kidsdocs@aap.org
www.aap.org

Reply to**Department of Federal Affairs**

Homer Building, Suite 400 N
601 13th St NW
Washington, DC 20005
Phone: 202/347-8600
Fax: 202/393-6137
E-mail: kids1st@aap.org

Executive Committee**President**

Thomas K. McInerney, MD, FAAP

President-Elect

James M. Perrin, MD, FAAP

Immediate Past President

Robert W. Block, MD, FAAP

Executive Director/CEO

Errol R. Alden, MD, FAAP

Board of Directors**District I**

Carole E. Allen, MD, FAAP
Arlington, MA

District II

Danielle Laraque, MD, FAAP
Brooklyn, NY

District III

David I. Bromberg, MD, FAAP
Frederick, MD

District IV

Francis E. Rushton, Jr, MD, FAAP
Beaufort, SC

District V

Marilyn J. Bull, MD, FAAP
Indianapolis, IN

District VI

Pamela K. Shaw, MD, FAAP
Kansas City, KS

District VII

Kenneth E. Matthews, MD, FAAP
College Station, TX

District VIII

Kyle Yasuda, MD, FAAP
Seattle, WA

District IX

Stuart A. Cohen, MD, MPH, FAAP
San Diego, CA

District X

Sara H. Goza, MD, FAAP
Fayetteville, GA

December 12, 2012

The Honorable Dennis Kucinich
2445 Rayburn House Office Building
Washington, DC 20515

Dear Representative Kucinich:

On behalf of the American Academy of Pediatrics (AAP), a non-profit professional organization of 60,000 primary care pediatricians, pediatric medical subspecialists, and pediatric surgical specialists dedicated to the health, safety and well-being of infants, children, adolescents, and young adults, I would like to share our support of H.R. 6358, the *Cell Phone Right to Know Act*.

The AAP strongly supports H.R. 6358's emphasis on examining the effects of radiofrequency (RF) energy on vulnerable populations, including children and pregnant women. In addition, we are pleased that the bill would require the consideration of those effects when developing maximum exposure standards. Children are disproportionately affected by environmental exposures, including cell phone radiation. The differences in bone density and the amount of fluid in a child's brain compared to an adult's brain could allow children to absorb greater quantities of RF energy deeper into their brains than adults. It is essential that any new standards for cell phones or other wireless devices be based on protecting the youngest and most vulnerable populations to ensure they are safeguarded throughout their lifetimes.

In addition, the AAP supports the product labeling requirements in H.R. 6358. These standards will ensure consumers can make informed choices in selecting mobile phone purchases. They will also enable parents to better understand the potential dangers of RF energy exposure and protect their children.

On July 24, the U.S. Government Accountability Office (GAO) published a report on federal cell phone radiation exposure limits and testing requirements. The GAO noted that the Federal Communications Commission's (FCC) most recent data indicates that the number of estimated mobile phone subscribers has grown from approximately 3.5 million in 1989 to approximately 289 million at the end of 2009. Cell phone use behaviors have also changed during that time. The quantity and duration of cell phone calls has increased, as has the amount of time people use mobile phones, while cell phone and wireless technology has undergone substantial changes. Many more people, especially adolescents and young adults, now use cell phones as their only phone line, and they begin using wireless phones at much younger ages.

Despite these dramatic changes in mobile phone technology and behavior, the FCC has not revisited the standard for cell phone radiation exposure since 1996. The current FCC standard for maximum radiation exposure levels is based on the heat emitted by mobile phones. These guidelines specify exposure limits for hand-held wireless devices in terms of the Specific Absorption Rate (SAR), which measures the rate the body absorbs radiofrequency (RF). The current allowable SAR limit is 1.6 watts per kilogram (W/kg), as averaged over one gram of tissue. Although wireless devices sold in the United States must ensure that they do not exceed the maximum allowable SAR limit when operating at the device's highest possible power level, concerns have been raised that long-term RF energy exposure at this level affects the brain and other tissues and may be connected to types of brain cancer, including glioma and meningioma.

In May 2011, the International Agency for Research on Cancer (IARC), the United Nations' World Health Organization's (WHO) agency promoting international cancer research collaboration, classified RF energy as "possibly carcinogenic to humans." In addition, the National Cancer Institute has stated that although studies have not definitively linked RF energy exposure from cell phones to cancer, more research is required to address rapidly changing cell phone technology and use patterns.

This and other research identified by the GAO demonstrates the need for further research on this issue, and makes clear that exposure standards should be reexamined.

The GAO concluded that the current exposure limits may not reflect the latest research on RF energy, and that current mobile phone testing requirements may not identify maximum RF energy exposure. The GAO proposed that the FCC formally reassess its limit and testing requirements to determine whether they are effective. The AAP commends the activities proposed under H.R. 6358, as they would address this research gap and improve consumer knowledge and safety. Establishing an expanded federal research program as the basis for exposure standards will ensure that consumer protections incorporate the latest research. Currently, the National Institute of Health (NIH), the only federal agency the GAO identified as directly funding research on this topic, provided approximately \$35 million from 2001 to 2011. Given this previous funding level, the AAP supports the \$50 million per fiscal year for seven years that H.R. 6358 would authorize.

The AAP appreciates your recognition of the need for new research and standards for mobile phone radiation, and is pleased to support H.R. 6358. For further assistance, please do not hesitate to contact Sonya Clay, Assistant Director, Department of Federal Affairs, at 202-347-8600 or sclay@aap.org.

Sincerely,

A handwritten signature in dark ink, reading "Thomas K. McInerney". The signature is written in a cursive, flowing style.

Thomas K. McInerney, MD, FAAP
President



INTERNET ARCHIVE
waybackmachine

2 captures
25 Oct 16 – 1 Dec 16

http://www.aappublications.org/news/2016/05/27/Cancer052716 Go

SEP OCT DEC Close
25
2015 2016 2017 Help

AAP responds to study showing link between cell phone radiation, tumors in rats

Melissa Jenco, News content editor

Some rats developed tumors after being exposed to cell phone radiation, according to preliminary results of a study released Thursday.

In light of the findings, the Academy continues to reinforce its recommendation that parents should limit use of cell phones by children and teens.

“They’re not toys. They have radiation that is emitted from them and the more we can keep it off the body and use (the phone) in other ways, it will be safer,”



said Jennifer A. Lowry, M.D., FAACT, FAAP, chair of the AAP Council on Environmental Health Executive Committee.

The [study by the National Toxicology Program](#), part of the National Institute of Environmental Health Sciences, exposed rats to radiofrequency radiation for nine hours a day for two years beginning in utero and compared them to rats that were not exposed. Some of the male rats developed malignant tumors in their hearts and brains while the control group did not, according to the report, which included only partial findings.

Dr. Lowry, chief for the Section of Clinical Toxicology at Children's Mercy Hospital, said it is difficult to translate the results in rats to humans, and exposure was heavier than it would be for most people.

In *Pediatric Environmental Health, 3rd Edition*, the Academy recommends “exposures can be reduced by encouraging children to use text messaging when possible, make only short and essential calls on cellular phones, use hands free kits and wired headsets and maintain the cellular phone an inch or more away from the head.” The book also warns against talking on the phone or texting while driving.

Copyright © 2016 American Academy of Pediatrics

We recommend

Study: 3% of young children have high blood lead levels
Melissa Jenco et al., AAP News , 2016

AAP Conference Preview: Environmental health expert to tackle tough questions
Carla Kemp et al., AAP News , 2016

NCE Preview: Environmental health expert to tackle tough questions
Carla Kemp et al., AAP News , 2016

Pesticides and Kids: An AAP Report
James Roberts, MD, MPH, et al., Medscape , 2013

AAP: No Amount of Lead Exposure Is Safe for Children
PracticeUpdate , 2016

Cell Phones and Cancer: Is There a Connection?
Bret Stetka, MD, et al., Medscape , 2011

Keeping Children Home Is Often Not Justified



Healthy Children (/English) > Safety & Prevention (<https://www.healthychildren.org/English/safety-prevention>) > All Around (<https://www.healthychildren.org/English/safety-prevention/all-around>) > Cell Phone Radiation & Children's Health: What Parents Need to Know

SAFETY & PREVENTION

Cell Phone Radiation & Children's Health: What Parents Need to Know

Children are not just little adults; their growing minds and bodies make them uniquely vulnerable to the effects of the environment around them, including cell phone radiation. Because technology is being adopted by children at younger ages than ever before, it's even more important to investigate if cell phone usage is a health hazard.



What is cell phone radiation, anyway?

There are two types of radiation: ionizing and non-ionizing.

- **Ionizing radiation** (e.g., x-rays, radon, sunlight) is high frequency (and high energy).
- **Non-ionizing** is low frequency (low energy) radiation.

Cell phones have non-ionizing radiation. Your phone sends radio frequency waves from its antenna to nearby cell towers. When you make a call, text, or use data, your phone receives radio frequency waves to its antenna from cell towers.

What does the latest research say?

Several studies have been done to find out if cell phone use can lead to cancer. These types of studies in people have not shown clear evidence of an increased cancer risk with cell phone use. While there was a slight increase in a type of brain tumor, called a glioma, in a small group of people who spent the most total time on cell phone calls in one study, other studies have not found this to be true.

In May 2016, the US National Toxicology Program, which is part of the National Institutes of Health (NIH), released partial findings from a two-year study (<http://biorxiv.org/content/early/2016/05/26/055699>) that exposed rats to the types of radio frequency radiation that cell phones give off and compared them with a non-exposed group. Some rats developed cancerous tumors after being exposed to the radiation—showing a potential connection between exposure to radiation and an increased risk of cancer.

A few words of caution about this study:

- This study was only done on rats. While rats can be good test subjects for medical research, they are not the same as humans. We do not yet know if the same results would occur in people.
- The rats were exposed to very large amounts of radiation—nine hours a day, seven days a week, for two years. This is far more than most people spend holding their cell phones.
- More male rats developed cancerous tumors after being exposed to the radiation than female rats. Some of the rats who developed tumors lived longer than the control group rats that were not exposed to radiation.
- The analysis of all of the data from this study is not yet complete.

Why is more research needed?

Parents should not panic over the latest research, but it can be used as a good reminder to limit both children's screen time (/English/family-life/Media/Pages/Media-Time-Family-Pledge.aspx) and exposure from cell phones and other devices emitting radiation from electromagnetic fields (EMF) (/English/safety-prevention/all-around/Pages/Electromagnetic-Fields-A-Hazard-to-Your-Health.aspx). Partial findings from studies like this one give scientists reason to look into the issue more. **The American Academy of Pediatrics (AAP) supports more research into how cell phone exposure affects human health long term, particularly children's health.**

How can we limit cell phone radiation for ourselves and our children?

The AAP reinforces its existing recommendations on limiting cell phone use for children and teenagers. The AAP also reminds parents that cell phones are not toys, and are not recommended for infants and toddlers (/English/family-life/Media/Pages/Tablets-and-Smartphones-Not-for-Babies.aspx) to play with.

Cell phone safety tips for families:

- Use text messaging when possible, and use cell phones in speaker mode or with the use of hands-free kits.
- When talking on the cell phone, try holding it an inch or more away from your head.
- Make only short or essential calls on cell phones.
- Avoid carrying your phone against the body like in a pocket, sock, or bra. Cell phone manufacturers can't guarantee that the amount of radiation you're absorbing will be at a safe level.
- Do not talk on the phone or text while driving (/English/ages-stages/teen/safety/Pages/Sample-Driving-Rules-Teens-Must-Follow.aspx). This increases the risk of automobile crashes.
- Exercise caution when using a phone or texting while walking or performing other activities. "Distracted walking" injuries are also on the rise.
- If you plan to watch a movie on your device, download it first, then switch to airplane mode while you watch in order to avoid unnecessary radiation exposure.
- Keep an eye on your signal strength (i.e. how many bars you have). The weaker your cell signal, the harder your phone has to work and the more radiation it gives off. It's better to wait until you have a stronger signal before using your device.
- Avoid making calls in cars, elevators, trains, and buses. The cell phone works harder to get a signal through metal, so the power level increases.
- Remember that cell phones are not toys or teething items.

Are there any regulations in place to limit cell phone radiation in the United States?

The Federal Communications Commission (FCC) decides how much radiation cell phones are allowed to give off in the US. Currently, the FCC limit is at 1.6 W/Kg. The FCC, however, has not revised the standard for cell phone radiation since 1996, and a lot has changed since then.

- There are now more cell phones in the United States than there are people.
- The number of cell phone calls per day, the length of each call, and the amount of time people use cell phones has increased.
- Cell phone and wireless technology have had huge changes over the years. For example, how many cell phone models have you had since 1996?

Another problem is that the cell phone radiation test used by the FCC is based on the devices' possible effect on large adults—not children. Children's skulls are thinner and can absorb more radiation.

Where the AAP stands:

The AAP supports the review of radiation standards for cell phones in an effort to protect children's health, reflect current cell phone use patterns, and provide meaningful consumer disclosure. Providing parents with information about any potential risks arms them with the information they need to make informed decisions for

their families. The AAP advocates for more research into how cell phone exposure affects human health long term, particularly children's health.

Additional Information & Resources:

- Cell Phones: What's the Right Age to Start? (/English/family-life/Media/Pages/Cell-Phones-Whats-the-Right-Age-to-Start.aspx)
- Parents of Young Children: Put Down Your Smartphones (/English/family-life/Media/Pages/Parents-of-Young-Children-Put-Down-Your-Smartphones.aspx)
- Cell Phones (<http://www.niehs.nih.gov/health/topics/agents/cellphones/>) (National Institute of Environmental Health Sciences)
- Cell Phones and Cancer Risk Fact Sheet (<http://www.cancer.gov/about-cancer/causes-prevention/risk/radiation/cell-phones-fact-sheet>) (National Cancer Institute)

Last Updated 6/13/2016

Source American Academy of Pediatrics (Copyright © 2016)

The information contained on this Web site should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

Contact Us (<https://www.healthychildren.org/english/pages/contact-us.aspx>)

About Us (<https://www.healthychildren.org/english/pages/about-aap.aspx>)

Privacy Policy (<https://www.healthychildren.org/english/pages/privacy-policy.aspx>)

Terms of Use (<https://www.healthychildren.org/english/pages/terms-of-use.aspx>)

Editorial Policy (<https://www.healthychildren.org/English/Pages/Editorial-Policy.aspx>)



ie HONcode standard for trustworthy health (<http://www.healthonnet.org/HONcode/Conduct.html>)

© Copyright 2016 American Academy of Pediatrics. All rights reserved.




Healthy Children (/English) > Safety & Prevention (<https://www.healthychildren.org/English/safety-prevention>) > All Around (<https://www.healthychildren.org/English/safety-prevention/all-around>) > Electromagnetic Fields: A Hazard to Your Health?

SAFETY & PREVENTION

Electromagnetic Fields: A Hazard to Your Health?

Household Appliances:

For most people, their highest magnetic field exposures come from using household appliances with motors, transformers, or heaters.



- If a parent is concerned about electric and magnetic field exposure from appliances, identify the major sources of exposure and limit a child's time near those appliances.
- Manufacturers have reduced magnetic field exposures from electric blankets (since 1990) and from computers (since the early 1990s).
- Because magnetic fields decline rapidly with increasing distance, an easy measure is to increase the distance between children and the appliance.

Power Lines:

A Massachusetts study published back in 1993 showed a significant association between proximity to power lines and depressive symptoms; that is, people who were able to see the towers from their house or yard were nearly 3 times more likely than those living farther away to experience depression. A Finnish study done a few years later confirmed a much higher risk of severe depression among those living within 100 yards of a power line.

There remains some degree of uncertainty in the literature on electric and magnetic field exposure and developing cancer (/English/health-issues/conditions/cancer/Pages/default.aspx). This uncertainty should be considered in the context of the low individual risk and the comparable environmental risks (eg, traffic accidents) in other locations.

Obtaining magnetic field measurements in the home sometimes will show that field levels are at approximately the average level despite proximity to the power line.

Cell Phones:

In recent years, concern has increased about exposure to radio frequency electromagnetic radiation emitted from cell phones and phone station antennae. An Egyptian study confirmed concerns that living nearby mobile phone base stations increased the risk for developing:

- Headaches (/English/health-issues/conditions/head-neck-nervous-system/pages/Headaches.aspx)
- Memory problems
- Dizziness (/English/health-issues/conditions/head-neck-nervous-system/Pages/Dizziness-and-Fainting-Spells.aspx)
- Depression (/English/health-issues/conditions/emotional-problems/pages/Childhood-Depression-What-Parents-Can-Do-To-Help.aspx)
- Sleep problems

Short-term exposure to these fields in experimental studies have not always shown negative effects, but this does not rule out cumulative damage from these fields, so larger studies over longer periods are needed to help understand who is at risk. In large studies, an association has been observed between symptoms and exposure to these fields in the everyday environment.

Last Updated 11/21/2015

Source Adapted from Pediatric Environmental Health, 3rd Edition (Copyright © American Academy of Pediatrics 2011)

The information contained on this Web site should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.



P.O. Box 58
Teton Village, WY 83025
www.ehtrust.org

Jennifer Lowry, MD
Chair of the AAP Council on Environmental Health
Director of MAPEHSU
Chief, Section of Toxicology
Pediatrician, Department of Pediatrics
Children's Mercy Hospital
jlowry@cmh.edu

October 27, 2016

Dear Dr. Jennifer Lowry,

We were delighted to learn that based on the cancer findings from the National Toxicology Program (NTP) study on cell phone radiofrequency radiation (RFR), the American Academy of Pediatrics has reconfirmed its recommendation to limit exposure of children and teenagers to cell phones and other devices that emit RFR. However, along with that recommendation were four statements that downplayed the significance of the results from the NTP study. We are referring to the [Healthy Children.org AAP webpage with Ten Cell Phone Safety Tips](#).

Our comments provided below are intended to provide clarification on the reliability of available data on cancer risks associated with exposure to cell phone RFR. Based on the accumulating scientific evidence of increased cancer risk from cell phone RFR, it is necessary that health agencies and individuals promote precautionary measures now rather than waiting for absolute proof of human harm.

Statement 1: *"While there was a slight increase in a type of brain tumor, called a glioma, in a small group of people who spent the most total time on cell phone calls in one study, other studies have not found this to be true."*

Response: In their evaluation of the cancer risk of radiofrequency radiation, an expert working group of the International Agency for Research on Cancer (IARC) noted that brain cancer risks were increased significantly after 10 years of use, and risk levels were greatest on the side of the head on which users held their cell phones. Risks of glioma and acoustic neuroma were increased significantly in the multicenter Interphone case-control study as well as in pooled case control studies of Northern European countries that were included in the Interphone study, and in case control studies by Hardell et al. in Sweden¹²³⁴⁵⁶⁷⁸. The classification of RFR as a possible human

¹ Schoemaker, M. J., Swerdlow, A. J., Ahlbom, A., Auvinen, A., Blaasaas, K. G., Cardis, E., ... & Klaeboe, L. (2005). Mobile phone use and risk of acoustic neuroma: results of the Interphone case-control study in five North European countries. *British Journal of Cancer*, [93\(7\), 842-848](#).

carcinogen by IARC was based on “positive associations observed between exposure to radiofrequency radiation from wireless phones and glioma, and acoustic neuroma,” and for which a causal relationship was considered to be credible⁸⁹. Those associations were not considered to represent “sufficient evidence of carcinogenicity” at that time because recall bias in the case-control studies could not be fully ruled out as a possible contributing factor.

Since the IARC classification additional published studies indicate an association with increased tumor formation¹⁰¹¹¹²¹³¹⁴.

Statement 2: “*This study (NTP) was only done on rats. While rats can be good test subjects for medical research, they are not the same as humans. We do not yet know if the same results would occur in people.*”

The findings of brain tumors (gliomas) and malignant Schwann cell tumors of the heart in the NTP study, as well as DNA damage in brain cells of exposed animals, present a major public health concern because these tumors occurred in the same types of cells that had been reported to develop into tumors (gliomas and acoustic neuromas) in epidemiological studies of adult cell phone users.

Carcinogenicity studies in rodents are useful for several important reasons: (1) animals and humans exhibit similarities in biological processes of disease induction (that is why animal models are used in preclinical trials of new pharmaceutical agents), (2) it is unethical to intentionally expose humans to agents in order to test for adverse health effects such as cancer, (3) every agent that is known to cause cancer in humans is carcinogenic in animals when

² Lahkola, A., Auvinen, A., Raitanen, J., Schoemaker, M. J., Christensen, H. C., Feychting, M., ... & Tynes, T. (2007). Mobile phone use and risk of glioma in 5 North European countries. *International Journal of Cancer*, [120\(8\), 1769-1775](#).

³ INTERPHONE Study Group. (2010). Brain tumour risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Int. J. Epidemiol*, [39\(3\), 675-94](#).

⁴ INTERPHONE Study Group. (2010). Supplementary Material - Brain tumour risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Int. J. Epidemiol*, [39\(3\), 675-94](#).

⁵ INTERPHONE Study Group. (2011). Acoustic neuroma risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Cancer Epidemiol*, [35, 453-64](#).

⁶ Cardis, E. et al. (2011). Risk of brain tumours in relation to estimated RF dose from mobile phones: results from five Interphone countries. *Occup. Environ. Med*, [68\(9\), 631-40](#).

⁷ Hardell L., Carlberg M., & Hansson M.K. (2011). Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol*, [38\(5\):1465-74](#).

⁸ Han, Y. Y., Kano, H., Davis, D. L., Niranjan, A., & Lunsford, L. D. (2009). Cell phone use and acoustic neuroma: the need for standardized questionnaires and access to industry data. *Surgical neurology*, [72\(3\), 216-222](#).

⁹ International Agency for Research on Cancer. (2011). IARC classifies radiofrequency electromagnetic fields as possibly carcinogenic to humans. *Press release*, [\(208\)](#).

⁹ IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. (2013). Non-ionizing radiation, Part 2: Radiofrequency electromagnetic fields. *IARC monographs on the evaluation of carcinogenic risks to humans/World Health Organization, International Agency for Research on Cancer*, [102\(2\), 1-460](#).

¹⁰ Coureau, G. et al. (2014). Mobile phone use and brain tumours in the CERENAT case-control study. *Occup Environ Med*, [71\(7\), 514-22](#).

¹¹ Lerchl, A., Klose, M., Grote, K., Wilhelm, A. F., Spathmann, O., Fiedler, T., ... & Clemens, M. (2015). Tumor promotion by exposure to radiofrequency electromagnetic fields below exposure limits for humans. *Biochemical and biophysical research communications*, [459\(4\), 585-590](#).

¹² Hardell, L., & Carlberg, M. (2015). Mobile phone and cordless phone use and the risk for glioma—Analysis of pooled case-control studies in Sweden, 1997–2003 and 2007–2009. *Pathophysiology*, [22\(1\), 1-13](#).

¹³ Hardell, L., Carlberg, M., Söderqvist, F., & Mild, K. H. (2013). Case-control study of the association between malignant brain tumours diagnosed between 2007 and 2009 and mobile and cordless phone use. *International Journal of Oncology*, [43\(6\), 1833-1845](#).

¹⁴ Hardell, L., & Carlberg, M. (2013). Using the Hill viewpoints from 1965 for evaluating strengths of evidence of the risk for brain tumors associated with use of mobile and cordless phones. *Reviews on environmental health*, [28\(2-3\), 97-106](#).

adequately tested (IARC, preamble), (4) almost one-third of human carcinogens were identified after carcinogenic effects were found in well-conducted animal studies, (5) animal studies can eliminate the need to wait for a high incidence of human cancers (which may clinically manifest as much as 30 years from time of first exposure) before implementing public health-protective strategies, and (6) the control of exposure conditions in animal studies can eliminate the potential impact of confounding factors on the interpretation of study results.

Statement 3: *“The rats were exposed to very large amounts of radiation—nine hours a day, seven days a week, for two years. This is far more than most people spend holding their cell phones.”*

Response: While the exposure limit to RFR by the Federal Communications Commission is 0.08 W/kg averaged over the whole body, the localized exposure limit is 1.6 W/kg averaged over any one gram of tissue. For cell phone users, body tissues located nearest to the phone’s antenna receive higher exposures than tissues located distant from the antenna. Thus, when an individual holds a cell phone next to his or her head, exposure to the brain will be much higher than exposures averaged over the whole body. When considering organ-specific risk (e.g., risk to the brain) from cell phones, the important measure of exposure is the 1.6 W/kg value. Cell phone manufacturers provide values for their phone’s emissions. Many cell phones emit radiation that can produce local doses near 1.6 W/kg. In the NTP study in which animals were exposed to 1.5, 3, and 6.0 W/kg RFR, exposures in the brain were within 10% of the whole body exposure levels. Therefore, with respect to exposures to the brain, exposures of rats to RFR were similar to or slightly higher than human exposures from cell phones held next to the head.

Experimental carcinogenicity studies are generally conducted in small groups of rodents (approximately 50 animals of each sex and species per exposure or control group), and incidence values of adverse effects are used to assess health risks to potentially millions of exposed people. While an increased incidence of 1% in an experimental study would not be statistically significant, such an increase or even an increase in brain cancer risk of 0.001% in the general population would be dreadful; this concern is particularly pertinent for cell phones as there are more than 250 million cell phone users in the US and more than 4 billion users worldwide. Thus, to identify a hazardous agent, exposure levels in animal studies are often much higher than human exposures, while lower doses are included for analyses of dose-response relationships and assessments of human health risks. The NTP study of RFR could not use exposure intensities much higher than that of cell phones in order to prevent any measurable increases in body temperature. Consequently, the duration of exposure was extended to nine hours a day to determine whether cell phone radiation could cause adverse health effects and to provide data to characterize dose-response relationships for any detected effect and to assess human risk.

Statement 4: *“More male rats developed cancerous tumors after being exposed to the radiation than female rats. Some of the rats who developed tumors lived longer than the control group rats that were not exposed to radiation.”*

While the incidence of brain tumors and schwannomas of the heart was greater in exposed male rats than in female rats, these rare and uncommon tumors were observed only in RFR exposed

animals of both sexes with none observed in the controls. In addition, pre-cancerous lesions (glial hyperplasia and Schwann cell hyperplasia) were observed only in RFR exposed male and female rats. Observing numerical differences in response between the sexes is common in animal carcinogenicity studies as well as in human populations. For example, brain cancer mortality rates are approximately 50% higher in men than in women, and for many human cancers (e.g., colon-rectal, liver, soft tissue including heart, kidney, non-Hodgkin lymphoma, etc.) the incidence and mortality rates are much higher in men than in women. The different response rate between male and female rats in the RFR study does not alter the relevance of the cancer findings from this study.

The criticism that exposed rats lived longer than control rats, which might have affected the tumor findings, is an inaccurate portrayal and interpretation of the data for at least two reasons. First, there was no statistical difference in survival between control male rats and the exposure group with the highest rate of gliomas and heart schwannomas (male rats exposed to CDMA modulated RFR at 6 W/kg). Second, no glial cell hyperplasias (potential pre-cancerous lesions) or heart schwannomas were observed in any control rat, even though glial cell hyperplasia was detected in a CDMA-exposed rat as early as week 58 and heart schwannomas were detected as early as week 70 in exposed rats during the 2-year study. Thus, survival was sufficient to detect tumors or pre-cancerous lesions in control male rats. The exclusive findings of these tumors and pre-cancerous lesions in exposed animals support the carcinogenic potential of RFR in living organisms.

We hope these comments are helpful to you as the AAP develops future recommendations to protect children from adverse effects of RFR. It is also important to note that actively used cell phones are not the exclusive source of exposure to RFR, other sources of daily exposures include cell phones powered on even when not communicating, Wi-Fi devices, cordless phones and cell towers. Babies, toddlers and preschoolers are handed iPads and tablets as toys to play games and watch movies on. Many young children engage in wireless streamed content through devices resting on their laps, yet parents are unaware such Wi-Fi connectivity results in radiofrequency exposure to their bodies.

For children, health risks may be greater than that for adults because of greater penetration and absorption of cell phone radiation in the brains of children and because the developing nervous system of children is more susceptible to tissue damaging agents.

Sincerely,

Ron Melnick PhD

Senior Toxicologist and Director of Special Programs in the Environmental Toxicology Program at the National Institute of Environmental Health Sciences (NIEHS), National Institutes of Health, now retired.

Devra Davis, PhD MPH

President and Founder [Environmental Health Trust](#)

Visiting Professor Hebrew University Hadassah Medical Center

Does Cell Phone Use Cause Brain Cancer? What the New Study Means For You

Groundbreaking study reveals the strongest link yet between cell phone radiation and cancer. Important advice for all consumers.

By Jeneen Interlandi

May 27, 2016

The [results of a new study](#) by the National Toxicology Program—the largest and most expensive study of its kind—show a link between cell phone radiation and cancer in rats.

For many people, these findings likely raise questions and concerns about the safety of devices that we now carry with us nearly all the time.

Consumer Reports health and safety experts, who have long been concerned about the potential risks of cell phones and urged precautions when using them, say the new study supports that caution.

"Consumers don't need to stop using their phones," says Michael Hansen, Ph.D., a senior scientist with Consumer Reports who has studied this issue for years. "But there are some simple, common-sense steps you can and should take to reduce your exposure."

Specifically, Consumer Reports recommends that you:

- Try to keep the cell phone away from your head and body. Keeping it an arm's distance away significantly reduces exposure to the low-level radiation it emits. This is particularly important when the cellular signal is weak—when

your phone has only one bar, for example—because phones may increase their power then to compensate.

- Text or video call when possible, because this allows you to hold the phone farther from your body.
- When speaking, use the speakerphone on your device or a [hands-free headset](#).
- Don't stow your phone in your pants or shirt pocket. Instead, carry it in a bag or use a belt clip.

Below, answers to other basic questions about the study and what it means for you and your family.

So What Did This New Study Find?

The [study](#) found that male rats had a higher incidence of two kinds of tumors when exposed to the same type of radiation emitted by cell phones.

The results are not conclusive, and the overall relevance to human cell phone use is something that's "not currently completely worked out," said John Bucher, Ph.D., associate director of the NTP, part of the National Institutes of Health.

But the [new report](#) adds weight to human epidemiological studies that have previously raised similar concerns, and when combined with those earlier studies, is poised to force a reconsidering among federal agencies of the potential risks posed by cell phones. "In my experience," Bucher said, "the people who have reviewed our findings agree with the findings."

A spokesman for CTIA, a trade group for the wireless industry, says "Numerous international and U.S. organizations, including the U.S. Food and Drug Administration, World Health Organization, and American Cancer Society, have determined that the already existing body of peer-reviewed and published studies shows that there are no established health effects from radio frequency signals used in cellphones."

Why Should I Be Worried About a Study Using Rats?

Animal studies are actually the gold standard for determining cancer risk, for several reasons.

For one, it is unethical to expose humans to suspected carcinogens in a lab setting.

Second, studies in animals such as rats and mice can be completed much more quickly than they can be in humans, simply because their lifespans are so much shorter than ours. For example, the new NIH study involved exposing the rodents to cell phone radiation for just two years.

Finally, animal studies can validate results of previous observational studies in humans. Those studies, which track large groups of people over time, can look for associations between how many hours people said they used cell phones every day and the incidence of cancers in those people, but they can't prove a cause and effect relationship. Laboratory studies in rats, showing that exposure to cell phone radiation can cause cancers compared to a similar non-exposed group of rats, give credence to the results of observational human studies, and point strongly to cause and effect.


What Do Studies in Humans Show?

The current animal studies are worrisome precisely because they do line up with the results of some previous observational studies in humans.

Last year, [Consumer Reports reviewed that research](#), focusing on five large population studies that investigated that question. Together the studies included more than a million people worldwide, comparing cell phone users with nonusers.

Three of the studies—one from Sweden, another from France, and a third that combined data from 13 countries—suggest a connection between heavy cell phone use and gliomas, the same kind of tumors detected in the new NIH study. Those tumors are usually cancerous and often deadly.

One of those studies also hinted at a link between cell phones and acoustic neuromas (noncancerous tumors); that kind of tumor is related to the second cancer detected in the current study, malignant schwannoma of the heart.



This MRI shows a possible glioma in a human brain. The NIH study suggests cell phones radiation could cause that kind of cancer in rats.

How Might Cell Phone Radiation Cause Cancer?

Scientists [previously thought](#) that the radiation from cell phones might damage cells by heating human tissue. At high power levels radiofrequency waves—the kind emitted by cell phones—can heat up water molecules. Since human tissue is mostly water, scientists hypothesized that those waves might cause damage by heating.

The Federal Communications Commission’s cell phone emission test—which all cell phones must pass before being allowed on the market—is based on that principle.

But in 2011, scientists at the NIH found that low level radiation, held close to the head, could alter brain cells without raising body temperatures. Likewise, in 2015, German researchers reported that the same type of radiation emitted by cell phones could promote the growth of brain tumors in mice without raising body temperatures.

The NTP study controlled for heating effects by making sure that the body temperatures of exposed rats did not increase by more than 1° C (1.9° F), suggesting that the cancers were triggered by some other mechanism.

Read our previous coverage about the potential dangers of radiation from [cell phones](#) and [CT scans and X-rays](#).

How Well Does the NTP Study Mimic Current Cell Phone Usage?

The study used specially designed chambers that allowed researchers to expose rodents to standardized doses of radiation. The rodents were exposed for nine hours total each day, at intervals of 10 minutes on, 10 minutes off, for two years.

The radiation frequencies and signal modulation used were the same used by 2G (GSM or CDMA) phones, which were standard when the study began. Newer cell phones use 3G (such as UMTS or CDMA-2000) or 4G (LTE), which may have lower power outputs and different signal modulation.

“These changes may be a critical difference in whether there is a hazard today,” says Consumer Reports’ Hansen. “But the study raises enough concern with the older technologies that we recommend an additional study be done with current technology.”

The rodents were exposed over their entire bodies. While that’s obviously different than the way humans use cell phones, the rodent results are still revealing, Hansen says.

“The reason we see schwannomas in the heart here, and not the auditory system, could be due to the fact that in rodents the heart is closer to the surface of the body,” he says. “What’s more important is that the cell type found in the heart in the NTP study is the same as in some brain tumors found in several human epidemiology studies.”

What Does Consumer Reports Think the Government and Industry Should Do Now?

The substantial questions and concerns raised by this and previous research regarding cell phones and cancer requires swift and decisive action by the government and industry. Specifically, Consumer Reports believes that:

- The National Institutes of Health should commission another animal study using current cell phone technology to determine if it poses the same risks as found in this new study.
- The Federal Communications Commission should update its requirements for testing the effect of cell phone radiation on human heads. The agency's current test is based on the devices' possible effect on large adults, though research suggests that children's thinner skulls mean they may absorb more radiation. The FCC should develop new tests that take into account the potential increased vulnerability of children.
- The Food and Drug Administration and the FCC should determine whether the maximum specific absorption rate of 1.6 W/kg over a gram of tissue is an adequate maximum limit of radiation from cell phones.
- The Centers for Disease Control and Prevention should repost its advice on the potential hazard of cell phone radiation and cautionary advice that was taken down in August 2014.
- Cell phone manufacturers should prominently display advice on steps that cell phone users can take to reduce exposure to cell phone radiation.

Are You Pregnant?

Protect Your Baby from Wireless Radiation!



Recent studies from Yale University show that exposure to wireless radiation can have profound effects on brain development, including symptoms of Attention Deficit/Hyperactivity Disorder (ADHD).

Now, doctors and researchers around the world are urging pregnant women to take simple steps to limit their exposure to wireless radiation.

For more information, including links to scientific studies and video interviews with medical doctors and public health experts, please visit our web site, **BabySafeProject.org**.

10 Tips for Reducing Your Exposure to Wireless Radiation

1. Avoid carrying your cell phone on your body (e.g. in a pocket or bra).
2. Avoid holding any wireless device against your body when in use.
3. Use your cell phone on speaker setting or with an “air tube” headset.
4. Avoid using your wireless device in cars, trains or elevators.
5. Avoid cordless phones, especially where you sleep.
6. Whenever possible, connect to the internet with wired cables.
7. When using Wi-Fi, connect only to download, then disconnect and disable Wi-Fi.
8. Avoid prolonged or direct exposure to Wi-Fi routers.
9. Unplug your home Wi-Fi router when not in use (e.g. at bedtime).
10. Sleep as far away from wireless utility meters (i.e. “smart” meters) as possible



The BabySafe Project
www.BabySafeProject.org
#KnowYourExposure

What's Your
Exposure?
Take the Quiz!



*What You Need to Know
About Wireless Radiation
and Your Baby*



The BabySafe
Project

INTRODUCTION

The beautiful and awe-inspiring process by which a few cells develop into a new life is truly miraculous. As if following some invisible instructions, the intricate and complex steps for the creation of a new life unfold, forming delicate organs and vital systems and making billions of important connections and patterns in the brain alone.

Your body is the first home for this new life, and pregnancy triggers some instructions for you as well. Your body provides a protected and warm space for your baby to grow, sharing important nutrients and preparing you to nourish your baby beginning immediately after birth.

Protecting your baby from any possible harm is a strong and natural instinct. Every stage of your baby's development will be affected by the choices you make and the environments in which you live. And those choices you make are at the same time becoming more difficult and more important.



Today, scientists and medical professionals around the world are working to understand more about how a developing human life is affected by environmental exposures in the food we eat, the air we breathe, the water we drink and the surroundings in which we live and work. Identifying and avoiding all of them can seem overwhelming, but knowing the risks and the steps you can take to reduce your exposure is something you can do.

While there are many things yet to be discovered about the important relationship between your developing baby and your own environmental exposures, we do know that your body can provide only a certain degree of protection. Another thing we know is that the impact can depend on the amount and timing of the exposure.

One focus of current research is wireless radiation, also referred to as microwave radiation or radio frequency radiation (RFR). Wireless radiation is emitted by the myriad of wireless devices we encounter every day. It was once thought to be relatively harmless. Decades ago, when exposure thresholds were established, the thermal or heating effects on human tissue were the main concern.

However, we now know that wireless radiation can cause non-thermal biological effects as well, including damage to cells and DNA, even at low levels. The International Agency for Research on Cancer (IARC) of the World Health Organization (WHO) recently classified wireless radiation as possibly carcinogenic to humans, and numerous animal studies have shown neurological effects, including behavioral disorders, that may lead to long-term health consequences. All of this suggests that taking certain precautions around wireless radiation is appropriate for our most vulnerable populations, including pregnant women.

“It is so simple to move your cell phone away from your abdomen when you're pregnant. Why not do it? There's essentially no downside to being cautious and protecting your baby.”

- Dr. Hugh Taylor, Yale University

Generally speaking, *proximity* is the most important factor in determining the amount of wireless radiation to which you and your baby are exposed. Radiation levels fall off dramatically as you distance yourself from the source.

Reducing Your Exposure to Wireless Radiation

- Avoid carrying your cell phone on your body (e.g. don't carry it in a pocket or bra).
- Avoid placing your cell phone, wireless laptop or tablet on your abdomen. Don't talk or text while holding a phone against your body.
- Talk on speaker setting or with an "air tube" headset.
- Avoid using your cell phone in cars, trains and elevators.
- Avoid using cordless phones (which also expose you to wireless radiation), especially where you sleep. Corded phones do not emit RFR.
- Streaming videos and playing games results in higher levels of RFR exposure. Download and then view or play with the device in "airplane" mode. Download your email and then disconnect from WiFi, allowing you to read and respond without being exposed.
- Whenever possible, connect wired cables to a router or modem and turn off the WiFi feature on the router, modem and your personal device(s).
- Unplug your home WiFi router when not in use (e.g. at bedtime).
- Avoid prolonged or direct exposure to WiFi routers at home, school or work. If your job requires you to use a computer, position yourself so that you are not sitting close to the router and not between the router and your computer.
- If your home or apartment or close neighbor has a wireless utility meter (i.e. a "smart" meter), sleep in a room as far away from it as possible.

MORE ABOUT WIRELESS RADIATION

Radio frequency waves have been used for more than a hundred years to carry signals from transmitting towers to distant receivers. This technology has informed and entertained millions of people around the world.

However, the technology offered today by the wireless industry puts powerful transmitters as well as receivers much closer to users of all ages than ever before. This two-way communication, and the increased radiation needed to support it, is reason for concern. Indeed, manufacturers of wireless devices warn consumers to keep their phones, tablets, baby monitors or other devices away from their bodies.

Consumer demand for connectivity everywhere has resulted in the installation of many more powerful local wireless transmitters and receivers, and now hundreds of thousands of rooftop, pole-mounted and tower transmitters (antennas) are placed in close proximity to private homes, apartments, schools, office buildings, retail and recreation areas. “Free WiFi” is commonly advertised to attract customers at bars, restaurants, hotels and coffee shops. Wireless routers in public spaces are very powerful because they are intended to power many laptops or tablets simultaneously.



This ubiquitous and ever-growing wireless world that we live in means that wireless radiation is all around us. But you can still make some personal choices that can reduce your exposure. As mentioned previously, keeping a safe distance from transmitters or antennas and keeping your personal wireless devices away from your body is relatively easy to do. The amount of time you spend using wireless devices is also important. Remember that exposure adds up over time.

We hope that you will take the time to learn more about wireless radiation by visiting our website, www.BabySafeProject.org.



“Most pregnant women understand that they should stay away from cigarette smoke, avoid alcohol and be careful about eating potentially contaminated foods of any kind. The weight of evidence clearly supports the need to protect pregnant women from wireless radiation exposures as well.”

- Dr. Devra Davis, University of California at Berkeley

MORE ABOUT THE SCIENCE

A recent study (Aldad, et al 2012) conducted at Yale University found that pregnant laboratory mice exposed to ordinary cell phone radiation produced offspring that were more hyperactive and had poorer memories compared to a control group that was not exposed.

Dr. Hugh Taylor, Chair of the Department of Obstetrics, Gynecology and Reproductive Sciences at Yale University School of Medicine and his team of researchers concluded that cell phone radiation had damaged neurons in the prefrontal cortex of the brain.

The work of the Yale researchers followed a steady progression of scientific studies that demonstrated health and behavioral effects from wireless radiation. Twenty years ago, a review of the scientific literature on radiofrequency/microwave radiation conducted by the U. S. Air Force Materiel Command (Bolen 1994) concluded that “behavior may be the most sensitive biological component to RF/Microwave radiation.” Scientists at the University of Washington demonstrated DNA breaks in brain cells of rats resulting from exposure to microwave radiation (Lai, et al 1995).

Over the next decade, numerous studies were conducted regarding the safety of RF radiation with varying results, and controversy remained about the relevance of animal studies to humans.

Of particular importance was the work of researchers at the University of Kentucky where they were able to show how exposure to wireless radiation can damage or even destroy brain cells (Zhao, et al 2007). Researchers in Samsun, Turkey published findings that rats prenatally exposed to cell phone radiation developed impaired learning and also showed damage to those parts of the brain involved in memory and learning (Inkinci, et al 2013). In yet another study, rats prenatally exposed to wireless radiation also had damaged spinal cords (Odaci, et al 2013).

Regarding human impacts of wireless radiation, UCLA researchers (Divan, et al 2008) studied 13,000 mothers and children and found that prenatal exposure to cell phones was

associated with a higher risk for behavioral problems and hyperactivity in children.

Scientists continue to conduct research on the human impact of wireless radiation exposure. While we wait for the scientific process to provide us with a deeper understanding of this issue, and for government agencies to adopt appropriate exposure thresholds, a precautionary approach to exposures, especially during pregnancy, seems warranted.

For a listing of scientific studies regarding wireless radiation and its biological effects, please visit our web site BabySafeProject.org.

Government agencies, professional societies and public health organizations around the world calling for further research on wireless radiation include:

United States Environmental Protection Agency (EPA)
American Academy of Environmental Medicine (AAEM)
National Cancer Institute (NCI)
American Academy of Pediatrics (AAP)
American Cancer Society (ACS)
National Institute of Environmental Health Sciences (NIEHS)
International Agency for Research on Cancer (IARC)
International Commission for Electromagnetic Safety (ICES)
European Parliament and European Environment Agency (EEA)

“Current FCC standards do not account for the unique vulnerability and use patterns specific to pregnant women and children. It is essential that any new standards for cell phones and other wireless devices be based on protecting the youngest and most vulnerable population to ensure they are safeguarded throughout their lifetimes.”

- American Academy of Pediatrics 2013

Joint Statement on Pregnancy and Wireless Radiation

We join together as physicians, scientists and educators to express our concerns about the risks that wireless radiation poses to pregnancy and to urge pregnant women to limit their exposures.

We recognize that the exquisite systems that direct the development of human life are vulnerable to environmental insults, and that even minute exposures during critical windows of development may have serious and life-long consequences.

We know that the scientific process demands a thorough and exhaustive examination of the possible impact of wireless radiation on health; however, we believe substantial evidence of risk, rather than absolute proof of harm, must be the trigger for action to protect public health.

We call on the research community to conduct more studies to identify the mechanisms by which a fetus could be affected by wireless radiation. We call on our elected leaders to support such research and to advance policies and regulations that limit exposures for pregnant women. We call on industry to implement and explore technologies and designs that will reduce radiation exposures until such research is carried out.

We affirm our role as health and science professionals to inform the public about the potential dangers associated with early-life exposures to wireless radiation, and invite all professionals engaged in obstetric, pediatric, and environmental health advocacy to join us in our quest to ensure the safety and health of future generations.

For a current list of medical doctors, researchers and health advocates who have signed the Joint Statement, please visit BabySafeProject.org

For more information about pregnancy and wireless radiation
and to obtain additional copies of this brochure please visit

BabySafeProject.org

#KnowYourExposure

Know Your Exposure



The BabySafe Project
www.BabySafeProject.org

*The BabySafe Project is a joint initiative of
Grassroots Environmental Education
and*

Environmental Health Trust

© 2014 Grassroots Environmental Education. All rights reserved.

Dec 8

----- Forwarded message -----

From: **Cecelia Doucette** <c2douce@gmail.com>

Date: Thu, Sep 8, 2016 at 4:09 PM

Subject: Please Protect Children from Wireless Radiation

To: ematsui@jhmi.edu, [rachel.hess-](mailto:rachel.hess-mutinda@maryland.gov)

mutinda@maryland.gov, emily@marylandpirg.org, guy.guzzzone@senate.state.md.us, amezu@msde.state.md.us, allison@mdehn.org, Angela.Angel@house.state.md.us, benoythomas@gmail.com, brandi.stocksdales@maryland.gov, [Varney-](mailto:Varney-Alvarado@dncd.state.md.us)

Alvarado@dncd.state.md.us, christina.church@maryland.gov, dbishai@jhsph.edu, dianna.abney@maryland.gov, gdiette@jhmi.edu, lornegarr@msn.com, bgitterman@aol.com, jicarella@msn.com, julianlevy@comcast.net, megan.latshaw@aphl.org, cliff.mitchell@maryland.gov, nancy.servatius@maryland.gov, nobot@cehn.org, rob.hofstetter@maryland.gov

Dear Maryland Children's Environmental Health and Protection Advisory Council Members,

Please allow me to introduce myself, my name is Cece Doucette and I live in Ashland, MA. In the summer of 2013 I learned all wireless and cellular devices operate off of high levels of airborne microwave radiation that can cause infertility, brain cancer, breast cancer, behavioral issues, DNA damage and more. Children and fetuses are especially vulnerable.

As former President of the Ashland Education Foundation and Grant Coordinator for Ashland Public Schools in Massachusetts, I helped bring wireless technology into our district so I felt morally obliged to investigate and thus I started the conversation in my community.

Our School Committee researched the history, science, manufacturers and lobbyists. Health advisories are buried in [the fine print](#) of each device so they realized most folks, themselves included, did not know wi-fi poses harm. Ashland Public Schools rolled out [Best Practices for Mobile Devices](#) to keep safer distances and turn off wi-fi when not in use. The Environmental Health Trust tells us we are the [first public school](#) in the nation to do so although [other countries](#) are already setting laws into place to protect their citizens.

While scientists strongly urge using only hard-wired connections, I was grateful our School Committee put precautions in place as a first step. I realized the rest of our citizens didn't know about the potential harm, and our public health officials didn't appear to either.

So, I bought a wi-fi radiation meter and measured the radiation coming from Senator Karen Spilka's cell phone and her district director's laptop. Both measured off the charts. Senator Spilka introduced [Senate Bill 1222 An Act creating a special commission to study the health impacts of electromagnetic fields](#). It has been reported out favorably by the Joint Committee on Public Health, and the Massachusetts Department of Public Health has now hired an epidemiologist who will be looking into microwave radiation from wireless technology.

Ashland Public Library has become the first public library in the nation to host a six-part film and discussion series on [Electromagnetic Radiation and Health](#) to help educate the public. I received a grant, and we are now the first in the nation to put a Radiofrequency Acoustimeter on loan in our library so residents can measure and remediate the wireless radiation in their spaces. Our cable station, WACA, also produced a [public service video](#) to teach folks how to use it. Other communities could do the same.

I also educated our Ashland Board of Health on technology safety, and am told the following 33-minute video is a good primer for those just joining the conversation. Each of the documents I discuss is included in the Comments section below the YouTube video:

<https://www.youtube.com/watch?v=9T-4Aj6gb7A>

In my home, we've run Ethernet cables to our laptops, turned off wi-fi, and disconnected the routers. We enjoy much faster, more reliable and safer connections now that we are hardwired again. We put our cell phones in Airplane Mode when not in use and check periodically for messages when we're out, and forward calls to our landline when we're home. We replaced the cordless DECT phones, whose base stations serve as mini-cell towers pulsing radiation 24/7, with corded landline phones. The utility "smart meter" is the only radiation producer at home we cannot control, and it also pulses 24/7. We will work with our department of public utilities to have the old analog meter reinstalled. These are important measures that all families should be encouraged to take. The video above explains the biological damage otherwise.

We are fortunate that our town leaders have cared enough to really examine this issue and not fall for the industry's creation of doubt while marketing their untested products. I hope you can find something in our efforts here that might help Maryland too.

I would appreciate it if you would include this letter as a formal submission in the comments section of your report Appendix.

Thank you for your time and consideration, and for all you do to protect children from environmental toxins. Please do not hesitate to reach out if I can provide additional information -- I know it is a lot to absorb this issue, especially when it is so unknown to most.

Warm regards,

Cecelia Doucette
31 Fatima Drive
Ashland, MA 01721
[508-881-3878](tel:508-881-3878)



3 August 2016

Petaluma City Schools
District Office
200 Douglas Street
Petaluma, California 94952

Dear Sirs/Madams:

I am a public health physician who served as the Co-Editor of the Bioinitiative Report, published in 2007 as a comprehensive review of the adverse health effects of radiofrequency electromagnetic fields.

There is strong and consistent evidence that excessive exposure to radiofrequency electromagnetic fields has adverse human health effects. Of particular concern is the clear evidence that children are more vulnerable than adults. The best-documented adverse effects are an increase in risk of cancer, but cancers do not appear immediately upon exposure but rather come years later. The National Toxicology Program has within the past couple of months reported that even rats exposed to radiofrequency radiation develop brain cancer! Within a school setting there is increasing evidence that excessive exposures reduce learning ability, which is the last thing one wants in a school. Some children will also develop a syndrome of electrohypersensitivity, where they get headaches and reduced ability to pay attention and learn. While these effects are not nearly as well documented as those relating to cancer, they are particularly important within a school. This is especially the case in a wireless computer classroom, where exposure can be very high. However there will be essentially no exposure in a wired computer classroom.

The exposure levels of the Federal Communications Commission are totally outdated and do not protect the health of the public, especially of children. I urge you to abandon any plans for wireless communication within schools. It is of course critical that all children have access to the Internet, but when this is done through wired connections they will not be exposed to excessive electromagnetic fields.

Yours sincerely,

David O. Carpenter, M.D.
Director, Institute for Health and the Environment
University at Albany

Studies that show **Cell Phone** Health Effects

Table of Contents

Over 700 Studies Showing Health Effects from Cell Phone Radio Frequency Radiation

Biological Effect	No of Studies	Page #
Cancer	57	2 to 29
DNA Damage	47	29 to 52
Effects on Pregnancy and the Fetus	29	52 to 67
Fertility and Reproductive Harm	55	67 to 94
Neurological Effects	44	94-114
Effects on Brain	104	114-161
Central Nervous System Effects	25	161-172
Hearing Effects	23	172-183
Effect on Eyes	11	183-188
Children Health Effects	20	188-197
Oxidative Stress	14	197-204
Cell Function Impairment	41	205-224
Effects on Hormones	11	224-229
Effects on the Heart	5	229-232
Effects on Glucose Tolerance	2	233
Effects on the Whole Body	10	233-237
Survival Effects	2	237-239
Medical Implants and Devices Effects	32	239-255
Cell Phone Addiction	2	255-256
Synergistic Effects With Other Agents	12	256-263
Inflammation	3	263-264
Effects on Behavior	10	264-269
Electro-Magnetic Hypersensitivity	6	269-272
Effects on Blood	13	272-278
Effects on Wellbeing	18	278-286
Effects on Glands	5	286-288
Effects on Sleep	19	288-297
Effects on Skin	10	298-302
Allergies Effect	4	302-304
Effects to Critical Organs	10	304-310
Mercury Release in Dental Amalgams	1	310
Effects on Other Living Things	14	310-323
Miscellaneous Effects	12	323-329
Driving Impairment Using a Cell Phone	31	329-343
Overlooked Effects in Science	7	343-347

Studies that show **Cell Phone** Health Effects

Cancer

Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. Int J Oncol. 38(5):1465-1474, 2011.

We studied the association between use of mobile and cordless phones and malignant brain tumours. Pooled analysis was performed of two case-control studies on patients with malignant brain tumours diagnosed during 1997-2003 and matched controls alive at the time of study inclusion and one case-control study on deceased patients and controls diagnosed during the same time period. Cases and controls or relatives to deceased subjects were interviewed using a structured questionnaire. Replies were obtained for 1,251 (85%) cases and 2,438 (84%) controls. The risk increased with latency period and cumulative use in hours for both mobile and cordless phones. Highest risk was found for the most common type of glioma, astrocytoma, yielding in the >10 year latency group for mobile phone use odds ratio (OR) = 2.7, 95% confidence interval (CI) = 1.9-3.7 and cordless phone use OR = 1.8, 95% CI = 1.2-2.9. In a separate analysis, these phone types were independent risk factors for glioma. The risk for astrocytoma was highest in the group with first use of a wireless phone before the age of 20; mobile phone use OR = 4.9, 95% CI = 2.2-11, cordless phone use OR = 3.9, 95% CI = 1.7-8.7. In conclusion, an increased risk was found for glioma and use of mobile or cordless phone. The risk increased with latency time and cumulative use in hours and was highest in subjects with first use before the age of 20.

Lonn S, Ahlbom A, Hall P, Feychting M. Mobile phone use and the risk of acoustic neuroma. Epidemiology. 15(6):653-659, 2004.

BACKGROUND:: Radiofrequency exposure from mobile phones is concentrated to the tissue closest to the handset, which includes the auditory nerve. If this type of exposure increases tumor risk, acoustic neuroma would be a potential concern. METHODS:: In this population-based case-control study we identified all cases age 20 to 69 years diagnosed with acoustic neuroma during 1999 to 2002 in certain parts of Sweden. Controls were randomly selected from the study base, stratified on age, sex, and residential area. Detailed information about mobile phone use and other environmental exposures was collected from 148 (93%) cases and 604 (72%) controls. RESULTS:: The overall odds ratio for acoustic neuroma associated with regular mobile phone use was 1.0 (95% confidence interval = 0.6-1.5). Ten years after the start of mobile phone use the estimates relative risk increased to 1.9 (0.9-4.1); when restricting to tumors on the same side of the head as the phone was normally used, the relative risk was 3.9 (1.6-9.5). CONCLUSIONS:: Our findings do not indicate an increased risk of acoustic neuroma related to short-term mobile phone use after a short latency period. However, our data suggest an increased risk of acoustic neuroma associated with mobile phone use of at least 10 years' duration.

Levis AG, Minicuci N, Ricci P, Gennaro V, Garbisa S. Mobile phones and head tumours.

Studies that show **Cell Phone** Health Effects

The discrepancies in cause-effect relationships in the epidemiological studies - how do they arise? *Environ Health.* 10:59, 2011.

BACKGROUND: Whether or not there is a relationship between use of mobile phones (analogue and digital cellars, and cordless) and head tumour risk (brain tumours, acoustic neuromas, and salivary gland tumours) is still a matter of debate; progress requires a critical analysis of the methodological elements necessary for an impartial evaluation of contradictory studies. **METHODS:** A close examination of the protocols and results from all case-control and cohort studies, pooled- and meta-analyses on head tumour risk for mobile phone users was carried out, and for each study the elements necessary for evaluating its reliability were identified. In addition, new meta-analyses of the literature data were undertaken. These were limited to subjects with mobile phone latency time compatible with the progression of the examined tumours, and with analysis of the laterality of head tumour localisation corresponding to the habitual laterality of mobile phone use. **RESULTS:** Blind protocols, free from errors, bias, and financial conditioning factors, give positive results that reveal a cause-effect relationship between long-term mobile phone use or latency and statistically significant increase of ipsilateral head tumour risk, with biological plausibility. Non-blind protocols, which instead are affected by errors, bias, and financial conditioning factors, give negative results with systematic underestimate of such risk. However, also in these studies a statistically significant increase in risk of ipsilateral head tumours is quite common after more than 10 years of mobile phone use or latency. The meta-analyses, our included, examining only data on ipsilateral tumours in subjects using mobile phones since or for at least 10 years, show large and statistically significant increases in risk of ipsilateral brain gliomas and acoustic neuromas. **CONCLUSIONS:** Our analysis of the literature studies and of the results from meta-analyses of the significant data alone shows an almost doubling of the risk of head tumours induced by long-term mobile phone use or latency.

Hardell L, Carlberg M, Hansson Mild K. Use of mobile phones and cordless phones is associated with increased risk for glioma and acoustic neuroma. Pathophysiology. 2012 Dec 20. pii: S0928-4680(12)00110-1. doi: 10.1016/j.pathophys.2012.11.001. [Epub ahead of print]

The International Agency for Research on Cancer (IARC) at WHO evaluation of the carcinogenic effect of RF-EMF on humans took place during a 24-31 May 2011 meeting at Lyon in France. The Working Group consisted of 30 scientists and categorised the radiofrequency electromagnetic fields from mobile phones, and from other devices that emit similar non-ionising electromagnetic fields (RF-EMF), as Group 2B, i.e., a 'possible', human carcinogen. The decision on mobile phones was based mainly on the Hardell group of studies from Sweden and the IARC Interphone study. We give an overview of current epidemiological evidence for an increased risk for brain tumours including a meta-analysis of the Hardell group and Interphone results for mobile phone use. Results for cordless phones are lacking in Interphone. The meta-analysis gave for glioma in the most exposed part of the brain, the temporal lobe, odds ratio (OR)=1.71, 95% confidence interval (CI)=1.04-2.81 in the ≥10 years (>10 years in the Hardell group)

Studies that show **Cell Phone** Health Effects

latency group. Ipsilateral mobile phone use ≥ 1640 h in total gave OR=2.29, 95% CI=1.56-3.37. The results for meningioma were OR=1.25, 95% CI=0.31-4.98 and OR=1.35, 95% CI=0.81-2.23, respectively. Regarding acoustic neuroma ipsilateral mobile phone use in the latency group ≥ 10 years gave OR=1.81, 95% CI=0.73-4.45. For ipsilateral cumulative use ≥ 1640 h OR=2.55, 95% CI=1.50-4.40 was obtained. Also use of cordless phones increased the risk for glioma and acoustic neuroma in the Hardell group studies. Survival of patients with glioma was analysed in the Hardell group studies yielding in the >10 years latency period hazard ratio (HR)=1.2, 95% CI=1.002-1.5 for use of wireless phones. This increased HR was based on results for astrocytoma WHO grade IV (glioblastoma multiforme). Decreased HR was found for low-grade astrocytoma, WHO grades I-II, which might be caused by RF-EMF exposure leading to tumour-associated symptoms and earlier detection and surgery with better prognosis. Some studies show increasing incidence of brain tumours whereas other studies do not. It is concluded that one should be careful using incidence data to dismiss results in analytical epidemiology. The IARC carcinogenic classification does not seem to have had any significant impact on governments' perceptions of their responsibilities to protect public health from this widespread source of radiation.

Hardell L, Carlberg M, Söderqvist F, Hansson Mild K. Meta-analysis of long-term mobile phone use and the association with brain tumours. *Int J Oncol.* 32(5):1097-1103, 2008.

We evaluated long-term use of mobile phones and the risk for brain tumours in case-control studies published so far on this issue. We identified ten studies on glioma and meta-analysis yielded OR = 0.9, 95% CI = 0.8-1.1. Latency period of ≥ 10 -years gave OR = 1.2, 95% CI = 0.8-1.9 based on six studies, for ipsilateral use (same side as tumour) OR = 2.0, 95% CI = 1.2-3.4 (four studies), but contralateral use did not increase the risk significantly, OR = 1.1, 95% CI = 0.6-2.0. Meta-analysis of nine studies on acoustic neuroma gave OR = 0.9, 95% CI = 0.7-1.1 increasing to OR = 1.3, 95% CI = 0.6-2.8 using ≥ 10 -years latency period (four studies). Ipsilateral use gave OR = 2.4, 95% CI = 1.1-5.3 and contra-lateral OR = 1.2, 95% CI = 0.7-2.2 in the ≥ 10 -years latency period group (three studies). Seven studies gave results for meningioma yielding overall OR = 0.8, 95% CI = 0.7-0.99. Using ≥ 10 -years latency period OR = 1.3, 95% CI = 0.9-1.8 was calculated (four studies) increasing to OR = 1.7, 95% CI = 0.99-3.1 for ipsilateral use and OR = 1.0, 95% CI = 0.3-3.1 for contralateral use (two studies). We conclude that this meta-analysis gave a consistent pattern of an association between mobile phone use and ipsilateral glioma and acoustic neuroma using ≥ 10 -years latency period.

Hardell LO, Carlberg M, Soderqvist F, Hansson Mild K, Morgan LL Long-term use of cellular phones and brain tumours - increased risk associated with use for > 10 years. *Occup Environ Med.* 64(9):626-632, 2007.

AIM: To evaluate brain tumour risk among long-term users of cellular telephones.
METHODS: One cohort study and 13 case-control studies were identified on this topic. Data were scrutinized for use of mobile phone for > 10 years and ipsilateral exposure if presented. RESULTS: The cohort study was of limit value due to methodological

Studies that show **Cell Phone** Health Effects

shortcomings in the study. Of the 13 case-control studies, 9 gave results for > 10 years use or latency period. Most of these results were based on low numbers. Clearly an association with acoustic neuroma was found in four studies with two- to three-fold increased risk in the group with at least 10 years use of a mobile phone. No risk was found in one study, but the tumour size was significantly larger among users. Five studies gave results for malignant brain tumours in that latency group. All gave increased OR especially for ipsilateral exposure. Highest OR = 5.4, 95 % CI = 3.0-9.6 was calculated for high-grade glioma and ipsilateral exposure in one study. CONCLUSIONS: Results from present studies on use of mobile phones for > 10 years give a consistent pattern of an increased risk for acoustic neuroma and glioma, most pronounced for high-grade glioma. The risk is highest for ipsilateral exposure.

Hardell L, Carlberg M, Cell and cordless phone risk for glioma - Analysis of pooled case-control studies in Sweden, 1997-2003 and 2007-2009, Pathophysiology (2014), Available online 29 October 2014. <http://dx.doi.org/10.1016/j.pathophys.2014.10.001>

We made a pooled analysis of 2 case-control studies on malignant brain tumours with patients diagnosed during 1997-2003 and 2007-2009. They were aged 20-80 years and 18-75 years, respectively, at the time of diagnosis. Only cases with histopathological verification of the tumour were included. Population-based controls, matched on age and gender, were used. Exposures were assessed by questionnaire. The whole reference group was used in the unconditional regression analysis adjusted for gender, age, year of diagnosis and socio-economic index.

In total 1,498 (89%) cases and 3,530 (87%) controls participated. Mobile phone use increased the risk of glioma, OR = 1.3, 95% CI = 1.1-1.6 overall, increasing to OR = 3.0, 95% CI = 1.7-5.2 in the > 25 year latency group. Use of cordless phones increased the risk to OR = 1.4, 95% CI = 1.1-1.7, with highest risk in the >15-20 year latency group yielding OR = 1.7, 95% CI = 1.1-2.5. The OR increased statistically significant both per 100 h of cumulative use, and per year of latency for mobile and cordless phone use. Highest ORs overall were found for ipsilateral mobile or cordless phone use, OR = 1.8, 95% CI = 1.4-2.2 and OR = 1.7, 95% CI = 1.3-2.1, respectively. The highest risk was found for glioma in the temporal lobe. First use of mobile or cordless phone before the age of 20 gave higher OR for glioma than in later age groups

Hardell L, Carlberg M. Use of Mobile and Cordless Phones and Survival of Patients with Glioma. Neuroepidemiology. 40(2):101-108, 2012.

Background: We analysed the survival of patients after glioma diagnosis in relation to the use of wireless phones. Methods: All cases diagnosed between 1997 and 2003 with a malignant brain tumour (n = 1,251) in our case-control studies were included and followed from the date of diagnosis to the date of death or until May 30, 2012. Results: For glioma, the use of wireless phones (mobile and cordless phones) gave a hazard ratio (HR) = 1.1 (95% confidence interval, CI = 0.9-1.2), with >10-year latency HR = 1.2 (95% CI = 1.002-1.5, p trend = 0.02). For astrocytoma grade I-II (low-grade), the results were, HR = 0.5 (95% CI = 0.3-0.9) and for astrocytoma grade IV (glioblastoma), HR = 1.1 (95% CI =

Studies that show **Cell Phone** Health Effects

0.95-1.4), with >10 year latency HR = 1.3 (95% CI = 1.03-1.7). In the highest tertile (>426 h) of cumulative use, HR = 1.2 (95% CI = 0.95-1.5) was found for glioblastoma. The results were similar for mobile and cordless phones. Conclusions: Decreased survival of glioma cases with long-term and high cumulative use of wireless phones was found. A survival disadvantage for astrocytoma grade IV, but a survival benefit for astrocytoma grade I-II was observed which could be due to exposure-related tumour symptoms leading to earlier diagnosis and surgery in that patient group.

Carlberg M, Hardell L. Decreased Survival of Glioma Patients with Astrocytoma Grade IV (Glioblastoma Multiforme) Associated with Long-Term Use of Mobile and Cordless Phones. *Int J Environ Res Public Health*. 11(10):10790-10805, 2014.

On 31 May 2011 the WHO International Agency for Research on Cancer (IARC) categorised radiofrequency electromagnetic fields (RF-EMFs) from mobile phones, and from other devices that emit similar non-ionising electromagnetic fields, as a Group 2B, i.e., a "possible", human carcinogen. A causal association would be strengthened if it could be shown that the use of wireless phones has an impact on the survival of glioma patients. We analysed survival of 1678 glioma patients in our 1997-2003 and 2007-2009 case-control studies. Use of wireless phones in the >20 years latency group (time since first use) yielded an increased hazard ratio (HR) = 1.7, 95% confidence interval (CI) = 1.2-2.3 for glioma. For astrocytoma grade IV (glioblastoma multiforme; n = 926) mobile phone use yielded HR = 2.0, 95% CI = 1.4-2.9 and cordless phone use HR = 3.4, 95% CI = 1.04-11 in the same latency category. The hazard ratio for astrocytoma grade IV increased statistically significant per year of latency for wireless phones, HR = 1.020, 95% CI = 1.007-1.033, but not per 100 h cumulative use, HR = 1.002, 95% CI = 0.999-1.005. HR was not statistically significant increased for other types of glioma. Due to the relationship with survival the classification of IARC is strengthened and RF-EMF should be regarded as human carcinogen requiring urgent revision of current exposure guidelines.

Sadetzki S, Chetrit A, Jarus-Hakak A, Cardis E, Deutch Y, Duvdevani S, Zultan A, Novikov I, Freedman L, Wolf M. Cellular phone use and risk of benign and malignant parotid gland tumors--a nationwide case-control study. *Am J Epidemiol*. 167(4):457-467, 2008.

The objective of this nationwide study was to assess the association between cellular phone use and development of parotid gland tumors (PGTs). The methods were based on the international INTERPHONE study that aimed to evaluate possible adverse effects of cellular phone use. The study included 402 benign and 58 malignant incident cases of PGTs diagnosed in Israel at age 18 years or more, in 2001-2003, and 1,266 population individually matched controls. For the entire group, no increased risk of PGTs was observed for ever having been a regular cellular phone user (odds ratio = 0.87; p = 0.3) or for any other measure of exposure investigated. However, analysis restricted to regular users or to conditions that may yield higher levels of exposure (e.g., heavy use in rural areas) showed consistently elevated risks. For ipsilateral use, the odds ratios in the highest category of cumulative number of calls and call time without use of hands-free

Studies that show **Cell Phone** Health Effects

devices were 1.58 (95% confidence interval: 1.11, 2.24) and 1.49 (95% confidence interval: 1.05, 2.13), respectively. The risk for contralateral use was not significantly different from 1. A positive dose-response trend was found for these measurements. Based on the largest number of benign PGT patients reported to date, our results suggest an association between cellular phone use and PGTs.

Richter ED, Berman T, Levy O. Brain cancer with induction periods of less than 10 years in young military radar workers. Arch Environ Health 57(4):270-272, 2002.

The authors have reported on 5 young patients who had brain tumors that appeared within 10 yr of initial occupational exposures to radar. Four of the patients were less than 30 yr of age when the diagnoses were initially made. Brief induction periods that follow high exposures in individual sentinel patients are a recognized indicator of impending group risk, and these periods call attention to the need for precautionary measures. Similarly, reports of short induction periods for brain cancer on the side of the head in which there has been prior use of cell phones may also indicate increased risk.

Hardell L, Carlberg M, Söderqvist F, Mild KH. Case-control study of the association between malignant brain tumours diagnosed between 2007 and 2009 and mobile and cordless phone use. Int J Oncol. 43(6):1833-1845, 2013.

Previous studies have shown a consistent association between long-term use of mobile and cordless phones and glioma and acoustic neuroma, but not for meningioma. When used these phones emit radiofrequency electromagnetic fields (RF-EMFs) and the brain is the main target organ for the handheld phone. The International Agency for Research on Cancer (IARC) classified in May, 2011 RF-EMF as a group 2B, i.e. a 'possible' human carcinogen. The aim of this study was to further explore the relationship between especially long-term (>10 years) use of wireless phones and the development of malignant brain tumours. We conducted a new case-control study of brain tumour cases of both genders aged 18-75 years and diagnosed during 2007-2009. One population-based control matched on gender and age (within 5 years) was used to each case. Here, we report on malignant cases including all available controls. Exposures on e.g. use of mobile phones and cordless phones were assessed by a self-administered questionnaire. Unconditional logistic regression analysis was performed, adjusting for age, gender, year of diagnosis and socio-economic index using the whole control sample. Of the cases with a malignant brain tumour, 87% (n=593) participated, and 85% (n=1,368) of controls in the whole study answered the questionnaire. The odds ratio (OR) for mobile phone use of the analogue type was 1.8, 95% confidence interval (CI)=1.04-3.3, increasing with >25 years of latency (time since first exposure) to an OR=3.3, 95% CI=1.6-6.9. Digital 2G mobile phone use rendered an OR=1.6, 95% CI=0.996-2.7, increasing with latency >15-20 years to an OR=2.1, 95% CI=1.2-3.6. The results for cordless phone use were OR=1.7, 95% CI=1.1-2.9, and, for latency of 15-20 years, the OR=2.1, 95% CI=1.2-3.8. Few participants had used a cordless phone for >20-25 years. Digital type of wireless phones (2G and 3G mobile phones, cordless phones) gave increased risk with latency >1-5 years, then a lower risk in the following latency groups, but again increasing risk with latency

Studies that show **Cell Phone** Health Effects

>15-20 years. Ipsilateral use resulted in a higher risk than contralateral mobile and cordless phone use. Higher ORs were calculated for tumours in the temporal and overlapping lobes. Using the meningioma cases in the same study as reference entity gave somewhat higher ORs indicating that the results were unlikely to be explained by recall or observational bias. This study confirmed previous results of an association between mobile and cordless phone use and malignant brain tumours. These findings provide support for the hypothesis that RF-EMFs play a role both in the initiation and promotion stages of carcinogenesis.

Carlberg M, Hardell L. On the association between glioma, wireless phones, heredity and ionising radiation. Pathophysiology. 19(4):243-252, 2012.

We performed two case-control studies on brain tumours diagnosed during 1 January 1997 to 30 June 2000 and 1 July 2000 to 31 December 2003, respectively. Living cases and controls aged 20-80 years were included. An additional study was performed on deceased cases with a malignant brain tumour using deceased controls. Pooled results for glioma yielded for ipsilateral use of mobile phone odds ratio (OR)=2.9, 95% confidence interval (CI)=1.8-4.7 in the >10 years latency group. The corresponding result for cordless phone was OR=3.8, 95% CI=1.8-8.1. OR increased statistically significant for cumulative use of wireless phones per 100h and per year of latency. For high-grade glioma ipsilateral use of mobile phone gave OR=3.9, 95% CI=2.3-6.6 and cordless phone OR=5.5, 95% CI=2.3-13 in the >10 years latency group. Heredity for brain tumour gave OR=3.4, 95% CI=2.1-5.5 for glioma. There was no interaction with use of wireless phones. X-ray investigation of the head gave overall OR=1.3, 95% CI=1.1-1.7 for glioma without interaction with use of wireless phones or heredity. In conclusion use of mobile and cordless phone increased the risk for glioma with highest OR for ipsilateral use, latency >10 years and third tertile of cumulative use in hours. In total, the risk was highest in the age group <20 years for first use of a wireless phone.

Hardell L, Carlberg M, Hansson Mild K. Mobile phone use and the risk for malignant brain tumors: A case-control study on deceased cases and controls. Neuroepidemiology. 35(2):109-114, 2010.

We investigated the use of mobile or cordless phones and the risk for malignant brain tumors in a group of deceased cases. Most previous studies have either left out deceased cases of brain tumors or matched them to living controls and therefore a study matching deceased cases to deceased controls is warranted. Recall error is one issue since it has been claimed that increased risks reported in some studies could be due to cases blaming mobile phones as a cause of the disease. This should be of less importance for deceased cases and if cancer controls are used. In this study brain tumor cases aged 20-80 years diagnosed during 1997-2003 that had died before inclusion in our previous studies on the same topic were included. Two control groups were used: one with controls that had died from another type of cancer than brain tumor and one with controls that had died from other diseases. Exposure was assessed by a questionnaire sent to the next-of-kin for both cases and controls. Replies were obtained for 346 (75%) cases, 343 (74%) cancer controls and 276 (60%) controls with other

Studies that show **Cell Phone** Health Effects

diseases. Use of mobile phones gave an increased risk, highest in the >10 years' latency group yielding odds ratio (OR) = 2.4, and 95% confidence interval (CI) = 1.4-4.1. The risk increased with cumulative number of lifetime hours for use, and was highest in the >2,000 h group (OR = 3.4, 95% CI = 1.6-7.1). No clear association was found for use of cordless phones, although OR = 1.7, 95% CI = 0.8-3.4 was found in the group with >2,000 h of cumulative use. This investigation confirmed our previous results of an association between mobile phone use and malignant brain tumors.

Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of two case-control studies on use of cellular and cordless telephones and the risk for malignant brain tumours diagnosed in 1997-2003. *Int Arch Occup Environ Health*. 79(8):630-639, 2006.

Objectives: To study the use of cellular and cordless telephones and the risk for malignant brain tumours. Methods: Two case-control studies on malignant brain tumours diagnosed during 1997-2003 included answers from 905 (90%) cases and 2,162 (89%) controls aged 20-80 years. We present pooled analysis of the results in the two studies. Results: Cumulative lifetime use for >2,000 h yielded for analogue cellular phones odds ratio (OR)=5.9, 95% confidence interval (CI)=2.5-14, digital cellular phones OR=3.7, 95% CI=1.7-7.7, and for cordless phones OR=2.3, 95% CI=1.5-3.6. Ipsilateral exposure increased the risk for malignant brain tumours; analogue OR=2.1, 95% CI=1.5-2.9, digital OR=1.8, 95% CI=1.4-2.4, and cordless OR=1.7, 95% CI=1.3-2.2. For high-grade astrocytoma using >10 year latency period analogue phones yielded OR=2.7, 95% CI=1.8-4.2, digital phones OR=3.8, 95% CI=1.8-8.1, and cordless phones OR=2.2, 95% CI=1.3-3.9. In the multivariate analysis all phone types increased the risk. Regarding digital phones OR=3.7, 95% CI=1.5-9.1 and cordless phones OR=2.1, 95% CI=0.97-4.6 were calculated for malignant brain tumours for subjects with first use <20 years of age, higher than in older persons. Conclusion: Increased risk was obtained for both cellular and cordless phones, highest in the group with >10 years latency period.

Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of two case-control studies on the use of cellular and cordless telephones and the risk of benign brain tumours diagnosed during 1997-2003. *Int J Oncol*. 28(2):509-518, 2006.

The use of cellular and cordless telephones and the risk of brain tumours is of concern since the brain is a high exposure area. We present the results of a pooled analysis of two case-control studies on benign brain tumours diagnosed during 1997-2003 including answers from 1,254 (88%) cases and 2,162 (89%) controls aged 20-80 years. For acoustic neuroma, the use of analogue cellular phones gave an odds ratio (OR) of 2.9 and a 95% confidence interval (CI) of 2.0-4.3; for digital cellular phones, OR=1.5; 95% CI=1.1-2.1; and for cordless telephones, OR=1.5, 95% CI=1.04-2.0. The highest OR was found for analogue phones with a latency period of >15 years; OR=3.8, 95% CI=1.4-10. Regarding meningioma, the results were as follows: for analogue phones, OR=1.3, 95% CI=0.99-1.7; for digital phones, OR=1.1, 95% CI=0.9-1.3; and for cordless phones, OR=1.1, 95% CI=0.9-1.4. In the multivariate analysis, a significantly increased risk of acoustic neuroma was found with the use of analogue phones.

Studies that show **Cell Phone** Health Effects

Richter E, Berman T, Ben-Michael E, Laster R, Westin JB, Cancer in Radar Technicians Exposed to Radiofrequency/Microwave Radiation: Sentinel Episodes. *Int J Occup Environ Health* 6(3):187-193, 2000.

Controversy exists concerning the health risks from exposures to radiofrequency/microwave irradiation (RF/MW). The authors report exposure-effect relationships in sentinel patients and their co-workers, who were technicians with high levels of exposure to RF/MW radiation. Information about exposures of patients with sentinel tumors was obtained from interviews, medical records, and technical sources. One patient was a member of a cohort of 25 workers with six tumors. The authors estimated relative risks for cancer in this group and latency periods for a larger group of self-reported individuals. Index patients with melanoma of the eye, testicular cancer, nasopharyngioma, non-Hodgkin's lymphoma, and breast cancer were in the 20-37-year age group. Information about work conditions suggested prolonged exposures to high levels of RF/MW radiation that produced risks for the entire body. Clusters involved many different types of tumors. Latency periods were extremely brief in index patients and a larger self-reported group. The findings suggest that young persons exposed to high levels of RF/MW radiation for long periods in settings where preventive measures were lax were at increased risk for cancer. Very short latency periods suggest high risks from high-level exposures. Calculations derived from a linear model of dose-response suggest the need to prevent exposures in the range of 10-100 $\mu\text{W}/\text{cm}^2$.

Hardell L, Carlberg M, Hansson Mild K, Eriksson M. Case-control study on the use of mobile and cordless phones and the risk for malignant melanoma in the head and neck region. *Pathophysiology*. 18(4):325-333, 2011.

The incidence of cutaneous malignant melanoma has increased during the last decades in Sweden as in many other countries. Besides of ultraviolet radiation and constitutional factors such as light-sensitive skin and poor ability to tan few risk factors are established. Some studies indicate that electromagnetic fields might be of concern. In this case-control study we assessed use of mobile and cordless phones in 347 cases with melanoma in the head and neck region and 1184 controls. These subjects constituted 82% and 80%, respectively, that answered the questionnaire. Overall no increased risk was found. However, in the most exposed area; temporal, cheek and ear, cumulative use >365h of mobile phone yielded in the >1-5-year latency group odds ratio (OR)=2.1, 95% confidence interval (CI)=0.7-6.1 and cordless phone use gave OR=2.1, 95% CI=1.1-3.8. Highest OR was calculated for first use of mobile or cordless phone before the age of 20 years regardless of anatomical localisation in the head and neck region. No interaction was found with established risk factors such as red, medium blond or fair hair colour, blue eyes, skin type I or II (never or sometimes tanned), severe sunburns as teenager or heredity. The results must be interpreted with caution due to low numbers and potential methodological shortcomings in a case-control study. However, the findings might be consistent with a late carcinogenic effect from microwaves, i.e. tumour promotion, but need to be confirmed.

Myung SK, Ju W, McDonnell DD, Lee YJ, Kazinets G, Cheng CT, Moskowitz JM. Mobile

Studies that show **Cell Phone** Health Effects

Phone Use and Risk of Tumors: A Meta-Analysis. J Clin Oncol. 27:5565-5572, 2009.

PURPOSE: Case-control studies have reported inconsistent findings regarding the association between mobile phone use and tumor risk. We investigated these associations using a meta-analysis. **METHODS:** We searched MEDLINE (PubMed), EMBASE, and the Cochrane Library in August 2008. Two evaluators independently reviewed and selected articles based on predetermined selection criteria. **RESULTS:** Of 465 articles meeting our initial criteria, 23 case-control studies, which involved 37,916 participants (12,344 patient cases and 25,572 controls), were included in the final analyses. Compared with never or rarely having used a mobile phone, the odds ratio for overall use was 0.98 for malignant and benign tumors (95% CI, 0.89 to 1.07) in a random-effects meta-analysis of all 23 studies. However, a significant positive association (harmful effect) was observed in a random-effects meta-analysis of eight studies using blinding, whereas a significant negative association (protective effect) was observed in a fixed-effects meta-analysis of 15 studies not using blinding. Mobile phone use of 10 years or longer was associated with a risk of tumors in 13 studies reporting this association (odds ratio = 1.18; 95% CI, 1.04 to 1.34). Further, these findings were also observed in the subgroup analyses by methodologic quality of study. Blinding and methodologic quality of study were strongly associated with the research group. **CONCLUSION:** The current study found that there is possible evidence linking mobile phone use to an increased risk of tumors from a meta-analysis of low-biased case-control studies. Prospective cohort studies providing a higher level of evidence are needed.

Hardell L, Mild KH, Carlberg M. Further aspects on cellular and cordless telephones and brain tumours. Int J Oncol 22(2):399-407, 2003.

We included in a case-control study on brain tumours and mobile and cordless telephones 1,617 patients aged 20-80 years of both sexes diagnosed during January 1, 1997 to June 30, 2000. They were alive at the study time and had histopathology verified brain tumour. One matched control to each case was selected from the Swedish Population Register. The study area was the Uppsala-Orebro, Stockholm, Linköping and Göteborg medical regions of Sweden. Exposure was assessed by a questionnaire that was answered by 1,429 (88%) cases and 1,470 (91%) controls. In total use of analogue cellular telephones gave an increased risk with odds ratio (OR)=1.3, 95% confidence interval (CI)=1.04-1.6, whereas digital and cordless phones did not overall increase the risk significantly. Ipsilateral use of analogue phones gave OR=1.7, 95% CI=1.2-2.3, digital phones OR=1.3, 95% CI=1.02-1.8 and cordless phones OR=1.2, 95% CI=0.9-1.6. The risk for ipsilateral use was significantly increased for astrocytoma for all studied phone types, analogue phones OR=1.8, 95% CI=1.1-3.2, digital phones OR=1.8, 95% CI=1.1-2.8, cordless phones OR=1.8, 95% CI=1.1-2.9. Use of a telephone on the opposite side of the brain was not associated with a significantly increased risk for brain tumours. Regarding anatomical area of the tumour and exposure to microwaves, the risk was increased for tumours located in the temporal area on the same side of the brain that was used during phone calls, significantly so for analogue cellular telephones OR=2.3, 95% CI=1.2-4.1. For acoustic neurinoma OR=4.4, 95% CI=2.1-9.2 was calculated among analogue

Studies that show **Cell Phone** Health Effects

cellular telephone users. When duration of use was analysed as a continuous variable in the total material, the risk increased per year for analogue phones with OR=1.04, 95% CI=1.01-1.08. For astrocytoma and ipsilateral use the trend was for analogue phones OR=1.10, 95% CI=1.02-1.19, digital phones OR=1.11, 95% CI=1.01-1.22, and cordless phones OR=1.09, 95% CI=1.01-1.19. There was a tendency of a shorter tumour induction period for ipsilateral exposure to microwaves than for contralateral, which may indicate a tumour promotor effect.

Hardell, L., Carlberg, M., Mild, K., 2005. Case-control study of the association between the use of cellular and cordless telephones and malignant brain tumors diagnosed during 2000-2003. Environ Res 100(2):232-241, 2006.

We performed a case-control study on the use of cellular and cordless telephones and the risk for brain tumors diagnosed during 2000-2003. We report the results for malignant brain tumors with data from 317 cases (88%) and 692 controls (84%). The use of analog cellular phones yielded odds ratio (OR) of 2.6 and a 95% confidence interval (CI) of 1.5-4.3, increasing to OR=3.5 and 95% CI=2.0-6.4 with a >10-year latency period. Regarding digital cellular telephones, the corresponding results were OR=1.9, 95% CI=1.3-2.7 and OR=3.6, 95% CI=1.7-7.5, respectively. Cordless telephones yielded OR=2.1, 95% CI=1.4-3.0, and with a >10-year latency period, OR=2.9, 95% CI=1.6-5.2. The OR increased with the cumulative number of hours of use and was highest for high-grade astrocytoma. A somewhat increased risk was also found for low-grade astrocytoma and other types of malignant brain tumors, although not significantly so. In multivariate analysis, all three phone types studied showed an increased risk.

Hardell L, Carlberg M, Hansson Mild K. Case-control study on cellular and cordless telephones and the risk for acoustic neuroma or meningioma in patients diagnosed 2000-2003. Neuroepidemiology 25:120-128, 2005.

We performed a case-control study on the use of cellular and cordless telephones and the risk for brain tumors. We report the results for benign brain tumors with data from 413 cases (89% response rate), 305 with meningioma, 84 with acoustic neuroma, 24 with other types and 692 controls (84% response rate). For meningioma, analogue phones yielded odds ratio (OR) = 1.7, 95% confidence interval (CI) = 0.97-3.0, increasing to OR = 2.1, 95% CI = 1.1-4.3 with a >10-year latency period. Also digital cellular phones and cordless phones increased the risk to some extent. For acoustic neuroma, analogue phones gave OR = 4.2, 95% CI = 1.8-10 increasing to OR = 8.4, 95% CI = 1.6-45 with a >15-year latency period, but based on low numbers. Digital phones yielded OR = 2.0, 95% CI = 1.05-3.8, whereas for cordless phones OR was not significantly increased. In the multivariate analysis, analogue phones represented a significant risk factor for acoustic neuroma.

Hardell L, Eriksson M, Carlberg M, Sundstrom C, Mild KH. Use of cellular or cordless telephones and the risk for non-Hodgkin's lymphoma. Int Arch Occup Environ Health. 78(8):625-632, 2005.

Objectives: To evaluate the use of cellular and cordless telephones as the risk factor for

Studies that show **Cell Phone** Health Effects

non-Hodgkin's lymphoma (NHL). Methods: Male and female subjects aged 18-74 years living in Sweden were included during a period from 1 December 1999 to 30 April 2002. Controls were selected from the national population registry. Exposure to different agents was assessed by questionnaire. Results: In total, 910 (91%) cases and 1016 (92%) controls participated. NHL of the B-cell type was not associated with the use of cellular or cordless telephones. Regarding T-cell NHL and >5 year latency period, the use of analogue cellular phones yielded: odds ratio (OR) = 1.46, 95%; confidence interval (CI) = 0.58-3.70, digital: OR=1.92, 95%; CI=0.77-4.80 and cordless phones: OR=2.47; CI=1.09-5.60. The corresponding results for certain, e.g. cutaneous and leukaemia, T-cell lymphoma for analogue phones were: OR=3.41, 95%; CI=0.78-15.0, digital: OR=6.12, 95%; CI=1.26-29.7 and cordless phones: OR=5.48, 95%; CI=1.26-23.9. Conclusions: The results indicate an association between T-cell NHL and the use of cellular and cordless telephones, however based on low numbers and must be interpreted with caution. Regarding B-cell NHL no association was found.

Hardell L, Carlberg M, Hansson Mild K. Use of cellular telephones and brain tumour risk in urban and rural areas. *Occup Environ Med* 62:390-394, 2005.

Aim: To investigate the association between the use of cellular or cordless telephones and the risk for brain tumours in different geographical areas, urban and rural. Methods: patients aged 20-80 years, living in the middle part of Sweden, and diagnosed between 1 January 1997 and 30 June 2000 were included. One control matched for sex and age in five year age groups was selected for each case. Use of different phone types was assessed by a questionnaire. Results: The number of participating cases was 1429; there were 1470 controls. An effect of rural living was most pronounced for digital cellular telephones. Living in rural areas yielded an odd ratio (OR) of 1.4 (95% CI 0.98 to 2.0), increasing to 3.2 (95% CI 1.2 to 8.4) with > 5 year latency time for digital phones. The corresponding Ors for living in urban areas were 0.9 (95% CI 0.8 to 1.2) and 0.9 (95% CI 0.6 to 1.4), respectively. This effect was most obvious for malignant brain tumours. Conclusion: In future studies, place of residence should be considered in assessment of exposure to microwaves from cellular telephone, although the results in this study must be interpreted with caution due to low numbers in some of the calculations.

Hardell L, Mild KH, Sandstrom M, Carlberg M, Hallquist A, Pahlson A. Vestibular schwannoma, tinnitus and cellular telephones. *Neuroepidemiol* 22:124-129, 2003.

Cases with tinnitus after using analogue cellular telephones are presented. An increased odds ratio of 3.45, 95% confidence interval (CI) 1.77-6.76, was found for vestibular schwannoma (VS) associated with the use of analogue cell phones. During the time period 1960-1998, the age-standardized incidence of VS in Sweden significantly increased yearly by +2.53% (CI 1.71-3.35). A significant increase in the incidence of VS was only found for the latter of the two time period 1960-1979 and 1980-1998. For all other brain tumors taken together, the incidence significantly increased yearly by +0.80% (CI 0.59-1.02) for the time period 1960-1998, although the increase was only significant for benign tumors other than VS during 1960-1979.

Studies that show **Cell Phone** Health Effects

Hardell L, Mild KH, Carlsberg M. Case-control study on the use of cellular and cordless phones and the risk for malignant brain tumours. Int. J. Radiat. Biol. 78:931-936, 2002.

Purpose: To investigate the use of cellular and cordless phones and the risk for malignant brain tumours. Materials and Methods: A case-control study was performed on 649 patients aged 20-80 years of both sexes with malignant brain tumour diagnosed from 1 January 1997 to 30 June 2000. All patients were alive during the time of the study and had histopathology verified brain tumours. One matched control to each case was selected from the Swedish Population Register. The study area was the Uppsala-Orebro, Stockholm, Linköping and Göteborg medical regions of Sweden. Results: Exposure was assessed by a questionnaire answered by 588 (91%) cases and 581 (90%) controls. Phone usage was defined as 'ever use' and usage starting 1 year before diagnosis was disregarded. Overall, no significantly increased risks were found: analogue cellular phones yielded an odds ratio (OR) = 1.13, 95% confidence interval (CI) = 0.82-1.57, digital cellular phones OR = 1.13, CI = 0.86-1.48, and cordless phones OR = 1.13, CI = 0.85-1.50. For ipsilateral (same side) radiofrequency exposure, analogue mobile phones gave OR = 1.85, CI = 1.16-2.96, for all malignant brain tumours. For astrocytoma, this risk was OR = 1.95, CI = 1.12-3.39. For all malignant brain tumours, digital mobile phones yielded OR = 1.59, CI = 1.05-2.41, and cordless phones yielded OR = 1.46, CI = 0.96-2.23, in the analysis of ipsilateral exposure. Conclusion: The ipsilateral use of an analogue cellular phone yielded a significantly increased risk for malignant brain tumours.

Hardell L, Hallquist A, Hansson Mild K, Carlberg M, Pahlson A, Lilja A. cellular and cordless telephones and the risk for brain tumours. Europ J Cancer Prevent 11:377-386, 2002.

Microwave exposure from the use of cellular telephones has been discussed in recent years as a potential risk factor for brain tumours. We included in a case-control study 1617 patients aged 20-80 years of both sexes with brain tumour diagnosed between 1 January 1997 and 30 June 2000. They were alive at the study time and had histopathologically verified brain tumour. One matched control to each case was selected from the Swedish Population Register. The study area was the Uppsala-Orebro, Stockholm, Linköping and Göteborg medical regions of Sweden. Exposure was assessed by a questionnaire that was answered by 1429 (88%) cases and 1470 (91%) controls. In total, use of analogue cellular telephones gave an increased risk with an odds ratio (OR) of 1.3 (95% confidence interval (CI) 1.02-1.6). With a tumour induction period of >10 years the risk increased further; OR 1.8 (95% CI 1.1-2.9). No clear association was found for digital or cordless telephones. With regard to the anatomical area of the tumour and exposure to microwaves, the risk was increased for tumours located in the temporal area on the same side of the brain that was used during phone calls; for analogue cellular telephones the OR was 2.5 (95% CI 1.3-4.9). Use of a telephone on the opposite side of the brain was not associated with an increased risk for brain tumours. With regard to different tumour types, the highest risk was for acoustic neuroma (OR 3.5, 95% CI 1.8-6.8) among analogue cellular telephone users.

Hardell L, Nasman A, Pahlson A, Hallquist A. Case-control study on radiology work,

Studies that show **Cell Phone** Health Effects

medical x-ray investigations, and use of cellular telephones as risk factors for brain tumors. Medscape General Medicine May 4, 2000.

Abstract. Context. Ionizing radiation is a well-established risk factor for brain tumors. During recent years, microwave exposure from the use of cellular telephones has been discussed as a potential risk factor. **Objective.** To determine risk factors for brain tumors. **Design.** A case-control study, with exposure assessed by questionnaires. **Participants.** A total of 233 currently living men and women, aged 20 to 80 years, were included. The case patients had histopathologically verified brain tumors and lived in the Uppsala-Orebro region (1994-1996) or the Stockholm region (1995-1996). Two matched controls to each case were selected from the Swedish Population Register. **Main Outcome Measures.** Ionizing radiation and use of cellular telephones as risk factors for brain tumors. **Results.** A total of 209 cases (90%) and 425 controls (91%) answered the questionnaire. Work as a physician yielded an odds ratio (OR) of 6.00, with a 95% confidence interval (CI) of 0.62 to 57.7. All three case patients had worked with fluoroscopy. Radiotherapy of the head and neck region yielded an OR of 3.61 (95% CI, 0.65-19.9). Medical diagnostic x-ray examination of the same area yielded an OR of 2.10 (95% CI, 1.25-3.53), with a tumor induction period of 5 years or more. Chemical industry work yielded an OR of 4.10 (95% CI, 1.25-13.4), and laboratory work yielded an OR of 3.21 (95% CI, 1.16-8.85). Ipsilateral use of cellular telephones increased the risk for tumors in the temporal, temporoparietal, and occipital lobes (OR, 2.42; 95% CI, 0.97-6.05), ie, the anatomic areas with highest exposure to microwaves from a mobile telephone. The result was further strengthened (OR, 2.62; 95% CI, 1.02-6.71) in a multivariate analysis that included laboratory work and medical diagnostic x-ray investigations of the head and neck. **Conclusion.** Exposure to ionizing radiation, work in laboratories, and work in the chemical industry increased the risk of brain tumors. Use of a cellular telephone was associated with an increased risk in the anatomic area with highest exposure.

Di Carlo A, White N, Guo F, Garrett P, Litovitz T. Chronic electromagnetic field exposure decreases HSP70 levels and lowers cytoprotection. J. Cell. Biochem. 84: 447-454, 2002.

Electromagnetic field (EMF) exposures have been shown to induce heat shock proteins (HSPs), which help to maintain the conformation of cellular proteins during periods of stress. We have previously reported that short-term exposure of chick embryos to either 60 Hz (extremely low frequency: ELF), or radio-frequency (RF: 915 MHz) EMFs induce protection against hypoxia. Experiments presented in the current report are based on a study in which long-term (4 days), continuous exposure to ELF-EMFs decreased protection against ultraviolet radiation. Based on this result, it was hypothesized that de-protection against hypoxia should also occur following long-term, continuous, or daily, repeated exposures to EMFs. To test this hypothesis, chick embryos were exposed to ELF-EMFs (8 μ T) continuously for 4 days, or to ELF or RF (3.5 mW incident power)- EMFs repeated daily (20, 30, or 60 min once or twice daily for 4 days). Several of the exposure protocols yielded embryos that had statistically significant decreases in protection against hypoxic stress (continuous and 30 or 60 min ELF twice

Studies that show **Cell Phone** Health Effects

daily; or 30 or 60 min once daily RF). This is consistent with our finding that following 4 days of ELF-EMF exposure, HSP70 levels decline by 27% as compared to controls. In addition, the superposition of ELF-EM noise, previously shown to minimize ELF-EMF induced hypoxia protection, inhibited hypoxia de-protection caused by long term, continuous ELF or daily, repeated RF exposures. This EMF-induced decrease in HSP70 levels and resulting decline in cytoprotection suggests a mechanism by which daily exposure (such as might be experienced by mobile phone users) could enhance the probability of cancer and other diseases.

Hardell, L, Nasman, A, Pahlson, A, Hallquist, A, Hansson Mild, K, Use of cellular telephones and the risk for brain tumours: A case-control study. Int J Oncol 15(1):113-116, 1999.

The use of cellular telephones has increased dramatically during the 1990's in the world. In the 1980's the analogue NMT system was used whereas the digital GSM system was introduced in early 1990's and is now the preferred system. Case reports of brain tumours in users initiated this case-control study on brain tumours and use of cellular telephones. Also other exposures were assessed. All cases, both males and females, with histopathologically verified brain tumour living in Uppsala-Orebro region (1994-96) and Stockholm region (1995-96) aged 20-80 at the time of diagnosis and alive at start of the study were included, 233 in total. Two controls to each case were selected from the Swedish Population Register matched for sex, age and study region. Exposure was assessed by questionnaires supplemented over the phone. The analyses were based on answers from 209 (90%) cases and 425 (91%) controls. Use of cellular telephone gave odds ratio (OR) = 0.98 with 95% confidence interval (CI) = 0.69-1.41. For the digital GSM system OR = 0.97, CI = 0.61-1.56 and for the analogue NMT system OR = 0.94, CI = 0.62-1.44 were calculated. Dose-response analysis and using different tumour induction periods gave similar results. Non-significantly increased risk was found for tumour in the temporal or occipital lobe on the same side as a cellular phone had been used, right side OR = 2.45, CI = 0.78-7.76, left side OR = 2.40, CI = 0.52-10.9 Increased risk was found only for use of the NMT system. For GSM use the observation time is still too short for definite conclusions. An increased risk for brain tumour in the anatomical area close to the use of a cellular telephone should be especially studied in the future.

Hardell L, Carlberg M. Mobile phones, cordless phones and the risk for brain tumours. Int J Oncol. 35(1):5-17, 2009.

The Hardell-group conducted during 1997-2003 two case control studies on brain tumours including assessment of use of mobile phones and cordless phones. The questionnaire was answered by 905 (90%) cases with malignant brain tumours, 1,254 (88%) cases with benign tumours and 2,162 (89%) population-based controls. Cases were reported from the Swedish Cancer Registries. Anatomical area in the brain for the tumour was assessed and related to side of the head used for both types of wireless phones. In the current analysis we defined ipsilateral use (same side as the tumour) as $\geq 50\%$ of the use and contralateral use (opposite side) as $< 50\%$ of the calling time. We report now further results for use of mobile and cordless phones. Regarding

Studies that show **Cell Phone** Health Effects

astrocytoma we found highest risk for ipsilateral mobile phone use in the >10 year latency group, OR=3.3, 95% CI=2.0-5.4 and for cordless phone use OR=5.0, 95% CI=2.3-11. In total, the risk was highest for cases with first use <20 years age, for mobile phone OR=5.2, 95% CI=2.2-12 and for cordless phone OR=4.4, 95% CI=1.9-10. For acoustic neuroma, the highest OR was found for ipsilateral use and >10 year latency, for mobile phone OR=3.0, 95% CI=1.4-6.2 and cordless phone OR=2.3, 95% CI=0.6-8.8. Overall highest OR for mobile phone use was found in subjects with first use at age <20 years, OR=5.0, 95% CI 1.5-16 whereas no association was found for cordless phone in that group, but based on only one exposed case. The annual age-adjusted incidence of astrocytoma for the age group >19 years increased significantly by +2.16%, 95% CI +0.25 to +4.10 during 2000-2007 in Sweden in spite of seemingly underreporting of cases to the Swedish Cancer Registry. A decreasing incidence was found for acoustic neuroma during the same period. However, the medical diagnosis and treatment of this tumour type has changed during recent years and underreporting from a single center would have a large impact for such a rare tumour.

Hardell L, Hansson Mild K, Pahlson A, Hallquist A, Ionizing radiation, cellular telephones and the risk of brain tumours. *Europ J Cancer Prevent* 10:523-529, 2001.

A case-control study on brain tumours included 233 patients aged 20-80 years and alive at the study time. They had histopathologically verified brain tumour and lived in the Upsala-Orebro region (1994-1996) or in Stockholm region (1995-1996). Two matched controls to each case were selected from the Swedish Population Register. Two hundred and nine cases (90%) and 425 controls (91%) answered the questionnaire. Results are presented for the whole study group, as given here, and for malignant and benign tumours separately. For workers in the chemical industry the odds ratio (OR) was 4.10, 95% confidence interval (95% CI) 1.25-13.4 and laboratory workers OR 3.21, 95%CI 1.16-8.85. Radiotherapy of the head and neck region gave OR 3.61, 95% CI 0.65-19.9. Medical diagnostic X-ray of the same area yielded OR 1.64, 95% CI 1.04-2.58. Work as a physician gave OR 6.00, 95% CI 0.62-57.7. All three cases had worked with fluoroscopy. Ipsilateral (same side) use of a cellular telephone increased the risk of tumours in the temporal, temporo-parietal and occipital areas, with OR 2.42, 95% CI 0.97-6.05 (i.e., the anatomical areas with highest exposure to microwaves from a mobile phone).

Hardell, L, Reizenstein, J, Johansson, B, Gertzen, H, Mild, KH, Angiosarcoma of the scalp and use of a cordless (portable) telephone. *Epidemiology* 10(6):785-786, 1999.

This is a case study of a 57-year old woman who was diagnosed in January 1999 with an angiosarcoma of the scalp on the left side of her head (2 cm above her ear). Angiosarcoma is a rare type of soft tissue sarcoma (STS). Since 1988, the patient had, on a regular basis (1 hr per day), used a cordless (portable) phone, always using her left ear. Starting in 1994, she also used a GSM (digital) mobile phone on her left ear (a few minutes per week). The exposure to microwaves from a cordless phone in this patient is of interest because she had exposure on a daily basis starting 10 years before the first clinical signs of her angiosarcoma. Moreover, the tumor developed in the anatomical area with the highest exposure to microwaves on the same side of the head that she

Studies that show **Cell Phone** Health Effects

had used the phone. She also reported no exposure to other know risk factors for STS.

Leszczynski D, Joenväärä S, Reivinen J, Kuokka R, Non-thermal activation of the hsp27/p38MAPK stress pathway by mobile phone radiation in human endothelial cells: Molecular mechanism for cancer- and blood-brain barrier-related effects.

Differentiation 70:120–129, 2002.

Abstract We have examined whether non-thermal exposures of cultures of the human endothelial cell line EA.hy926 to 900 MHz GSM mobile phone microwave radiation could activate stress response. Results obtained demonstrate that 1-hour non-thermal exposure of EA.hy926 cells changes the phosphorylation status of numerous, yet largely unidentified, proteins. One of the affected proteins was identified as heat shock protein-27 (hsp27). Mobile phone exposure caused a transient increase in phosphorylation of hsp27, an effect which was prevented by SB203580, a specific inhibitor of p38 mitogen-activated protein kinase (p38MAPK). Also, mobile phone exposure caused transient changes in the protein expression levels of hsp27 and p38MAPK. All these changes were non-thermal effects because, as determined using temperature probes, irradiation did not alter the temperature of cell cultures, which remained throughout the irradiation period at $37\pm0.3^{\circ}\text{C}$. Changes in the overall pattern of protein phosphorylation suggest that mobile phone radiation activates a variety of cellular signal transduction pathways, among them the hsp27/p38MAPK stress response pathway. Based on the known functions of hsp27, we put forward the hypothesis that mobile phone radiation-induced activation of hsp27 may (i) facilitate the development of brain cancer by inhibiting the cytochrome c/caspase-3 apoptotic pathway and (ii) cause an increase in blood-brain barrier permeability through stabilization of endothelial cell stress fibers. We postulate that these events, when occurring repeatedly over a long period of time, might become a health hazard because of the possible accumulation of brain tissue damage. Furthermore, our hypothesis suggests that other brain damaging factors may co-participate in mobile phone radiation-induced effects.

Hansson Mild K, Hardell L, Carlberg M. Pooled analysis of two Swedish case-control studies on the use of mobile and cordless telephones and the risk of brain tumours diagnosed during 1997-2003. *Int J Occup Saf Ergon* 13(1):63-71, 2007.

Here we present the pooled analysis of 2 case-control studies on the association of brain tumours with mobile phone use. Use of analogue cellular phones increased the risk for acoustic neuroma by 5%, 95% confidence interval (CI) = 2-9% per 100 hrs of use. The risk increased for astrocytoma grade III-IV with latency period with highest estimates using >10-year time period from first use of these phone types. The risk increased per one year of use of analogue phones by 10%, 95% CI = 6-14%, digital phones by 11%, 95% CI = 6-16%, and cordless phones by 8%, 95% CI = 5-12%. For all studied phone types OR for brain tumours, mainly acoustic neuroma and malignant brain tumours, increased with latency period, especially for astrocytoma grade III-IV.

Lourencini da Silva R , Albano F, Lopes dos Santos LR , Tavares AD, Felzenszwalb I, The effect of electromagnetic field exposure on the formation of DNA lesions. *Redox Rep*

Studies that show **Cell Phone** Health Effects

5(5):299-301, 2000.

In an attempt to determine whether electromagnetic field (EMF) exposure might lead to DNA damage, we exposed SnCl₂-treated pBR322 plasmids to EMF and analysed the resulting conformational changes using agarose gel electrophoresis. An EMF-dependent potentiation of DNA scission (i.e. the appearance of relaxed plasmids) was observed. In confirmation of this, plasmids pre-exposed to EMF also were less capable of transforming *Escherichia coli*. The results indicate that EMF, in the presence of a transition metal, is capable of causing DNA damage. These observations support the idea that EMF, probably through secondary generation of reactive oxygen species, can be clastogenic and provide a possible explanation for the observed correlation between EMF exposure and the frequency of certain types of cancers in humans.

Muscat JE, Malkin MG, Shore RE, Thompson S, Neugut AL, Stellman SD, Bruce J. Handheld cellular telephones and risk of acoustic neuroma. *Neurology* 58:1304-1306, 2002.

The hypothesis that intracranial energy deposition from handheld cellular telephones causes acoustic neuroma was tested in an epidemiologic study of 90 patients and 86 control subjects. The relative risk was 0.9 ($p = 0.07$) and did not vary significantly by the frequency, duration, and lifetime hours of use. In patients who used cellular telephones, the tumor occurred more often on the contralateral than ipsilateral side of the head. Further efforts should focus on potentially longer induction periods.

Mashevich M, Folkman D, Kesar A, Barbul A, Korenstein R, Jerby E, Avivi L, Exposure of human peripheral blood lymphocytes to electromagnetic fields associated with cellular phones leads to chromosomal instability. *Bioelectromagnetics* 24:82-90, 2003.

Whether exposure to radiation emitted from cellular phones poses a health hazard is at the focus of current debate. We have examined whether in vitro exposure of human peripheral blood lymphocytes (PBL) to continuous 830 MHz electromagnetic fields causes losses and gains of chromosomes (aneuploidy), a major "somatic mutation" leading to genomic instability and thereby to cancer. PBL were irradiated at different average absorption rates (SAR) in the range of 1.6-8.8 W/kg for 72 hr in an exposure system based on a parallel plate resonator at temperatures ranging from 34.5-37.5 °C. The averaged SAR and its distribution in the exposed tissue culture flask were determined by combining measurements and numerical analysis based on a finite element simulation code. A linear increase in chromosome 17 aneuploidy was observed as a function of the SAR value, demonstrating that this radiation has a genotoxic effect. The SAR dependent aneuploidy was accompanied by an abnormal mode of replication of the chromosome 17 region engaged in segregation (repetitive DNA arrays associated with the centromere), suggesting that epigenetic alterations are involved in the SAR dependent genetic toxicity. Control experiments (i.e., without any RF radiation) carried out in the temperature range of 34.5-38.5 °C showed that elevated temperature is not associated with either the genetic or epigenetic alterations observed following RF radiation - the increased levels of aneuploidy and the modification in replication of the centromeric DNA arrays. These findings indicate that the genotoxic effect of the

Studies that show **Cell Phone** Health Effects

electromagnetic radiation is elicited via a non-thermal pathway. Moreover, the fact that aneuploidy is a phenomenon known to increase the risk for cancer, should be taken into consideration in future evaluation of exposure guidelines.

Akhavan-Sigari R, Baf MM, Ariabod V, Rohde V, Rahighi S. Connection between Cell Phone use, p53 Gene Expression in Different Zones of Glioblastoma Multiforme and Survival Prognoses. Rare Tumors. 2014 Aug 8;6(3):5350. doi: 10.4081/rt.2014.5350.

The aim of this paper is to investigate p53 gene expression in the central and peripheral zones of glioblastoma multiforme using a real-time reverse transcription polymerase chain reaction (RT-PCR) technique in patients who use cell phones ≥ 3 hours a day and determine its relationship to clinicopathological findings and overall survival. Sixty-three patients (38 males and 25 females), diagnosed with glioblastoma multiforme (GBM), underwent tumor resection between 2008 and 2011. Patient ages ranged from 25 to 88 years, with a mean age of 55. The levels of expression of p53 in the central and peripheral zone of the GBM were quantified by RT-PCR. Data on p53 gene expression from the central and peripheral zone, the related malignancy and the clinicopathological findings (age, gender, tumor location and size), as well as overall survival, were analyzed. Forty-one out of 63 patients (65%) with the highest level of cell phone use (≥ 3 hours/day) had higher mutant type p53 expression in the peripheral zone of the glioblastoma; the difference was statistically significant ($P=0.034$). Results from the present study on the use of mobile phones for ≥ 3 hours a day show a consistent pattern of increased risk for the mutant type of p53 gene expression in the peripheral zone of the glioblastoma, and that this increase was significantly correlated with shorter overall survival time. The risk was not higher for ipsilateral exposure. We found that the mutant type of p53 gene expression in the peripheral zone of the glioblastoma was increased in 65% of patients using cell phones ≥ 3 hours a day.

Ozgur E, Guler G, Kismali G, Seyhan N. Mobile Phone Radiation Alters Proliferation of Hepatocarcinoma Cells. Cell Biochem Biophys. 2014 May 11. [Epub ahead of print]

This study investigated the effects of intermittent exposure (15 min on, 15 min off for 1, 2, 3, or 4 h, at a specific absorption rate of 2 W/kg) to enhanced data rates for global system for mobile communication evolution-modulated radiofrequency radiation (RFR) at 900- and 1,800-MHz frequencies on the viability of the Hepatocarcinoma cells (Hep G2). Hep G2 cell proliferation was measured by a colorimetric assay based on the cleavage of the tetrazolium salt WST-1 by mitochondrial dehydrogenases in viable cells. Cell injury was evaluated by analyzing the levels of lactate dehydrogenase (LDH) and glucose released from lysed cells into the culture medium. Morphological observation of the nuclei was carried out by 4',6-diamidino-2-phenylindole (DAPI) staining using fluorescence microscopy. In addition, TUNEL assay was performed to confirm apoptotic cell death. It was observed that cell viability, correlated with the LDH and glucose levels, changed according to the frequency and duration of RFR exposure. Four-hour exposure produced more pronounced effects than the other exposure durations. 1,800-MHz RFR had a larger impact on cell viability and Hep G2 injury than the RFR at 900 MHz. Morphological observations also supported the biochemical results indicating that most

Studies that show **Cell Phone** Health Effects

of the cells showed irregular nuclei pattern determined by using the DAPI staining, as well as TUNEL assay which shows DNA damage especially in the cells after 4 h of exposure to 1,800-MHz RFR. Our results indicate that the applications of 900- and 1,800-MHz (2 W/kg) RFR cause to decrease in the proliferation of the Hep G2 cells after 4 h of exposure. Further studies will be conducted on other frequency bands of RFR and longer duration of exposure.

Zeng QL, Weng Y, Chen GD, Lu DQ, Chiang H, Xu ZP [Effects of GSM 1800 MHz radiofrequency electromagnetic fields on protein expression profile of human breast cancer cell MCF-7.] Zhonghua Yu Fang Yi Xue Za Zhi. 40(3):153-158, 2006. [Article in Chinese]

OBJECTIVE: To study the effects of GSM 1800 MHz radiofrequency electromagnetic fields (RF EMF) exposure on protein expression profile of human breast cancer cell line (MCF-7), as to exploring the possible effects on normal cell physiological function.
METHODS: MCF-7 cells were continuously or intermittently (5 minutes field on followed by 10 minutes off) exposed to RF EMF for different duration (1 hour, 3 hours, 6 hours, 12 hours, or 24 hours) at an average specific absorption rate (SAR) of 3.5 W/kg. The extracted proteins were separated by 2-dimensional electrophoresis and the protein-spot distribution of the silver-stained gels was analyzed by using PDQuest software 7.1. Each experiment was repeated three times. **RESULTS:** On the average, around 1100 proteins were detected using pH 4 - 7 IPG strip. There were no differential proteins found under continuous exposure at SAR of 3.5 W/kg for 6 hours. Under other exposure conditions, we found various differentially expressed proteins in exposure groups as compared with the sham-exposed controls. Especially in 3 hours intermittent exposure and 12 hours continuous exposure, eighteen and seven differential proteins were detected, respectively. The categories and functions of these differentially expressed proteins were analyzed by searching of SWISS-PROT protein database, which suggested that these proteins should be related to the functions of biosynthesis, signal transduction, and DNA damage and repair. **CONCLUSIONS:** Data indicated that the protein expression changes induced by RF radiation might depend on exposure duration and mode. Many biological processes might be affected by RF exposure.

Marinelli F, La Sala D, Ciccio G, Cattini L, Trimarchi C, Putti S, Zamparelli A, Giuliani L, Tomassetti G, Cinti C. Exposure to 900 MHz electromagnetic field induces an unbalance between pro-apoptotic and pro-survival signals in T-lymphoblastoid leukemia CCRF-CEM cells. J Cell Physiol. 198(2):324-332, 2004.

It has been recently established that low-frequency electromagnetic field (EMFs) exposure induces biological changes and could be associated with increased incidence of cancer, while the issue remains unresolved as to whether high-frequency EMFs can have hazardous effect on health. Epidemiological studies on association between childhood cancers, particularly leukemia and brain cancer, and exposure to low- and high-frequency EMF suggested an etiological role of EMFs in inducing adverse health effects. To investigate whether exposure to high-frequency EMFs could affect in vitro cell survival, we cultured acute T-lymphoblastoid leukemia cells (CCRF-CEM) in the

Studies that show **Cell Phone** Health Effects

presence of unmodulated 900 MHz EMF, generated by a transverse electromagnetic (TEM) cell, at various exposure times. We evaluated the effects of high-frequency EMF on cell growth rate and apoptosis induction, by cell viability (MTT) test, FACS analysis and DNA ladder, and we investigated pro-apoptotic and pro-survival signaling pathways possibly involved as a function of exposure time by Western blot analysis. At short exposure times (2-12 h), unmodulated 900 MHz EMF induced DNA breaks and early activation of both p53-dependent and -independent apoptotic pathways while longer continuous exposure (24-48 h) determined silencing of pro-apoptotic signals and activation of genes involved in both intracellular (Bcl-2) and extracellular (Ras and Akt1) pro-survival signaling. Overall our results indicate that exposure to 900 MHz continuous wave, after inducing an early self-defense response triggered by DNA damage, could confer to the survivor CCRF-CEM cells a further advantage to survive and proliferate.

Kundi M. The controversy about a possible relationship between mobile phone use and cancer. Cien Saude Colet. 15(5):2415-2430, 2010.

Over the last decade, mobile phone use increased to almost 100% prevalence in many countries. Evidence for potential health hazards accumulated in parallel by epidemiologic investigations has raised controversies about the appropriate interpretation and the degree of bias and confounding responsible for reduced or increased risk estimates. Overall, 33 epidemiologic studies were identified in the peer-reviewed literature, mostly (25) about brain tumors. Methodologic considerations revealed that three important conditions for epidemiologic studies to detect an increased risk are not met: no evidence-based exposure metric is available; the observed duration of mobile phone use is generally still too low; no evidence-based selection of end points among the grossly different types of neoplasias is possible because of lack of etiologic hypotheses. The overall evidence speaks in favor of an increased risk, but its magnitude cannot be assessed at present because of insufficient information on long-term use.

Repacholi, MH, Basten, A, Gebiski, V, Noonan, D, Finnie, J, Harris, AW, Lymphomas in E mu-Pim1 transgenic mice exposed to pulsed 900 MHz electromagnetic fields. Radiat Res 147(5):631-640, 1997.

Whether radiofrequency (RF) fields are carcinogenic is controversial; epidemiological data have been inconclusive and animal tests limited. The aim of the present study was to determine whether long-term exposure to pulse-modulated RF fields similar to those used in digital mobile telecommunications would increase the incidence of lymphoma in E mu-Pim1 transgenic mice, which are moderately predisposed to develop lymphoma spontaneously. One hundred female E mu-Pim1 mice were sham-exposed and 101 were exposed for two 30-min periods per day for up to 18 months to plane-wave fields of 900 MHz with a pulse repetition frequency of 217 Hz and a pulse width of 0.6 ms. Incident power densities were 2.6-13 W/m² and specific absorption rates were 0.008-4.2 W/kg, averaging 0.13-1.4 W/kg. Lymphoma risk was found to be significantly higher in the exposed mice than in the controls (OR = 2.4. P = 0.006, 95% CI = 1.3-4.5). Follicular

Studies that show **Cell Phone** Health Effects

lymphomas were the major contributor to the increased tumor incidence. Thus long-term intermittent exposure to RF fields can enhance the probability that mice carrying a lymphomagenic oncogene will develop lymphomas. We suggest that such genetically cancer-prone mice provide an experimental system for more detailed assessment of dose-response relationships for risk of cancer after RF-field exposure.

Hruby R, Neubauer G, Kuster N, Frauscher M Study on potential effects of "902-MHz GSM-type Wireless Communication Signals" on DMBA-induced mammary tumours in Sprague-Dawley rats. *Mutat Res.* 649(1-2):34-44, 2008.

The aim of the study was to detect whether long-term exposure to "902-MHz GSM-type Wireless Communication Signals" ("radio-frequency (RF)-exposure") would affect 7,12-dimethylbenz(a)anthracene (DMBA)-induced mammary tumours in female Sprague-Dawley rats. Five hundred female rats were each given a single oral dose of 17mg DMBA per kg body weight (bw) at an age of 46-48 days. Three groups of 100 animals each were RF-exposed (902MHz; crest factor 8; pulse width=0.57ms) from the next day onwards to normal whole-body averaged doses (expressed as specific absorption rate, SAR) of 0.4, 1.3 or 4.0W/kg bw (low/mid/high-dose group) for 4h/d, 5d/week, during 6 months. A sham-exposed and a cage-control group remained without RF-exposure ($<0.01\text{mW/kg}$). Animals were weekly weighed and palpated for mammary tumours; all mammary glands were examined histopathologically. There were several statistically significant differences between RF-exposed groups and the sham-exposed group, as follows: All RF-exposed groups had, at different times, significantly more palpable tissue masses. There were fewer animals with benign neoplasms, but more with malignant tumours in the high-dose group. In addition, there were more adenocarcinomas in the low-dose group, more malignant neoplasms in the low- and high-dose groups, more animals with adenocarcinomas in the high-dose group, and fewer animals with fibroadenomas in the low- and mid-dose groups. The cage-control group had, when compared with the sham-exposed group, statistically significantly more palpable tissue masses, more benign and also more malignant neoplasms. The cage-control group had in most aspects the highest incidence and malignancy of neoplasms among all groups. None of the above findings in RF-exposed animals produced a clear dose-response relation and the responses of the cage-control group were either similar to or stronger than those of any of the RF-exposed group. The significant differences between the sham-exposed animals and one or more RF-exposed groups may be interpreted as evidence of an effect of RF-exposure. In the context of the results of the cage-control group, in the light of controversial results reported in the literature, and given the fact that the DMBA-mammary tumour model is known to be prone to high variations in the results, it is the authors' opinion that the differences between the groups are rather incidental ones.

Yang L, Hao D, Wang M, Zeng Y, Wu S, Zeng Y. Cellular Neoplastic Transformation Induced by 916 MHz Microwave Radiation. *Cell Mol Neurobiol.*32(6):1039-1046, 2012.

There has been growing concern about the possibility of adverse health effects resulting from exposure to microwave radiations, such as those emitted by mobile phones. The purpose of this study was to investigate the cellular neoplastic transformation effects of

Studies that show **Cell Phone** Health Effects

electromagnetic fields. 916 MHz continuous microwave was employed in our study to simulate the electromagnetic radiation of mobile phone. NIH/3T3cells were adopted in our experiment due to their sensitivity to carcinogen or cancer promoter in environment. They were divided randomly into one control group and three microwave groups. The three microwave groups were exposed to 916 MHz EMF for 2 h per day with power density of 10, 50, and 90 w/m², respectively, in which 10 w/m² was close to intensity near the antenna of mobile phone. The morphology and proliferation of NIH/3T3cells were examined and furthermore soft agar culture and animal carcinogenesis assay were carried out to determine the neoplastic promotion. Our experiments showed NIH/3T3cells changed in morphology and proliferation after 5-8 weeks exposure and formed clone in soft agar culture after another 3-4 weeks depending on the exposure intensity. In the animal carcinogenesis study, lumps developed on the back of SCID mice after being inoculated into exposed NIH/3T3cells for more than 4 weeks. The results indicate that microwave radiation can promote neoplastic transformation of NIH/3T3cells.

Trivino Pardo JC, Grimaldi S, Taranta M, Naldi I, Cinti C. Microwave electromagnetic field regulates gene expression in T-lymphoblastoid leukemia CCRF-CEM cell line exposed to 900 MHz. *Electromagn Biol Med.* 31(1):1-18, 2012.

Electric, magnetic, and electromagnetic fields are ubiquitous in our society, and concerns have been expressed regarding possible adverse effects of these exposures. Research on Extremely Low-Frequency (ELF) magnetic fields has been performed for more than two decades, and the methodology and quality of studies have improved over time. Studies have consistently shown increased risk for childhood leukemia associated with ELF magnetic fields. There are still inadequate data for other outcomes. More recently, focus has shifted toward Radio Frequencies (RF) exposures from mobile telephony. There are no persuasive data suggesting a health risk, but this research field is still immature with regard to the quantity and quality of available data. This technology is constantly changing and there is a need for continued research on this issue. To investigate whether exposure to high-frequency electromagnetic fields (EMF) could induce adverse health effects, we cultured acute T-lymphoblastoid leukemia cells (CCRF-CEM) in the presence of 900 MHz MW-EMF generated by a transverse electromagnetic (TEM) cell at short and long exposure times. We evaluated the effect of high-frequency EMF on gene expression and we identified functional pathways influenced by 900 MHz MW-EMF exposure.

Hartikka H, Heinävaara S, Mäntylä R, Kähärä V, Kurtio P, Auvinen A. Mobile phone use and location of glioma: A case-case analysis. *Bioelectromagnetics.* 30(3):176-182, 2009.

We assessed a new approach for evaluating the glioma risk among users of mobile phones to focus on the part of the brain most heavily exposed to radiofrequency electromagnetic fields from mobile phones. The tumor midpoint was defined from radiological imaging. A case-case analysis with 99 gliomas was performed using logistic regression. The exposed cases were those with the tumor mid-point within 4.6 cm from

Studies that show **Cell Phone** Health Effects

the line between the mouth and the external meatus of the ear, representing the most likely location of the mobile phone (the source of exposure). Alternative analyses based on various indicators of mobile phone use as the outcome were also carried out. The majority of cases were regular mobile phone users. A slightly higher proportion of gliomas among mobile phone users than non-users occurred within 4.6 cm from the presumed location of the mobile phone (28% vs. 14%). Modestly elevated odds ratios were observed for several indicators of mobile phone use, but without an exposure gradient. The highest odds ratios were found for contralateral and short-term use. Our results, though limited by the small sample size, demonstrate that detailed information on tumor location allows evaluation of the risk related to the most heavily exposed part of the brain, representing direct evaluation of the possible local carcinogenic effects of the radiofrequency fields. However, field strength varies between users and over time also within a given anatomic site, due to the output power of the phone. Collaborative analysis of a larger sample is planned.

Salahaldin AH, Bener A. Long-term and frequent cellular phone use and risk of acoustic neuroma. *Int Tinnitus J.* 12(2):145-148, 2006.

Human exposure to radio frequency radiation has increased dramatically during recent years from widespread use of mobile phones, and in some studies this exposure has been linked to the development of acoustic neuroma. The aim of our study was to describe the epidemiology of acoustic neuroma in a newly developed country, Qatar. We reviewed all cases of acoustic neuroma registered at the Hamad Medical Corporation during the period 2004-2005. We collected and assessed the sociodemographic information, presenting complaints, audiological evaluation, and laboratory investigations. During the study period, we diagnosed acoustic neuroma in 13 patients (10 women, median age 55 years, and 3 men, median age 49 years). Most of the cell phones were used daily for an average of 14 times per day (range, 8-20 times) and had been used for the duration of more than 5 years, with the exception of 3 patients who had used the cell phone excessively (> 20 minutes per call more than five times daily) owing to the nature of their jobs. The total incidence rate for Qatar was found to be 17.2 per million population. In conclusion, the incidence of acoustic neuroma in Qatar is slightly higher than that in other countries. Despite the presence of facilities in Qatar, no proper screening and management protocol is available. This study highlights the need for the development and implementation of a national registry plan whereby effective care services can be delivered and high-risk groups can be targeted.

de Vocht F, Hannam K, Buchan I. Environmental risk factors for cancers of the brain and nervous system: the use of ecological data to generate hypotheses. *Occup Environ Med.* 2013 Jan 23. [Epub ahead of print]

BACKGROUND: There is a public health need to balance timely generation of hypotheses with cautious causal inference. For rare cancers this is particularly challenging because standard epidemiological study designs may not be able to elucidate causal factors in an early period of newly emerging risks. Alternative methodologies need to be considered for generating and shaping hypotheses prior to definitive investigation. OBJECTIVES: To

Studies that show **Cell Phone** Health Effects

evaluate whether open-access databases can be used to explore links between potential risk factors and cancers at an ecological level, using the case study of brain and nervous system cancers as an example. **METHODS:** National age-adjusted cancer incidence rates were obtained from the GLOBOCAN 2008 resource and combined with data from the United Nations Development Report and the World Bank list of development indicators. Data were analysed using multivariate regression models. **RESULTS:** Cancer rates, potential confounders and environmental risk factors were available for 165 of 208 countries. 2008 national incidences of brain and nervous system cancers were associated with continent, gross national income in 2008 and Human Development Index Score. The only exogenous risk factor consistently associated with higher incidence was the penetration rate of mobile/cellular telecommunications subscriptions, although other factors were highlighted. According to these ecological results the latency period is at least 11-12 years, but probably more than 20 years. Missing data on cancer incidence and for other potential risk factors prohibit more detailed investigation of exposure-response associations and/or explore other hypotheses. **CONCLUSIONS:** Readily available ecological data may be underused, particularly for the study of risk factors for rare diseases and those with long latencies. The results of ecological analyses in general should not be overinterpreted in causal inference, but equally they should not be ignored where alternative signals of aetiology are lacking.

Czyz J, GuanK, ZengQ, NikolovaT, MeisterA, SchönbornF, SchudererJ, KusterN, WobusAM, High frequency electromagnetic fields (GSM signals) affect gene expression levels in tumor suppressor p53-deficient embryonic stem cells. Bioelectromagnetics 25:296-307, 2004.

Effects of electromagnetic fields (EMF) simulating exposure to the Global System for Mobile Communications (GSM) signals were studied using pluripotent embryonic stem (ES) cells in vitro. Wild-type ES cells and ES cells deficient for the tumor suppressor p53 were exposed to pulse modulated EMF at 1.71 GHz, lower end of the uplink band of GSM 1800, under standardized and controlled conditions, and transcripts of regulatory genes were analyzed during in vitro differentiation. Two dominant GSM modulation schemes (GSM-217 and GSM-Talk), which generate temporal changes between GSM-Basic (active during talking phases) and GSM-DTX (active during listening phases thus simulating a typical conversation), were applied to the cells at and below the basic safety limits for local exposures as defined for the general public by the International Commission on Nonionizing Radiation Protection (ICNIRP). GSM-217 EMF induced a significant upregulation of mRNA levels of the heat shock protein, hsp70 of p53-deficient ES cells differentiating in vitro, paralleled by a low and transient increase of c-jun, c-myc, and p21 levels in p53-deficient, but not in wild-type cells. No responses were observed in either cell type after EMF exposure to GSM-Talk applied at similar slot-averaged specific absorption rates (SAR), but at lower time-averaged SAR values. Cardiac differentiation and cell cycle characteristics were not affected in embryonic stem and embryonic carcinoma cells after exposure to GSM-217 EMF signals. Our data indicate that the genetic background determines cellular responses to GSM modulated EMF.

Studies that show **Cell Phone** Health Effects

Brautbar N. Rapid development of brain tumors in 2 cellular phone testers. Arch Environ Health. 58(9):606, 2003.

Two case reports. Comment on [Richter ED, Berman T, Levy O. Brain cancer with induction periods of less than 10 years in young military radar workers. Arch Environ Health. 2002 Jul-Aug;57(4):270-272.]

Auvinen A, Hietanen M, Luukkonen R, Koskela R-S, Brain tumors and salivary gland cancers among cellular telephone users Epidemiology 13:356-359, 2002.

Background. Possible risk of cancer associated with use of cellular telephones has lately been a subject of public debate. Methods. We conducted a register-based, case-control study on cellular phone use and cancer. The study subjects were all cases of brain tumor (N = 398) and salivary gland cancer (N = 34) diagnosed in Finland in 1996, with five controls per case. Results. Cellular phone use was not associated with brain tumors or salivary gland cancers overall, but there was a weak association between gliomas and analog cellular phones. Conclusions. A register-based approach has limited value in risk assessment of cellular phone use owing to lack of information on exposure.

Berg G, Spallek J, Schuz J, Schlehofer B, Bohler E, Schlaefer K, Hettinger I, Kunna-Grass K, Wahrendorf J, Blettner M. Occupational exposure to radio frequency/microwave radiation and the risk of brain tumors: Interphone Study Group, Germany. Am J Epidemiol.164(6):538-548, 2006.

It is still under debate whether occupational exposure to radio frequency/microwave electromagnetic fields (RF/MW-EMF) contributes to the development of brain tumors. This analysis examined the role of occupational RF/MW-EMF exposure in the risk of glioma and meningioma. A population-based, case-control study including 381 meningioma cases, 366 glioma cases, and 1,494 controls aged 30-69 years was performed in three German regions in 2000-2003. An exposure matrix for occupational activity was constructed by using information on RF/MW-EMF exposure collected in a computer-assisted personal interview. "High" exposure was defined as an occupational exposure that may exceed the RF/MW-EMF exposure limits for the general public recommended by the International Commission on Non-Ionizing Radiation Protection. Multiple conditional logistic regressions were performed separately for glioma and meningioma. No significant association between occupational exposure to RF/MW-EMF and brain tumors was found. For glioma, the adjusted odds ratio for highly exposed persons compared with persons not highly exposed was 1.21 (95% confidence interval: 0.69, 2.13); for meningioma, it was 1.34 (95% confidence interval: 0.64, 2.81). However, the slight increase in risk observed with increasing duration of exposure merits further research with larger sample sizes.

Caraglia M, Marra M, Mancinelli F, D'Ambrosio G, Massa R, Giordano A, Budillon A, Abbruzzese A, Bismuto E. Electromagnetic fields at mobile phone frequency induce apoptosis and inactivation of the multi-chaperone complex in human epidermoid cancer cells. J Cell Physiol. 204(2):539-548, 2005.

The exposure to non-thermal microwave electromagnetic field (MW-EMF) at 1.95 MHz,

Studies that show **Cell Phone** Health Effects

a frequency used in mobile communication, affects the refolding kinetics of eukaryotic proteins (Mancinelli et al., 2004). On these basis we have evaluated the in vivo effect of MW-EMF in human epidermoid cancer KB cells. We have found that MW-EMF induces time-dependent apoptosis (45% after 3 h) that is paralleled by an about 2.5-fold decrease of the expression of ras and Raf-1 and of the activity of ras and Erk-1/2. Although also the expression of Akt was reduced its activity was unchanged likely as a consequence of the increased expression of its upstream activator PI3K. In the same experimental conditions an about 2.5-fold increase of the ubiquitination of ras and Raf-1 was also found and the addition for 12 h of proteasome inhibitor lactacystin at 10 microM caused an accumulation of the ubiquitinated isoforms of ras and Raf-1 and counteracted the effects of MW-EMF on ras and Raf-1 expression suggesting an increased proteasome-dependent degradation induced by MW-EMF. The exposure of KB cells to MW-EMF induced a differential activation of stress-dependent pathway with an increase of JNK-1 activity and HSP70 and 27 expression and with a reduction of p38 kinase activity and HSP90 expression. The overexpression of HSP90 induced by transfection of KB cells with a plasmid encoding for the factor completely antagonized the apoptosis and the inactivation of the ras --> Erk-dependent survival signal induced by MW-EMF. Conversely, the inhibition of Erk activity induced by 12 h exposure to 10 mM Mek-1 inhibitor U0126 antagonized the effects induced by HSP90 transfection on apoptosis caused by MW-EMF. In conclusion, these results demonstrate for the first time that MW-EMF induces apoptosis through the inactivation of the ras --> Erk survival signaling due to enhanced degradation of ras and Raf-1 determined by decreased expression of HSP90 and the consequent increase of proteasome dependent degradation.

Heikkinen P, Kosma VM, Alhonen L, Huuskonen H, Komulainen H, Kumlin T, Laitinen JT, Lang S, Puranen L, Juutilainen J. Effects of mobile phone radiation on UV-induced skin tumourigenesis in ornithine decarboxylase transgenic and non-transgenic mice. Int J Radiat Biol 79(4):221-233, 2003.

Purpose: The effects of low-level radiofrequency radiation (RFR) on ultraviolet (UV)-induced skin tumorigenesis were evaluated in ornithine decarboxylase (ODC) and non-transgenic mice. Materials and methods: Transgenic female mice over-expressing the human ODC gene and their non-transgenic littermates (20 animals in the cage control group, and 45-49 animals in the other groups) were exposed for 52 weeks to UV radiation or a combination of UV radiation and pulsed RFR. The UV dose was 240 Jm(-2) (1.2 x human minimum erythral dose) delivered three times a week. One group of animals was exposed to Digital Advanced Mobile Phone System (DAMPS)-type RFR, the other group to Global System for Mobile (GSM)-type RFR at a nominal average specific absorption rate of 0.5 W kg(-1), 1.5 h day(-1), for 5 days a week. The skin was carefully palpated weekly for macroscopic tumours. Histopathological analyses of all skin lesions and of a specified dorsal skin area were performed on all animals. Results: UV exposure resulted in development of macroscopic skin tumours in 11.5 and 36.8% of non-transgenic and transgenic animals, respectively. The RFR exposures did not give a statistically significant effect on the development of skin tumours in either transgenic or

Studies that show Cell Phone Health Effects

non-transgenic animals, or in combined analysis, but tumour development appeared slightly accelerated especially in non-transgenic animals. No effects of RFR exposures were found on excretion of 6-hydroxymelatonin sulphate into urine or on polyamine levels in dorsal skin. Conclusion: RFR exposures did not significantly enhance skin tumourigenesis. However, the slightly accelerated tumour development may warrant further evaluation.

DNA Damage

Franzellitti S, Valbonesi P, Ciancaglini N, Biondi C, Contin A, Bersani F, Fabbri E. Transient DNA damage induced by high-frequency electromagnetic fields (GSM 1.8 GHz) in the human trophoblast HTR-8/SVneo cell line evaluated with the alkaline comet assay. *Mutat Res.* 683(1-2):35-42, 2010.

One of the most controversial issue regarding high-frequency electromagnetic fields (HF-EMF) is their putative capacity to affect DNA integrity. This is of particular concern due to the increasing use of HF-EMF in communication technologies, including mobile phones. Although epidemiological studies report no detrimental effects on human health, the possible disturbance generated by HF-EMF on cell physiology remains controversial. In addition, the question remains as to whether cells are able to compensate their potential effects. We have previously reported that a 1-h exposure to amplitude-modulated 1.8 GHz sinusoidal waves (GSM-217 Hz, SAR=2 W/kg) largely used in mobile telephony did not cause increased levels of primary DNA damage in human trophoblast HTR-8/SVneo cells. Nevertheless, further investigations on trophoblast cell responses after exposure to GSM signals of different types and durations were considered of interest. In the present work, HTR-8/SVneo cells were exposed for 4, 16 or 24h to 1.8 GHz continuous wave (CW) and different GSM signals, namely GSM-217 Hz and GSM-Talk (intermittent exposure: 5 min field on, 10 min field off). The alkaline comet assay was used to evaluate primary DNA damages and/or strand breaks due to uncompleted repair processes in HF-EMF exposed samples. The amplitude-modulated signals GSM-217 Hz and GSM-Talk induced a significant increase in comet parameters in trophoblast cells after 16 and 24h of exposure, while the un-modulated CW was ineffective. However, alterations were rapidly recovered and the DNA integrity of HF-EMF exposed cells was similar to that of sham-exposed cells within 2h of recovery in the absence irradiation. Our data suggest that HF-EMF with a carrier frequency and modulation scheme typical of the GSM signal may affect the DNA integrity.

Hekmat A, Saboury AA, Moosavi-Movahedi AA. The toxic effects of mobile phone radiofrequency (940MHz) on the structure of calf thymus DNA. *Ecotoxicol Environ Saf.* 2012 Nov 16. pii: S0147-6513(12)00368-5. doi: 10.1016/j.ecoenv.2012.10.016. [Epub ahead of print]

Currently, the biological effects of nonionizing electromagnetic fields (EMFs) including radiofrequency (RF) radiation have been the subject of numerous experimental and theoretical studies. The aim of this study is to evaluate the possible biological effects of

Studies that show **Cell Phone** Health Effects

mobile phone RF (940MHz, 15V/m and SAR=40mW/kg) on the structure of calf thymus DNA (ct DNA) immediately after exposure and 2h after 45min exposure via diverse range of spectroscopic instruments. The UV-vis and circular dichroism (CD) experiments depict that mobile phone EMFs can remarkably cause disturbance on ct DNA structure. In addition, the DNA samples, immediately after exposure and 2h after 45min exposure, are relatively thermally unstable compared to the DNA solution, which was placed in a small shielded box (unexposed ct DNA). Furthermore, the exposed DNA samples (the DNA samples that were exposed to 940MHz EMF) have more fluorescence emission when compared with the unexposed DNA, which may have occurred attributable to expansion of the exposed DNA structure. The results of dynamic light scattering (DLS) and zeta potential experiments demonstrate that RF-EMFs lead to increment in the surface charge and size of DNA. The structure of DNA immediately after exposure is not significantly different from the DNA sample 2h after 45min exposure. In other words, the EMF-induced conformational changes are irreversible. Collectively, our results reveal that 940MHz can alter the structure of DNA. The displacement of electrons in DNA by EMFs may lead to conformational changes of DNA and DNA disaggregation. Results from this study could have an important implication on the health effects of RF-EMFs exposure. In addition, this finding could proffer a novel strategy for the development of next generation of mobile phone.

Sekeroğlu V, Akar A, Sekeroğlu ZA. Cytotoxic and genotoxic effects of high-frequency electromagnetic fields (GSM 1800 MHz) on immature and mature rats. *Ecotoxicol Environ Saf.* 80:140-144, 2012.

We investigated the cytogenotoxic effects of high frequency electromagnetic fields (HF-EMF) for 45 day and the effect of a recovery period of 15 day after exposure to EMF on bone marrow cells of immature and mature rats. The animals in treatment groups were exposed to 1800 MHz EMF at SAR of 0.37 W/kg and 0.49 W/kg for 2h/day for 45 day. Two recovery groups were kept for a recovery period of 15 day without EMF after exposure to HF-EMF. Two control groups for both immature and mature rats were also included. Significant differences were also observed in chromosome aberrations (CA), micronucleus (MN) frequency, mitotic index (MI) and ratio of polychromatic erythrocytes (PCEs) in all treatment groups. The cytogenotoxic damage was more remarkable in immature rats and, the recovery period did not improve this damage in immature rats. Because much higher and irreversible cytogenotoxic damage was observed in immature rats than in mature rats, further studies are needed to understand effects of EMF on DNA damage and DNA repair, and to determine safe limits for environment and human, especially for children.

Tomruk A, Guler G, Dincel AS. The influence of 1800 MHz GSM-like signals on hepatic oxidative DNA and lipid damage in nonpregnant, pregnant, and newly born rabbits. *Cell Biochem Biophys.* 56(1):39-47, 2010.

The aim of our study is to evaluate the possible biological effects of whole-body 1800 MHz GSM-like radiofrequency (RF) radiation exposure on liver oxidative DNA damage and lipid peroxidation levels in nonpregnant, pregnant New Zealand White rabbits, and

Studies that show **Cell Phone** Health Effects

in their newly borns. Eighteen nonpregnant and pregnant rabbits were used and randomly divided into four groups which were composed of nine rabbits: (i) Group I (nonpregnant control), (ii) Group II (nonpregnant-RF exposed), (iii) Group III (pregnant control), (iv) Group IV (pregnant-RF exposed). Newborns of the pregnant rabbits were also divided into two groups: (v) Group V (newborns of Group III) and (vi) Group VI (newborns of Group III). 1800 MHz GSM-like RF radiation whole-body exposure (15 min/day for a week) was applied to Group II and Group IV. No significant differences were found in liver 8 OHdG/10(6) dG levels of exposure groups (Group II and Group IV) compared to controls (Group I and Group III). However, in Group II and Group IV malondialdehyde (MDA) and ferrous oxidation in xylene orange (FOX) levels were increased compared to Group I ($P < 0.05$, Mann-Whitney). No significant differences were found in liver tissue of 8 OHdG/10(6) dG and MDA levels between Group VI and Group V ($P > 0.05$, Mann-Whitney) while liver FOX levels were found significantly increased in Group VI with respect to Group V ($P < 0.05$, Mann-Whitney). Consequently, the whole-body 1800 MHz GSM-like RF radiation exposure may lead to oxidative destruction as being indicators of subsequent reactions that occur to form oxygen toxicity in tissues.

Schwarz C, Kratochvil E, Pilger A, Kuster N, Adlkofer F, Rüdiger HW. Radiofrequency electromagnetic fields (UMTS, 1,950 MHz) induce genotoxic effects in vitro in human fibroblasts but not in lymphocytes. *Int Arch Occup Environ Health*. 81(6):755-767, 2008.

OBJECTIVE: Universal Mobile Telecommunication System (UMTS) was recently introduced as the third generation mobile communication standard in Europe. This was done without any information on biological effects and genotoxic properties of these particular high-frequency electromagnetic fields. This is disconcerting, because genotoxic effects of the second generation standard Global System for Mobile Communication have been reported after exposure of human cells in vitro. **METHODS:** Human cultured fibroblasts of three different donors and three different short-term human lymphocyte cultures were exposed to 1,950 MHz UMTS below the specific absorption rate (SAR) safety limit of 2 W/kg. The alkaline comet assay and the micronucleus assay were used to ascertain dose and time-dependent genotoxic effects. Five hundred cells per slide were visually evaluated in the comet assay and comet tail factor (CTF) was calculated. In the micronucleus assay 1,000 binucleated cells were evaluated per assay. The origin of the micronuclei was determined by fluorescence labeled anticentromere antibodies. All evaluations were performed under blinded conditions. **RESULTS:** UMTS exposure increased the CTF and induced centromere-negative micronuclei (MN) in human cultured fibroblasts in a dose and time-dependent way. Incubation for 24 h at a SAR of 0.05 W/kg generated a statistically significant rise in both CTF and MN ($P = 0.02$). At a SAR of 0.1 W/kg the CTF was significantly increased after 8 h of incubation ($P = 0.02$), the number of MN after 12 h ($P = 0.02$). No UMTS effect was obtained with lymphocytes, either unstimulated or stimulated with Phytohemagglutinin. **CONCLUSION:** UMTS exposure may cause genetic alterations in some but not in all human cells in vitro.

Studies that show **Cell Phone** Health Effects

Pesnya DS, Romanovsky AV. Comparison of cytotoxic and genotoxic effects of plutonium-239 alpha particles and mobile phone GSM 900 radiation in the Allium cepa test. *Mutat Res.* 2012 Oct 8. pii: S1383-5718(12)00291-4. doi: 10.1016/j.mrgentox.2012.08.010. [Epub ahead of print]

The goal of this study was to compare the cytotoxic and genotoxic effects of plutonium-239 alpha particles and GSM 900 modulated mobile phone (model Sony Ericsson K550i) radiation in the Allium cepa test. Three groups of bulbs were exposed to mobile phone radiation during 0 (sham), 3 and 9h. A positive control group was treated during 20min with plutonium-239 alpha-radiation. Mitotic abnormalities, chromosome aberrations, micronuclei and mitotic index were analyzed. Exposure to alpha-radiation from plutonium-239 and exposure to modulated radiation from mobile phone during 3 and 9h significantly increased the mitotic index. GSM 900 mobile phone radiation as well as alpha-radiation from plutonium-239 induced both clastogenic and aneugenic effects. However, the aneugenic activity of mobile phone radiation was more pronounced. After 9h of exposure to mobile phone radiation, polyploid cells, three-groups metaphases, amitoses and some unspecified abnormalities were detected, which were not registered in the other experimental groups. Importantly, GSM 900 mobile phone radiation increased the mitotic index, the frequency of mitotic and chromosome abnormalities, and the micronucleus frequency in a time-dependent manner. Due to its sensitivity, the A. cepa test can be recommended as a useful cytogenetic assay to assess cytotoxic and genotoxic effects of radiofrequency electromagnetic fields.

Zhijian C, Xiaoxue L, Wei Z, Yezhen L, Jianlin L, Deqiang L, Shijie C, Lifan J, Jiliang H. Studying the protein expression in human B lymphoblastoid cells exposed to 1.8-GHz (GSM) radiofrequency radiation (RFR) with protein microarray. *Biochem Biophys Res Commun.* 433(1):36-39, 2013.

In the present study, the protein microarray was used to investigate the protein expression in human B-cell lymphoblastoid cells intermittently exposed to 1.8-GHz GSM radiofrequency radiation (RFR) at the specific absorption rate (SAR) of 2.0W/kg for 24h. The differential expression of 27 proteins was found, which were related to DNA damage repair, apoptosis, oncogenesis, cell cycle and proliferation (ratio >1.5-fold, $P < 0.05$). The results validated with Western blot assay indicated that the expression of RPA32 was significantly down-regulated ($P < 0.05$) while the expression of p73 was significantly up-regulated in RFR exposure group ($P < 0.05$). Because of the crucial roles of those proteins in DNA repair and cell apoptosis, the results of present investigation may explain the biological effects of RFR on DNA damage/repair and cell apoptosis.

Souza LD, Cerqueira ED, Meireles JR. Assessment of nuclear abnormalities in exfoliated cells from the oral epithelium of mobile phone users. *Electromagn Biol Med.* 2013 May 28. [Epub ahead of print]

Transmission and reception of mobile telephony signals take place through electromagnetic wave radiation, or electromagnetic radiofrequency fields, between the mobile terminal and the radio base station. Based on reports in the literature on

Studies that show **Cell Phone Health Effects**

adverse effects from exposure to this type of radiation, the objective of this study was to evaluate the genotoxic and cytotoxic potential of such exposure, by means of the micronucleus test on exfoliated cells from the oral epithelium. The sample included 45 individuals distributed in 3 groups according to the amount of time in hours per week (t) spent using mobile phones: group I, $t > 5$ h; group II, $t > 1$ h and ≤ 5 h; and group III, $t \leq 1$ h. Cells from the oral mucosa were analyzed to assess the numbers of micronuclei, broken egg structures and degenerative nuclear abnormalities indicative of apoptosis (condensed chromatin, karyorrhexis and pyknosis) or necrosis (karyolysis in addition to these changes). The occurrences of micronuclei and degenerative nuclear abnormalities did not differ between the groups, but the number of broken egg (structures that may be associated with gene amplification) was significantly greater in the individuals in group I ($p < 0.05$).

Esmekaya MA, Aytekin E, Ozgur E, Güler G, Ergun MA, Omeroğlu S, Seyhan N. Mutagenic and morphologic impacts of 1.8GHz radiofrequency radiation on human peripheral blood lymphocytes (hPBLs) and possible protective role of pre-treatment with Ginkgo biloba (EGb 761). Sci Total Environ. 410-411:59-64, 2011.

The mutagenic and morphologic effects of 1.8GHz Global System for Mobile Communications (GSM) modulated RF (radiofrequency) radiation alone and in combination with Ginkgo biloba (EGb 761) pre-treatment in human peripheral blood lymphocytes (hPBLs) were investigated in this study using Sister Chromatid Exchange (SCE) and electron microscopy. Cell viability was assessed with 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide (MTT) reduction assay. The lymphocyte cultures were exposed to GSM modulated RF radiation at 1.8GHz for 6, 8, 24 and 48h with and without EGb 761. We observed morphological changes in pulse-modulated RF radiated lymphocytes. Longer exposure periods led to destruction of organelle and nucleus structures. Chromatin change and the loss of mitochondrial crista occurred in cells exposed to RF for 8h and 24h and were more pronounced in cells exposed for 48h. Cytoplasmic lysis and destruction of membrane integrity of cells and nuclei were also seen in 48h RF exposed cells. There was a significant increase ($p < 0.05$) in SCE frequency in RF exposed lymphocytes compared to sham controls. EGb 761 pre-treatment significantly decreased SCE from RF radiation. RF radiation also inhibited cell viability in a time dependent manner. The inhibitory effects of RF radiation on the growth of lymphocytes were marked in longer exposure periods. EGb 761 pre-treatment significantly increased cell viability in RF+EGb 761 treated groups at 8 and 24h when compared to RF exposed groups alone. The results of our study showed that RF radiation affects cell morphology, increases SCE and inhibits cell proliferation. However, EGb 761 has a protective role against RF induced mutagenity. We concluded that RF radiation induces chromosomal damage in hPBLs but this damage may be reduced by EGb 761 pre-treatment.

Diem E, Schwarz C, Adlkofer F, Jahn O, Rudiger H. Non-thermal DNA breakage by mobile-phone radiation (1800MHz) in human fibroblasts and in transformed GFSH-R17 rat granulosa cells in vitro. Mutat Res. 583:178-183, 2005.

Studies that show **Cell Phone Health Effects**

Cultured human diploid fibroblasts and cultured rat granulosa cells were exposed to intermittent and continuous radiofrequency electromagnetic fields (RF-EMF) used in mobile phones, with different specific absorption rates (SAR) and different mobile-phone modulations. DNA strand breaks were determined by means of the alkaline and neutral comet assay. RF-EMF exposure (1800MHz; SAR 1.2 or 2W/kg; different modulations; during 4, 16 and 24h; intermittent 5min on/10min off or continuous wave) induced DNA single- and double-strand breaks. Effects occurred after 16h exposure in both cell types and after different mobile-phone modulations. The intermittent exposure showed a stronger effect in the comet assay than continuous exposure. Therefore we conclude that the induced DNA damage cannot be based on thermal effects.

Yao K, Wu W, Wang K, Ni S, Ye P, Yu Y, Ye J, Sun L. Electromagnetic noise inhibits radiofrequency radiation-induced DNA damage and reactive oxygen species increase in human lens epithelial cells. Mol Vis. 14:964-969, 2008.

PURPOSE: The goal of this study was to investigate whether superposing of electromagnetic noise could block or attenuate DNA damage and intracellular **reactive oxygen species (ROS)** increase of cultured human lens epithelial cells (HLECs) induced by acute exposure to 1.8 GHz radiofrequency field (RF) of the Global System for Mobile Communications (GSM). METHODS: An sXc-1800 RF exposure system was used to produce a GSM signal at 1.8 GHz (217 Hz amplitude-modulated) with the specific absorption rate (SAR) of 1, 2, 3, and 4 W/kg. After 2 h of intermittent exposure, the ROS level was assessed by the fluorescent probe, 2',7'-dichlorodihydrofluorescein diacetate (DCFH-DA). DNA damage to HLECs was examined by alkaline comet assay and the phosphorylated form of histone variant H2AX (gammaH2AX) foci formation assay. RESULTS: After exposure to 1.8 GHz RF for 2 h, HLECs exhibited significant intracellular ROS increase in the 2, 3, and 4 W/kg groups. RF radiation at the SAR of 3 W/kg and 4 W/kg could induce significant DNA damage, examined by alkaline comet assay, which was used to detect mainly **single strand breaks (SSBs)**, while no statistical difference in double strand breaks (DSBs), evaluated by gammaH2AX foci, was found between RF exposure (SAR: 3 and 4 W/kg) and sham exposure groups. When RF was superposed with 2 muT electromagnetic noise could block RF-induced ROS increase and DNA damage. CONCLUSIONS: DNA damage induced by 1.8 GHz radiofrequency field for 2 h, which was mainly SSBs, may be associated with the increased ROS production. Electromagnetic noise could block RF-induced ROS formation and DNA damage.

Zhang SZ, Yao GD, Lu DQ, Chiang H, Xu ZP. [Effect of 1.8 GHz radiofrequencyelectromagnetic fields on gene expression of rat neurons]. Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi. 26(8):449-452, 2008. [Article in Chinese]

OBJECTIVE: To investigate the changes of gene expression in rat neuron induced by 1.8 GHz radiofrequencyelectromagnetic fields (RF EMF) to screen for RF EMF-responsive genes and the effect of different exposure times and modes on the gene expression in neuron. METHODS: Total RNA was extracted immediately and purified from the primary culture of neurons after intermittent exposed or sham-exposed to a frequency of 1.8

Studies that show **Cell Phone** Health Effects

GHz RF EMF for 24 hours at an average special absorption rate (SAR) of 2 W/kg. Affymetrix Rat Neurobiology U34 array was applied to investigate the changes of gene expression in rat neuron. Differentially expressed genes (Egr-1, Mbp and Plp) were further confirmed by semi-quantitative reverse transcription polymerase chain reaction (RT PCR). The expression levels of Egr-1, Mbp and Plp were observed at different exposure times (6, 24 h) and modes (intermittent and continuous exposure). RESULTS: Among 1200 candidate genes, 24 up-regulated and 10 down-regulated genes were found by using Affymetrix microarray suite software 5.0 which are associated with multiple cellular functions (cytoskeleton, signal transduction pathway, metabolism, etc.) after functional classification. Under 24 h and 6 h intermittent exposure, Egr-1 and Plp in experiment groups showed statistic significance ($P < 0.05$) compared with the control groups, while expression of Mbp did not change significantly ($P > 0.05$). After 24 h continuous exposure, Egr-1 and Mbp in experiment groups showed statistic significance ($P < 0.05$) compared with the control group, while expression of Plp did not change significantly ($P > 0.05$). Under the same exposure mode 6 h, expression of all the 3 genes did not change significantly. Different times (6, 24 h) and modes (intermittent and continuous exposure) of exposure exerted remarkable different influences on the expression of Egr-1, Mbp, Plp genes ($P < 0.01$). CONCLUSION: The changes of many genes transcription were involved in the effect of 1.8 GHz RF EMF on rat neurons; Down-regulation of Egr-1 and up-regulation of Mbp, Plp indicated the negative effects of RF EMF on neurons; The effect of RF intermittent exposure on gene expression was more obvious than that of continuous exposure; The effect of 24 h RF exposure (both intermittent and continuous) on gene expression was more obvious than that of 6 h (both intermittent and continuous).

Zhao R, Zhang S, Xu Z, Ju L, Lu D, Yao G. Studying gene expression profile of rat neuron exposed to 1800MHz radiofrequency electromagnetic fields with cDNA microassay. *Toxicology*. 235(3):167-175, 2007.

A widespread use of mobile phone (MP) evokes a growing concern for their possible adverse effects on human, especially the brain. Gene expression is a unique way of characterizing how cells and organism adapt to changes in the external environment, so the aim of this investigation was to determine whether 1800 MHz radiofrequency electromagnetic fields (RF EMF) can influence the gene expression of neuron. Affymetrix Rat Neurobiology U34 array was applied to investigate the changes of gene expression in rat neuron after exposed to the pulsed RF EMF at a frequency of 1800 MHz modulated by 217 Hz which is commonly used in MP. Among 1200 candidate genes, 24 up-regulated genes and 10 down-regulated genes were identified after 24-h intermittent exposure at an average special absorption rate (SAR) of 2 W/kg, which are associated with multiple cellular functions (cytoskeleton, signal transduction pathway, metabolism, etc.) after functional classification. The results were further confirmed by quantitative real-time polymerase chain reaction (RT PCR). The present results indicated that the gene expression of rat neuron could be altered by exposure to RF EMF under our experimental conditions.

Studies that show **Cell Phone** Health Effects

Prohofsky EW. RF absorption involving biological macromolecules.

Bioelectromagnetics. 25(6):441-451, 2004.

The fundamental intramolecular frequency of a globular protein can be obtained from the measurements of acoustic velocities of bulk protein matter. This lowest frequency for common size molecules is shown to be above several hundred GHz. All modes below this frequency would then be intermolecular modes or bulk modes of the molecule and surrounding matter or tissue. The lowest frequency modes of an extended DNA double helix are also shown to be bulk modes because of interaction with water. Only DNA modes, whose frequency is well above 4 GHz, can be intrahelical modes, that is, confined to the helix rather than in the helix plus surroundings. Near 4 GHz, they are heavily damped and, therefore, not able to resonantly absorb. Modes that absorb radio frequency (RF) below this frequency are bulk modes of the supporting matter. Bulk modes rapidly thermalize all absorbed energy. The implication of these findings for the possibility of athermal RF effects is considered. The applicability of these findings for other biological molecules is discussed.

Cam ST, Seyhan N. Single-strand DNA breaks in human hair root cells exposed to mobile phone radiation. Int J Radiat Biol.88(5):420-424, 2012.

Abstract. Purpose: To analyze the short term effects of radiofrequency radiation (RFR) exposure on genomic deoxyribonucleic acid (DNA) of human hair root cells. Subjects and methods: Hair samples were collected from 8 healthy human subjects immediately before and after using a 900-MHz GSM (Global System for Mobile Communications) mobile phone for 15 and 30 minutes. Single-strand DNA breaks of hair root cells from the samples were determined using the 'comet assay'. Results: The data showed that talking on a mobile phone for 15 or 30 minutes significantly increased ($p < .05$) single-strand DNA breaks in cells of hair roots close to the phone. Comparing the 15-min and 30-min data using the paired t-test also showed that significantly more damages resulted after 30 minutes than after 15 minutes of phone use. Conclusions: A short-term exposure (15 and 30 minutes) to RFR (900-MHz) from a mobile phone caused a significant increase in DNA single-strand breaks in human hair root cells located around the ear which is used for the phone calls.

Campisi A, Gulino M, Acquaviva R, Bellia P, Raciti G, Grasso R, Musumeci F, Vanella A, Triglia A. Reactive oxygen species levels and DNA fragmentation on astrocytes in primary culture after acute exposure to low intensity microwave electromagnetic field. Neurosci Lett.31 473(1):52-55, 2010.

The exposure of primary rat neocortical astroglial cell cultures to acute electromagnetic fields (EMF) in the microwave range was studied. Differentiated astroglial cell cultures at 14 days in vitro were exposed for 5, 10, or 20min to either 900MHz continuous waves or 900MHz waves modulated in amplitude at 50Hz using a sinusoidal waveform and 100% modulation index. The strength of the electric field (rms value) at the sample position was 10V/m. No change in cellular viability evaluated by MTT test and lactate dehydrogenase release was observed. A significant increase in ROS levels and DNA fragmentation was found only after exposure of the astrocytes to modulated EMF for

Studies that show **Cell Phone** Health Effects

20min. No evident effects were detected when shorter time intervals or continuous waves were used. The irradiation conditions allowed the exclusion of any possible thermal effect. Our data demonstrate, for the first time, that even acute exposure to low intensity EMF induces ROS production and DNA fragmentation in astrocytes in primary cultures, which also represent the principal target of modulated EMF. Our findings also suggest the hypothesis that the effects could be due to hyperstimulation of the glutamate receptors, which play a crucial role in acute and chronic brain damage. Furthermore, the results show the importance of the amplitude modulation in the interaction between EMF and neocortical astrocytes.

Gandhi G, Anita Genetic damage in mobile phone users: some preliminary findings. Ind J Hum Genet 11(2): 99-104, 2005.

BACKGROUND: The impact of microwave (MW)/radio frequency radiation (RFR) on important biological parameters is probably more than a simply thermal one. Exposure to radio frequency (RF) signals generated by the use of cellular telephones have increased dramatically and reported to affect physiological, neurological, cognitive and behavioural changes and to induce, initiate and promote carcinogenesis. Genotoxicity of RFR has also been reported in various test systems after in vitro and/or in vivo exposure but none in mobile phone users. **AIMS:** In the present study, DNA and chromosomal damage investigations were carried out on the peripheral blood lymphocytes of individuals using mobile phones, being exposed to MW frequency ranging from 800 to 2000 MHz. **METHODS:** DNA damage was assessed using the single cell gel electrophoresis assay and aneugenic and clastogenic damage by the in vivo capillary blood micronucleus test (MNT) in a total of 24 mobile phone users. **RESULTS:** Mean comet tail length (26.76 ± 0.054 mm; 39.75% of cells damaged) in mobile phone users was highly significant from that in the control group. The in vivo capillary blood MNT also revealed highly significant (0.25) frequency of micronucleated (MNd) cells. **CONCLUSIONS:** These results highlight a correlation between mobile phone use (exposure to RFR) and genetic damage and require interim public health actions in the wake of widespread use of mobile telephony.

Franzellitti S, Valbonesi P, Ciancaglini N, Biondi C, Contin A, Bersani F, Fabbri E. Transient DNA damage induced by high-frequency electromagnetic fields (GSM 1.8 GHz) in the human trophoblast HTR-8/SVneo cell line evaluated with the alkaline comet assay. Mutat Res. 683(1-2):35-42, 2010.

One of the most controversial issue regarding high-frequency electromagnetic fields (HF-EMF) is their putative capacity to affect DNA integrity. This is of particular concern due to the increasing use of HF-EMF in communication technologies, including mobile phones. Although epidemiological studies report no detrimental effects on human health, the possible disturbance generated by HF-EMF on cell physiology remains controversial. In addition, the question remains as to whether cells are able to compensate their potential effects. We have previously reported that a 1-h exposure to amplitude-modulated 1.8 GHz sinusoidal waves (GSM-217 Hz, SAR=2 W/kg) largely used in mobile telephony did not cause increased levels of primary DNA damage in human

Studies that show **Cell Phone** Health Effects

trophoblast HTR-8/SVneo cells. Nevertheless, further investigations on trophoblast cell responses after exposure to GSM signals of different types and durations were considered of interest. In the present work, HTR-8/SVneo cells were exposed for 4, 16 or 24h to 1.8 GHz continuous wave (CW) and different GSM signals, namely GSM-217 Hz and GSM-Talk (intermittent exposure: 5 min field on, 10 min field off). The alkaline comet assay was used to evaluate primary DNA damages and/or strand breaks due to uncompleted repair processes in HF-EMF exposed samples. The amplitude-modulated signals GSM-217 Hz and GSM-Talk induced a significant increase in comet parameters in trophoblast cells after 16 and 24h of exposure, while the un-modulated CW was ineffective. However, alterations were rapidly recovered and the DNA integrity of HF-EMF exposed cells was similar to that of sham-exposed cells within 2h of recovery in the absence irradiation. Our data suggest that HF-EMF with a carrier frequency and modulation scheme typical of the GSM signal may affect the DNA integrity.

Wu W, Yao K, Wang KJ, Lu DQ, He JL, Xu LH, Sun WJ. [Blocking 1800 MHz mobile phone radiation-induced reactive oxygen species production and DNA damage in lens epithelial cells by noise magnetic fields.] Zhejiang Da Xue Xue Bao Yi Xue Ban. 37(1):34-38, 2008. [Article in Chinese]

OBJECTIVE: To investigate whether the exposure to the electromagnetic noise can block reactive oxygen species (ROS) production and DNA damage of lens epithelial cells induced by 1800 MHz mobile phone radiation. **METHODS:** The DCFH-DA method and comet assay were used respectively to detect the intracellular ROS and DNA damage of cultured human lens epithelial cells induced by 4 W/kg 1800 MHz mobile phone radiation or/and 2microT electromagnetic noise for 24 h intermittently. **RESULT:** 1800 MHz mobile phone radiation at 4 W/kg for 24 h increased intracellular ROS and DNA damage significantly ($P<0.05$). However, the ROS level and DNA damage of mobile phone radiation plus noise group were not significant enhanced ($P>0.05$) as compared to sham exposure group. **Conclusion:** Electromagnetic noise can block intracellular ROS production and DNA damage of human lens epithelial cells induced by 1800 MHz mobile phone radiation.

Ji S, Oh E, Sul D, Choi JW, Park H, Lee E. DNA Damage of Lymphocytes in Volunteers after 4 hours Use of Mobile Phone. J Prev Med Public Health. 37(4):373-380, 2004.

OBJECTIVES: There has been gradually increasing concern about the adverse health effects of electromagnetic radiation originating from cell phones which are widely used in modern life. Cell phone radiation may affect human health by increasing free radicals of human blood cells. This study has been designed to identify DNA damage of blood cells by electromagnetic radiation caused by cell phone use. **METHODS:** This study investigated the health effect of acute exposure to commercially available cell phones on certain parameters such as an indicator of DNA damage for 14 healthy adult volunteers. Each volunteer during the experiment talked over the cell phone with the keypad facing the right side of the face for 4 hours. The single cell gel electrophoresis assay (Comet assay), which is very sensitive in detecting the presence of DNA strand-breaks and alkali-labile damage in individual cells, was used to assess peripheral blood

Studies that show **Cell Phone** Health Effects

cells (T-cells, B-cells, granulocytes) from volunteers before and after exposure to cell phone radiation. The parameters of Comet assay measured were Olive Tail Moment and Tail DNA %. **RESULTS:** The Olive Tail Moment of B-cells and granulocytes and Tail DNA % of B-cells and granulocytes were increased by a statistically significant extent after 4-hour use of a cell phone compared with controls. **CONCLUSIONS:** It is concluded that cell phone radiation caused the DNA damage during the 4 hours of experimental condition. Nonetheless, this study suggested that cell phone use may increase DNA damage by electromagnetic radiation and other contributing factors.

Buttiglione M, Roca L, Montemurno E, Vitiello F, Capozzi V, Cibelli G. Radiofrequency radiation (900 MHz) induces Egr-1 gene expression and affects cell-cycle control in human neuroblastoma cells. J Cell Physiol. 213(3):759-767, 2007.

Many environmental signals, including ionizing radiation and UV rays, induce activation of Egr-1 gene, thus affecting cell growth and apoptosis. The paucity and the controversial knowledge about the effect of electromagnetic fields (EMF) exposure of nerve cells prompted us to investigate the bioeffects of radiofrequency (RF) radiation on SH-SY5Y neuroblastoma cells. The effect of a modulated RF field of 900 MHz, generated by a wire patch cell (WPC) antenna exposure system on Egr-1 gene expression, was studied as a function of time. Short-term exposures induced a transient increase in Egr-1 mRNA level paralleled with activation of the MAPK subtypes ERK1/2 and SAPK/JNK. The effects of RF radiations on cell growth rate and apoptosis were also studied. Exposure to RF radiation had an anti-proliferative activity in SH-SY5Y cells with a significant effect observed at 24 h. RF radiation impaired cell cycle progression, reaching a significant G2-M arrest. In addition, the appearance of the sub-G1 peak, a hallmark of apoptosis, was highlighted after a 24-h exposure, together with a significant decrease in mRNA levels of Bcl-2 and survivin genes, both interfering with signaling between G2-M arrest and apoptosis. Our results provide evidence that exposure to a 900 MHz-modulated RF radiation affect both Egr-1 gene expression and cell regulatory functions, involving apoptosis inhibitors like Bcl-2 and survivin, thus providing important insights into a potentially broad mechanism for controlling in vitro cell viability.

Tice RR, Hook GG, Donner M, McRee DI, Guy AW. Genotoxicity of radiofrequency signals. I. Investigation of DNA damage and micronuclei induction in cultured human blood cells. Bioelectromagnetics 23:113-126, 2002.

As part of a comprehensive investigation of the potential genotoxicity of radiofrequency (RF) signals emitted by cellular telephones, in vitro studies evaluated the induction of DNA and chromosomal damage in human blood leukocytes and lymphocytes, respectively. The signals were voice modulated 837 MHz produced by an analog signal generator or by a time division multiple access (TDMA) cellular telephone, 837 MHz generated by a code division multiple access (CDMA) cellular telephone (not voice modulated), and voice modulated 1909.8 MHz generated by a global system of mobile communication (GSM)-type personal communication systems (PCS) cellular telephone. DNA damage (strand breaks/alkali labile sites) was assessed in leukocytes using the alkaline (pH>13) single cell gel electrophoresis (SCG) assay. Chromosomal damage was evaluated in lymphocytes mitogenically stimulated to divide

Studies that show **Cell Phone** Health Effects

postexposure using the cytochalasin B-binucleate cell micronucleus assay. Cells were exposed at $37\pm 1^\circ\text{C}$, for 3 or 24 h at average specific absorption rates (SARs) of 1.0-10.0 W/kg. Exposure for either 3 or 24 h did not induce a significant increase in DNA damage in leukocytes, nor did exposure for 3 h induce a significant increase in micronucleated cells among lymphocytes. However, exposure to each of the four RF signal technologies for 24 h at an average SAR of 5.0 or 10.0 W/kg resulted in a significant and reproducible increase in the frequency of micronucleated lymphocytes. The magnitude of the response (approximately four fold) was independent of the technology, the presence or absence of voice modulation, and the frequency (837 vs. 1909.8 MHz). This research demonstrates that, under extended exposure conditions, RF signals at an average SAR of at least 5.0 W/kg are capable of inducing chromosomal damage in human lymphocytes.

Fucic A, Garaj-Vrhovac V, Skara M, Dimitrovic B, X-rays, microwaves and vinyl chloride monomer: their clastogenic and aneugenic activity, using the micronucleus assay on human lymphocytes. *Mutat Res* 282(4):265-271, 1992.

Chromosome aberration assays, sister-chromatid exchange techniques and micronucleus assays are commonly used methods for biomonitoring genetic material damaged by chemical or physical agents. On the other hand, their aneugenic activity, which can lead to hypoploidy and may also be associated with carcinogenesis, has not been thoroughly investigated. In our study we chose the micronucleus assay with a new mathematical approach to separate clastogenic from aneugenic activity of three well-known mutagens (vinyl chloride monomer, X-rays and microwaves) on the genome of human somatic cells. The comparison of frequencies of size distribution of micronuclei in the lymphocytes of humans exposed to each of these three mutagens showed that X-rays and microwaves were preferentially clastogens while vinyl chloride monomer showed aneugenic activity as well. Microwaves possess some mutagenic characteristics typical of chemical mutagens.

Sun LX, Yao K, Jiang H, He JL, Lu DQ, Wang KJ, Li HW [DNA damage and repair induced by acute exposure of microwave from mobile phone on cultured human lens epithelial cells] *Zhonghua Yan Ke Za Zhi*. 42(12):1084-1088, 2006. [Article in Chinese]

OBJECTIVE: To investigate the effects of acute exposure of low-power 217 Hz modulated 1.8 GHz microwave radiation on the DNA damage of human lens epithelial cells (hLECs) and repair. METHODS: Cultured hLECs were exposed to 217 Hz modulated 1.8 GHz microwave radiation at SAR (specific absorption rate) of 1.0, 2.0, 3.00 and 4.0 W/kg for 2 hours in an sXc-1800 incubator and irradiate system, the DNA single strand breaks were detected with comet assay (single-cell gel electrophoresis) in sham-irradiated cells and irradiated cells incubated for varying periods: 0, 30 and 60 minutes after irradiation. Images of comets were digitized and analyzed using an Imagine-pro plus software, and the indexes used in this study were tail length (TL) and tail moment (TM). BrdU was added into the medium with additional one hour incubation after radiation, the cell proliferation rate was determined using a BrdU-kit. RESULTS: The difference of DNA-breaks between the exposure and sham exposure groups induced by 1.0 and 2.0 W/kg irradiation were not significant in each time points ($P > 0.05$); there were significant

Studies that show **Cell Phone** Health Effects

difference in both groups at the exposure dose of 3.0 and 4.0 W/kg immediately and at the time of 30 minutes after irradiation ($P < 0.01$); if the radiation exposure time was beyond one hour no differences were able to be detected in 3.0 W/kg group ($P > 0.05$) compared with control, but the evidence of significant DNA damage still existed in 4.0 W/kg group at the same time point. Cell proliferation rate had no significant difference when the application of SAR was ≤ 3.0 W/kg ($P > 0.05$), however the cell proliferation was decreased significantly at the dose of 4.0 W/kg irradiation ($P < 0.01$). **CONCLUSIONS:** No effective DNA damage was induced using comet assay after 2 hours irradiation of 1.8 GHz microwave on hLECs at the dose SAR ≤ 3.0 W/kg. 4.0 W/kg irradiation caused significantly DNA damage and inhibition of hLECs proliferation.

Zhao TY, Zou SP, Knapp PE Exposure to cell phone radiation up-regulates apoptosis genes in primary cultures of neurons and astrocytes. *Neurosci Lett.* 412(1):34-38, 2007.

The health effects of cell phone radiation exposure are a growing public concern. This study investigated whether expression of genes related to cell death pathways are dysregulated in primary cultured neurons and astrocytes by exposure to a working Global System for Mobile Communication (GSM) cell phone rated at a frequency of 1900MHz. Primary cultures were exposed to cell phone emissions for 2h. We used array analysis and real-time RT-PCR to show up-regulation of caspase-2, caspase-6 and Asc (apoptosis associated speck-like protein containing a card) gene expression in neurons and astrocytes. Up-regulation occurred in both "on" and "stand-by" modes in neurons, but only in "on" mode in astrocytes. Additionally, astrocytes showed up-regulation of the Bax gene. The effects are specific since up-regulation was not seen for other genes associated with apoptosis, such as caspase-9 in either neurons or astrocytes, or Bax in neurons. The results show that even relatively short-term exposure to cell phone radiofrequency emissions can up-regulate elements of apoptotic pathways in cells derived from the brain, and that neurons appear to be more sensitive to this effect than astrocytes.

Hou Q, Wang M, Wu S, Ma X, An G, Liu H, Xie F. Oxidative changes and apoptosis induced by 1800-MHz electromagnetic radiation in NIH/3T3 cells. *Electromagn Biol Med.* 2014 Mar 25. [Epub ahead of print]

To investigate the potential adverse effects of mobile phone radiation, we studied reactive oxygen species (ROS), DNA damage and apoptosis in mouse embryonic fibroblasts (NIH/3T3) after intermittent exposure (5 min on/10 min off, for various durations from 0.5 to 8 h) to an 1800-MHz GSM-talk mode electromagnetic radiation (EMR) at an average specific absorption rate of 2 W/kg. A 2',7'-dichlorofluorescein diacetate fluorescence probe was used to detect intracellular ROS levels, immunofluorescence was used to detect γ H2AX foci as a marker for DNA damage, and flow cytometry was used to measure apoptosis. Our results showed a significant increase in intracellular ROS levels after EMR exposure and it reached the highest level at an exposure time of 1 h ($p < 0.05$) followed by a slight decrease when the exposure continued for as long as 8 h. No significant effect on the number of γ H2AX was detected after EMR exposure. The percentage of late-apoptotic cells in the EMR-exposed group

Studies that show **Cell Phone** Health Effects

was significantly higher than that in the sham-exposed groups ($p < 0.05$). These results indicate that an 1800-MHz EMR enhances ROS formation and promotes apoptosis in NIH/3T3 cells.

Zhang DY, Xu ZP, Chiang H, Lu DQ, Zeng QL. [Effects of GSM 1800 MHz radiofrequency electromagnetic fields on DNA damage in Chinese hamster lung cells.] Zhonghua Yu Fang Yi Xue Za Zhi. 40(3):149-152, 2006. [Article in Chinese]

OBJECTIVE: To study the effects of GSM 1800 MHz radiofrequency electromagnetic fields (RF EMF) on DNA damage in Chinese hamster lung (CHL) cells. **METHODS:** The cells were intermittently exposed or sham-exposed to GSM 1800 MHz RF EMF (5 minutes on/10 minutes off) at a special absorption rate (SAR) of 3.0 W/kg for 1 hour or 24 hours. Meanwhile, cells exposed to 2-acetaminofluorene, a DNA damage agent, at a final concentration of 20 mg/L for 2 hours were used as positive control. After exposure, cells were fixed by using 4% paraformaldehyde and processed for phosphorylated form of H2AX (gammaH2AX) immunofluorescence measurement. The primary antibody used for immunofluorescence was mouse monoclonal antibody against gammaH2AX and the secondary antibody was fluorescein isothiocyanate (FITC)-conjugated goat anti-mouse IgG. Nuclei were counterstained with 4, 6-diamidino-2-phenylindole (DAPI). The gammaH2AX foci and nuclei were visualized with an Olympus AX70 fluorescent microscope. Image Pro-Plus software was used to count the gammaH2AX foci in each cell. For each exposure condition, at least 50 cells were selected to detect gammaH2AX foci. Cells were classified as positive when more than five foci were detected. The percentage of gammaH2AX foci positive cells was adopted as the index of DNA damage. **RESULTS:** The percentage of gammaH2AX foci positive cell of 1800 MHz RF EMF exposure for 24 hours ($37.9 \pm 8.6\%$) or 2-acetylaminofluorene exposure ($50.9 \pm 9.4\%$) was significantly higher compared with the sham-exposure ($28.0 \pm 8.4\%$). However, there was no significant difference between the sham-exposure and RF EMF exposure for 1 hour ($31.8 \pm 8.7\%$). **CONCLUSION:** 1800 MHz RF EMF (SAR, 3.0 W/kg) for 24 hours might induce DNA damage in CHL cells.

Zhao R, Zhang SZ, Yao GD, Lu DQ, Jiang H, Xu ZP.[Effect of 1.8 GHz radiofrequency electromagnetic fields on the expression of microtubule associated protein 2 in rat neurons]Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi. 24(4):222-225, 2006. [Article in Chinese]

OBJECTIVE: To investigate the changes of gene expression in rat neurons induced by 1.8 GHz radiofrequency electromagnetic fields (RF EMF) and to screen for the RF EMF-responsive genes. **METHODS:** Newly-born SD rats in 24 hours were sacrificed to obtain cortex and hippocampus neurons. The cells were divided randomly into two groups: the experiment group (the irradiation group) and the control group (the false irradiation group). In the irradiation group, after twelve days' culture, neurons were exposed to 1.8 GHz RF EMF modulated by 217 Hz at a specific absorption rate (SAR) of 2 W/kg for 24 hours (5 minutes on/10 minutes off) while in the false control group, the neurons were put in the same waveguide as in the irradiation group, but were not exposed to any irradiation. The total RNA was isolated and purified immediately after exposure. The

Studies that show **Cell Phone** Health Effects

affymetrix rat neurobiology U34 assay was used for detecting the changes in gene expression profile according to the manufacturer's instruction. RF EMF-responsive candidate gene was confirmed by using ribonuclease protection assay (RPA). RESULTS: Among 1200 candidate genes, the expression levels of 34 genes were up or down regulated. Microtubule associated protein 2 (Map2) gene was selected as the candidate and subjected to further analysis. RPA data clearly revealed that Map2 was statistically significantly up-regulated after neurons were exposed to the RF EMF ($P < 0.05$). CONCLUSION: The modulation of gene expression and function of Map2 as a neuron specific cytoskeleton protein is crucial to maintain the normal framework and function of neurons. The finding that 1.8 GHz RF EMF exposure increases the expression of Map2 might indicate some unknown effects of RF EMF on neurons.

Goswami PC, Albee LD, Parsian AJ, Baty JD, Moros EG, Pickard WF, Roti Roti JL, Hunt CR, Proto-oncogene mRNA levels and activities of multiple transcription factors in C3H 10T 1/2 murine embryonic fibroblasts exposed to 835.62 and 847.74 MHz cellular phone communication frequency radiation. Radiat Res 151(3):300-309, 1999.

This study was designed to determine whether two differently modulated radiofrequencies of the type generally used in cellular phone communications could elicit a general stress response in a biological system. The two modulations and frequencies studied were a frequency-modulated continuous wave (FMCW) with a carrier frequency of 835.62 MHz and a code division multiple-access (CDMA) modulation centered on 847.74 MHz. Changes in proto-oncogene expression, determined by measuring Fos, Jun, and Myc mRNA levels as well as by the DNA-binding activity of the AP1, AP2 and NF-kappaB transcription factors, were used as indicators of a general stress response. The effect of radiofrequency exposure on proto-oncogene expression was assessed (1) in exponentially growing C3H 10T 1/2 mouse embryo fibroblasts during their transition to plateau phase and (2) during transition of serum-deprived cells to the proliferation cycle after serum stimulation. Exposure of serum-deprived cells to 835.62 MHz FMCW or 847.74 MHz CDMA microwaves (at an average specific absorption rate, SAR, of 0.6 W/kg) did not significantly change the kinetics of proto-oncogene expression after serum stimulation. Similarly, these exposures did not affect either the Jun and Myc mRNA levels or the DNA-binding activity of AP1, AP2 and NF-kappaB in exponential cells during transit to plateau-phase growth. Therefore, these results suggest that the radiofrequency exposure is unlikely to elicit a general stress response in cells of this cell line under these conditions. However, statistically significant increases (approximately 2-fold, $P = 0.001$) in Fos mRNA levels were detected in exponential cells in transit to the plateau phase and in plateau-phase cells exposed to 835.62MHz FMCW microwaves. For 847.74 MHz CDMA exposure, the increase was 1.4-fold ($P = 0.04$). This increase in Fos expression suggests that expression of specific genes could be affected by radiofrequency exposure.

Trosić I, Pavčić I, Milković-Kraus S, Mladinić M, Zeljezić D. Effect of electromagnetic radiofrequency radiation on the rats' brain, liver and kidney cells measured by comet assay. Coll Antropol. 35(4):1259-1264, 2011.

Studies that show Cell Phone Health Effects

The goal of study was to evaluate DNA damage in rat's renal, liver and brain cells after in vivo exposure to radiofrequency/microwave (Rf/Mw) radiation of cellular phone frequencies range. To determine DNA damage, a single cell gel electrophoresis/comet assay was used. Wistar rats (male, 12 week old, approximate body weight 350 g) (N = 9) were exposed to the carrier frequency of 915 MHz with Global System Mobile signal modulation (GSM), power density of 2.4 W/m², whole body average specific absorption rate SAR of 0.6 W/kg. The animals were irradiated for one hour/day, seven days/week during two weeks period. The exposure set-up was Gigahertz Transversal Electromagnetic Mode Cell (GTEM--cell). Sham irradiated controls (N = 9) were apart of the study. The body temperature was measured before and after exposure. There were no differences in temperature in between control and treated animals. Comet assay parameters such as the tail length and tail intensity were evaluated. In comparison with tail length in controls (13.5 +/- 0.7 microm), the tail was slightly elongated in brain cells of irradiated animals (14.0 +/- 0.3 microm). The tail length obtained for liver (14.5 +/- 0.3 microm) and kidney (13.9 +/- 0.5 microm) homogenates notably differs in comparison with matched sham controls (13.6 +/- 0.3 microm) and (12.9 +/- 0.9 microm). Differences in tail intensity between control and exposed animals were not significant. The results of this study suggest that, under the experimental conditions applied, repeated 915 MHz irradiation could be a cause of DNA breaks in renal and liver cells, but not affect the cell genome at the higher extent compared to the basal damage.

Gandhi G, Singh P. Cytogenetic damage in mobile phone users: preliminary data. Int J Hum Genet 5(4):259-265, 2005.

Mobile telephones, sometimes called cellular (cell) phones or handies, are now an integral part of modern life. The mobile phone handsets are low-powered radiofrequency transmitters, emitting maximum powers in the range of 0.2 to 0.6 watts. Scientific concerns have increased sufficiently over the possible hazard to health from using cell phones. The reported adverse health effects include physiological, behavioural and cognitive changes as well as tumour formation and genetic damage. However findings are controversial and no consensus exists. Genotoxicity has been observed either in lower organisms or in vitro studies. The aim of the present study hence was to detect any cytogenetic damage in mobile phone users by analysing short term peripheral lymphocyte cultures for chromosomal aberrations and the buccal mucosal cells for micronuclei (aneugenicity and clastogenicity). The results revealed increased number of micronucleated buccal cells and cytological abnormalities in cultured lymphocytes indicating the genotoxic response from mobile phone use.

Tiwari R, Lakshmi NK, Surender V, Rajesh AD, Bhargava SC, Ahuja YR. Combinative exposure effect of radio frequency signals from CDMA mobile phones and aphidicolin on DNA integrity. Electromagn Biol Med. 27(4):418-425, 2008.

The aim of present study is to assess DNA integrity on the effect of exposure to a radio frequency (RF) signal from Code Division Multiple Access (CDMA) mobile phones. Whole blood samples from six healthy male individuals were exposed for RF signals from a CDMA mobile phone for 1 h. Alkaline comet assay was performed to assess the DNA

Studies that show **Cell Phone** Health Effects

damage. The combinative exposure effect of the RF signals and APC at two concentrations on DNA integrity was studied. DNA repair efficiency of the samples was also studied after 2 h of exposure. The RF signals and APC (0.2 microg/ml) alone or in synergism did not have any significant DNA damage as compared to sham exposed. However, univariate analysis showed that DNA damage was significantly different among combinative exposure of RF signals and APC at 0.2 microg/ml ($p < 0.05$) and at 2 microg/ml ($p < 0.02$). APC at 2 microg/ml concentration also showed significant damage levels ($p < 0.05$) when compared to sham exposed. DNA repair efficiency also varied in a significant way in combinative exposure sets ($p < 0.05$). From these results, it appears that the repair inhibitor APC enhances DNA breaks at 2 microg/ml concentration and that the damage is possibly repairable. Thus, it can be inferred that the in vitro exposure to RF signals induces reversible DNA damage in synergism with APC.

Sykes PJ, McCallum BD, Bangay MJ, Hooker AM, Morley AA. Effect of Exposure to 900 MHz Radiofrequency Radiation on Intrachromosomal Recombination in pKZ1 Mice. *Radiat Res* 156(5):495-502, 2001.

Radiofrequency (RF) radiation emitted from mobile phones is not considered to be directly genotoxic, but it may have downstream effects on cellular DNA. We studied the effect of 4 W/kg pulsed 900 MHz RF radiation on somatic intrachromosomal recombination in the spleen in the pKZ1 recombination mutagenesis model. Somatic intrachromosomal recombination inversion events were detected in spleen tissue of pKZ1 mice by histochemical staining for E. coli beta-galactosidase protein in cells in which the lacZ transgene has undergone an inversion event. pKZ1 mice were exposed daily for 30 min to plane-wave fields of 900 MHz with a pulse repetition frequency of 217 Hz and a pulse width of 0.6 ms for 1, 5 or 25 days. Three days after the last exposure, spleen sections were screened for DNA inversion events. There was no significant difference between the control and treated groups in the 1- and 5-day exposure groups, but there was a significant reduction in inversions below the spontaneous frequency in the 25-day exposure group. This observation suggests that exposure to RF radiation can lead to a perturbation in recombination frequency which may have implications for recombination repair of DNA. The biological significance of a reduction below the spontaneous frequency is not known. The number of mice in each treatment group in this study was small ($n = 10$ or $n = 20$). Therefore, repetition of this study with a larger number of animals is required to confirm these observations.

Gajski G, Garaj-Vrhovac V. Radioprotective effects of honeybee venom (*Apis mellifera*) against 915-MHz microwave radiation-induced DNA damage in wistar rat lymphocytes: in vitro study. *Int J Toxicol*. 28(2):88-98, 2009.

The aim of this study is to investigate the radioprotective effect of bee venom against DNA damage induced by 915-MHz microwave radiation (specific absorption rate of 0.6 W/kg) in Wistar rats. Whole blood lymphocytes of Wistar rats are treated with 1 microg/mL bee venom 4 hours prior to and immediately before irradiation. Standard and formamidopyrimidine-DNA glycosylase (Fpg)-modified comet assays are used to assess basal and oxidative DNA damage produced by reactive oxygen species. Bee

Studies that show **Cell Phone** Health Effects

venom shows a decrease in DNA damage compared with irradiated samples. Parameters of Fpg-modified comet assay are statistically different from controls, making this assay more sensitive and suggesting that oxidative stress is a possible mechanism of DNA damage induction. Bee venom is demonstrated to have a radioprotective effect against basal and oxidative DNA damage. Furthermore, bee venom is not genotoxic and does not produce oxidative damage in the low concentrations used in this study.

Furtado-Filho OV, Borba JB, Dallegrave A, Pizzolato TM, Henriques JA, Moreira JC, Saffi J. Effect of 950 MHz UHF electromagnetic radiation on biomarkers of oxidative damage, metabolism of UFA and antioxidants in the livers of young rats of different ages. Int J Radiat Biol. 2013 Jul 25. [Epub ahead of print]

Purpose: To assess the effect of 950 MHz ultra-high-frequency electromagnetic radiation (UHF EMR) on biomarkers of oxidative damage, as well as to verify the concentration of unsaturated fatty acids (UFA) and the expression of the catalase in the livers of rats of different ages. Materials and methods: Twelve rats were equally divided into two groups as controls (CR) and exposed (ER), for each age (0, 6, 15 and 30 days). Radiation exposure lasted half an hour per day for up to 51 days (21 days of gestation and 6, 15 or 30 days of life outside the womb). The specific absorption rate (SAR) ranged from 1.3-1.0 W/kg. The damage to lipids, proteins and DNA was verified by thiobarbituric acid reactive substances (TBARS), protein carbonyls and comets, respectively. UFA were determined by gas chromatography with a flame ionization detector. The expression of catalase was by Western blotting. Results: The neonates had low levels of TBARS and concentrations of UFA after exposure. There was no age difference in the accumulation of protein carbonyls for any age. The DNA damage of ER 15 or 30 days was different. The exposed neonates exhibited lower expression of catalase. Conclusions: 950 MHz UHF EMR does not cause oxidative stress (OS), and it is not genotoxic to the livers of neonates or those of 6 and 15 day old rats, but it changes the concentrations of polyunsaturated fatty acid (PUFA) in neonates. For rats of 30 days, no OS, but it is genotoxic to the livers of ER to total body irradiation.

Donnellan M, McKenzie DR, French PW, Effects of exposure to electromagnetic radiation at 835 MHz on growth, morphology and secretory characteristics of a mast cell analogue, RBL-2H3. Cell Biol Int 21:427-439, 1997.

A mast cell line, RBL-2H3, was exposed to 835 MHz for 20 minutes, three times per day for 7 days at a power density of 8.1 ± 3 mW/cm². From day 4 onwards, it was observed that the rate of DNA synthesis and cell replication increased, that actin distribution and cell morphology became altered, and the amount of beta-hexosaminidase (a marker of granule secretion) released in response to a calcium ionophore was significantly enhanced, in comparison to unexposed cultures. There were no effects seen on levels of cytoskeletal protein synthesis or of beta-actin mRNA. Morphological changes persisted following subculture for at least 7 days in the absence of further exposure. It is hypothesized that effects of exposure to an electromagnetic field at 835 MHz may be mediated via a signal transduction pathway.

Studies that show **Cell Phone** Health Effects

Sun LX, Yao K, He JL, Lu DQ, Wang KJ, Li HW. [Effect of acute exposure to microwave from mobile phone on DNA damage and repair of cultured human lens epithelial cells in vitro.] Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi. 24(8):465-467, 2006.

[Article in Chinese]

OBJECTIVE: To investigate the DNA damage of human lens epithelial cells (LECs) caused by acute exposure to low-power 217 Hz modulated 1.8 GHz microwave radiation and DNA repair. **METHODS:** Cultured LECs were exposed to 217 Hz modulated 1.8 GHz microwave radiation at SAR (specific absorption rate) of 0, 1, 2, 3 and 4 W/kg for 2 hours in an sXc-1800 incubator and irradiate system. The DNA single strand breaks were detected with comet assay in sham-irradiated cells and irradiated cells incubated for varying periods: 0, 30, 60, 120 and 240 min after irradiation. Images of comets were digitized and analyzed using an Imagine-pro plus software, and the indexes used in this study were tail length (TL) and tail moment (TM). **RESULTS:** The difference in DNA-breaks between the exposure and sham exposure groups induced by 1 and 2 W/kg irradiation was not significant at every detect time ($P > 0.05$). As for the dosage of 3 and 4 W/kg there was difference in both group immediately after irradiation ($P < 0.01$). At the time of 30 min after irradiation the difference went on at both group ($P < 0.01$). However, the difference disappeared after one hour's incubation in 3 W/kg group ($P > 0.05$), and existed in 4 W/kg group. **CONCLUSION:** No or repairable DNA damage was observed after 2 hour irradiation of 1.8 GHz microwave on LECs when $SAR \leq 3$ W/kg. The DNA damages caused by 4 W/kg irradiation were irreversible.

Belyaev IY, Hillert L, Protopopova M, Tamm C, Malmgren LO, Persson BR, Selivanova G, Harms-Ringdahl M. 915 MHz microwaves and 50 Hz magnetic field affect chromatin conformation and 53BP1 foci in human lymphocytes from hypersensitive and healthy persons. Bioelectromagnetics. 26(3):173-184, 2005.

We used exposure to microwaves from a global system for mobile communication (GSM) mobile phone (915 MHz, specific absorption rate (SAR) 37 mW/kg) and power frequency magnetic field (50 Hz, 15 μ T peak value) to investigate the response of lymphocytes from healthy subjects and from persons reporting hypersensitivity to electromagnetic field (EMF). The hypersensitive and healthy donors were matched by gender and age and the data were analyzed blind to treatment condition. The changes in chromatin conformation were measured with the method of anomalous viscosity time dependencies (AVTD). 53BP1 protein, which has been shown to colocalize in foci with DNA double strand breaks (DSBs), was analyzed by immunostaining in situ. Exposure at room temperature to either 915 MHz or 50 Hz resulted in significant condensation of chromatin, shown as AVTD changes, which was similar to the effect of heat shock at 41 degrees C. No significant differences in responses between normal and hypersensitive subjects were detected. Neither 915 MHz nor 50 Hz exposure induced 53BP1 foci. On the contrary, a distinct decrease in background level of 53BP1 signaling was observed upon these exposures as well as after heat shock treatments. This decrease correlated with the AVTD data and may indicate decrease in accessibility of 53BP1 to antibodies because of stress-induced chromatin condensation. Apoptosis was determined by morphological changes and by apoptotic fragmentation of DNA as

Studies that show **Cell Phone** Health Effects

analyzed by pulsed-field gel electrophoresis (PFGE). No apoptosis was induced by exposure to 50 Hz and 915 MHz microwaves. In conclusion, 50 Hz magnetic field and 915 MHz microwaves under specified conditions of exposure induced comparable responses in lymphocytes from healthy and hypersensitive donors that were similar but not identical to stress response induced by heat shock.

Belyaev IY, Koch CB, Terenius O, Roxstrom-Lindquist K, Malmgren LO, H Sommer W, Salford LG, Persson BR. Exposure of rat brain to 915 MHz GSM microwaves induces changes in gene expression but not double stranded DNA breaks or effects on chromatin conformation. *Bioelectromagnetics*.27(4):295-306,2006.

We investigated whether exposure of rat brain to microwaves (MWs) of global system for mobile communication (GSM) induces DNA breaks, changes in chromatin conformation and in gene expression. An exposure installation was used based on a test mobile phone employing a GSM signal at 915 MHz, all standard modulations included, output power level in pulses 2 W, specific absorption rate (SAR) 0.4 mW/g. Rats were exposed or sham exposed to MWs during 2 h. After exposure, cell suspensions were prepared from brain samples, as well as from spleen and thymus. For analysis of gene expression patterns, total RNA was extracted from cerebellum. Changes in chromatin conformation, which are indicative of stress response and genotoxic effects, were measured by the method of anomalous viscosity time dependencies (AVTD). DNA double strand breaks (DSBs) were analyzed by pulsed-field gel electrophoresis (PFGE). Effects of MW exposure were observed on neither conformation of chromatin nor DNA DSBs. Gene expression profiles were obtained by Affymetrix U34 GeneChips representing 8800 rat genes and analyzed with the Affymetrix Microarray Suite (MAS) 5.0 software. In cerebellum from all exposed animals, 11 genes were upregulated in a range of 1.34-2.74 fold and one gene was downregulated 0.48-fold ($P < .0025$). The induced genes encode proteins with diverse functions including neurotransmitter regulation, blood-brain barrier (BBB), and melatonin production. The data shows that GSM MWs at 915 MHz did not induce PFGE-detectable DNA double stranded breaks or changes in chromatin conformation, but affected expression of genes in rat brain cells

Phillips, J.L., Ivaschuk, O., Ishida-Jones, T., Jones, R.A., Campbell-Beachler, M. and Haggren, W. DNA damage in Molt-4 T- lymphoblastoid cells exposed to cellular telephone radiofrequency fields in vitro. *Bioelectrochem. Bioenerg.* 45:103-110, 1998. Molt-4 T-lymphoblastoid cells have been exposed to pulsed signals at cellular telephone frequencies of 813.5625 MHz (iDEN signal) and 836.55 MHz (TDMA signal). These studies were performed at low SAR (average = 2.4 and 24 microwatt/g for iDEN and 2.6 and 26 microwatt/g for TDMA) in studies designed to look for athermal RF effects. The alkaline comet, or single cell gel electrophoresis, assay was employed to measure DNA single-strand breaks in cell cultures exposed to the radiofrequency (RF) signal as compared to concurrent sham-exposed cultures. Tail moment and comet extent were calculated as indicators of DNA damage. Statistical differences in the distribution of values for tail moment and comet extent between exposed and control cell cultures were evaluated with the SKolmogorov-Smirnoff distribution test. Data points for all

Studies that show **Cell Phone** Health Effects

experiments of each exposure condition were pooled and analyzed as single groups. It was found that: 1) exposure of cells to the iDEN signal at an SAR of 2.4 microwatt/g for 2 h or 21 h significantly decreased DNA damage; 2) exposure of cells to the TDMA signal at an SAR of 2.6 microwatt/g for 2 h and 21 h significantly decreased DNA damage; 3) exposure of cells to the iDEN signal at an SAR of 24 microwatt/g for 2 h and 21 h significantly increased DNA damage; 4) exposure of cells to the TDMA signal at an SAR of 26 microwatt/g for 2 h significantly decreased DNA damage. The data indicate a need to study the effects of exposure to RF signals on direct DNA damage and on the rate at which DNA damage is repaired.

Belyaev IY, Markovà E, Hillert L, Malmgren LO, Persson BR. Microwaves from UMTS/GSM mobile phones induce long-lasting inhibition of 53BP1/gamma-H2AX DNA repair foci in human lymphocytes. *Bioelectromagnetics*.30(2):129-141, 2009

We have recently described frequency-dependent effects of mobile phone microwaves (MWs) of global system for mobile communication (GSM) on human lymphocytes from persons reporting hypersensitivity to electromagnetic fields and healthy persons. Contrary to GSM, universal global telecommunications system (UMTS) mobile phones emit wide-band MW signals. Hypothetically, UMTS MWs may result in higher biological effects compared to GSM signal because of eventual "effective" frequencies within the wideband. Here, we report for the first time that UMTS MWs affect chromatin and inhibit formation of DNA double-strand breaks co-localizing 53BP1/gamma-H2AX DNA repair foci in human lymphocytes from hypersensitive and healthy persons and confirm that effects of GSM MWs depend on carrier frequency. Remarkably, the effects of MWs on 53BP1/gamma-H2AX foci persisted up to 72 h following exposure of cells, even longer than the stress response following heat shock. The data are in line with the hypothesis that the type of signal, UMTS MWs, may have higher biological efficiency and possibly larger health risk effects compared to GSM radiation emissions. No significant differences in effects between groups of healthy and hypersensitive subjects were observed, except for the effects of UMTS MWs and GSM-915 MHz MWs on the formation of the DNA repair foci, which were different for hypersensitive ($P < 0.02[53BP1]/0.01[\text{gamma-H2AX}]$) but not for control subjects ($P > 0.05$). The non-parametric statistics used here did not indicate specificity of the differences revealed between the effects of GSM and UMTS MWs on cells from hypersensitive subjects and more data are needed to study the nature of these differences.

Belyaev IY. Radiation-induced DNA repair foci: Spatio-temporal aspects of formation, application for assessment of radiosensitivity and biological dosimetry. *Mutat Res*. 704(1-3):132-141, 2010.

Several proteins involved in DNA repair and DNA damage signaling have been shown to produce discrete foci in response to ionizing radiation. These foci are believed to co-localize to DSB and referred to as ionizing radiation-induced foci (IRIF) or DNA repair foci. Recent studies have revealed that some residual IRIF remain in cells for a relatively long time after irradiation, and have indicated a possible correlation between radiosensitivity of cells and residual IRIF. Remarkably, residual foci are significantly larger in size than

Studies that show **Cell Phone** Health Effects

the initial foci. Increase in the size of IRIF with time upon irradiation has been found in various cell types and has partially been correlated with dynamics and fusion of initial foci. Although it is admitted that the number of IRIF reflect that of DSB, several studies report a lack of correlation between kinetics for IRIF and DSB and a lack of co-localization between DSB repair proteins. These studies suggest that some proportion of residual IRIF that depend on cell type, dose, and post-irradiation time may represent alternations in chromatin structure after DSB have been repaired or misrepaired. While precise functions of residual foci are presently unknown, their possible link to remaining chromatin alternations, nuclear matrix, apoptosis, delayed repair and misrejoining of DSB, activity of several kinases, phosphatases, and checkpoint signaling has been suggested. Another intriguing possibility is that some of DNA repair foci may mark break-points at chromosomal aberrations (CA). While this possibility has not been confirmed substantially, the residual foci seem to be useful for biological dosimetry and estimation of individual radiosensitivity in radiotherapy of cancer.

Dawe AS, Smith B, Thomas DW, Greedy S, Vasic N, Gregory A, Loader B, de Pomerai DI. A small temperature rise may contribute towards the apparent induction by microwaves of heat-shock gene expression in the nematode *Caenorhabditis Elegans*. Bioelectromagnetics.27(2):88-97, 2006.

We have previously reported that low intensity microwave exposure (0.75-1.0 GHz CW at 0.5 W; SAR 4-40 mW/kg) can induce an apparently non-thermal heat-shock response in *Caenorhabditis elegans* worms carrying hsp16-1::reporter genes. Using matched copper TEM cells for both sham and exposed groups, we can detect only modest reporter induction in the latter exposed group (15-20% after 2.5 h at 26 degrees C, rising to approximately 50% after 20 h). Traceable calibration of our copper TEM cell by the National Physical Laboratory (NPL) reveals significant power loss within the cell (8.5% at 1.0 GHz), accompanied by slight heating of exposed samples (approximately 0.3 degrees C at 1.0 W). Thus, exposed samples are in fact slightly warmer (by ≤ 0.2 degrees C at 0.5 W) than sham controls. Following NPL recommendations, our TEM cell design was modified with the aim of reducing both power loss and consequent heating. In the modified silver-plated cell, power loss is only 1.5% at 1.0 GHz, and sample warming is reduced to approximately 0.15 degrees C at 1.0 W (i.e., ≤ 0.1 degrees C at 0.5 W). Under sham:sham conditions, there is no difference in reporter expression between the modified silver-plated TEM cell and an unmodified copper cell. However, worms exposed to microwaves (1.0 GHz and 0.5 W) in the silver-plated cell also show no detectable induction of reporter expression relative to sham controls in the copper cell. Thus, the 20% "microwave induction" observed using two copper cells may be caused by a small temperature difference between sham and exposed conditions. In worms incubated for 2.5 h at 26.0, 26.2, and 27.0 degrees C with no microwave field, there is a consistent and significant increase in reporter expression between 26.0 and 26.2 degrees C (by approximately 20% in each of the six independent runs), but paradoxically expression levels at 27.0 degrees C are similar to those seen at 26.0 degrees C. This surprising result is in line with other evidence pointing towards complex regulation of hsp16-1 gene expression across the sub-heat-shock range of 25-27.5 degrees C in C.

Studies that show **Cell Phone** Health Effects

elegans. We conclude that our original interpretation of a non-thermal effect of microwaves cannot be sustained; at least part of the explanation appears to be thermal.

P, Tattersall J, Jones D, Candido P, Non-thermal heat-shock response to microwaves, Nature 405:417-418, 2000.

Nematode worms (*C. elegans*) exposed overnight to 750-MHz microwaves at a SAR of 0.001 W/kg showed an increased in heat shock proteins (HSPs). (Heat shock proteins are induced in most organisms by adverse conditions (such as heat or toxins) that cause damage to cellular proteins, acting as molecular chaperones to rescue damaged proteins). The authors give several arguments that the microwave-induced effect on HSPs is non-thermal and suggest that 'current exposure limits for microwave equipment may need to be reconsidered.

de Pomerai DI, Dawe A, DjerbibL, Allan, Brunt G, Daniells C. Growth and maturation of the nematode *Caenorhabditis elegans* following exposure to weak microwave fields. Enzyme Microbial Tech 30:73-79, 2002.

Prolonged exposure to weak microwave fields (750-1000 MHz, 0.5 W) at 25°C induces a heat-shock response in transgenic *C. elegans* strains carrying *hsp16* reporter genes [1]. A comparable response to heat alone requires a substantially higher temperature of 28°C, suggesting that microwave heating of worms or of the system as a whole might provide a sufficient explanation, although this can be ruled out by indirect arguments [1]. Here we investigate two further biological consequences of prolonged microwave exposure at 25°C in synchronised cultures of wild-type worm larvae, namely alterations in (i) growth rate (GR) and (ii) the proportion of worms later maturing into egg-bearing adults (MP). Both of these parameters are significantly increased following microwave exposure (GR by 8-11%, and MP by 28-40%), whereas both are significantly decreased (GR by 10% and MP almost abolished) after mild heat treatment at 28°C for the same period. It follows that the biological consequences of microwave exposure are opposite to, and therefore incompatible with, those attributable to mild heating. This evidence does not in itself necessitate a non-thermal mechanism, but does eliminate explanations that invoke the bulk heating of tissues by microwaves. This latter, however, remains the sole basis for current regulations governing microwave exposure.

de Pomerai DI, Smith B, Dawe A, North K, Smith T, Archer DB, Duce IR, Jones D, Candido EP. Microwave radiation can alter protein conformation without bulk heating. FEBS Lett 543(1-3):93-97, 2003.

Exposure to microwave radiation enhances the aggregation of bovine serum albumin in vitro in a time- and temperature-dependent manner. Microwave radiation also promotes amyloid fibril formation by bovine insulin at 60 degrees C. These alterations in protein conformation are not accompanied by measurable temperature changes, consistent with estimates from field modelling of the specific absorbed radiation (15-20 mW kg⁻¹). Limited denaturation of cellular proteins could explain our previous observation that modest heat-shock responses are induced by microwave exposure in *Caenorhabditis elegans*. We also show that heat-shock responses both to heat and

Studies that show **Cell Phone** Health Effects

microwaves are suppressed after RNA interference ablating heat-shock factor function.

Tkalec M, Stambuk A, Srut M, Malarić K, Klobučar GI. Oxidative and genotoxic effects of 900 MHz electromagnetic fields in the earthworm *Eisenia fetida*. Ecotoxicol Environ Saf. 90:7-12, 2013.

Accumulating evidence suggests that exposure to radiofrequency electromagnetic field (RF-EMF) can have various biological effects. In this study the oxidative and genotoxic effects were investigated in earthworms *Eisenia fetida* exposed in vivo to RF-EMF at the mobile phone frequency (900 MHz). Earthworms were exposed to the homogeneous RF-EMF at field levels of 10, 23, 41 and 120 V m⁻¹ for a period of 2h using a Gigahertz Transversal Electromagnetic (GTEM) cell. At the field level of 23 V m⁻¹ the effect of longer exposure (4h) and field modulation (80% AM 1 kHz sinusoidal) was investigated as well. All exposure treatments induced significant genotoxic effect in earthworms coelomocytes detected by the Comet assay, demonstrating DNA damaging capacity of 900 MHz electromagnetic radiation. Field modulation additionally increased the genotoxic effect. Moreover, our results indicated the induction of antioxidant stress response in terms of enhanced catalase and glutathione reductase activity as a result of the RF-EMF exposure, and demonstrated the generation of lipid and protein oxidative damage. Antioxidant responses and the potential of RF-EMF to induce damage to lipids, proteins and DNA differed depending on the field level applied, modulation of the field and duration of *E. fetida* exposure to 900 MHz electromagnetic radiation. Nature of detected DNA lesions and oxidative stress as the mechanism of action for the induction of DNA damage are discussed.

Effects on Pregnancy and Fetus

Luo Q, Jiang Y, Jin M, Xu J, Huang HF. Proteomic Analysis on the Alteration of Protein Expression in the Early-Stage Placental Villous Tissue of Electromagnetic Fields Associated With Cell Phone Exposure. Reprod Sci. 2013 Feb 18. [Epub ahead of print]

Background:To explore the possible adverse effects and search for cell phone electromagnetic field (EMF)-responsive proteins in human early reproduction, a proteomics approach was employed to investigate the changes in protein expression profile induced by cell phone EMF in human chorionic tissues of early pregnancy in vivo.**Methods:**Volunteer women about 50 days pregnant were exposed to EMF at the average absorption rate of 1.6 to 8.8 W/kg for 1 hour with the irradiation device placed 10 cm away from the umbilicus at the midline of the abdomen. The changes in protein profile were examined using 2-dimensional electrophoresis (2-DE).**Results:**Up to 15 spots have yielded significant change at least 2- to 2.5-folds up or down compared to sham-exposed group. Twelve proteins were identified- procollagen-proline, eukaryotic translation elongation factor 1 delta, chain D crystal structure of human vitamin D-binding protein, thioredoxin-like 3, capping protein, isocitrate dehydrogenase 3 alpha, calumenin, Catechol-O-methyltransferase protein, proteinase inhibitor 6 (PI-6; SerpinB6) protein, 3,2-trans-enoyl-CoA isomerase protein, chain B human erythrocyte 2,3-

Studies that show **Cell Phone** Health Effects

bisphosphoglycerate mutase, and nucleoprotein. Conclusion: Cell phone EMF might alter the protein profile of chorionic tissue of early pregnancy, during the most sensitive stage of the embryos. The exposure to EMF may cause adverse effects on cell proliferation and development of nervous system in early embryos. Furthermore, 2-DE coupled with mass spectrometry is a promising approach to elucidate the effects and search for new biomarkers for environmental toxic effects.

Odacı E, Unal D, Mercantepe T, Topal Z, Hancı H, Türedi S, Erol H, Mungan S, Kaya H, Colakoğlu S. Pathological effects of prenatal exposure to a 900 MHz electromagnetic field on the 21-day-old male rat kidney. Biotech Histochem. 2014 Aug 27:1-9. [Epub ahead of print]

We investigated the effects on kidney tissue of 900 megahertz (MHz) EMF applied during the prenatal period. Pregnant rats were exposed to 900 MHz EMF, 1 h/day, on days 13-21 of pregnancy; no procedure was performed on control group pregnant rats or on mothers or newborns after birth. On postnatal day 21, kidney tissues of male rat pups from both groups were examined by light and electron microscopy. Malondialdehyde (MDA), superoxide dismutase (SOD), catalase (CAT) and glutathione levels also were investigated. Light microscopy revealed some degenerative changes in the tubule epithelium, small cystic formations in the primitive tubules and large cysts in the cortico-medullary or medullary regions in the experimental group. Electron microscopy revealed a loss of peritubular capillaries and atypical parietal layer epithelial cells in the experimental group. Biochemical analysis showed significantly increased MDA levels in the experimental group and decreased SOD and CAT levels. EMF applied during the prenatal period can caused pathological changes in kidney tissue in 21-day-old male rats owing to oxidative stress and decreased antioxidant enzyme levels.

Liu XY, Bian XM, Han JX, Cao ZJ, Fan GS, Zhang C, Zhang WL, Zhang SZ, Sun XG. [Risk factors in the living environment of early spontaneous abortion pregnant women] Zhongguo Yi Xue Ke Xue Yuan Xue Bao. 29(5):661-664, 2007. [Article in Chinese]

OBJECTIVE: To study the relationship between early spontaneous abortion and living environment, and explore the risk factors of spontaneous abortion. METHODS: We conducted analysis based on the interview of 200 spontaneous abortion cases and the matched control (age +/- 2 years) by using multifactor Logistic regression analysis. RESULTS: The proportions of watching TV > or =10 hours/week, operating computer > or =45 hours/week, using copycat, microwave oven and mobile phone, electromagnetism equipment near the dwell or work place, e. g. switch room < or =50 m and launching tower < or =500 m in the cases are significantly higher than those in the controls in single factor analysis (all P < 0.05). After adjusted the effect of other risk factors by multifactor analysis, using microwave oven and mobile phone, contacting abnormal smell of fitment material > or =3 months, having emotional stress during the first term of pregnancy and spontaneous abortion history were significantly associated with risk of spontaneous abortion. The odds ratios of these risk factors were 2.23 and 4.63, respectively. CONCLUSION: Using microwave oven and mobile phone, contacting

Studies that show **Cell Phone** Health Effects

abnormal smell of fitment material > or =3 months, having emotional stress during the first term of pregnancy, and spontaneous abortion history are risk factors of early spontaneous abortion.

Rezk AY, Abdulqawi K, Mustafa RM, Abo El-Azm TM, Al-Inany H. Fetal and neonatal responses following maternal exposure to mobile phones. *Saudi Med J.* 29(2):218-223, 2008.

OBJECTIVE: To study fetal and neonatal heart rate (HR) and cardiac output (COP), following acute maternal exposure to electromagnetic fields (EMF) emitted by mobile phones. **METHODS:** The present study was carried out at Benha University Hospital and El-Shorouq Hospital, Cairo, Egypt, from October 2003 to March 2004. Ninety women with uncomplicated pregnancies aged 18-33 years, and 30 full term healthy newborn infants were included. The pregnant mothers were exposed to EMF emitted by mobile telephones while on telephone-dialing mode for 10 minutes during pregnancy and after birth. The main outcome were measurements of fetal and neonatal HR and COP.

RESULTS: A statistical significant increase in fetal and neonatal HR, and statistical significant decrease in stroke volume and COP before and after use of mobile phone were noted. All these changes are attenuated with increase in gestational age.

CONCLUSION: Exposure of pregnant women to mobile phone significantly increase fetal and neonatal HR, and significantly decreased the COP.

Pyrpasopoulou A, Kotoula V, Cheva A, Hytioglou P, Nikolakaki E, Magras IN, Xenos TD, Tsiboukis TD, Karkavelas G. Bone morphogenetic protein expression in newborn rat kidneys after prenatal exposure to radiofrequency radiation. *Bioelectromagnetics* 25(3):216-227, 2004.

Effects of nonthermal radiofrequency radiation (RFR) of the global system of mobile communication (GSM) cellular phones have been as yet mostly studied at the molecular level in the context of cellular stress and proliferation, as well as neurotransmitter production and localization. In this study, a simulation model was designed for the exposure of pregnant rats to pulsed GSM-like RFR (9.4 GHz), based on the different resonant frequencies of man and rat. The power density applied was 5 microW/cm², in order to avoid thermal electromagnetic effects as much as possible. Pregnant rats were exposed to RFR during days 1-3 postcoitum (p.c.) (embryogenesis, pre-implantation) and days 4-7 p.c. (early organogenesis, peri-implantation). Relative expression and localization of bone morphogenetic proteins (BMP) and their receptors (BMPR), members of a molecular family currently considered as major endocrine and autocrine morphogens and known to be involved in renal development, were investigated in newborn kidneys from RFR exposed and sham irradiated (control) rats. Semi-quantitative duplex RT-PCR for BMP-4, -7, BMPR-IA, -IB, and -II showed increased BMP-4 and BMPR-IA, and decreased BMPR-II relative expression in newborn kidneys. These changes were statistically significant for BMP-4, BMPR-IA, and -II after exposure on days 1-3 p.c. (P <.001 each), and for BMP-4 and BMPR-IA after exposure on days 4-7 p.c. (P <.001 and P =.005, respectively). Immunohistochemistry and in situ hybridization (ISH) showed aberrant expression and localization of these molecules at the histological level.

Studies that show **Cell Phone** Health Effects

Our findings suggest that GSM-like RFR interferes with gene expression during early gestation and results in aberrations of BMP expression in the newborn. These molecular changes do not appear to affect renal organogenesis and may reflect a delay in the development of this organ. The differences of relative BMP expression after different time periods of exposure indicate the importance of timing for GSM-like RFR effects on embryonic development.

İkinci A, Odacı E, Yıldırım M, Kaya H, Akça M, Hancı H, Aslan A, Sönmez OF, Baş O. The Effects of Prenatal Exposure to a 900 Megahertz Electromagnetic Field on Hippocampus Morphology and Learning Behavior in Rat Pups. NeuroQuantology. 11(4):582-590, 2013.

The purpose of this study was to examine the effect on hippocampus morphology and learning behavior in rat pups following prenatal exposure to a 900 megahertz (MHz) electromagnetic field (EMF). Female Sprague Dawley rats weighing 180-250 g were left to mate with males. The following day, pregnant rats identified as such by the vaginal smear test were divided into two groups, control (n=3) and EMF (n=3). No procedures were performed on the control group. The rats in the EMF group were exposed to 900 MHz EMF on days 13 to 21 of pregnancy, for 1 h a day. Female rat pups were removed from their mothers at 22 days old. We then established two newborn rat groups, a 13 member control group and a 10 member EMF group. Radial arm maze and passive avoidance tests were used to measure rat pups' learning and memory performance. All rats were decapitated on the postnatal 32nd day. Routine histological procedures were performed on the brain tissues, and sections were stained with Cresyl fast violet. The radial arm maze (p=0.007) and passive avoidance (p=0.032) tests were administered to both groups under identical conditions, and compromised learning behavior was determined in the EMF group rats. Morphological compromise was also determined in the EMF group sections. Our results show that the application of a 900 MHz EMF in the prenatal period adversely affected female pups' learning behavior and also resulted in histopathological changes appearing in the hippocampus.

Ulubay M, Yahyazadeh A, Deniz OG, Kıvrak EG, Altunkaynak BZ, Erdem G, Kaplan S. Effects of prenatal 900 MHz electromagnetic field exposures on the histology of rat kidney. Int J Radiat Biol. 2014 Aug 1:1-25. [Epub ahead of print]

Purpose: To research the harmful effects of prenatal exposure of 900 megahertz (MHz) electromagnetic field (EMF) on kidneys of four-week-old male rats and to determine protective effects of melatonin (MEL) and omega-3 (ω -3). Materials and methods: Twenty-one Wistar albino rats were randomly placed into seven groups as follows: control (Cont), Sham, MEL, ω -3, EMF, EMF+MEL and EMF+ ω -3. After mating, three groups (EMF, EMF+MEL, EMF+ ω -3) were exposed to an EMF. In the fourth week subsequent to parturition, six rats were randomly chosen from each group. Mean volume of kidneys and renal cortices, the total number of glomeruli and basic histological structure of kidney were evaluated by stereological and light microscopical methods, respectively. Results: Stereological results determined the mean volume of the kidneys and cortices were significantly increased in EMF-exposed groups compared

Studies that show **Cell Phone** Health Effects

to the Cont group. However, EMF-unexposed groups were not significantly modified compared to the Cont group. Additionally, the total number of glomeruli was significantly higher in EMF-unexposed groups compared to the Cont group. Alternatively, the number of glomeruli in EMF-exposed groups was decreased compared to the Cont group. Conclusions: Prenatal exposure of rat kidneys to 900 MHz EMF resulted in increased total kidney volume and decreased the numbers of glomeruli. Moreover, MEL and ω -3 prevented adverse effects of EMF on the kidneys.

Köktürk S, Yardimoglu M, Celikozlu SD, Dolanbay EG, Cimbiz A. Effect of *Lycopersicon esculentum* extract on apoptosis in the rat cerebellum, following prenatal and postnatal exposure to an electromagnetic field. *Exp Ther Med.* 6(1):52-56, 2013.

The expansion of mobile phone technology has raised concerns regarding the effect of 900-MHz electromagnetic field (EMF) exposure on the central nervous system. At present, the developing human brain is regularly exposed to mobile telephones, pre- and postnatally. Several studies have demonstrated the acute effects of EMF exposure during pre- or postnatal periods; however, the chronic effects of EMF exposure are less understood. Thus, the aim of the present study was to determine the chronic effects of EMF on the pre- and postnatal rat cerebellum. The control group was maintained in the same conditions as the experimental groups, without the exposure to EMF. In the EMF1 group, the rats were exposed to EMF during pre- and postnatal periods (until postnatal day 80). In the EMF2 group, the rats were also exposed to EMF pre- and postnatally; in addition, however, they were provided with a daily oral supplementation of *Lycopersicon esculentum* extract (~2 g/kg). The number of caspase-3-labeled Purkinje neurons and granule cells present in the rats in the control and experimental groups were then counted. The neurodegenerative changes were studied using cresyl violet staining, and these changes were evaluated. In comparison with the control animals, the EMF1 group demonstrated a significant increase in the number of caspase-3-labeled Purkinje neurons and granule cells present in the cerebellum ($P < 0.001$). However, in comparison with the EMF1 group, the EMF2 group exhibited significantly fewer caspase-3-labeled Purkinje neurons and granule cells in the cerebellum. In the EMF1 group, the Purkinje neurons were revealed to have undergone dark neuron degenerative changes. However, the presence of dark Purkinje neurons was reduced in the EMF2 group, compared with the EMF1 group. The results indicated that apoptosis and neurodegeneration in rats exposed to EMF during pre- and postnatal periods may be reduced with *Lycopersicon esculentum* extract therapy.

Tillmann T, Ernst H, Streckert J, Zhou Y, Taugner F, Hansen V, Dasenbrock C. Indication of cocarcinogenic potential of chronic UMTS-modulated radiofrequency exposure in an ethylnitrosourea mouse model. *nt J Radiat Biol.* 86(7):529-541, 2010.

PURPOSE: To evaluate putative effects on tumour susceptibility in mice exposed to a UMTS (universal mobile telecommunications system) test signal for up to 24 months, commencing with embryo-fetal exposure. **MATERIAL AND METHODS:** Animals were exposed to UMTS fields with intensities of 0, 4.8, and 48 W/m², the low-dose group (4.8 W/m²) was subjected to additional prenatal ethylnitrosourea treatment (40 mg

Studies that show **Cell Phone** Health Effects

ENU/kg body weight).RESULTS: The high-level UMTS exposure (48 W/m(2)), the sham exposure, and the cage control groups showed comparable tumour incidences in the protocol organs. In contrast, the ENU-treated group UMTS-exposed at 4.8 W/m(2) displayed an enhanced lung tumour rate and an increased incidence of lung carcinomas as compared to the controls treated with ENU only. Furthermore, tumour multiplicity of the lung carcinomas was increased and the number of metastasising lung tumours was doubled in the ENU/UMTS group as compared to the ENU control group.CONCLUSION: This pilot study indicates a cocarcinogenic effect of lifelong UMTS exposure (4.8 W/m(2)) in female B6C3F1 descendants subjected to pretreatment with ethylnitrosourea.

Türedi S, Hancı H, Topal Z, Unal D, Mercantepe T, Bozkurt I, Kaya H, Odacı E. The effects of prenatal exposure to a 900-MHz electromagnetic field on the 21-day-old male rat heart. Electromagn Biol Med. 2014 Aug 28:1-8. [Epub ahead of print]

Abstract The growing spread of mobile phone use is raising concerns about the effect on human health of the electromagnetic field (EMF) these devices emit. The purpose of this study was to investigate the effects on rat pup heart tissue of prenatal exposure to a 900 megahertz (MHz) EMF. For this purpose, pregnant rats were divided into experimental and control groups. Experimental group rats were exposed to a 900 MHz EMF (1 h/d) on days 13-21 of pregnancy. Measurements were performed with rats inside the exposure box in order to determine the distribution of EMF intensity. Our measurements showed that pregnant experimental group rats were exposed to a mean electrical field intensity of 13.77 V/m inside the box (0.50 W/m²). This study continued with male rat pups obtained from both groups. Pups were sacrificed on postnatal day 21, and the heart tissues were extracted. Malondialdehyde, superoxide dismutase and catalase values were significantly higher in the experimental group rats, while glutathione values were lower. Light microscopy revealed irregularities in heart muscle fibers and apoptotic changes in the experimental group. Electron microscopy revealed crista loss and swelling in the mitochondria, degeneration in myofibrils and structural impairments in Z bands. Our study results suggest that exposure to EMF in the prenatal period causes oxidative stress and histopathological changes in male rat pup heart tissue.

Aldad TS, Gan G, Gao XB, Taylor HS.Fetal radiofrequency radiation exposure from 800-1900 mhz-rated cellular telephones affects neurodevelopment and behavior in mice.Sci Rep. 2:312, 2012.

Neurobehavioral disorders are increasingly prevalent in children, however their etiology is not well understood. An association between prenatal cellular telephone use and hyperactivity in children has been postulated, yet the direct effects of radiofrequency radiation exposure on neurodevelopment remain unknown. Here we used a mouse model to demonstrate that in-utero radiofrequency exposure from cellular telephones does affect adult behavior. Mice exposed in-utero were hyperactive and had impaired memory as determined using the object recognition, light/dark box and step-down assays. Whole cell patch clamp recordings of miniature excitatory postsynaptic currents

Studies that show **Cell Phone** Health Effects

(mEPSCs) revealed that these behavioral changes were due to altered neuronal developmental programming. Exposed mice had dose-responsive impaired glutamatergic synaptic transmission onto layer V pyramidal neurons of the prefrontal cortex. We present the first experimental evidence of neuropathology due to in-utero cellular telephone radiation. Further experiments are needed in humans or non-human primates to determine the risk of exposure during pregnancy.

Barteri M, Pala A, Rotella S. Structural and kinetic effects of mobile phone microwaves on acetylcholinesterase activity. Biophys Chem. 113(3):245-253, 2005.

The present study provides evidence that "in vitro" simple exposure of an aqueous solution of electric eel acetylcholinesterase (EeAChE; EC 3.1.1.7.) to cellular phone emission alters its enzymatic activity. This paper demonstrates, by combining different experimental techniques, that radio frequency (RF) radiations irreversibly affect the structural and biochemical characteristics of an important CNS enzyme. These results were obtained by using a commercial cellular phone to reproduce the reality of the human exposition. This experimental procedure provided surprising effects collected practically without experimental errors because they were obtained comparing native and irradiated sample of the same enzyme solution. Although these results cannot be used to conclude whether exposure to RF during the use of cellular phone can lead to any hazardous health effect, they may be a significant first step towards further verification of these effects on other "ex vivo" or "in vivo" biological systems.

Cetin H, Nazıroğlu M, Celik O, Yüksel M, Pastacı N, Ozkaya MO. Liver antioxidant stores protect the brain from electromagnetic radiation (900 and 1800 MHz)-induced oxidative stress in rats during pregnancy and the development of offspring. J Matern Fetal Neonatal Med. 2014 Mar 3. [Epub ahead of print]

Objectives: The present study determined the effects of mobile phone (900 and 1800 MHz)-induced electromagnetic radiation (EMR) exposure on oxidative stress in the brain and liver as well as the element levels in growing rats from pregnancy to 6 weeks of age. **Methods:** Thirty-two rats and their offspring were equally divided into 3 different groups: the control, 900 MHz, and 1800 MHz groups. The 900 MHz and 1800 MHz groups were exposed to EMR for 60 min/day during pregnancy and neonatal development. At the 4th, 5th, and 6th weeks of the experiment, brain samples were obtained. **Results:** Brain and liver glutathione peroxidase (GSH-Px) activities, as well as liver vitamin A and β -carotene concentrations decreased in the EMR groups, although brain iron, vitamin A, and β -carotene concentrations increased in the EMR groups. In the 6th week, selenium concentrations in the brain decreased in the EMR groups. There were no statistically significant differences in glutathione, vitamin E, chromium, copper, magnesium, manganese, and zinc concentrations between the 3 groups. **Conclusion:** EMR-induced oxidative stress in the brain and liver was reduced during the development of offspring. Mobile phone-induced EMR could be considered as a cause of oxidative brain and liver injury in growing rats.

Odaci E, Bas O, Kaplan S. Effects of prenatal exposure to a 900 MHz electromagnetic

Studies that show **Cell Phone** Health Effects

field on the dentate gyrus of rats: a stereological and histopathological study.Brain Res. 1238:224-229, 2008.

Electromagnetic fields (EMFs) inhibit the formation and differentiation of neural stem cells during embryonic development. In this study, the effects of prenatal exposure to EMF on the number of granule cells in the dentate gyrus of 4-week-old rats were investigated. This experiment used a control (Cont) group and an EMF exposed (EMF) group (three pregnant rats each group). The EMF group consisted of six offspring (n=6) of pregnant rats that were exposed to an EMF of up to 900 megahertz (MHz) for 60 min/day between the first and last days of gestation. The control group consisted of five offspring (n=5) of pregnant rats that were not treated at all. The offspring were sacrificed when they were 4 weeks old. The numbers of granule cells in the dentate gyrus were analyzed using the optical fractionator technique. The results showed that prenatal EMF exposure caused a decrease in the number of granule cells in the dentate gyrus of the rats (P<0.01). This suggests that prenatal exposure to a 900 MHz EMF affects the development of the dentate gyrus granule cells in the rat hippocampus. Cell loss might be caused by an inhibition of granule cell neurogenesis in the dentate gyrus.

Odacı E, İkinci A, Yıldırım M, Kaya H, Akça M, Hancı H, Sönmez OF, Aslan A, Okuyan M, Baş O. The Effects of 900 Megahertz Electromagnetic Field Applied in the Prenatal Period on Spinal Cord Morphology and Motor Behavior in Female Rat Pups. NeuroQuantology 11:573-581, 2013.

This study investigated the effect of a 900 megahertz (MHz) electromagnetic field (EMF) applied in the prenatal period on the spinal cord and motor behavior of female rat pups. Beginning of the study, female Sprague Dawley rats (180–250 g) were left to mate with male rats. Rats identified as pregnant were then divided into control (n=3) and EMF groups (n=3). The EMF group was exposed to 1-h 900 MHz EMF daily between days 13 and 21 of pregnancy. At 21 days old, rat pups were removed from their mothers and divided into two newborn rat groups, control (n=13) and EMF (n=10). The rotarod test was applied to the rat pups to assess motor functions and the open field test to evaluate locomotor activity. On day 32 of the study, the rat pups were decapitated, and the spinal cord in the upper thoracic region was removed. Following routine histological tests, they were stained with Cresyl fast violet. Rotarod test results revealed a significant increase in EMF group rat pups' motor functions (p=0.037). However, no difference was observed in the open field test results (p>0.05). In the EMF group' rat pups, we observed pathological changes in the spinal cord. On the basis of our results, 900 MHz EMF applied in the prenatal period affected spinal cord development. This effect was observed in the form of pathological changes in the spinal cord of rat pups, and it may be that these pathological changes led to an increase in rat pups' motor activities.

Jing J, Yuhua Z, Xiao-qian Y, Rongping J, Dong-mei G, Xi C. The influence of microwave radiation from cellular phone on fetal rat brain. Electromagn Biol Med. 31(1):57-66, 2012.

The increasing use of cellular phones in our society has brought focus on the potential detrimental effects to human health by microwave radiation. The aim of our study was

Studies that show **Cell Phone** Health Effects

to evaluate the intensity of oxidative stress and the level of neurotransmitters in the brains of fetal rats chronically exposed to cellular phones. The experiment was performed on pregnant rats exposed to different intensities of microwave radiation from cellular phones. Thirty-two pregnant rats were randomly divided into four groups: CG, GL, GM, and GH. CG accepted no microwave radiation, GL group radiated 10 min each time, GM group radiated 30 min, and GH group radiated 60 min. The 3 experimental groups were radiated 3 times a day from the first pregnant day for consecutively 20 days, and on the 21st day, the fetal rats were taken and then the contents of superoxide dismutase (SOD), glutathione peroxidase (GSH-Px), malondialdehyde (MDA), noradrenaline (NE), dopamine (DA), and 5-hydroxyindole acetic acid (5-HT) in the brain were assayed. Compared with CG, there were significant differences ($P < 0.05$) found in the contents of SOD, GSH-Px, and MDA in GM and GH; the contents of SOD and GSH-Px decreased and the content of MDA increased. The significant content differences of NE and DA were found in fetal rat brains in GL and GH groups, with the GL group increased and the GH group decreased. Through this study, we concluded that receiving a certain period of microwave radiation from cellular phones during pregnancy has certain harm on fetal rat brains.

Dasdag S; Akdag MZ; Ayyıldız O, Demirtas OC, Yayla M, Sert C. Do cellular phones alter blood parameters and birth weight of rats? Electromag Biol Med. 19:107-113, 2000.

The present study aimed to investigate the effects of microwaves (MW) emitted by cellular phones (CPs) on peripheral blood parameters and birth weights of rats. Thirty-six albino rats were divided into four groups, male ($n = 6$) and female sham-exposed groups ($n = 12$) and male ($n = 6$) and female experimental groups ($n = 12$). No blood parameters differed following exposure ($p > 0.05$). The birth weight of offspring in the experimental group was significantly lower than in the sham-exposed group ($p < 0.001$). No significant differences were observed between rectal temperatures of rats in the sham and experimental groups ($p > 0.05$). The specific absorption rate (SAR) was found to be 0.155 W/kg for the experimental groups. All parameters investigated were normal in the next generation of rats ($p > 0.05$).

Fragopoulou AF, Koussoulakos SL, Margaritis LH. Cranial and postcranial skeletal variations induced in mouse embryos by mobile phone radiation. Pathophysiology. 17(3):169-177, 2010.

This study focuses on foetal development following mild daily exposure of pregnant mice to near field electromagnetic radiation emitted by a mobile phone. The investigation was motivated by the fact that the potentially hazardous electromagnetic radiation emitted by mobile phones is currently of tremendous public interest. Physically comparable pregnant mice were exposed to radiofrequency radiation GSM 900MHz emitted by a mobile phone. Within 5h after birth most cubs were fixed followed by double staining in toto, and conventional paraffin histology. Other cubs remained with their mothers until teeth eruption. Structural development was assessed by examining newborns for the presence of anomalies and/or variations in soft tissues

Studies that show **Cell Phone** Health Effects

and skeletal anatomy. Electromagnetic radiofrequency exposed newborns, externally examined, displayed a normal phenotype. Histochemical and histological studies, however, revealed variations in the exposed fetuses with respect to control ones concerning the ossification of cranial bones and thoracic cage ribs, as well as displacement of Meckelian cartilage. Littermates examined after teeth eruption displayed normal phenotypes. It is concluded that mild exposure to mobile phone radiation may affect, although transiently, mouse foetal development at the ossification level. The developmental variations observed could be explained by considering the different embryonic origin and mode of ossification of the affected skeletal elements.

Ferreira AR, Knakievicz T, de Bittencourt Pasquali MA, Gelain DP, Dal-Pizzol F, Fernandez CE, de Almeida de Salles AA, Ferreira HB, Moreira JC. Ultra high frequency-electromagnetic field irradiation during pregnancy leads to an increase in erythrocytes micronuclei incidence in rat offspring. Life Sci. 80(1)43-50, 2006.

Mobile telephones and their base stations are an important ultra high frequency-electromagnetic field (UHF-EMF) source and their utilization is increasing all over the world. Epidemiological studies suggested that low energy UHF-EMF emitted from a cellular telephone may cause biological effects, such as DNA damage and changes on oxidative metabolism. An in vivo mammalian cytogenetic test, the micronucleus (MN) assay, was used to investigate the occurrence of chromosomal damage in erythrocytes from rat offspring exposed to a non-thermal UHF-EMF from a cellular phone during their embryogenesis; the irradiated group showed a significant increase in MN occurrence. In order to investigate if UHF-EMF could also alter oxidative parameters in the peripheral blood and in the liver - an important hematopoietic tissue in rat embryos and newborns - we also measured the activity of antioxidant enzymes, quantified total sulfhydryl content, protein carbonyl groups, thiobarbituric acid-reactive species and total non-enzymatic antioxidant defense. No significant differences were found in any oxidative parameter of offspring blood and liver. The average number of pups in each litter has also not been significantly altered. Our results suggest that, under our experimental conditions, UHF-EMF is able to induce a genotoxic response in hematopoietic tissue during the embryogenesis through an unknown mechanism.

Haghani M, Shabani M, Moazzami K. Maternal mobile phone exposure adversely affects the electrophysiological properties of Purkinje neurons in rat offspring. Neuroscience. 2013 Jul 29. pii: S0306-4522(13)00643-X. doi: 10.1016/j.neuroscience.2013.07.049. [Epub ahead of print]

Electromagnetic field (EMF) radiations emitted from mobile phones may cause structural damage to neurons. With the increased usage of mobile phones worldwide, concerns about their possible effects on the nervous system are rising. In the present study, we aimed to elucidate the possible effects of prenatal EMF exposure on the cerebellum of offspring Wistar rats. Rats in EMF group were exposed to 900 MHz Pulse-EMF irradiation for six hours per day during all gestation period. Ten offspring's per each group were evaluated for behavioral and electrophysiological evaluations. Cerebellum - related behavioral dysfunctions were analyzed using motor learning and cerebellum-

Studies that show **Cell Phone** Health Effects

dependent functional tasks (Accelerated Rotarod, Hanging and Open field tests). Whole cell- patch clamp recordings were used for electrophysiological evaluations. The results of the present study failed to show any behavioral abnormalities in rats exposed to chronic EMF radiation. However, whole cell patch clamp recordings revealed decreased neuronal excitability of Purkinje cells in rats exposed to EMF. The most prominent changes included afterhyperpolarization amplitude, spike frequency, half width and first spike latency. In conclusion, the results of the present study show that prenatal EMF exposure results in altered electrophysiological properties of Purkinje neurons. However, these changes may not be severe enough to alter the cerebellum-dependent functional tasks.

Col-Araz N. Evaluation of factors affecting birth weight and preterm birth in southern Turkey. J Pak Med Assoc. 2013 Apr;63(4):459-62.

OBJECTIVE: To identify factors affecting birth weight and pre-term birth, and to find associations with electromagnetic devices such as television, computer and mobile phones. **METHODS:** The study was conducted in Turkey at Gazintep University, Faculty of Medicine's Outpatient Clinic at the Paediatric Ward. It comprised 500 patients who presented at the clinic from May to December 2009. All participants were administered a questionnaire regarding their pregnancy history. SPSS 13 was used for statistical analysis. **RESULTS:** In the study, 90 (19%) patients had pre-term birth, and 64 (12.9%) had low birth weight rate Birth weight was positively correlated with maternal age and baseline maternal weight ($r = 0.115$, $p < 0.010$; $r = 0.168$, $p < 0.000$, respectively). Pre-term birth and birth weight less than 2500g were more common in mothers with a history of disease during pregnancy ($p < 0.046$ and $p < 0.008$, respectively). The habit of watching television and using mobile phones and computer by mothers did not demonstrate any relationship with birth weight. Mothers who used mobile phones or computers during pregnancy had more deliveries before 37 weeks ($p < 0.018$, $p < 0.034$; respectively). Similarly, pregnancy duration was shorter in mothers who used either mobile phone or computers during pregnancy ($p < 0.005$, $p < 0.048$, respectively). **CONCLUSION:** Mobile phones and computers may have an effect on pre-term birth.

Orhan Baş, Osman Fikret Sönmez, Ali Aslan, Ayşe İkinci, Hatice Hancı, Mehmet Yıldırım, Haydar Kaya, Metehan Akça, Ersan Odacı. Pyramidal Cell Loss in the Cornu Ammonis of 32-day-old Female Rats Following Exposure to a 900 Megahertz Electromagnetic Field During Prenatal Days 13–21. NeuroQuantology 11:591-599, 2013.

The number of studies reporting that the electromagnetic field (EMF) emitted by mobile phones affects human health is increasing by the day. In previous studies we reported that a 900 megahertz (MHz) EMF applied throughout the prenatal period reduced the number of pyramidal cells in the cornu ammonis of rat pups in the postnatal period. In this study we investigated the effect of a 900 MHz EMF applied on days 13-21 of the prenatal period on the number of pyramidal cells in the cornu ammonis of rat pups in the postnatal period. For that purpose, pregnant rats were divided into experimental and control groups. Experimental group pregnant rats were exposed to the effect of a

Studies that show **Cell Phone** Health Effects

900 MHz EMF on days 13-21 of pregnancy. No procedure was applied to the control group. Newborn female rat pups were added to the study, and no procedure was performed on these after birth. Five newborn female rats were obtained from the experimental group and six from the control group. All female rat pups were decapitated on the postnatal 32nd day, and histological procedures were performed on the brain tissues. Sections were stained with Cresyl fast violet. The optical dissector technique was used to estimate the total number of pyramidal cells in the cornu ammonis. Sections of cornu ammonis were subjected to histopathological evaluations. Our results showed that exposure to 900 MHz EMF during prenatal days 13-21 led to a significant decrease in the number of pyramidal cells in the cornu ammonis of the experimental group female rat pups ($P<0.05$). Histopathological examination revealed picnotic cells in the cornu ammonis in experimental female rat pups. The pyramidal cell loss in the cornu ammonis may therefore be attributed to exposure to 900 MHz EMF in days 13-21 of the prenatal period.

Ayşe İkinci, Ersan Odacı, Mehmet Yıldırım, Haydar Kaya, Metehan Akça, Hatice Hancı, Ali Aslan, Osman Fikret Sönmez, Orhan Baş. The Effects of Prenatal Exposure to a 900 Megahertz Electromagnetic Field on Hippocampus Morphology and Learning Behavior in Rat Pups. *NeuroQuantology*. 11(4):582-590. December 2013.

The purpose of this study was to examine the effect on hippocampus morphology and learning behavior in rat pups following prenatal exposure to a 900 megahertz (MHz) electromagnetic field (EMF). Female Sprague Dawley rats weighing 180-250 g were left to mate with males. The following day, pregnant rats identified as such by the vaginal smear test were divided into two groups, control ($n=3$) and EMF ($n=3$). No procedures were performed on the control group. The rats in the EMF group were exposed to 900 MHz EMF on days 13 to 21 of pregnancy, for 1 h a day. Female rat pups were removed from their mothers at 22 days old. We then established two newborn rat groups, a 13 member control group and a 10 member EMF group. Radial arm maze and passive avoidance tests were used to measure rat pups' learning and memory performance. All rats were decapitated on the postnatal 32nd day. Routine histological procedures were performed on the brain tissues, and sections were stained with Cresyl fast violet. The radial arm maze ($p=0.007$) and passive avoidance ($p=0.032$) tests were administered to both groups under identical conditions, and compromised learning behavior was determined in the EMF group rats. Morphological compromise was also determined in the EMF group sections. Our results show that the application of a 900 MHz EMF in the prenatal period adversely affected female pups' learning behavior and also resulted in histopathological changes appearing in the hippocampus.

Ozgur E, Kismali G, Guler G, Akcay A, Ozkurt G, Sel T, Seyhan N. Effects of Prenatal and Postnatal Exposure to GSM-Like Radiofrequency on Blood Chemistry and Oxidative Stress in Infant Rabbits, an Experimental Study. *Cell Biochem Biophys*. 2013 Mar 24. [Epub ahead of print]

Studies that show **Cell Phone** Health Effects

We aimed to investigate the potential hazardous effects of prenatal and/or postnatal exposure to 1800 MHz GSM-like radiofrequency radiation (RFR) on the blood chemistry and lipid peroxidation levels of infant rabbits. A total of 72 New Zealand female and male white rabbits aged 1-month were used. Thirty-six female and 36 male were divided into four groups which were composed of nine infants: (i) Group 1 were the sham exposure (control), (ii) Group 2 were exposed to RFR, 15 min daily for 7 days in the prenatal period (between 15th and 22nd days of the gestational period) (prenatal exposure group). (iii) Group 3 were exposed to RFR 15 min/day (14 days for male, whereas 7 days for female) after they reached 1-month of age (postnatal exposure group). (iv) Group 4 were exposed to RFR for 15 min daily during 7 days in the prenatal period (between 15th and 22nd days of the gestational period) and 15 min/day (14 days for male, whereas 7 days for female) after they reached 1-month of age (prenatal and postnatal exposure group). Results showed that serum lipid peroxidation level in both female and male rabbits changed due to the RFR exposure. However, different parameters of the blood biochemistry were affected by exposure in male and female infants. Consequently, the whole-body 1800 MHz GSM-like RFR exposure may lead to oxidative stress and changes on some blood chemistry parameters. Studies on RFR exposure during prenatal and postnatal periods will help to establish international standards for the protection of pregnant and newborns from environmental RFR.

Guler G, Tomruk A, Ozgur E, Seyhan N. The effect of radiofrequency radiation on DNA and lipid damage in non-pregnant and pregnant rabbits and their newborns. *Gen Physiol Biophys.* 29(1):59-66, 2010.

The concerns of people on possible adverse health effects of radiofrequency radiation (RFR) generated from mobile phones as well as their supporting transmitters (base stations) have increased markedly. RFR effect on oversensitive people, such as pregnant women and their developing fetuses, and older people is another source of concern that should be considered. In this study, oxidative DNA damage and lipid peroxidation levels in the brain tissue of pregnant and non-pregnant New Zealand White rabbits and their newborns exposed to RFR were investigated. Thirteen-month-old rabbits were studied in four groups as non-pregnant-control, non-pregnant-RFR exposed, pregnant-control and pregnant-RFR exposed. They were exposed to RFR (1800 MHz GSM; 14 V/m as reference level) for 15 min/day during 7 days. Malondialdehyde (MDA) and 8-hydroxy-2'-deoxyguanosine (8-OHdG) levels were analyzed. MDA and 8-OHdG levels of non-pregnant and pregnant-RFR exposed animals significantly increased with respect to controls ($p < 0.001$, Mann-Whitney test). No difference was found in the newborns ($p > 0.05$, Mann-Whitney). There exist very few experimental studies on the effects of RFR during pregnancy. It would be beneficial to increase the number of these studies in order to establish international standards for the protection of pregnant women from RFR.

Larsen AI, Olsen J, Svane O, Gender-specific reproductive outcome and exposure to high-frequency electromagnetic radiation among physiotherapists. *Scand J Work Environ Health* 17(5):324-329, 1991.

Studies that show **Cell Phone** Health Effects

The aim of this case-referent study was to investigate reproductive hazards other than congenital malformations after exposure to high-frequency electromagnetic radiation. Cases and referents were sampled from a cohort of pregnancies of members of the Union of Danish Physiotherapists through linkage of the union file with national medical registers. Case groups were spontaneous abortions and children with low birth-weight prematurity, and stillbirth/death within one year. Exposure to high-frequency electromagnetic radiation before and during pregnancy was assessed through telephone interviews. As referents to the 270 cases, 316 pregnancies were randomly sampled. A total of 8.4% did not participate. Only 23.5% of the children born by the highly exposed mothers were boys. This value is a statistically significantly altered gender ratio showing a dose-response pattern. High-frequency electromagnetic radiation was furthermore associated with low birthweight, but only for male newborns. The other outcomes were not statistically significantly associated with exposure to high-frequency electromagnetic radiation.

Kismali G, Ozgur E, Guler G, Akcay A, Sel T, Seyhan N. The influence of 1800 MHz GSM-like signals on blood chemistry and oxidative stress in non-pregnant and pregnant rabbits. *Int J Radiat Biol.* 88(5):414-419, 2012.

PURPOSE: Environmental electromagnetic fields originate from man-made sources, such as mobile phones and base stations, and have led to increasing public concern about their possible adverse health effects. We aimed to investigate the possible effects of radiofrequency radiation (RFR) generated from these devices on oversensitive animals, such as pregnant rabbits. MATERIALS AND METHODS: In the present study, the effects of whole body 1800 MHz Global System for Mobile Communications (GSM)-like RFR exposure for 15 min/day for seven days on blood chemistry and lipid peroxidation levels in both non-pregnant and pregnant New Zealand White rabbits were investigated. Thirteen-month-old rabbits were studied in the following four groups: Non-pregnant control, non-pregnant RFR-exposed, pregnant control and pregnant RFR-exposed. RESULTS: Lipid peroxidation, namely malondialdehyde (MDA) levels, did not change after RFR exposure. However, blood chemistry parameters, such as cholesterol (CHO), total protein (TP), albumin (ALB), uric acid, creatinin and creatine kinase (CK) and creatine kinase-myocardial band isoenzyme (CK-MB) changed due to both pregnancy and RFR exposure. CONCLUSION: Our investigations have been shown that no indication for oxidative stress was detected in the blood of pregnant rabbits upon RF exposure at specific conditions employed in the present study. Minor changes in some blood chemistry parameters were detected but CK-MB and CK increases were found remarkable. Studies on RFR exposure during pregnancy will help establish international standards for the protection of pregnant women from environmental RFR.

Gao X, Luo R, Ma B, Wang H, Liu T, Zhang J, Lian Z, Cui X. [Interference of vitamin E on the brain tissue damage by electromagnetic radiation of cell phone in pregnant and fetal rats]. *Wei Sheng Yan Jiu.* 2013 Jul;42(4):642-646.

[Article in Chinese]

OBJECTIVE: To investigate the interference of vitamin E on brain tissue damage by electromagnetic radiation of cell phone in pregnant and fetal rats. METHODS: 40

Studies that show **Cell Phone** Health Effects

pregnant rats were randomly divided into five groups (positive control, negative control, low, middle and high dosage of vitamin E groups). The low, middle and high dosage of vitamin E groups were supplemented with 5, 15 and 30 mg/ml vitamin E respectively since the first day of pregnancy. And the negative control group and the positive control group were given peanut oil without vitamin E. All groups except for the negative control group were exposed to 900MHz intensity of cell phone radiation for one hour each time, three times per day for 21 days. After accouchement, the right hippocampus tissue of fetal rats in each group was taken and observed under electron microscope. The vitality of superoxide dismutase (SOD) and glutathione peroxidase (GSH-Px), and the content of malondialdehyde (MDA) in pregnant and fetal rats' brain tissue were tested. RESULTS: Compared with the negative control group, the chondriosomes in neuron and neuroglia of brain tissues was swelling, mild edema was found around the capillary, chromatin was concentrated and collected, and bubbles were formed in vascular endothelial cells (VEC) in the positive fetal rat control group, whereas the above phenomenon was un-conspicuous in the middle and high dosage of vitamin E groups. We can see uniform chromatin, abundant mitochondrion, rough endoplasmic reticulum and free ribosomes in the high dosage group. The apoptosis has not found in all groups' sections. In the antioxidase activity analysis, compared with the negative control group, the vitality of SOD and GSH-Px significantly decreased and the content of MDA significantly increased both in the pregnant and fetal rats positive control group ($P < 0.05$). In fetal rats, the vitality of SOD and GSH-Px significantly increased in the brain tissues of all three different vitamin E dosages groups when compared with the positive control group, and the content of MDA was found significantly decreased in both middle and high dosage of vitamin E groups ($P < 0.05$). The same results have also been found in high dosage pregnant rat group, but in middle dosage group only SOD activity was found increased with significance ($P < 0.05$). With the dosage increase of vitamin E, the vitality of SOD and GSH-Px was increasing and the content of MDA was decreasing. CONCLUSION: Under the experimental dosage, vitamin E has certain interference on damage of antioxidant capacity and energy metabolism induced by electromagnetic radiation of cell phone in pregnant rats and fetal rats.

Batellier F, Couty I, Picard D, Brillard JP. Effects of exposing chicken eggs to a cell phone in "call" position over the entire incubation period. *Theriogenology*.69(6):737-745,2008

The aim of the present study was to assess the effects of exposing fertile chicken eggs to a cell phone repeatedly calling a ten-digit number at 3-min intervals over the entire period of incubation. A pre-experiment was performed first to adjust incubation conditions in an experimental chamber devoid of metallic content and without automatic turning until the overall performance of hatchability was reproducible in the absence of the cell phone. The experimental period consisted of a series of 4 incubations referred to as "replicates". For each replicate, one batch of 60 eggs was exposed to the immediate environment (≤ 25 cm) of a cell phone in the "call" position (exposed group), while another batch of 60 eggs, 1.5m away from the exposed group and also in the incubation chamber, was exposed to a similar cell phone in the "off"

Studies that show **Cell Phone** Health Effects

position (sham group). For each replicate, 2 other groups each of 60 eggs were also incubated, one in a standard mini-incubator ("Control I" group) and the second in a standard medium size incubator ("Control II" group). Temperature, relative humidity and electromagnetic fields in the experimental chamber were permanently monitored over the entire experiment. A significantly higher percentage of embryo mortality was observed in the "exposed" compared to the "sham" group in 2 of the 4 replicates ($p < .05$). In comparison with control groups, additional embryo mortality in the exposed group occurred mainly between Days 9 and 12 of incubation but a causal relationship between the intensity of the electric field and embryo mortality could not be established.

Sarapultseva EI, Igolkina JV, Tikhonov VN, Dubrova YE. THE IN VIVO EFFECTS OF LOW-INTENSITY RADIOFREQUENCY FIELDS ON THE MOTOR ACTIVITY OF PROTOZOA. Int J Radiat Biol. 2013 Nov 25. [Epub ahead of print]

Purpose: To analyze the direct and transgenerational effects of exposure to low-dose 1 GHz (mobile phone/wireless telecommunication range) and 10 GHz (radar/satellite communication range) radiofrequency electromagnetic fields (RF-EMF) on the motility of ciliates *Spirostomum ambiguum*. Materials and Methods: *S. ambiguum* were exposed to 1 GHz and 10 GHz RF-EMF with power flux densities (PD) ranging from 0.05 to 0.5 W/m² over a period of time from 0.05 to 10 h. The motility of directly exposed ciliates and their non-exposed progeny across 10-15 generations was measured. Results: Exposure to 0.1 W/m² of either 1 or 10 GHz RF-EMF resulted in a significant decrease in the motility. The dose of exposure capable of altering the mobility of ciliates was inversely correlated with the flux density of RF-EMF. The motility of the non-exposed progeny of ciliates irradiated with 0.1 W/m² of 10 GHz RF-EMF remained significantly compromised, at least, across 10-15 generations, thus indicating the presence of transgenerational effects. Conclusions: The results of our study show that low-dose exposure to RF-EMF can significantly affect the motility of irradiated ciliates and their non-exposed offspring, thus providing further insights into the unknown mechanisms underlying the in vivo effects of RF-EMF.

Effects on Fertility and Reproductive Organs

Wdowiak A, Wdowiak L, Wiktor H. Evaluation of the effect of using mobile phones on male fertility. Ann Agric Environ Med. 14(1):169-172, 2007.

The problem of the lack of offspring is a phenomenon concerning approximately 15% of married couples in Poland. Infertility is defined as inability to conceive after a year of sexual intercourses without the use of contraceptives. In half of the cases the causative factor is the male. Males are exposed to the effect of various environmental factors, which may decrease their reproductive capabilities. A decrease in male fertility is a phenomenon which occurs within years, which may suggest that one of the reasons for the decrease in semen parameters is the effect of the development of techniques in the

Studies that show **Cell Phone** Health Effects

surrounding environment. A hazardous effect on male fertility may be manifested by a decrease in the amount of sperm cells, disorders in their mobility, as well as structure. The causative agents may be chemical substances, ionizing radiation, stress, as well as electromagnetic waves. The objective of the study was the determination of the effect of the usage of cellular phones on the fertility of males subjected to marital infertility therapy. The following groups were selected from among 304 males covered by the study: Group A: 99 patients who did not use mobile phones, Group B: 157 males who have used GSM equipment sporadically for the period of 1-2 years, and Group C: 48 people who have been regularly using mobile phone for more than 2 years. In the analysis of the effect of GSM equipment on the semen it was noted that an increase in the percentage of sperm cells of abnormal morphology is associated with the duration of exposure to the waves emitted by the GSM phone. It was also confirmed that a decrease in the percentage of sperm cells in vital progressing motility in the semen is correlated with the frequency of using mobile phones.

Agarwal A, Deepinder F, Sharma RK, Ranga G, Li J. Effect of cell phone usage on semen analysis in men attending infertility clinic: an observational study. Fertil Steril.89(1):124-128, 2008.

OBJECTIVE: To investigate the effect of cell phone use on various markers of semen quality. DESIGN: Observational study. SETTING: Infertility clinic. PATIENT(S): Three hundred sixty-one men undergoing infertility evaluation were divided into four groups according to their active cell phone use: group A: no use; group B: <2 h/day; group C: 2-4 h/day; and group D: >4 h/day. INTERVENTION(S): None. MAIN OUTCOME MEASURE(S): Sperm parameters (volume, liquefaction time, pH, viscosity, sperm count, motility, viability, and morphology). RESULT(S): The comparisons of mean sperm count, motility, viability, and normal morphology among four different cell phone user groups were statistically significant. Mean sperm motility, viability, and normal morphology were significantly different in cell phone user groups within two sperm count groups. The laboratory values of the above four sperm parameters decreased in all four cell phone user groups as the duration of daily exposure to cell phones increased. CONCLUSION(S): Use of cell phones decrease the semen quality in men by decreasing the sperm count, motility, viability, and normal morphology. The decrease in sperm parameters was dependent on the duration of daily exposure to cell phones and independent of the initial semen quality.

Mailankot M, Kunnath AP, Jayalekshmi H, Koduru B, Valsalan R. Radio frequency electromagnetic radiation (RF-EMR) from GSM (0.9/1.8GHz) mobile phones induces oxidative stress and reduces sperm motility in rats. Clinics (Sao Paulo). 64(6):561-565, 2009.

INTRODUCTION: Mobile phones have become indispensable in the daily lives of men and women around the globe. As cell phone use has become more widespread, concerns have mounted regarding the potentially harmful effects of RF-EMR from these devices. OBJECTIVE: The present study was designed to evaluate the effects of RF-EMR from mobile phones on free radical metabolism and sperm quality. MATERIALS AND

Studies that show **Cell Phone** Health Effects

METHODS: Male albino Wistar rats (10-12 weeks old) were exposed to RF-EMR from an active GSM (0.9/1.8 GHz) mobile phone for 1 hour continuously per day for 28 days. Controls were exposed to a mobile phone without a battery for the same period. The phone was kept in a cage with a wooden bottom in order to address concerns that the effects of exposure to the phone could be due to heat emitted by the phone rather than to RF-EMR alone. Animals were sacrificed 24 hours after the last exposure and tissues of interest were harvested. **RESULTS:** One hour of exposure to the phone did not significantly change facial temperature in either group of rats. No significant difference was observed in total sperm count between controls and RF-EMR exposed groups. However, rats exposed to RF-EMR exhibited a significantly reduced percentage of motile sperm. Moreover, RF-EMR exposure resulted in a significant increase in lipid peroxidation and low GSH content in the testis and epididymis. **CONCLUSION:** Given the results of the present study, we speculate that RF-EMR from mobile phones negatively affects semen quality and may impair male fertility.

Rago R, Salacone P, Caponecchia L, Sebastianelli A, Marcucci I, Calogero AE, Condorelli R, Vicari E, Morgia G, Favilla V, Cimino S, Arcoria AF, La Vignera S. The semen quality of the mobile phone users. J Endocrinol Invest. 36(11):970-974, 2013.

BACKGROUND: The increased use of mobile phones, the media's attention for general health, and the increase of idiopathic male infertility suggest to investigate the possible consequences of an excessive use of mobile phones on semen quality. **AIM:** To evaluate the conventional and some of the main biofunctional sperm parameters in healthy men according to the different use of the mobile phone. **SUBJECTS AND METHODS:** All the enrolled subjects in this study were divided into four groups according to their active cell phone use: group A= no use (no.=10 subjects); group B= <2 h/day (no.=16); group C= 2-4 h/day (no.=17); and group D= >4 h/day (no.=20). Among the subjects of the group D (>4 h/day), a further evaluation was made between the "trousers users"(no.=12) and "shirt users"(no.=8), and they underwent semen collection to evaluate conventional and biofunctional sperm parameters (density, total count, morphology, progressive motility, apoptosis, mitochondrial membrane potential, chromatin compaction, DNA fragmentation). **RESULTS:** None of the conventional sperm parameters examined were significantly altered. However, the group D and the trousers users showed a higher percentage of sperm DNA fragmentation compared to other groups. **CONCLUSION:** These results suggest that the sperm DNA fragmentation could represent the only parameter significantly altered in the subjects who use the mobile phone for more than 4 h/day and in particular for those who use the device in the pocket of the trousers.

Qin F, Zhang J, Cao H, Guo W, Chen L, Shen O, Sun J, Yi C, Li J, Wang J, Tong J. Circadian alterations of reproductive functional markers in male rats exposed to 1800-MHz radiofrequency field. Chronobiol Int. 2013 Oct 11. [Epub ahead of print]

In this study, we explored the circadian effects of daily radiofrequency field (RF) exposure on reproductive functional markers in adult male Sprague-Dawley rats. Animals in circadian rhythm (as indicated by melatonin measurements), were

Studies that show **Cell Phone** Health Effects

divided into several groups and exposed to 1800 MHz RF at 205 $\mu\text{W}/\text{cm}^2$ power density (specific absorption rate 0.0405 W/kg) for 2 h/day for 32 days at different zeitgeber time (ZT) points, namely, ZT0, ZT4, ZT8, ZT12, ZT16 and ZT20. Sham-exposed animals were used as controls in the study. From each rat, testicular and epididymis tissues were collected and assessed for testosterone levels, daily sperm production and sperm motility, testis marker enzymes γ -GT and ACP, cytochrome P450 side-chain cleavage (p450cc) mRNA expression, and steroidogenic acute regulatory protein (StAR) mRNA expression. Via these measurements, we confirmed the existence of circadian rhythms in sham-exposed animals. However, rats exposed to RF exhibited a disruption of circadian rhythms, decreased testosterone levels, lower daily sperm production and sperm motility, down-regulated activity of γ -GT and ACP, as well as altered mRNA expression of cytochrome P450 and StAR. All of these observations were more pronounced when rats were exposed to RF at ZT0. Thus, our findings indicate potential adverse effects of RF exposure on male reproductive functional markers, in terms of both the daily overall levels as well as the circadian rhythmicity.

Sepehrimanesh M, Saeb M, Nazifi S, Kazemipour N, Jelodar G, Saeb S. Impact of 900 MHz electromagnetic field exposure on main male reproductive hormone levels: a *Rattus norvegicus* model. Int J Biometeorol. 2013 Dec 20. [Epub ahead of print]

This work analyzes the effects of radiofrequency-electromagnetic field (RF-EMF) exposure on the reproductive system of male rats, assessed by measuring circulating levels of FSH, LH, inhibin B, activin B, prolactin, and testosterone. Twenty adult male Sprague-Dawley rats (180 ± 10 g) were exposed to 900 MHz RF-EMF in four equal separated groups. The duration of exposure was 1, 2, and 4 h/day over a period of 30 days and sham-exposed animals were kept under the same environmental conditions as the exposed group except with no RF-EMF exposure. Before the exposure, at 15 and 30 days of exposure, determination of the abovementioned hormone levels was performed using ELISA. At the end of the experiment, FSH and LH values of the long time exposure (LTE) group were significantly higher than the sham-exposed group ($p < 0.05$). Serum activin B and prolactin in the LTE group showed significant increase and inhibin B showed significant decrease than sham and short time exposed (STE) groups after 30 days RF-EMF exposure ($p < 0.05$). Also, a significant decrease in serum testosterone levels in the LTE group was found compared to short and moderate time exposed (MTE) groups after 30 days RF-EMF exposure ($p < 0.05$). Results suggest that reproductive hormone levels are disturbed as a result of RF-EMF exposure and it may possibly affect reproductive functions. However, testosterone and inhibin B concentrations as a fertility marker and spermatogenesis were decreased significantly.

Zalata, A., A. Z. El-Samanoudy, D. Shaalan, Y. El-Baiomy, and T. Mostafa. In vitro effect of cell phone radiation on motility, DNA fragmentation and clusterin gene expression of sperm. Int J Fertil Steril, In Press. Published online ahead of print.

Background: Use of cellular phones that emits radiofrequency electromagnetic field (RF-EMF) has been increased exponentially and became a part of everyday life. This study

Studies that show **Cell Phone** Health Effects

aimed to investigate the effects of RF-EMF radiation emitted from cellular phones on sperm motility variables, sperm DNA fragmentation and clusterin (CLU) gene expression. **Materials and Methods:** 124 semen samples were grouped into; normozoospermia (N, n=26), asthenozoospermia (A, n=32), asthenoteratozoospermia (AT, n=31) and oligoasthenoteratozoospermia (OAT, n=35). Semen samples were divided into two aliquots; samples not exposed to cell phone and samples exposed to cell phone radiation (850 MHz, maximum power < 1 watt; SAR 1.46 W/kg at 10 cm distance) for 1 hr. Before and immediately after exposure both aliquots were subjected to assessment of sperm motility, acrosin activity, sperm DNA fragmentation and CLU gene expression. Statistical differences were analyzed using paired t-student test for comparisons where $P < 0.05$ was set as significant. **Results:** There was significant decrease in sperm motility, sperm linear velocity, sperm linearity index, sperm acrosin activity and significant increase in sperm DNA fragmentation percent, CLU gene expression and CLU protein levels in the exposed semen samples to RF-EMF compared with non- exposed samples in $OAT > AT > A > N$ groups ($P < 0.05$). **Conclusions:** Cell phone emissions have a negative impact on exposed sperm motility indices, sperm acrosin activity, sperm DNA fragmentation and CLU gene expression especially in OAT cases.

Guney M, Ozguner F, Oral B, Karahan N, Mungan T. 900 MHz radiofrequency-induced histopathologic changes and oxidative stress in rat endometrium: protection by vitamins E and C. *Toxicol Ind Health*. 23(7):411-420, 2007.

There are numerous reports on the effects of electromagnetic radiation (EMR) in various cellular systems. Mechanisms of adverse effects of EMR indicate that reactive oxygen species (ROS) may play a role in the biological effects of this radiation. The aims of this study were to examine 900 MHz mobile phone-induced oxidative stress that promotes production of ROS and to investigate the role of vitamins E and C, which have antioxidant properties, on endometrial tissue against possible 900 MHz mobile phone-induced endometrial impairment in rats. The animals were randomly grouped (eight each) as follows: 1) Control group (without stress and EMR, Group I), 2) sham-operated rats stayed without exposure to EMR (exposure device off, Group II), 3) rats exposed to 900 MHz EMR (EMR group, Group III) and 4) a 900 MHz EMR exposed + vitamin-treated group (EMR + Vit group, Group IV). A 900 MHz EMR was applied to EMR and EMR + Vit group 30 min/day, for 30 days using an experimental exposure device. Endometrial levels of nitric oxide (NO, an oxidant product) and malondialdehyde (MDA, an index of lipid peroxidation), increased in EMR exposed rats while the combined vitamins E and C caused a significant reduction in the levels of NO and MDA. Likewise, endometrial superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GSH-Px) activities decreased in EMR exposed animals while vitamins E and C caused a significant increase in the activities of these antioxidant enzymes. In the EMR group histopathologic changes in endometrium, diffuse and severe apoptosis was present in the endometrial surface epithelial and glandular cells and the stromal cells. Diffuse eosinophilic leucocyte and lymphocyte infiltration were observed in the endometrial stroma whereas the combination of vitamins E and C caused a significant decrease in these effects of EMR. It is concluded that oxidative endometrial damage plays an

Studies that show **Cell Phone** Health Effects

important role in the 900 MHz mobile phone-induced endometrial impairment and the modulation of oxidative stress with vitamins E and C reduces the 900 MHz mobile phone-induced endometrial damage both at biochemical and histological levels.

Agarwal A, Desai NR, Makker K, Varghese A, Mouradi R, Sabanegh E, Sharma R. Effects of radiofrequency electromagnetic waves (RF-EMW) from cellular phones on human ejaculated semen: an in vitro pilot study. Fertil Steril. 92(4) 1318-1325, 2009.

OBJECTIVE: To evaluate effects of cellular phone radiofrequency electromagnetic waves (RF-EMW) during talk mode on unprocessed (neat) ejaculated human semen. DESIGN: Prospective pilot study. SETTING: Center for reproductive medicine laboratory in tertiary hospital setting. SAMPLES: Neat semen samples from normal healthy donors (n = 23) and infertile patients (n = 9). INTERVENTION(S): After liquefaction, neat semen samples were divided into two aliquots. One aliquot (experimental) from each patient was exposed to cellular phone radiation (in talk mode) for 1 h, and the second aliquot (unexposed) served as the control sample under identical conditions. MAIN OUTCOME MEASURE(S): Evaluation of sperm parameters (motility, viability), reactive oxygen species (ROS), total antioxidant capacity (TAC) of semen, ROS-TAC score, and sperm DNA damage. RESULT(S): Samples exposed to RF-EMW showed a significant decrease in sperm motility and viability, increase in ROS level, and decrease in ROS-TAC score. Levels of TAC and DNA damage showed no significant differences from the unexposed group. CONCLUSION(S): Radiofrequency electromagnetic waves emitted from cell phones may lead to oxidative stress in human semen. We speculate that keeping the cell phone in a trouser pocket in talk mode may negatively affect spermatozoa and impair male fertility.

Susa M, Pavčić I. Djelovanje Radiofrekvencijskog Elektromagnetskog Zracenja na Spermatogenez u Sisavaca. Arh Hig Rada Toksikol. 58(4):449-459, 2007.

This article reviews studies about the effects of radiofrequency electromagnetic (RF EM) fields on male reproductive system and reproductive health in mammals. According to current data, there are almost 4 million active mobile phone lines in Croatia while this number has risen to 2 billion in the world. Increased use of mobile technology raises scientific and public concern about possible hazardous effects of RF fields on human health. The effects of radiofrequencies on reproductive health and consequences for the offspring are still mainly unknown. A number of in vivo and in vitro studies indicated that RF fields could interact with charged intracellular macromolecular structures. Results of several laboratory studies on animal models showed how the RF fields could affect the mammalian reproductive system and sperm cells. Inasmuch as, in normal physiological conditions spermatogenesis is a balanced process of division, maturation and storage of cells, it is particularly vulnerable to the chemical and physical environmental stimuli. Especially sensitive could be the cytoskeleton, composed of charged proteins; actin, intermedial filaments and microtubules. Cytoskeleton is a functional and structural part of the cell that has important role in the sperm motility, and is actively involved in the morphologic changes that occur during mammalian spermiogenesis.

Studies that show **Cell Phone** Health Effects

Yan JG, Agresti M, Bruce T, Yan YH, Granlund A, Matloub HS. Effects of cellular phone emissions on sperm motility in rats. *Fertil Steril*.88(4):957-64, 2007.

OBJECTIVE: To evaluate the effects of cellular phone emissions on rat sperm cells.
DESIGN: Classic experimental. SETTING: Animal research laboratory. SUBJECTS: Sixteen 3-month-old male Sprague-Dawley rats, weighing 250-300 g. INTERVENTION(S): Rats in the experimental group were exposed to two 3-hour periods of daily cellular phone emissions for 18 weeks; sperm samples were then collected for evaluation. MAIN OUTCOME MEASURE(S): Evaluation of sperm motility, sperm cell morphology, total sperm cell number, and mRNA levels for two cell surface adhesion proteins. RESULT(S): Rats exposed to 6 hours of daily cellular phone emissions for 18 weeks exhibited a significantly higher incidence of sperm cell death than control group rats through chi-squared analysis. In addition, abnormal clumping of sperm cells was present in rats exposed to cellular phone emissions and was not present in control group rats. CONCLUSION(S): These results suggest that carrying cell phones near reproductive organs could negatively affect male fertility.

Salama N, Kishimoto T, Kanayama HO, Kagawa S. Effects of exposure to a mobile phone on sexual behavior in adult male rabbit: an observational study. *Int J Impot Res*.22(2):127-33, 2010.

The accumulating effects of exposure to electromagnetic radiation emitted by a conventional mobile phone (MP) on male sexual behaviour have not yet been analyzed. Therefore, we studied these effects in 18 male rabbits that were randomly divided into phone and control groups. Six female teasers were taken successively to the male's cage and the copulatory behavior was recorded. Serum total testosterone, dopamine and cortisol were evaluated. The animals of the phone group were exposed to MPs (800 MHz) in a standby position for 8 h daily for 12 weeks. At the end of the study, the copulatory behavior and hormonal assays were re-evaluated. Mounts without ejaculation were the main mounts in the phone group and its duration and frequency increased significantly compared with the controls, whereas the reverse was observed in its mounts with ejaculation. Ejaculation frequency dropped significantly, biting/grasping against teasers increased notably and mounting latency in accumulated means from the first to the fourth teasers were noted in the phone group. The hormonal assays did not show any significant differences between the study groups. Therefore, the pulsed radiofrequency emitted by a conventional MP, which was kept on a standby position, could affect the sexual behavior in the rabbit.

Salama N, Kishimoto T, Kanayama HO. Effects of exposure to a mobile phone on testicular function and structure in adult rabbit. *Int J Androl*.33(1):88-94, 2010.

Summary The accumulating effects of exposure to electromagnetic radiation emitted by a conventional mobile phone (standby position) on the testicular function and structure are not yet fully investigated. To study these effects longitudinally, a total of 24 adult male rabbits were randomly and equally divided into three groups. Rabbits in the first (phone) group were exposed, in specially designed cages, to radio frequency emitted from the mobile phone (800 MHz) in a standby position opposite to that of testes for 8 h

Studies that show **Cell Phone** Health Effects

daily for 12 weeks. The second group consisted of the stress controls which were kept in the same kind of cages to appreciate any cage-induced anxiety. The third group included the ordinary controls which were kept in the conventional roomy cages. Semen analysis and sperm function tests (viability, hypo-osmotic swelling and acridine orange) were conducted weekly. Histological testicular sections and serum total testosterone were also evaluated. A drop in the sperm concentration appeared in the phone group at week 6. This became statistically significant at week 8, compared with the two control (stress and ordinary) groups (133, 339 and 356 x 10⁶/mL, respectively) and to the initial sperm count (341 x 10⁶/mL) of this group. Motile sperm population showed similarity amongst the three study groups until week 10 when it declined significantly, and thereafter in the phone and stress control groups, with more significant decline in the phone animals (50, 61 and 72.4%, respectively). Histological examination showed also a significant decrease in the diameter of seminiferous tubules in the phone group vs. the stress and ordinary controls (191 μ m vs. 206 and 226 μ m, respectively). The other study points did not show any difference. In conclusion, low intensity pulsed radio frequency emitted by a conventional mobile phone kept in the standby position could affect the testicular function and structure in the adult rabbit.

Sepehrimanesh M, Kazemipour N, Saeb M, Nazifi S. Analysis of rat testicular proteome following 30-days exposure to 900 MHz electromagnetic field radiation.

Electrophoresis. 2014 Aug 21. doi: 10.1002/elps.201400273. [Epub ahead of print]

The use of electromagnetic field (EMF) generating apparatuses such as cell phones is increasing, and has caused an interest in the investigations of its effects on human health. We analyzed proteome in preparations from the whole testis in adult male Sprague-Dawley rats exposed for 1, 2 or 4 h/d for 30 consecutive days to 900 MHz EMF radiation, simulating a range of possible human cell phone use. Subjects were sacrificed immediately after the end of the experiment and testes fractions were solubilized and separated via high resolution 2-dimensional electrophoresis, and gel patterns were scanned, digitized and processed. Thirteen of the proteins which found only in sham or in exposure groups were identified by MALDI-TOF/TOF-MS. Among them, heat shock proteins, superoxide dismutase, peroxiredoxin-1 and other proteins related to misfolding of proteins and/or stress were identified. These results demonstrate significant effects of radio-frequency modulated electromagnetic fields (RF-EMF) exposure on proteome, particularly in protein species in the rodent testis, and suggest that a 30 d exposure to EMF radiation induces non-thermal stress in testicular tissue. The functional implication of the identified proteins was discussed.

Aitken RJ, Bennetts LE, Sawyer D, Wiklendt AM, King BV. Impact of radio frequency electromagnetic radiation on DNA integrity in the male germline. Inter J Androl 28:171-179, 2005.

Concern has arisen over human exposures to radio frequency electromagnetic radiation (RFEMR), including a recent report indicating that regular mobile phone use can negatively impact upon human semen quality. These effects would be particularly serious if the biological effects of RFEMR included the induction of DNA damage in male

Studies that show **Cell Phone** Health Effects

germ cells. In this study, mice were exposed to 900 MHz RFEMR at a specific absorption rate of approximately 90 mW/kg inside a waveguide for 7 days at 12 h per day. Following exposure, DNA damage to caudal epididymal spermatozoa was assessed by quantitative PCR (QPCR) as well as alkaline and pulsed-field gel electrophoresis. The treated mice were overtly normal and all assessment criteria, including sperm number, morphology and vitality were not significantly affected. Gel electrophoresis revealed no gross evidence of increased single- or double-DNA strand breakage in spermatozoa taken from treated animals. However, a detailed analysis of DNA integrity using QPCR revealed statistically significant damage to both the mitochondrial genome ($p < 0.05$) and the nuclear β -globin locus ($p < 0.01$). This study suggests that while RFEMR does not have a dramatic impact on male germ cell development, a significant genotoxic effect on epididymal spermatozoa is evident and deserves further investigation.

Tas M, Dasdag S, Akdag MZ, Cirit U, Yegin K, Seker U, Ozmen MF, Eren LB. Long-term effects of 900 MHz radiofrequency radiation emitted from mobile phone on testicular tissue and epididymal semen quality. Electromagn Biol Med. 2013 Jun 19. [Epub ahead of print]

Abstract The purpose of this study is to bridge this gap by investigating effects of long term 900 MHz mobile phone exposure on reproductive organs of male rats. The study was carried out on 14 adult Wistar Albino rats by dividing them randomly into two groups (n: 7) as sham group and exposure group. Rats were exposed to 900 MHz radiofrequency (RF) radiation emitted from a GSM signal generator. Point, 1 g and 10 g specific absorption rate (SAR) levels of testis and prostate were found as 0.0623 W/kg, 0.0445 W/kg and 0.0373 W/kg, respectively. The rats in the exposure group were subject to RF radiation 3 h per day (7 d a week) for one year. For the sham group, the same procedure was applied, except the generator was turned off. At the end of the study, epididymal sperm concentration, progressive sperm motility, abnormal sperm rate, all-genital organs weights and testis histopathology were evaluated. Any differences were not observed in sperm motility and concentration ($p > 0.05$). However, the morphologically normal spermatozoa rates were found higher in the exposure group ($p < 0.05$). Although histological examination showed similarity in the seminiferous tubules diameters in both groups, tunica albuginea thickness and the Johnsen testicular biopsy score were found lower in the exposure group ($p < 0.05$, $p < 0.0001$). In conclusion, we claim that long-term exposure of 900 MHz RF radiation alter some reproductive parameters. However, more supporting evidence and research is definitely needed on this topic.

Al-Ali BM, Patzak J, Fischereder K, Pummer K, Shamloul R. Cell phone usage and erectile function. Cent European J Urol. 66(1):75-77, 2013. INTRODUCTION: The objective of this pilot study was to report our experience concerning the effects of cell phone usage on erectile function (EF) in men. MATERIAL AND METHODS: We recruited 20 consecutive men complaining of erectile dysfunction (ED) for at least six months (Group A), and another group of 10 healthy men with no complaints of ED (Group B). Anamnesis, basic laboratory investigations, and clinical examinations were performed.

Studies that show **Cell Phone** Health Effects

All men completed the German version of the Sexual Health Inventory for Men (SHIM) for evaluation of the International Index of Erectile Function (IIEF), as well as another questionnaire designed by our clinicians that assessed cell phone usage habits. RESULTS: There was no significant difference between both groups regarding age, weight, height, and total testosterone (Table 1). The SHIM scores of Group A were significantly lower than that of Group B, 11.2 ± 5 and 24.2 ± 2.3 , respectively. Total time spent talking on the cell phone per week was not significantly higher in Group A over B, 17.6 ± 11.1 vs. 12.5 ± 7 hours. Men with ED were found to carry their 'switched on' cell phones for a significantly longer time than those without ED, 4.4 ± 3.6 vs. 1.8 ± 1 hours per day. CONCLUSIONS: We found a potential correlation with cell phone usage and a negative impact on EF. Further large-scale studies confirming our initial data and exploring the mechanisms involved in this phenomenon are recommended.

Fejes I, Za Vaczki Z, Szollosi J, Kolosza R S, Daru J, Kova Cs L, Pa L **Als there a relationship between cell phone use and semen quality? Arch Androl. 51(5):385-393, 2005.**

This study was conducted to determine a possible relationship between regular cell phone use and different human semen attributes. The history-taking of men in our university clinic was supplemented with questions concerning cell phone use habits, including possession, daily standby position and daily transmission times. Semen analyses were performed by conventional methods. Statistics were calculated with SPSS statistical software. A total of 371 were included in the study. The duration of possession and the daily transmission time correlated negatively with the proportion of rapid progressive motile sperm ($r = -0.12$ and $r = -0.19$, respectively), and positively with the proportion of slow progressive motile sperm ($r = 0.12$ and $r = 0.28$, respectively). The low and high transmitter groups also differed in the proportion of rapid progressive motile sperm (48.7% vs. 40.6%). The prolonged use of cell phones may have negative effects on the sperm motility characteristics.

Falzone N, Huyser C, Becker P, Leszczynski D, Franken DR. **The effect of pulsed 900-MHz GSM mobile phone radiation on the acrosome reaction, head morphometry and zona binding of human spermatozoa. Int J Androl. 34(1):20-26, 2011.**

Summary Several recent studies have indicated that radiofrequency electromagnetic fields (RF-EMF) have an adverse effect on human sperm quality, which could translate into an effect on fertilization potential. This study evaluated the effect of RF-EMF on sperm-specific characteristics to assess the fertilizing competence of sperm. Highly motile human spermatozoa were exposed for 1 h to 900-MHz mobile phone radiation at a specific absorption rate of 2.0 W/kg and examined at various times after exposure. The acrosome reaction was evaluated using flow cytometry. The radiation did not affect sperm propensity for the acrosome reaction. Morphometric parameters were assessed using computer-assisted sperm analysis. Significant reduction in sperm head area ($9.2 \pm 0.7 \mu\text{m}^2$ vs. $18.8 \pm 1.4 \mu\text{m}^2$) and acrosome percentage of the head area ($21.5 \pm 4\%$ vs. $35.5 \pm 11.4\%$) was reported among exposed sperm compared with unexposed controls. Sperm-zona binding was assessed directly after exposure using the

Studies that show **Cell Phone** Health Effects

hemizona assay. The mean number of zona-bound sperm of the test hemizona and controls was 22.8 ± 12.4 and 31.8 ± 12.8 ($p < 0.05$), respectively. This study concludes that although RF-EMF exposure did not adversely affect the acrosome reaction, it had a significant effect on sperm morphometry. In addition, a significant decrease in sperm binding to the hemizona was observed. These results could indicate a significant effect of RF-EMF on sperm fertilization potential.

Fejes I, Za Vaczki Z, Szollosi J, Kolosza R S, Daru J, Kova Cs L, Pa L **Als there a relationship between cell phone use and semen quality? Arch Androl. 51(5):385-393, 2005.**

This study was conducted to determine a possible relationship between regular cell phone use and different human semen attributes. The history-taking of men in our university clinic was supplemented with questions concerning cell phone use habits, including possession, daily standby position and daily transmission times. Semen analyses were performed by conventional methods. Statistics were calculated with SPSS statistical software. A total of 371 were included in the study. The duration of possession and the daily transmission time correlated negatively with the proportion of rapid progressive motile sperm ($r = -0.12$ and $r = -0.19$, respectively), and positively with the proportion of slow progressive motile sperm ($r = 0.12$ and $r = 0.28$, respectively). The low and high transmitter groups also differed in the proportion of rapid progressive motile sperm (48.7% vs. 40.6%). The prolonged use of cell phones may have negative effects on the sperm motility characteristics.

Erogul O, Oztas E, Yildirim I, Kir T, Aydur E, Komesli G, Irkilata HC, Irmak MK, Peker AF. **Effects of electromagnetic radiation from a cellular phone on human sperm motility: an in vitro study. Arch Med Res. 37(7):840-843, 2006.**

BACKGROUND: There has been growing public concern on the effects of electromagnetic radiation (EMR) emitted by cellular phones on human health. Many studies have recently been published on this topic. However, possible consequences of the cellular phone usage on human sperm parameters have not been investigated adequately. METHODS: A total number of 27 males were enrolled in the study. The semen sample obtained from each participant was divided equally into two parts. One of the specimens was exposed to EMR emitted by an activated 900 MHz cellular phone, whereas the other was not. The concentration and motility of the specimens were compared to analyze the effects of EMR. Assessment of sperm movement in all specimens was performed using four criteria: (A) rapid progressive, (B) slow progressive, (C) nonprogressive, (D) no motility. RESULTS: Statistically significant changes were observed in the rapid progressive, slow progressive and no-motility categories of sperm movement. EMR exposure caused a subtle decrease in the rapid progressive and slow progressive sperm movement. It also caused an increase in the no-motility category of sperm movement. There was no statistically significant difference in the sperm concentration between two groups. CONCLUSIONS: These data suggest that EMR emitted by cellular phone influences human sperm motility. In addition to these acute adverse effects of EMR on sperm motility, long-term EMR exposure may lead to

Studies that show **Cell Phone** Health Effects

behavioral or structural changes of the male germ cell. These effects may be observed later in life, and they are to be investigated more seriously.

De Iuliis GN, Newey RJ, King BV, Aitken RJ. Mobile phone radiation induces reactive oxygen species production and DNA damage in human spermatozoa in vitro. *PLoS One*. 4(7):e6446, 2009.

BACKGROUND: In recent times there has been some controversy over the impact of electromagnetic radiation on human health. The significance of mobile phone radiation on male reproduction is a key element of this debate since several studies have suggested a relationship between mobile phone use and semen quality. The potential mechanisms involved have not been established, however, human spermatozoa are known to be particularly vulnerable to oxidative stress by virtue of the abundant availability of substrates for free radical attack and the lack of cytoplasmic space to accommodate antioxidant enzymes. Moreover, the induction of oxidative stress in these cells not only perturbs their capacity for fertilization but also contributes to sperm DNA damage. The latter has, in turn, been linked with poor fertility, an increased incidence of miscarriage and morbidity in the offspring, including childhood cancer. In light of these associations, we have analyzed the influence of RF-EMR on the cell biology of human spermatozoa in vitro. PRINCIPAL FINDINGS: Purified human spermatozoa were exposed to radio-frequency electromagnetic radiation (RF-EMR) tuned to 1.8 GHz and covering a range of specific absorption rates (SAR) from 0.4 W/kg to 27.5 W/kg. In step with increasing SAR, motility and vitality were significantly reduced after RF-EMR exposure, while the mitochondrial generation of reactive oxygen species and DNA fragmentation were significantly elevated ($P < 0.001$). Furthermore, we also observed highly significant relationships between SAR, the oxidative DNA damage bio-marker, 8-OH-dG, and DNA fragmentation after RF-EMR exposure. CONCLUSIONS: RF-EMR in both the power density and frequency range of mobile phones enhances mitochondrial reactive oxygen species generation by human spermatozoa, decreasing the motility and vitality of these cells while stimulating DNA base adduct formation and, ultimately DNA fragmentation. These findings have clear implications for the safety of extensive mobile phone use by males of reproductive age, potentially affecting both their fertility and the health and wellbeing of their offspring.

Falzone N, Huyser C, Becker P, Leszczynski D, Franken DR. The effect of pulsed 900-MHz GSM mobile phone radiation on the acrosome reaction, head morphometry and zona binding of human spermatozoa. *Int J Androl*. 34(1):20-26, 2011.

Summary Several recent studies have indicated that radiofrequency electromagnetic fields (RF-EMF) have an adverse effect on human sperm quality, which could translate into an effect on fertilization potential. This study evaluated the effect of RF-EMF on sperm-specific characteristics to assess the fertilizing competence of sperm. Highly motile human spermatozoa were exposed for 1 h to 900-MHz mobile phone radiation at a specific absorption rate of 2.0 W/kg and examined at various times after exposure. The acrosome reaction was evaluated using flow cytometry. The radiation did not affect sperm propensity for the acrosome reaction. Morphometric parameters were assessed

Studies that show **Cell Phone** Health Effects

using computer-assisted sperm analysis. Significant reduction in sperm head area ($9.2 \pm 0.7 \mu\text{m}^2$ vs. $18.8 \pm 1.4 \mu\text{m}^2$) and acrosome percentage of the head area ($21.5 \pm 4\%$ vs. $35.5 \pm 11.4\%$) was reported among exposed sperm compared with unexposed controls. Sperm-zona binding was assessed directly after exposure using the hemizona assay. The mean number of zona-bound sperm of the test hemizona and controls was 22.8 ± 12.4 and 31.8 ± 12.8 ($p < 0.05$), respectively. This study concludes that although RF-EMF exposure did not adversely affect the acrosome reaction, it had a significant effect on sperm morphometry. In addition, a significant decrease in sperm binding to the hemizona was observed. These results could indicate a significant effect of RF-EMF on sperm fertilization potential.

Kesari KK, Kumar S, Nirala J, Siddiqui MH, Behari J. Biophysical Evaluation of Radiofrequency Electromagnetic Field Effects on Male Reproductive Pattern. *Cell Biochem Biophys*. 65(2): 85-96, 2013 .

There are possible hazardous health effects of exposure to radiofrequency electromagnetic radiations emitted from mobile phone on the human reproductive pattern. It is more effective while keeping mobile phones in pocket or near testicular organs. Present review examines the possible concern on radio frequency radiation interaction and biological effects such as enzyme induction, and toxicological effects, including genotoxicity and carcinogenicity, testicular cancer, and reproductive outcomes. Testicular infertility or testicular cancer due to mobile phone or microwave radiations suggests an increased level of reactive oxygen species (ROS). Though generation of ROS in testis has been responsible for possible toxic effects on physiology of reproduction, the reviews of last few decades have well established that these radiations are very harmful and cause mutagenic changes in reproductive pattern and leads to infertility. The debate will be focused on bio-interaction mechanism between mobile phone and testicular cancer due to ROS formation. This causes the biological damage and leads to several changes like decreased sperm count, enzymatic and hormonal changes, DNA damage, and apoptosis formation. In the present review, physics of mobile phone including future research on various aspects has been discussed.

Salama N, Kishimoto T, Kanayama HO, Kagawa S. The mobile phone decreases fructose but not citrate in rabbit semen: a longitudinal study. *Syst Biol Reprod Med*. 55(5):181-187, 2009.

The negative impact of mobile phones on sperm motility has been previously described. Both fructose and citrate are important components in semen that facilitate sperm motility. To date, no studies have investigated the effect of exposure to electromagnetic radiation emitted from the mobile phone on their levels.. Therefore, a longitudinal study using the adult rabbit as a model was undertaken. A total of 30 adult male rabbits were randomly divided into three groups. The first (phone) group was placed in specially designed cages, and exposed to radio frequency emitted from a mobile phone (900 MHz) kept in standby mode and positioned adjacent to the genitalia for 8 h daily for 12 weeks. The other two groups served as controls; the stress group which was housed in

Studies that show **Cell Phone** Health Effects

the same kind of cages to evaluate any cage-induced anxiety, and the control group which was housed in the conventional roomy cages. Semen samples were retrieved weekly. Sperm motility and viability, semen fructose and citrate, and serum testosterone were measured. Histological sections from the prostatic complex, ampulla, and vesicular gland were evaluated. A significant drop in both fructose levels (257 +/- 11.6 vs. 489 +/- 8.4 mg %, the baseline level) and number of motile sperms (50 vs. 72%) was observed in the phone group at the 10th week. However, no correlation was found between the two values. The stress control animals showed a similar but significantly less decline in motility. No significant changes in citrate levels or other study parameters were seen in the three animal groups throughout the study. In conclusion, the pulsed radio frequency emitted by the mobile phone kept in the standby position longitudinally affected sperm motility and fructose but not citrate levels in rabbit semen.

Kesari KK, Behari J. Evidence for mobile phone radiation exposure effects on reproductive pattern of male rats: Role of ROS. *Electromagn Biol Med.* 31(3):213-22, 2012.

The relationship between radiofrequency electromagnetic fields emitted from mobile phone and infertility is a matter of continuing debate. It is postulated that these radiations may affect the reproduction pattern by targeting biochemistry of sperm. In an attempt to expedite the issue, 70 days old Wistar rats (n = 6) were exposed to mobile phone radiofrequency (RF) radiation for 2 h per day for 45 days and data compared with sham exposed (n = 6) group. A significant decrease (P < 0.05) in the level of testosterone and an increase in caspase-3 activity were found in the RF-exposed animals. Distortions in sperm head and mid piece of sperm mitochondrial sheath were also observed as captured by Transmission Electron Microscope (TEM). In addition, progeny from RF-exposed rats showed significant decreases in number and weight as compared with that of sham-exposed animals. A reduction in testosterone, an increase in caspase-3, and distortion in spermatozoa could be caused by overproduction of reactive oxygen species (ROS) in animals under mobile phone radiation exposure. Our findings on these biomarkers are clear indications of possible health implications of repeated exposure to mobile phone radiation.

Kesari KK, Kumar S, Behari J. Mobile phone usage and male infertility in Wistar rats. *Indian J Exp Biol.* 48(10):987-992, 2010.

A significant decrease in protein kinase C and total sperm count along with increased apoptosis were observed in male Wistar rats exposed to mobile phone frequencies (2 h/day x 35 days at 0.9 W/kg specific absorption rate). The results suggest that a reduction in protein kinase activity may be related to overproduction of reactive oxygen species (ROS) under microwave field exposure. Decrease in sperm count and an increase in apoptosis may be causative factor due to mobile radiation exposure leading to infertility.

Studies that show **Cell Phone** Health Effects

Kesari KK, Kumar S, Behari J. Effects of Radiofrequency Electromagnetic Wave Exposure from Cellular Phones on the Reproductive Pattern in Male Wistar Rats. *Appl Biochem Biotechnol*. 164(4):546-559, 2011.

The present study investigates the effect of free radical formation due to mobile phone exposure and effect on fertility pattern in 70-day-old male Wistar rats (sham exposed and exposed). Exposure took place in Plexiglas cages for 2 h a day for 35 days to mobile phone frequency. The specific absorption rate was estimated to be 0.9 W/kg. An analysis of antioxidant enzymes glutathione peroxidase ($P < 0.001$) and superoxide dismutase ($P < 0.007$) showed a decrease, while an increase in catalase ($P < 0.005$) was observed. Malondialdehyde ($P < 0.003$) showed an increase and histone kinase ($P = 0.006$) showed a significant decrease in the exposed group. Micronuclei also show a significant decrease ($P < 0.002$) in the exposed group. A significant change in sperm cell cycle of G(0)-G(1) ($P = 0.042$) and G(2)/M ($P = 0.022$) were recorded. Generation of free radicals was recorded to be significantly increased ($P = 0.035$). Our findings on antioxidant, malondialdehyde, histone kinase, micronuclei, and sperm cell cycle are clear indications of an infertility pattern, initiated due to an overproduction of reactive oxygen species. It is concluded that radiofrequency electromagnetic wave from commercially available cell phones might affect the fertilizing potential of spermatozoa.

Nazıroğlu M, Yüksel M, Köse SA, Özkaya MO. Recent reports of Wi-Fi and mobile phone-induced radiation on oxidative stress and reproductive signaling pathways in females and males. *J Membr Biol*. 246(12):869-875, 2013.

Environmental exposure to electromagnetic radiation (EMR) has been increasing with the increasing demand for communication devices. The aim of the study was to discuss the mechanisms and risk factors of EMR changes on reproductive functions and membrane oxidative biology in females and males. It was reported that even chronic exposure to EMR did not increase the risk of reproductive functions such as increased levels of neoantigens abort. However, the results of some studies indicate that EMR induced endometriosis and inflammation and decreased the number of follicles in the ovarium or uterus of rats. In studies with male rats, exposure caused degeneration in the seminiferous tubules, reduction in the number of Leydig cells and testosterone production as well as increases in luteinizing hormone levels and apoptotic cells. In some cases of male and female infertility, increased levels of oxidative stress and lipid peroxidation and decreased values of antioxidants such as melatonin, vitamin E and glutathione peroxidase were reported in animals exposed to EMR. In conclusion, the results of current studies indicate that oxidative stress from exposure to Wi-Fi and mobile phone-induced EMR is a significant mechanism affecting female and male reproductive systems. However, there is no evidence to this date to support an increased risk of female and male infertility related to EMR exposure.

Ozlem Nisbet H, Nisbet C, Akar A, Cevik M, Karayigit MO. Effects of exposure to electromagnetic field (1.8/0.9 GHz) on testicular function and structure in growing rats. *Res Vet Sci*. 93(2):1001-1005, 2012.

Studies that show **Cell Phone** Health Effects

The aim of our study was to evaluate the possible effects of whole-body electromagnetic field (EMF) exposure on reproduction in growing male rats. Male albino Wistar rats (2 days old) were exposed to EMF 1800 and 900 MHz for 2 h continuously per day for 90 days. Sham control was kept under similar conditions except that the field was not applied for the same period. After blood samples were collected, the animals were sacrificed 24 h after the last exposure and the tissues of interest were harvested. The mean plasma total testosterone showed similarity among the two study groups and was significantly higher than the sham control rats. The percentage of epididymal sperm motility was significantly higher in the 1800 MHz group ($P < 0.05$). The morphologically normal spermatozoa rates were higher and the tail abnormality and total percentage abnormalities were lower in the 900 MHz group ($P < 0.05$). Histopathologic parameters in the 1800 MHz group were significantly higher ($P < 0.05$). In conclusion, the present study indicated that exposure to electromagnetic wave caused an increase in testosterone level, epididymal sperm motility (forward), and normal sperm morphology of rats. As a consequences, 1800 and 900 MHz EMF could be considered to be a cause of precocious puberty in growing rats.

Meo SA, Al-Drees AM, Husain S, Khan MM, Imran MB. Effects of mobile phone radiation on serum testosterone in Wistar albino rats. Saudi Med J. 31(8):869-873, 2010.

OBJECTIVE: To investigate the effects of electromagnetic field radiation generated by mobile phones on serum testosterone levels in Wistar albino rats. **METHODS:** This experimental interventional control study was conducted in the Department of Physiology, College of Medicine, King Saud University, Riyadh, Saudi Arabia during the period December 2006 to April 2008. A total of 34 male Albino rats [Wistar strain], 2 months of age, weighing 150-160 gm were used for the experiment. These animals were divided into 3 groups. The first group containing 6 rats was assigned as a control group. The second group containing 14 rats was exposed to mobile phone radiation for 30 minutes daily and the third group containing 14 rats was exposed to mobile phone radiation for 60 minutes daily for the total period of 3 months. At the end of experimental period, blood was collected into the container, and serum testosterone was analyzed using double-antibody radioimmunoassay method by Coat-A-Count. **RESULTS:** Exposure to mobile phone radiation for 60 minutes/day for the total period of 3 months significantly decrease the serum testosterone level [$p = 0.028$] in Wistar Albino rats compared to their matched control. **CONCLUSION:** Long-term exposure to mobile phone radiation leads to reduction in serum testosterone levels. Testosterone is a primary male gender hormone and any change in the normal levels may be devastating for reproductive and general health.

Gorpinchenko I, Nikitin O, Banyra O, Shulyak A. The influence of direct mobile phone radiation on sperm quality. Cent European J Urol. 67(1):65-71, 2014.

INTRODUCTION: It is impossible to imagine a modern socially-active man who does not use mobile devices and/or computers with Wi-Fi function. The effect of mobile phone radiation on male fertility is the subject of recent interest and investigations. The aim of

Studies that show **Cell Phone** Health Effects

this study was to investigate the direct in vitro influence of mobile phone radiation on sperm DNA fragmentation and motility parameters in healthy subjects with normozoospermia. **MATERIAL AND METHODS:** 32 healthy men with normal semen parameters were selected for the study. Each sperm sample was divided into two equal portions (A and B). Portions A of all involved men were placed for 5 hours in a thermostat, and portions B were placed into a second thermostat for the same period of time, where a mobile phone in standby/talk mode was placed. After 5 hours of incubation the sperm samples from both thermostats were re-evaluated regarding basic motility parameters. The presence of DNA fragmentation in both A and B portions of each sample was determined each hour using a standard sperm chromatin dispersion test. **RESULTS:** The number of spermatozoa with progressive movement in the group, influenced by electromagnetic radiation, is statistically lower than the number of spermatozoa with progressive movement in the group under no effect of the mobile phone. The number of non-progressive movement spermatozoa was significantly higher in the group, which was influenced by cell phone radiation. The DNA fragmentation was also significantly higher in this group. **CONCLUSIONS:** A correlation exists between mobile phone radiation exposure, DNA-fragmentation level and decreased sperm motility.

Ghanbari M, Mortazavi SB, Khavanin A, Khazaei M. The Effects of Cell Phone Waves (900 MHz-GSM Band) on Sperm Parameters and Total Antioxidant Capacity in Rats. Int J Fertil Steril. 7(1):21-28, 2013.

BACKGROUND: There is tremendous concern regarding the possible adverse effects of cell phone microwaves. Contradictory results, however, have been reported for the effects of these waves on the body. In the present study, the effect of cell phone microwaves on sperm parameters and total antioxidant capacity was investigated with regard to the duration of exposure and the frequency of these waves. **MATERIALS AND METHODS:** This experimental study was performed on 28 adult male Wistar rats (200-250 g). The animals were randomly assigned to four groups (n=7): i. control; ii. two-week exposure to cell phone-simulated waves; iii. three-week exposure to cell phonesimulated waves; and iv. two-week exposure to cell phone antenna waves. In all groups, sperm analysis was performed based on standard methods and we determined the mean sperm total antioxidant capacity according to the ferric reducing ability of plasma (FRAP) method. Data were analyzed by one-way ANOVA followed by Tukey's test using SPSS version 16 software. **RESULTS:** The results indicated that sperm viability, motility, and total antioxidant capacity in all exposure groups decreased significantly compared to the control group ($p<0.05$). Increasing the duration of exposure from 2 to 3 weeks caused a statistically significant decrease in sperm viability and motility ($p<0.05$). **CONCLUSION:** Exposure to cell phone waves can decrease sperm viability and motility in rats. These waves can also decrease sperm total antioxidant capacity in rats and result in oxidative stress.

Ozguner M, Koyu A, Cesur G, Ural M, Ozguner F, Gokcimen A, Delibas N. Biological and morphological effects on the reproductive organ of rats after exposure to

Studies that show **Cell Phone** Health Effects

electromagnetic field. Saudi Med J. 26(3):405-410, 2005.

OBJECTIVE: The biological effect of electromagnetic field (EMF) emitted from mobile phones is a current debate and still a controversial issue. Therefore, little is known on the possible adverse effects on reproduction as mobile phone bio-effects are only a very recent concern. The aim of this experimental study was to determine the biological and morphological effects of 900 MHz radiofrequency (RF) EMF on rat testes. **METHODS:** The study was performed in the Physiology and Histology Research Laboratories of Suleyman Demirel University, Faculty of Medicine, Isparta, Turkey in May 2004. Twenty adult male Sprague-Dawley rats weighing 270 - 320 gm were randomized into 2 groups of 10 animals: Group I (control group) was not exposed to EMF and Group II (EMF group) was exposed to 30 minutes per day, 5 days a week for 4 weeks to 900 MHz EMF. Testes tissues were submitted for histologic and morphologic examination. Testicular biopsy score count and the percentage of interstitial tissue to the entire testicular tissue were registered. Serum testosterone, plasma luteinizing hormone (LH) and follicle stimulating hormone (FSH) levels were assayed biochemically. **RESULTS:** The weight of testes, testicular biopsy score count and the percentage of interstitial tissue to the entire testicular tissue were not significantly different in EMF group compared to the control group. However, the diameter of the seminiferous tubules and the mean height of the germinal epithelium were significantly decreased in EMF group ($p<0.05$). There was a significant decrease in serum total testosterone level in EMF group ($p<0.05$). Therefore, there was an insignificant decrease in plasma LH and FSH levels in EMF group compared to the control group ($p>0.05$). **CONCLUSION:** The biological and morphological effects resulting from 900 MHz RF EMF exposure lends no support to suggestions of adverse effect on spermatogenesis, and on germinal epithelium. Therefore, testicular morphologic alterations may possibly be due to hormonal changes.

Meo SA, Arif M, Rashied S, Khan MM, Vohra MS, Usmani AM, Imran MB, Al-Drees AM. Hypospermatogenesis and spermatozoa maturation arrest in rats induced by mobile phone radiation. J Coll Physicians Surg Pak. 21(5):262-265, 2011.

Abstract. Objective: To determine the morphological changes induced by mobile phone radiation in the testis of Wistar albino rats. Study Design: Cohort study. Place and Duration of Study: Department of Physiology, College of Medicine, King Saud University, Riyadh, Saudi Arabia, from April 2007 to June 2008. Methodology: Forty male Wistar albino rats were divided in three groups. First group of eight served as the control. The second group [group B, $n=16$] was exposed to mobile phone radiation for 30 minutes/day and the third group [group C, $n=16$] was exposed to mobile phone radiation for 60 minutes/day for a total period of 3 months. Morphological changes in the testes induced by mobile phone radiations were observed under a light microscope. Results: Exposure to mobile phone radiation for 60 minutes/day caused 18.75% hypospermatogenesis and 18.75% maturation arrest in the testis of albino rats compared to matched controls. However, no abnormal findings were observed in albino rats that were exposed to mobile phone radiation for 30 minutes/day for a total period of 3 months. Conclusion: Long-term exposure to mobile phone radiation can cause hypospermatogenesis and maturation arrest in the spermatozoa in the testis of Wistar

Studies that show **Cell Phone** Health Effects

albino rats.

Meo SA, Arif M, Rashied S, Khan MM, Vohra MS, Usmani AM, Imran MB, Al-Drees AM. Hypospermatogenesis and spermatozoa maturation arrest in rats induced by mobile phone radiation. J Coll Physicians Surg Pak. 21(5):262-265, 2011.

OBJECTIVE: To determine the morphological changes induced by mobile phone radiation in the testis of Wistar albino rats. STUDY DESIGN: Cohort study. Place and Duration of Study: Department of Physiology, College of Medicine, King Saud University, Riyadh, Saudi Arabia, from April 2007 to June 2008. METHODOLOGY: Forty male Wistar albino rats were divided in three groups. First group of eight served as the control. The second group [group B, n=16] was exposed to mobile phone radiation for 30 minutes/day and the third group [group C, n=16] was exposed to mobile phone radiation for 60 minutes/day for a total period of 3 months. Morphological changes in the testes induced by mobile phone radiations were observed under a light microscope. RESULTS: Exposure to mobile phone radiation for 60 minutes/day caused 18.75% hypospermatogenesis and 18.75% maturation arrest in the testis of albino rats compared to matched controls. However, no abnormal findings were observed in albino rats that were exposed to mobile phone radiation for 30 minutes/day for a total period of 3 months. CONCLUSION: Long-term exposure to mobile phone radiation can cause hypospermatogenesis and maturation arrest in the spermatozoa in the testis of Wistar albino rats.

Gutschi T, Mohamad Al-Ali B, Shamloul R, Pummer K, Trummer H. Impact of cell phone use on men's semen parameters. Andrologia. 43(5):312-316, 2011.

The objective of the present retrospective study was to report our experience concerning the effects of cell phone usage on semen parameters. We examined 2110 men attending our infertility clinic from 1993 to October 2007. Semen analysis was performed in all patients. Serum free testosterone (T), follicle stimulating hormone (FSH), luteinising hormone (LH) and prolactin (PRL) were collected from all patients. The information on cell phone use of the patients was recorded and the subjects were divided into two groups according to their cell phone use: group A: cell phone use (n = 991); group B: no use (n = 1119). Significant difference was observed in sperm morphology between the two groups. In the patients of group A, 68.0% of the spermatozoa featured a pathological morphology compared to only 58.1% in the subjects of group B. Patients with cell phone usage showed significantly higher T and lower LH levels than those who did not use cell phone. No significant difference between the two groups was observed regarding FSH and PRL values. Our results showed that cell phone use negatively affects sperm quality in men. Further studies with a careful design are needed to determine the effect of cell phone use on male fertility.

Gul A, Celebi H, Uğraş S. The effects of microwave emitted by cellular phones on ovarian follicles in rats. Arch Gynecol Obstet. 280(5):729-733, 2009.

OBJECTIVE: The aim of this study was to investigate whether there were any toxic effects of microwaves of cellular phones on ovaries in rats. METHODS: In this study, 82

Studies that show **Cell Phone** Health Effects

female pups of rats, aged 21 days (43 in the study group and 39 in the control group) were used. Pregnant rats in the study group were exposed to mobile phones that were placed beneath the polypropylene cages during the whole period of pregnancy. The cage was free from all kinds of materials, which could affect electromagnetic fields. A mobile phone in a standby position for 11 h and 45 min was turned on to speech position for 15 min every 12 h and the battery was charged continuously. On the 21st day after the delivery, the female rat pups were killed and the right ovaries were removed. The volumes of the ovaries were measured and the number of follicles in every tenth section was counted. RESULTS: The analysis revealed that in the study group, the number of follicles was lower than that in the control group. The decreased number of follicles in pups exposed to mobile phone microwaves suggest that intrauterine exposure has toxic effects on ovaries. CONCLUSION: We suggest that the microwaves of mobile phones might decrease the number of follicles in rats by several known and, no doubt, countless unknown mechanisms.

Lukac N, Massanyi P, Roychoudhury S, Capcarova M, Tvrda E, Knazicka Z, Kolesarova A, Danko J. In vitro effects of radiofrequency electromagnetic waves on bovine spermatozoa motility. J Environ Sci Health A Tox Hazard Subst Environ Eng. 46(12):1417-1423, 2011.

In this study the effects of 1800 MHz GSM-like radiofrequency electromagnetic waves (RF-EMW) exposure on bovine semen was monitored. The experimental samples were analyzed in vitro in four time periods (0, 30, 120 and 420 min) and compared with unexposed samples (control). Spermatozoa motility was determined by computer assisted semen analyzer (CASA). Evaluation of the percentage of motile spermatozoa showed significant ($P < 0.001$) decrease in experimental groups after 120 and 420 min of culture when exposed to microwaves, in comparison to control. Similar spermatozoa motility inhibition was detected for the percentage of progressively motile spermatozoa, too. Average path distance decreased significantly ($p < 0.001$) in experimental groups after 30 and 420 min of culture. Path velocity increased in the experimental groups exposed to RF-EMW after 30 minutes of culture, but subsequently decreased after 420 min of culture, in comparison to control. This indicates a possible initial stimulation and subsequent velocity inhibition of bovine spermatozoa under RF-EMW exposure. Changes in spermatozoa motility were also detected for some fine parameters, too. A significant decrease ($P < 0.001$) was noted for amplitude of lateral head displacement in the experimental group after 420 minutes of culture. Detailed in vitro motility analysis of bovine spermatozoa exposed to microwave radiation suggested that the parameters of path and velocity at the beginning of the culture significantly increase, but after longer culture (420 minutes) a significant decrease occur in the experimental group as compared to control. In general, results of this experiment indicate a negative time-dependent effect of 1800 MHz RF-EMW radiation on bovine spermatozoa motility.

Kilgallon SJ, Simmons LW. Image content influences men's semen quality. Biol Lett 1:252-255, 2005.

Studies that show **Cell Phone** Health Effects

There is increasing evidence from non-human animals that males adjust their ejaculation expenditure according to the risk of sperm competition. In this study we show that, after controlling for lifestyle factors known to influence semen quality, human males viewing images depicting sperm competition had a higher percentage of motile sperm in their ejaculates. Many lifestyle variables were confirmed to influence semen quality, including the recent suggestion that storage of mobile phones close to the testes can decrease semen quality.

Liu C, Gao P, Xu SC, Wang Y, Chen CH, He MD, Yu ZP, Zhang L, Zhou Z. Mobile phone radiation induces mode-dependent DNA damage in a mouse spermatocyte-derived cell line: a protective role of melatonin. Int J Radiat Biol. 2013 Aug 19. [Epub ahead of print]

Purpose: To evaluate whether exposure to mobile phone radiation (MPR) can induce DNA damage in male germ cells. Materials and methods: A mouse spermatocyte-derived GC-2 cell line was exposed to a commercial mobile phone handset once every 20 minutes in standby, listen, dialed or dialing modes for 24 h. DNA damage was determined using an alkaline comet assay. Results: The levels of DNA damage were significantly increased following exposure to MPR in the listen, dialed and dialing modes. Moreover, there were significantly higher increases in the dialed and dialing modes than in the listen mode. Interestingly, these results were consistent with the radiation intensities of these modes. However, the DNA damage effects of MPR in the dialing mode were efficiently attenuated by melatonin pretreatment. Conclusions: These results regarding mode-dependent DNA damage have important implications for the safety of inappropriate mobile phone use by males of reproductive age and also suggest a simple preventive measure, keeping our body from mobile phones as far away as possible, not only during conversations but during "dialed" and "dialing" operation modes as well. Since the "dialed" mode is actually part of the standby mode, mobile phones should be kept at a safe distance from our body even during standby operation. Furthermore, the protective role of melatonin suggests that it may be a promising pharmacological candidate for preventing mobile phone use-related reproductive impairments.

Liu K, Zhang G, Wang Z, Liu Y, Dong J, Dong X, Liu J, Cao J, Ao L, Zhang S. The protective effect of autophagy on mouse spermatocyte derived cells exposure to 1800MHz radiofrequency electromagnetic radiation. Toxicol Lett. 2014 May 8. pii: S0378-4274(14)00195-7. doi: 10.1016/j.toxlet.2014.05.004. [Epub ahead of print]

The increasing exposure to radiofrequency (RF) radiation emitted from mobile phone use has raised public concern regarding the biological effects of RF exposure on the male reproductive system. Autophagy contributes to maintaining intracellular homeostasis under environmental stress. To clarify whether RF exposure could induce autophagy in the spermatocyte, mouse spermatocyte-derived cells (GC-2) were exposed to 1800MHz Global System for Mobile Communication (GSM) signals in GSM-Talk mode at specific absorption rate (SAR) values of 1 w/kg, 2w/kg or 4w/kg for 24h, respectively. The results indicated that the expression of LC3-II increased in a dose- and time-

Studies that show **Cell Phone** Health Effects

dependent manner with RF exposure, and showed a significant change at the SAR value of 4w/kg. The autophagosome formation and the occurrence of autophagy were further confirmed by GFP-LC3 transient transfection assay and transmission electron microscopy (TEM) analysis. Furthermore, the conversion of LC3-I to LC3-II was enhanced by co-treatment with Chloroquine (CQ), indicating autophagic flux could be enhanced by RF exposure. Intracellular ROS levels significantly increased in a dose- and time-dependent manner after cells were exposed to RF. Pretreatment with anti-oxidative NAC obviously decreased the conversion of LC3-I to LC3-II and attenuated the degradation of p62 induced by RF exposure. Meanwhile, phosphorylated extracellular-signal-regulated kinase (ERK) significantly increased after RF exposure at the SAR value of 2w/kg and 4w/kg. Moreover, we observed that RF exposure did not increase the percentage of apoptotic cells, but inhibition of autophagy could increase the percentage of apoptotic cells. These findings suggested that autophagy flux could be enhanced by 1800MHz GSM exposure (4w/kg), which is mediated by ROS generation. Autophagy may play an important role in preventing cells from apoptotic cell death under RF exposure stress.

Liu C, Duan W, Xu S, Chen C, He M, Zhang L, Yu Z, Zhou Z. Exposure to 1800 MHz radiofrequency electromagnetic radiation induces oxidative DNA base damage in a mouse spermatocyte-derived cell line. Toxicol Lett 218(1): 2-9, 2013.

Whether exposure to radiofrequency electromagnetic radiation (RF-EMR) emitted from mobile phones can induce DNA damage in male germ cells remains unclear. In this study, we conducted a 24 h intermittent exposure (5 min on and 10 min off) of a mouse spermatocyte-derived GC-2 cell line to 1800 MHz Global System for Mobile Communication (GSM) signals in GSM-Talk mode at specific absorption rates (SAR) of 1 W/kg, 2 W/kg or 4 W/kg. Subsequently, through the use of formamidopyrimidine DNA glycosylase (FPG) in a modified comet assay, we determined that the extent of DNA migration was significantly increased at a SAR of 4 W/kg. Flow cytometry analysis demonstrated that levels of the DNA adduct 8-oxoguanine (8-oxoG) were also increased at a SAR of 4 W/kg. These increases were concomitant with similar increases in the generation of reactive oxygen species (ROS); these phenomena were mitigated by co-treatment with the antioxidant α -tocopherol. However, no detectable DNA strand breakage was observed by the alkaline comet assay. Taking together, these findings may imply the novel possibility that RF-EMR with insufficient energy for the direct induction of DNA strand breaks may produce genotoxicity through oxidative DNA base damage in male germ cells.

Hancı H, Odacı E, Kaya H, Aliyazıcıoğlu Y, Turan I, Demir S, Colakoğlu S. The effect of prenatal exposure to 900-MHz electromagnetic field on the 21-old-day rat testicle. Reprod Toxicol. 42:203-209, 2013.

The aim of this study was to investigate the effect of exposure to a 900-MHz electromagnetic field (EMF) in the prenatal term on the 21-old-day rat testicle. Pregnant rats were divided into control (CG) and EMF (EMFG) groups. EMFG was exposed to 900-MHz EMF during days 13-21 of pregnancy. Newborn CG rats were obtained from the CG

Studies that show **Cell Phone** Health Effects

and newborn EMFG (NEMFG) rats from the EMFG. Testicles were extracted at postnatal day 21. Lipid peroxidation and DNA oxidation levels, apoptotic index and histopathological damage scores were compared. NEMFG rats exhibited irregularities in seminiferous tubule basal membrane and epithelium, immature germ cells in the lumen, and a decreased diameter in seminiferous tubules and thickness of epithelium. Apoptotic index, lipid peroxidation and DNA oxidation were higher in NEMFG rats than in NCG. 21-day-old rat testicles exposed to 900-MHz EMF in the prenatal term may be adversely affected, and this effect persists after birth.

Falzone N, Huyser C, Fourie F, Toivo T, Leszczynski D, Franken D. In vitro effect of pulsed 900 MHz GSM radiation on mitochondrial membrane potential and motility of human spermatozoa. *Bioelectromagnetics*.29(4):268-276, 2008.

Ejaculated, density purified, human spermatozoa were exposed to pulsed 900 MHz GSM mobile phone radiation at two specific absorption rate levels (SAR 2.0 and 5.7 W/kg) and compared with controls over time. Change in sperm mitochondrial membrane potential was analysed using flow cytometry. Sperm motility was determined by computer assisted sperm analysis (CASA). There was no effect of pulsed 900 MHz GSM radiation on mitochondrial membrane potential. This was also the case for all kinematic parameters assessed at a SAR of 2.0 W/kg. However, over time, the two kinematic parameters straight line velocity (VSL) and beat-cross frequency (BCF) were significantly impaired (P < 0.05) after the exposure at SAR 5.7 W/kg and no exposure by time interaction was present. This result should not be ascribed to thermal effects, due to the cooling methods employed in the RF chamber and temperature control within the incubator.

Dasdag, S, Ketani, MA, Akdag, Z, Ersay, AR, Sar,i I, Demirtas ,OC, Celik, MS, Whole-body microwave exposure emitted by cellular phones and testicular function of rats. *Urol Res* 27(3):219-223, 1999.

This study investigated whether there are adverse effects due to microwave exposure emitted by cellular phones in male rats. Eighteen Wistar Albino rats were separated into three groups, a sham group and two experimental groups. The rats were confined in Plexiglas cages and cellular phones were placed 0.5 cm under the cages. In the first experimental group, cellular phones were in standby position for 2 h. In the second experimental group, phones were turned to the speech position three times each for 1 min duration over 2 h. Rats in the first and second experimental groups were exposed to microwaves emitted by phones for 2 h/day for a duration of 1 month. After the last exposure the rats were killed. Brain, eyes, ears, liver, heart, lungs, stomach, kidneys, testes, small and large intestines and skin of the rats were observed histologically. The decrease of epididymal sperm counts in the speech groups were not found to be significant (P > 0.05). Differences in terms of normal and abnormal sperm forms were not observed (P > 0.05). Histological changes were especially observed in the testes of rats of the speech groups. Seminiferous tubular diameter of rat testes in the standby and speech groups was found to be lower than the sham group (P < 0.05). Rectal temperatures of rats in the speech group were found to be higher than the sham and standby groups (P < 0.05). The rectal temperatures of rats before

Studies that show **Cell Phone** Health Effects

and after exposure were also found to be significantly higher in the speech group ($P < 0.05$). Specific absorption rate (SAR) was determined as 0.141 W/kg.

Chen L, Qin F, Chen Y, Sun J, Tong J. [Chronotoxicity of 1800 MHz microwave radiation on sex hormones and spermatogenesis in male mice]. Wei Sheng Yan Jiu. **43(1):110-115, 2014.** [Article in Chinese]

OBJECTIVE: To study the chronotoxicity of 1800 MHz microwave radiation on the male reproductive system. METHODS: Sixty healthy male C57 mice with circadian rhythm in a 12:12 h light-dark photoperiod were divided into false radiation group (Sham) and microwave radiation (MR) group exposed to 1800 MHz RF at 208 microW/cm² power (SAR: 0.2221 W/kg) density at different zeitgeber times of a day (ZT01:00, ZT05:00, ZT09 : 00, ZT13: 00, ZT17 : 00, ZT21 : 00) for continuous 32 days with 2 h/d. The testicular sperm head was counted with a microscope, and serum testosterone (T) and estradiol (E2) levels were measured by ELISA method. RESULTS: Compared with the sham group, microwave radiation induced reduced level in testicular sperm head count and serum testosterone, while the level of serum estradiol increased. Also, the circadian rhythms of testicular sperm head count and estradiol disappeared after the microwave radiation. CONCLUSION: 1800 MHZ microwave radiation may disturb the level as well as circadian rhythmicity of the reproductive functions in male mice.

Celik S, Aridogan IA, Izol V, Erdoğan S, Polat S, Doran S. An Evaluation of the effects of long-term cell phone use on the testes via light and electron microscope analysis. Urology. **79(2):346-350, 2012**

Abstract. OBJECTIVE: To investigate whether the low-intensity electromagnetic waves transmitted by cell phones cause histopathological or ultrastructural changes in the testes of rats. MATERIALS AND METHODS: Wistar-Kyoto male rats were placed into either a control group or a group that was exposed to an electromagnetic field (EMF). Two cell phones with Specific Absorption Rate values of 1.58 were placed and left off in cages that housed 15 rats included in the control group, and four cell phones were placed and left on in cages that housed 30 rats included in the experimental group. After 3 months, weights, seminiferous tubule diameters, and spermatogenic cell conditions of all testes of the rats were evaluated. One half of each testis was examined also under an electron microscope. RESULTS: No significant differences were observed between the testis weights, seminiferous tubule diameters, and histopathological evaluations between rats that had and had not been exposed to EMF. Electron microscope analysis revealed that the membrana propria thickness and the collagen fiber contents were increased and the capillary veins extended in the experimental group. Common vacuolization in the cytoplasm of the Sertoli cells, growth of electron-dense structures, and existence of large lipid droplets were noted as the remarkable findings of this study.

Al-Damegh MA. Rat testicular impairment induced by electromagnetic radiation from a conventional cellular telephone and the protective effects of the antioxidants vitamins C and E. Clinics (Sao Paulo). **67(7):785-792, 2012.**

OBJECTIVE: The aim of this study was to investigate the possible effects of electromagnetic

Studies that show **Cell Phone** Health Effects

radiation from conventional cellular phone use on the oxidant and antioxidant status in rat blood and testicular tissue and determine the possible protective role of vitamins C and E in preventing the detrimental effects of electromagnetic radiation on the testes. **MATERIALS AND METHODS:** The treatment groups were exposed to an electromagnetic field, electromagnetic field plus vitamin C (40 mg/kg/day) or electromagnetic field plus vitamin E (2.7 mg/kg/day). All groups were exposed to the same electromagnetic frequency for 15, 30, and 60 min daily for two weeks. **RESULTS:** There was a significant increase in the diameter of the seminiferous tubules with a disorganized seminiferous tubule sperm cycle interruption in the electromagnetism-exposed group. The serum and testicular tissue conjugated diene, lipid hydroperoxide, and catalase activities increased 3-fold, whereas the total serum and testicular tissue glutathione and glutathione peroxidase levels decreased 3-5 fold in the electromagnetism-exposed animals. **CONCLUSION:** Our results indicate that the adverse effect of the generated electromagnetic frequency had a negative impact on testicular architecture and enzymatic activity. This finding also indicated the possible role of vitamins C and E in mitigating the oxidative stress imposed on the testes and restoring normality to the testes.

Azadi Oskouyi E, Rajaei F, Safari Variani A, Sarokhani MR, Javadi A. Effects of microwaves (950 MHz mobile phone) on morphometric and apoptotic changes of rabbit epididymis. Andrologia. 2014 Jul 25. doi: 10.1111/and.12321. [Epub ahead of print]

The effect of mobile phone radiation on human reproduction system is still a matter of debate. In this study, 18 male rabbits were randomly divided into two experimental groups and one control group. Experimental groups received simulated microwaves with the frequency of 950 MHz and the output power of 3 and 6 watts for 2 weeks, 2 h a day. After a week of rest, the microscopic slides from the quada of the excised epididymis were prepared. Then, the diameter of epididymis, the height of epithelium and the number of apoptotic cells in epithelium in study groups were determined. The data were compared using spss software and one-way anova test. The epithelial height and diameter of the epididymis in 3 watt and 6 watt groups had a significant decrease compared to the control group ($P < 0.001$), while the testosterone level only in 6 watt group was significantly decreased compared to control group. The rate of apoptosis in the epithelial cells of the epididymis had a significant increase only in 6 watt group compared to the control group ($P < 0.001$). This study showed that the microwaves with the frequency of 950 MHz can have negative impacts on morphometric and apoptotic changes of rabbit epididymis.

Baste V, Riise T, Moen BE. Radiofrequency electromagnetic fields; male infertility and sex ratio of offspring. Eur J Epidemiol.23(5):369-377,2008.

Concern is growing about exposure to electromagnetic fields and male reproductive health. The authors performed a cross-sectional study among military men employed in the Royal Norwegian Navy, including information about work close to equipment emitting radiofrequency electromagnetic fields, one-year infertility, children and sex of

Studies that show **Cell Phone** Health Effects

the offspring. Among 10,497 respondents, 22% had worked close to high-frequency aeriols to a "high" or "very high" degree. Infertility increased significantly along with increasing self-reported exposure to radiofrequency electromagnetic fields. In a logistic regression, odds ratio (OR) for infertility among those who had worked closer than 10 m from high-frequency aeriols to a "very high" degree relative to those who reported no work near high-frequency aeriols was 1.86 (95% confidence interval (CI): 1.46-2.37), adjusted for age, smoking habits, alcohol consumption and exposure to organic solvents, welding and lead. Similar adjusted OR for those exposed to a "high", "some" and "low" degree were 1.93 (95% CI: 1.55-2.40), 1.52 (95% CI: 1.25-1.84), and 1.39 (95% CI: 1.15-1.68), respectively. In all age groups there were significant linear trends with higher prevalence of involuntary childlessness with higher self-reported exposure to radiofrequency fields. However, the degree of exposure to radiofrequency radiation and the number of children were not associated. For self-reported exposure both to high-frequency aeriols and communication equipment there were significant linear trends with lower ratio of boys to girls at birth when the father reported a higher degree of radiofrequency electromagnetic exposure.

Panagopoulos DJ, Margaritis LH. The identification of an intensity 'window' on the bioeffects of mobile telephony radiation. *Int J Radiat Biol.* 86(5):358-366, 2010.

Erratum in *Int J Radiat Biol.* 2010 Sep;86(9):809.

PURPOSE: The increased bioactivity 'windows' of GSM 900 and 1800 MHz radiations, (Global System for Mobile telecommunications) revealed recently by us and published in this issue, manifesting themselves as a maximum decrease in the reproductive capacity of the insect *Drosophila melanogaster*, were examined to discover whether they depend on the intensity of radiation-fields. **METHODS:** In each experiment, one group of insects were exposed to the GSM 900 or 1800 radiation at 30 or 20 cm distances, respectively, from the antenna of a mobile phone, where the bioactivity 'window' appears for each type of radiation and another group was exposed at 8 or 5 cm, respectively, behind a metal grid, shielding both microwave radiation and the extremely low frequency (ELF) electric and magnetic fields for both types of radiation in a way that radiation and field intensities were roughly equal between the two groups. Then the effect on reproductive capacity was compared between groups for each type of radiation. **RESULTS:** The decrease in the reproductive capacity did not differ significantly between the two groups. **CONCLUSIONS:** The bioactivity window seems to be due to the intensity of radiation-field (10 microW/cm²), 0.6-0.7 V/m) at 30 or 20 cm from the GSM 900 or 1800 mobile phone antenna, respectively

Panagopoulos DJ, Margaritis LH. The effect of exposure duration on the biological activity of mobile telephony radiation. *Mutat Res*699(1-2):17-22,2010.

In the present experiments we studied the effects of different durations of a single, (continuous), daily exposure, ranging from 1min up to 21min, to the two established systems of digital mobile telephony radiation that are commonly used in Europe, viz. GSM 900MHz (Global System for Mobile telecommunications) and DCS 1800MHz (Digital Cellular System - referred to also as GSM 1800MHz), on a well-tested biological

Studies that show **Cell Phone** Health Effects

model, the reproductive capacity of the insect *Drosophila melanogaster*. The insects were exposed to each type of radiation at an intensity of about 10mW/cm², corresponding to a distance of 20cm or 30cm from the antenna of a DCS 1800 or a GSM 900 mobile phone handset, respectively. At these distances the bioactivity of mobile telephony radiation was found to be at a maximum due to the existence of a "window" of increased bioactivity around this value, as we have proposed recently [1-4]. The results show that the reproductive capacity decreases almost linearly with increasing exposure duration to both GSM 900 and DCS 1800 radiation, suggesting that short-term exposures to these radiations have cumulative effects on living organisms. Additionally, our results show again that GSM 900MHz radiation is slightly more bioactive than DCS 1800MHz radiation, at the same exposure durations and under equal radiation intensities, as shown in our previous experiments [5].

Panagopoulos DJ. Effect of microwave exposure on the ovarian development of *Drosophila melanogaster*. *Cell Biochem Biophys*. 63(2):121-132, 2012.

In the present experiments the effect of GSM radiation on ovarian development of virgin *Drosophila melanogaster* female insects was studied. Newly emerged adult female flies were collected and divided into separate identical groups. After the lapse of certain number of hours-different for each group-the insects (exposed and sham-exposed) were dissected and their intact ovaries were collected and photographed under an optical microscope with the same magnification. The size of the ovaries was compared between exposed and sham-exposed virgin female insects, during the time needed for the completion of oogenesis and maturation of the first eggs in the ovarioles. Immediately after the intact ovaries were photographed, they were further dissected into individual ovarioles and treated for TUNEL and acridine-orange assays to determine the degree of DNA damage in the egg chamber cells. The study showed that the ovarian size of the exposed insects is significantly smaller than that of the corresponding sham-exposed insects, due to destruction of egg chambers by the GSM radiation, after DNA damage and consequent cell death induction in the egg chamber cells of the virgin females as shown in previous experiments on inseminated females. The difference in ovarian size between sham-exposed and exposed virgin female flies becomes most evident 39-45 h after eclosion when the first eggs within the ovaries are at the late vitellogenic and post-vitellogenic stages (mid-late oogenesis). More than 45 h after eclosion, the difference in ovarian size decreases, as the first mature eggs of the sham-exposed insects are leaving the ovaries and are laid.

Forgacs Z, Somosy Z, Kubinyi G, Bakos J, Hudak A, Surjan A, Thuroczy G. Effect of whole-body 1800MHz GSM-like microwave exposure on testicular steroidogenesis and histology in mice. *Reprod Toxicol*.22(1):111-117, 2006.

The aim of our study was to evaluate the possible effects of whole-body 1800MHz GSM-like microwave exposure on male reproduction. After repeated exposure of mice to microwaves at 0.018-0.023W/kg whole-body specific energy absorption rate (SAR) an elevated serum testosterone level was measured, but no microwave exposure related histopathological alteration could be detected in the reproductive organs. The in vitro

Studies that show **Cell Phone** Health Effects

steroidogenic response of 48h Leydig cell cultures obtained from exposed animals did not differ from the controls, suggesting that Leydig cells were not the primary targets of the applied microwave exposure or direct action of microwaves on Leydig cells was temporary only. In exposed animals the red blood cell count and volume of packed red cells were also increased. Further investigations are required to clarify the mechanism of action of the applied microwave exposure on male mice, as well as to establish the biological significance of the observed phenomena.

Neurological Effects

Ntzouni MP, Stamatakis A, Stylianopoulou F, Margaritis LH. Short-term memory in mice is affected by mobile phone radiation. *Pathophysiology*. 18(3):193-199, 2011.

Abstract. The effects of mobile phone electromagnetic fields (EMFs) were studied on a non-spatial memory task (Object Recognition Task - ORT) that requires entorhinal cortex function. The task was applied to three groups of mice *Mus musculus* C57BL/6 (exposed, sham-exposed and control) combined with 3 different radiation exposure protocols. In the first protocol designated "acute exposure", mice 45 days old (PND45 - postnatal day 45) were exposed to mobile phone (MP) radiation (SAR value 0.22W/kg) during the habituation, the training and the test sessions of the ORT, but not during the 10min inter-trial interval (ITI) where consolidation of stored object information takes place. On the second protocol designated "chronic exposure-I", the same mice were exposed for 17 days for 90min/per day starting at PND55 to the same MP radiation. ORT recognition memory was performed at PND72 with radiation present only during the ITI phase. In the third protocol designated "chronic exposure-II", mice continued to be exposed daily under the same conditions up to PND86 having received radiation for 31 days. One day later the ORT test was performed without irradiation present in any of the sessions. The ORT-derived discrimination indices in all three exposure protocols revealed a major effect on the "chronic exposure-I" suggesting a possible severe interaction of EMF with the consolidation phase of recognition memory processes. This may imply that the primary EMF target may be the information transfer pathway connecting the entorhinal-parahippocampal regions which participate in the ORT memory task.

Yan JG, Agresti M, Zhang LL, Yan Y, Matloub HS. Upregulation of specific mRNA levels in rat brain after cell phone exposure. *Electromagn Biol Med*. 27(2):147-154, 2008.

Adult Sprague-Dawley rats were exposed to regular cell phones for 6 h per day for 126 days (18 weeks). RT-PCR was used to investigate the changes in levels of mRNA synthesis of several injury-associated proteins. Calcium ATPase, Neural Cell Adhesion Molecule, Neural Growth Factor, and Vascular Endothelial Growth Factor were evaluated. The results showed statistically significant mRNA up-regulation of these proteins in the brains of rats exposed to cell phone radiation. These results indicate that relative chronic exposure to cell phone microwave radiation may result in cumulative injuries that could eventually lead to clinically significant neurological damage.

Studies that show **Cell Phone** Health Effects

Hocking B, Westerman R, Neurological abnormalities associated with Mobile phone use. Occup Med 50: 366-368, 2000.

Dysaesthesiae of the scalp after mobile phone use have been previously reported but the pathological basis of these symptoms has been unclear. We report finding a neurological abnormality in a patient after prolonged use of a mobile phone. He had permanent unilateral dysaesthesiae of the scalp, slight loss of sensation, and abnormalities on current perception threshold testing of cervical and trigeminal nerves. A neurologist found no other disease. The implications regarding health effects of mobile phones and radio-frequency radiation is discussed.

Deshmukh PS, Banerjee BD, Abegaonkar MP, Megha K, Ahmed RS, Tripathi AK, Mediratta PK. Effect of low level microwave radiation exposure on cognitive function and oxidative stress in rats. Indian J Biochem Biophys. 50(2):114-119, 2013.

Use of wireless communicating devices is increasing at an exponential rate in present time and is raising serious concerns about possible adverse effects of microwave (MW) radiation emitted from these devices on human health. The present study aimed to evaluate the effects of 900 MHz MW radiation exposure on cognitive function and oxidative stress in blood of Fischer rats. Animals were divided into two groups (6 animals/group): Group I (MW-exposed) and Group II (Sham-exposed). Animals were subjected to MW exposure (Frequency 900 MHz; specific absorption rate 8.4738×10^{-5} W/kg) in Gigahertz transverse electromagnetic cell (GTEM) for 30 days (2 h/day, 5 days/week). Subsequently, cognitive function and oxidative stress parameters were examined for each group. Results showed significant impairment in cognitive function and increase in oxidative stress, as evidenced by the increase in levels of MDA (a marker of lipid peroxidation) and protein carbonyl (a marker of protein oxidation) and unaltered GSH content in blood. Thus, the study demonstrated that low level MW radiation had significant effect on cognitive function and was also capable of leading to oxidative stress.

Maier R, Greter SE, Maier N. Effects of pulsed electromagnetic fields on cognitive processes - a pilot study on pulsed field interference with cognitive regeneration. Acta Neurol Scand. 110(1):46-52, 2004.

BACKGROUND: Due to the ubiquitous use of cellular phones much has been speculated on secondary effects of electromagnetic irradiation emitted by those. Additionally, several studies have reported vegetative alterations as well as effects on the neuronal and molecular levels in humans. Here, using a psycho-physiological test paradigm, we examined effects of exposure to pulsed electromagnetic fields on cognitive performance. MATERIALS AND METHODS: In 11 volunteers, we tested cognitive processing under field exposure (GSM standard) and under field-free conditions. To examine the hypothesized effect of pulsed fields, we applied an auditory discrimination task and determined the participant's current 'Order Threshold' value. Following a first test cycle, the volunteers had to relax for 50 min while being, or not, exposed to pulsed electromagnetic fields. Subsequently, the test was repeated. Data acquired before and after the resting phase were compared from both experimental conditions. RESULTS:

Studies that show **Cell Phone** Health Effects

We found that nine of the 11 test participants (81.8%) showed worse results in their auditory discrimination performance upon field exposure as compared with control conditions. Group data comparison revealed a statistical significance of $P = 0.0105$. CONCLUSION: We could show that the participants' cognitive performance was impaired after exposure to pulsed electromagnetic fields. With regard to this finding, we recommend that the use of cellular phones should be restricted generally and in particular in respect of physical hazard of high-risk groups, e.g. elderly, children and ill people.

Maier R, Greter SE, Maier N. Effects of pulsed electromagnetic fields on cognitive processes - a pilot study on pulsed field interference with cognitive regeneration. Acta Neurol Scand. 110(1):46-52, 2004.

BACKGROUND: Due to the ubiquitous use of cellular phones much has been speculated on secondary effects of electromagnetic irradiation emitted by those. Additionally, several studies have reported vegetative alterations as well as effects on the neuronal and molecular levels in humans. Here, using a psycho-physiological test paradigm, we examined effects of exposure to pulsed electromagnetic fields on cognitive performance. MATERIALS AND METHODS: In 11 volunteers, we tested cognitive processing under field exposure (GSM standard) and under field-free conditions. To examine the hypothesized effect of pulsed fields, we applied an auditory discrimination task and determined the participant's current 'Order Threshold' value. Following a first test cycle, the volunteers had to relax for 50 min while being, or not, exposed to pulsed electromagnetic fields. Subsequently, the test was repeated. Data acquired before and after the resting phase were compared from both experimental conditions. RESULTS: We found that nine of the 11 test participants (81.8%) showed worse results in their auditory discrimination performance upon field exposure as compared with control conditions. Group data comparison revealed a statistical significance of $P = 0.0105$. CONCLUSION: We could show that the participants' cognitive performance was impaired after exposure to pulsed electromagnetic fields. With regard to this finding, we recommend that the use of cellular phones should be restricted generally and in particular in respect of physical hazard of high-risk groups, e.g. elderly, children and ill people.

Hocking B, Westerman R. Neurological effects of radiofrequency radiation. Occup Med (Lond) 53(2):123-127, 2003.

BACKGROUND: The health effects of radiofrequency radiation (RFR) and the adequacy of the safety standards are a subject of debate. One source of human data is case reports regarding peripheral neurological effects of RFR, mainly noxious sensations or dysaesthesiae. Aim To investigate health effects, neurophysiological mechanisms and safety levels for RFR. METHODS: We conducted a literature search for case reports and case series associated with mobile phone technology as well as other RFR sources using specific search terms on PubMed. RESULTS: We identified 11 original articles detailing case reports or case series and matching the search criteria. Five of the identified papers were written by at least one of the authors (B.H. or R.W.). CONCLUSIONS: Cases have

Studies that show **Cell Phone** Health Effects

arisen after exposure to much of the radiofrequency range. In some cases, symptoms are transitory but lasting in others. After very high exposures, nerves may be grossly injured. After lower exposures, which may result in dysaesthesia, ordinary nerve conduction studies find no abnormality but current perception threshold studies have found abnormalities. Only a small proportion of similarly exposed people develop symptoms. The role of modulations needs clarification. Some of these observations are not consistent with the prevailing hypothesis that all health effects of RFR arise from thermal mechanisms.

Nittby H, Grafström G, Tian DP, Malmgren L, Brun A, Persson BR, Salford LG, Eberhardt J. Cognitive impairment in rats after long-term exposure to GSM-900 mobile phone radiation. *Bioelectromagnetics*. 29:219-232, 2008.

Considering the frequent use of mobile phones, we have directed attention to possible implications on cognitive functions. In this study we investigated in a rat model the long-term effects of protracted exposure to Global System for Mobile Communication-900 MHz (GSM-900) radiation. Out of a total of 56 rats, 32 were exposed for 2 h each week for 55 weeks to radio-frequency electromagnetic radiation at different SAR levels (0.6 and 60 mW/kg at the initiation of the experimental period) emitted by a (GSM-900) test phone. Sixteen animals were sham exposed and eight animals were cage controls, which never left the animal house. After this protracted exposure, GSM-900 exposed rats were compared to sham exposed controls. Effects on exploratory behaviour were evaluated in the open-field test, in which no difference was seen. Effects on cognitive functions were evaluated in the episodic-like memory test. In our study, GSM exposed rats had impaired memory for objects and their temporal order of presentation, compared to sham exposed controls ($P = 0.02$). Detecting the place in which an object was presented was not affected by GSM exposure. Our results suggest significantly reduced memory functions in rats after GSM microwave exposure ($P = 0.02$).

Croft R, Chandler J, Burgess A, Barry R, Williams J, Clarke A. Acute mobile phone operation affects neural function in humans. *Clin Neurophysiol* 113(10):1623, 2002.

OBJECTIVES: Mobile phones (MP) are used extensively and yet little is known about the effects they may have on human physiology. There have been conflicting reports regarding the relation between MP use and the electroencephalogram (EEG). The present study suggests that this conflict may be due to methodological differences such as exposure durations, and tests whether exposure to an active MP affects EEG as a function of time. METHODS: Twenty-four subjects participated in a single-blind fully counterbalanced cross-over design, where both resting EEG and phase-locked neural responses to auditory stimuli were measured while a MP was either operating or turned off. RESULTS: MP exposure altered resting EEG, decreasing 1-4Hz activity (right hemisphere sites), and increasing 8-12Hz activity as a function of exposure duration (midline posterior sites). MP exposure also altered early phase-locked neural responses, attenuating the normal response decrement over time in the 4-8Hz band, decreasing the response in the 12-30Hz band globally and as a function of time, and increasing midline frontal and lateral posterior responses in the 30-45Hz band. CONCLUSIONS:

Studies that show **Cell Phone** Health Effects

Active MPs affect neural function in humans and do so as a function of exposure duration. The temporal nature of this effect may contribute to the lack of consistent results reported in the literature.

Eliyahu I, Luria R, Hareuveny R, Margaliot M, Meiran N, Shani G. Effects of radiofrequency radiation emitted by cellular telephones on the cognitive functions of humans. *Bioelectromagnetics*.27(2):119-126, 2006.

The present study examined the effects of exposure to Electromagnetic Radiation emitted by a standard GSM phone at 890 MHz on human cognitive functions. This study attempted to establish a connection between the exposure of a specific area of the brain and the cognitive functions associated with that area. A total of 36 healthy right-handed male subjects performed four distinct cognitive tasks: spatial item recognition, verbal item recognition, and two spatial compatibility tasks. Tasks were chosen according to the brain side they are assumed to activate. All subjects performed the tasks under three exposure conditions: right side, left side, and sham exposure. The phones were controlled by a base station simulator and operated at their full power. We have recorded the reaction times (RTs) and accuracy of the responses. The experiments consisted of two sections, of 1 h each, with a 5 min break in between. The tasks and the exposure regimes were counterbalanced. The results indicated that the exposure of the left side of the brain slows down the left-hand response time, in the second-later-part of the experiment. This effect was apparent in three of the four tasks, and was highly significant in only one of the tests. The exposure intensity and its duration exceeded the common exposure of cellular phone users.

Narayanan SN, Kumar RS, Potu BK, Nayak S, Mailankot M. Spatial memory performance of Wistar rats exposed to mobile phone. *Clinics*. 64(3):231-234, 2009.

INTRODUCTION: With the tremendous increase in number of mobile phone users world wide, the possible risks of this technology have become a serious concern. OBJECTIVE: We tested the effects of mobile phone exposure on spatial memory performance. MATERIALS AND METHODS: Male Wistar rats (10-12 weeks old) were exposed to 50 missed calls/day for 4 weeks from a GSM (900/1800 MHz) mobile phone in vibratory mode (no ring tone). After the experimental period, the animals were tested for spatial memory performance using the Morris water maze test. RESULTS: Both phone exposed and control animals showed a significant decrease in escape time with training. Phone exposed animals had significantly (approximately 3 times) higher mean latency to reach the target quadrant and spent significantly (approximately 2 times) less time in the target quadrant than age- and sex-matched controls. CONCLUSION: Mobile phone exposure affected the acquisition of learned responses in Wistar rats. This in turn points to the poor spatial navigation and the object place configurations of the phone-exposed animals.

Schneider J, Stangassinger M. Nonthermal Effects of Lifelong High-Frequency Electromagnetic Field Exposure on Social Memory Performance in Rats. *Behav Neurosci*. 2014 Jul 7. [Epub ahead of print]

Studies that show **Cell Phone** Health Effects

We are today surrounded almost constantly by high-frequency electromagnetic fields (EMFs) from mobile communications base stations. To date, however, there has been little concern regarding nonthermal effects of EMFs on cognition. In the present study, male and female rats were subjected to continuous far-field exposure to a frequency of 900-MHz (Global System for Mobile Communications [GSM]) or 1.966-GHz (Universal Mobile Telecommunications System [UMTS]) at 0.4 W/kg. Memory performance of adult EMF-exposed and sham-exposed female rats (at 6 months of age) and male rats (at 3 and 6 months of age) was tested using a social discrimination procedure. For this procedure, a target juvenile male was introduced to the subject's home cage for 4 min (Trial 1). After 30 min, the same target animal and a novel juvenile male were simultaneously presented to the subject for 4 min (Trial 2). Differences in sniffing duration to the familiar and novel target rats during Trial 2 were used to assess memory performance. EMF-exposed females exhibited no differences in sniffing duration compared with controls. In contrast, the sniffing durations of EMF-exposed males at 3 months of age were significantly affected. At 6 months of age, GSM-, but not UMTS-, exposed male adults showed a memory performance deficit. These findings provide new insight into the nonthermal effects of long-term high-frequency EMF exposure on memory.

Li H, Peng R, Wang C, Qiao S, Yong-Zou, Gao Y, Xu X, Wang S, Dong J, Zuo H, Li-Zhao, Zhou H, Wang L, Hu X. Alterations of cognitive function and 5-HT system in rats after long term microwave exposure. *Physiol Behav.* 2014 Dec 24. pii: S0031-9384(14)00663-5. doi: 10.1016/j.physbeh.2014.12.039. [Epub ahead of print]

The increased use of microwaves raises concerns about its impact on health including cognitive function in which neurotransmitter system plays an important role. In this study, we focused on the serotonergic system and evaluated the long term effects of chronic microwave radiation on cognition and correlated items. Wistar rats were exposed or sham exposed to 2.856GHz microwaves with the average power density of 5, 10, 20 or 30mW/cm² respectively for 6min three times a week up to 6weeks. At different time points after the last exposure, spatial learning and memory function, morphology structure of the hippocampus, electroencephalogram (EEG) and neurotransmitter content (amino acid and monoamine) of rats were tested. Above results raised our interest in serotonin system. Tryptophan hydroxylase 1 (TPH1) and monoamine oxidase (MAO), two important rate-limiting enzymes in serotonin synthesis and metabolic process respectively, were detected. Expressions of serotonin receptors including 5-HT_{1A}, 2A, 2C receptors were measured. We demonstrated that chronic exposure to microwave (2.856GHz, with the average power density of 5, 10, 20 and 30mW/cm²) could induce dose-dependent deficit of spatial learning and memory in rats accompanied with inhibition of brain electrical activity, the degeneration of hippocampus neurons, and the disturbance of neurotransmitters, among which the increase of 5-HT occurred as the main long-term change that the decrease of its metabolism partly contributed to. Besides, the variations of 5-HT_{1A}R and 5-HT_{2C}R expressions were also indicated. The results suggested that in the long-term way,

Studies that show **Cell Phone** Health Effects

chronic microwave exposure could induce cognitive deficit and 5-HT system may be involved in it.

Ntzouni MP, Skouroliahou A, Kostomitsopoulos N, Margaritis LH. Transient and cumulative memory impairments induced by GSM 1.8 GHz cell phone signal in a mouse model. Electromagn Biol Med. 2013 Jan 15. [Epub ahead of print]

This study was designed to investigate the transient and cumulative impairments in spatial and non-spatial memory of C57Bl/6J mice exposed to GSM 1.8 GHz signal for 90 min daily by a typical cellular (mobile) phone at a specific absorption rate value of 0.11 W/kg. Free-moving male mice 2 months old were irradiated in two experimental protocols, lasting for 66 and for 148 days respectively. Each protocol used three groups of animals (n = 8 each for exposed, sham exposed and controls) in combination with two behavioural paradigms, the object recognition task and the object location task sequentially applied at different time points. One-way analysis of variance revealed statistically significant impairments of both types of memory gradually accumulating, with more pronounced effects on the spatial memory. The impairments persisted even 2 weeks after interruption of the 8 weeks daily exposure, whereas the memory of mice as detected by both tasks showed a full recovery approximately 1 month later. Intermittent every other day exposure for 1 month had no effect on both types of memory. The data suggest that visual information processing mechanisms in hippocampus, perirhinal and entorhinal cortex are gradually malfunctioning upon long-term daily exposure, a phenotype that persists for at least 2 weeks after interruption of radiation, returning to normal memory performance levels 4 weeks later. It is postulated that cellular repair mechanisms are operating to eliminate the memory affecting molecules. The overall contribution of several possible mechanisms to the observed cumulative and transient impairments in spatial and non-spatial memory is discussed.

Vecchio F, Babiloni C, Ferreri F, Curcio G, Fini R, Del Percio C, Rossini PM. Mobile phone emission modulates interhemispheric functional coupling of EEG alpha rhythms. Eur J Neurosci. 25(6):1908-1913, 2007.

We tested the working hypothesis that electromagnetic fields from mobile phones (EMFs) affect interhemispheric synchronization of cerebral rhythms, an important physiological feature of information transfer into the brain. Ten subjects underwent two electroencephalographic (EEG) recordings, separated by 1 week, following a crossover double-blind paradigm in which they were exposed to a mobile phone signal (global system for mobile communications; GSM). The mobile phone was held on the left side of the subject head by a modified helmet, and orientated in the normal position for use over the ear. The microphone was orientated towards the corner of the mouth, and the antenna was near the head in the parietotemporal area. In addition, we positioned another similar phone (but without battery) on the right side of the helmet, to balance the weight and to prevent the subject localizing the side of GSM stimulation (and consequently lateralizing attention). In one session the exposure was real (GSM) while in the other it was Sham; both sessions lasted 45 min. Functional interhemispheric connectivity was modelled using the analysis of EEG spectral coherence between frontal, central and parietal electrode pairs. Individual EEG rhythms of interest were

Studies that show **Cell Phone** Health Effects

delta (about 2-4 Hz), theta (about 4-6 Hz), alpha 1 (about 6-8 Hz), alpha 2 (about 8-10 Hz) and alpha 3 (about 10-12 Hz). Results showed that, compared to Sham stimulation, GSM stimulation modulated the interhemispheric frontal and temporal coherence at alpha 2 and alpha 3 bands. The present results suggest that prolonged mobile phone emission affects not only the cortical activity but also the spread of neural synchronization conveyed by interhemispherical functional coupling of EEG rhythms.

Vecchio F, Babiloni C, Ferreri F, Buffo P, Cibelli G, Curcio G, Dijkman SV, Melgari JM, Giambattistelli F, Rossini PM. Mobile phone emission modulates inter-hemispheric functional coupling of EEG alpha rhythms in elderly compared to young subjects. Clin Neurophysiol. 121(2):163-171, 2010.

OBJECTIVE: It has been reported that GSM electromagnetic fields (GSM-EMFs) of mobile phones modulate - after a prolonged exposure - inter-hemispheric synchronization of temporal and frontal resting electroencephalographic (EEG) rhythms in normal young subjects [Vecchio et al., 2007]. Here we tested the hypothesis that this effect can vary on physiological aging as a sign of changes in the functional organization of cortical neural synchronization. **METHODS:** Eyes-closed resting EEG data were recorded in 16 healthy elderly subjects and 5 young subjects in the two conditions of the previous reference study. The GSM device was turned on (45min) in one condition and was turned off (45min) in the other condition. Spectral coherence evaluated the inter-hemispheric synchronization of EEG rhythms at the following bands: delta (about 2-4Hz), theta (about 4-6Hz), alpha 1 (about 6-8Hz), alpha 2 (about 8-10Hz), and alpha 3 (about 10-12Hz). The aging effects were investigated comparing the inter-hemispheric EEG coherence in the elderly subjects vs. a young group formed by 15 young subjects (10 young subjects of the reference study; Vecchio et al., 2007). **RESULTS:** Compared with the young subjects, the elderly subjects showed a statistically significant ($p<0.001$) increment of the inter-hemispheric coherence of frontal and temporal alpha rhythms (about 8-12Hz) during the GSM condition. **CONCLUSIONS:** These results suggest that GSM-EMFs of a mobile phone affect inter-hemispheric synchronization of the dominant (alpha) EEG rhythms as a function of the physiological aging. **SIGNIFICANCE:** This study provides further evidence that physiological aging is related to changes in the functional organization of cortical neural synchronization.

Hocking B, Westerman R. Neurological changes induced by a mobile phone. Occup Med (Lond) 52(7):413-415, 2002.

Dysaesthesiae of the scalp after mobile phone use have been previously reported, but the basis for this has not been clear. We report a case of a 34-year-old journalist who complained of symptoms associated with use of a mobile phone. She agreed to a provocation study with her phone. Current perception threshold testing before and after exposure showed marked changes in the C-fibre nerves of the affected area compared with the opposite side. The case is supportive of a neurological basis for some cases of dysaesthesiae associated with mobile phone use.

Curcio G, Ferrara M, Moroni F, D'Inzeo G, Bertini M, De Gennaro L. Is the brain

Studies that show **Cell Phone** Health Effects

influenced by a phone call? An EEG study of resting wakefulness. Neurosci Res. 53(3):265-270, 2005.

We recorded the resting electroencephalogram of 20 healthy subjects in order to investigate the effect of electromagnetic field (EMF) exposure on EEG waking activity and its temporal development. The subjects were randomly assigned to two groups and exposed, in double-blind conditions, to a typical mobile phone signal (902.40MHz, modulated at 217Hz, with an average power of 0.25W) before or during the EEG recording session. The results show that, under real exposure as compared to baseline and sham conditions, EEG spectral power was influenced in some bins of the alpha band. This effect was greater when the EMF was on during the EEG recording session than before it. The present data lend further support to the idea that pulsed high-frequency electromagnetic fields can affect normal brain functioning, also if no conclusions can be drawn about the possible health effects.

Vecchio F, Tombini M, Buffo P, Assenza G, Pellegrino G, Benvenga A, Babiloni C, Rossini PM. Mobile phone emission increases inter-hemispheric functional coupling of electroencephalographic alpha rhythms in epileptic patients. Int J Psychophysiol.84(2):164-171, 2012.

It has been reported that GSM electromagnetic fields (GSM-EMFs) of mobile phones modulate - after a prolonged exposure - inter-hemispheric synchronization of temporal and frontal resting electroencephalographic (EEG) rhythms in normal young and elderly subjects (Vecchio et al., 2007, 2010). Here we tested the hypothesis that this can be even more evident in epileptic patients, who typically suffer from abnormal mechanisms governing synchronization of rhythmic firing of cortical neurons. Eyes-closed resting EEG data were recorded in ten patients affected by focal epilepsy in real and sham exposure conditions. These data were compared with those obtained from 15 age-matched normal subjects of the previous reference studies. The GSM device was turned on (45min) in the "GSM" condition and was turned off (45min) in the other condition ("sham"). The mobile phone was always positioned on the left side in both patients and control subjects. Spectral coherence evaluated the inter-hemispheric synchronization of EEG rhythms at the following frequency bands: delta (about 2-4Hz), theta (about 4-6Hz), alpha1 (about 6-8Hz), alpha2 (about 8-10Hz), and alpha3 (about 10-12Hz). The effects on the patients were investigated comparing the inter-hemispheric EEG coherence in the epileptic patients with the control group of subjects evaluated in the previous reference studies. Compared with the control subjects, epileptic patients showed a statistically significant higher inter-hemispheric coherence of temporal and frontal alpha rhythms (about 8-12Hz) in the GSM than "Sham" condition. These results suggest that GSM-EMFs of mobile phone may affect inter-hemispheric synchronization of the dominant (alpha) EEG rhythms in epileptic patients. If confirmed by future studies on a larger group of epilepsy patients, the modulation of the inter-hemispheric alpha coherence due to the GSM-EMFs could have clinical implications and be related to changes in cognitive-motor function.

Balikci K, Cem Ozcan I, Turgut-Balik D, Balik HH. A survey study on some neurological

Studies that show **Cell Phone** Health Effects

symptoms and sensations experienced by long term users of mobile phones. Pathol Biol (Paris). 53(1):30-34, 2005.

A survey study was conducted to investigate the possible effects of mobile phone on headache, dizziness, extreme irritation, shaking in the hands, speaking falteringly, forgetfulness, neuro-psychological discomfort, increase in the carelessness, decrease of the reflex and clicking sound in the ears. There is no effect on dizziness, shaking in hands, speaking falteringly and neuro-psychological discomfort, but some statistical evidences are found that mobile phone may cause headache, extreme irritation, increase in the carelessness, forgetfulness, decrease of the reflex and clicking sound in the ears.

Vecsei Z, Csathó A, Thuróczy G, Hernádi I. Effect of a single 30 min UMTS mobile phone-like exposure on the thermal pain threshold of young healthy volunteers. Bioelectromagnetics. 2013 Jun 20. doi: 10.1002/bem.21801. [Epub ahead of print]

One of the most frequently investigated effects of radiofrequency electromagnetic fields (RF EMFs) on the behavior of complex biological systems is pain sensitivity. Despite the growing body of evidence of EMF-induced changes in pain sensation, there is no currently accepted experimental protocol for such provocation studies for the healthy human population. In the present study, therefore, we tested the effects of third generation Universal Mobile Telecommunications System (UMTS) RF EMF exposure on the thermal pain threshold (TPT) measured on the surface of the fingers of 20 young adult volunteers. The protocol was initially validated with a topical capsaicin treatment. The exposure time was 30 min and the genuine (or sham) signal was applied to the head through a patch antenna, where RF EMF specific absorption rate (SAR) values were controlled and kept constant at a level of 1.75 W/kg. Data were obtained using randomized, placebo-controlled trials in a double-blind manner. Subjective pain ratings were tested blockwise on a visual analogue rating scale (VAS). Compared to the control and sham conditions, the results provide evidence for intact TPT but a reduced desensitization effect between repeated stimulations within the individual blocks of trials, observable only on the contralateral side for the genuine UMTS exposure. Subjective pain perception (VAS) data indicated marginally decreased overall pain ratings in the genuine exposure condition only. The present results provide pioneering information about human pain sensation in relation to RF EMF exposure and thus may contribute to cover the existing gap between safety research and applied biomedical science targeting the potential biological effects of environmental RF EMFs.

Luria R, Eliyahu I, Hareuveny R, Margaliot M, Meiran N. Cognitive effects of radiation emitted by cellular phones: the influence of exposure side and time. Bioelectromagnetics. 30(3):198-204, 2009.

This study examined the time dependence effects of exposure to radiofrequency radiation (RFR) emitted by standard GSM cellular phones on the cognitive functions of humans. A total of 48 healthy right-handed male subjects performed a spatial working memory task (that required either a left-hand or a right-hand response) while being exposed to one of two GSM phones placed at both sides of the head. The subjects were

Studies that show **Cell Phone** Health Effects

randomly divided into three groups. Each group was exposed to one of three exposure conditions: left-side of the head, right-side, or sham-exposure. The experiment consisted of 12 blocks of trials. Response times (RTs) and accuracy of the responses were recorded. It was found that the average RT of the right-hand responses under left-side exposure condition was significantly longer than those of the right-side and sham-exposure groups averaged together during the first two time blocks. These results confirmed the existence of an effect of exposure on RT, as well as the fact that exposure duration (together with the responding hand and the side of exposure) may play an important role in producing detectable RFR effects on performance. Differences in these parameters might be the reason for the failure of certain studies to detect or replicate RFR effects.

Maskey D, Kim MJ. Immunohistochemical Localization of Brain-derived Neurotrophic Factor and Glial Cell Line-derived Neurotrophic Factor in the Superior Olivary Complex of Mice after Radiofrequency Exposure. Neuroscience Letters. 564:78-82, 2014.

Raising health concerns about the biological effects from radiofrequency exposure, even with conflicting results, has prompted calls for formulation of a guideline of the biological safety level. Given the close proximity between a mobile phone and the ear, it has been suggested that the central auditory system may be detrimentally influenced by radiofrequency exposure. In the auditory system, neurotrophins are important in the regulation of neuron survival, especially mammalian cochlear neurons. Neurotrophic factors like brain-derived neurotrophic factor (BDNF) and glial-derived neurotrophic factor (GDNF) present in the auditory system are responsible for the maintenance of auditory neurons. BDNF and GDNF may protect against acoustic trauma and prevent from hearing defect. The present study applied radiofrequency at a specific absorption rate (SAR) of 1.6 W/kg (E1.6) or 0 W/kg group to determine the distribution of BDNF and GDNF in the nuclei of superior olivary complex (SOC). In the E1.6 group, significant decrements of BDNF immunoreactivity (IR) were noted in the lateral superior olive, medial superior olive, superior paraolivary nucleus and medial nucleus of the trapezoid body. GDNF IR was also significantly decreased ($p < 0.001$) in all SOC nuclei of the E1.6 group. The decrease in the IR of these neurotrophic factors in the SOC of the E1.6 group suggests a detrimental effect of RF exposure in the auditory nuclei.

Qin F, Yuan H, Nie J, Cao Y, Tong J. [Effects of nano-selenium on cognition performance of mice exposed in 1800 MHz radiofrequency fields]. Wei Sheng Yan Jiu. 43(1):16-21, 2014. [Article in Chinese]

OBJECTIVE: To study the effects of nano-selenium (NSe) on cognition performance of mice exposed to 1800 MHz radiofrequency fields (RF). METHODS: Male mice were randomly divided into four groups, control and nano-Se low, middle and high dose groups (L, M, H). Each group was sub-divided into three groups, RF 0 min, RF 30 min and RF 120 min. Nano-se solution (2, 4 and 8 microg/ml) were administered to mice of L, M, H groups by intra-gastric injection respectively, 0.5 ml/d for 50 days, the conctral group were administered with distilled water. At the 21st day, the mice in RF subgroup were exposed to 208 microW/cm² 1800 MHz radiofrequency fields (0, 30 and 120 min/d

Studies that show **Cell Phone** Health Effects

respectively) for 30 days. The cognitive ability of the mice were tested with Y-maze. Further, the levels of MDA, GABA, Glu, Ach and the activities of CAT and GSH-Px in cerebra were measured. RESULTS: Significant impairments in learning and memory ($P < 0.05$) were observed in the RF 120 min group, and with reduction of the Ach level and the activities of CAT and GSH-Px and increase of the content of GABA, Glu and MDA in cerebrum. NSe enhanced cognitive performance of RF mice, decreased GABA, Glu and MDA levels, increased Ach levels, GSH-Px and CAT activities. CONCLUSION: NSe could improve cognitive impairments of mice exposed to RF, the mechanism of which might involve the increasing antioxidation, decreasing free radical content and the changes of cerebra neurotransmitters.

Fragopoulou AF, Miltiadous P, Stamatakis A, Stylianopoulou F, Koussoulakos SL, Margaritis LH. Whole body exposure with GSM 900MHz affects spatial memory in mice. Pathophysiology. 17(3):179-187,2010.

Extended work has been performed worldwide on the effects of mobile phone radiation upon rats' cognitive functions, however there is great controversy to the existence or not of deficits. The present work has been designed in order to test the effects of mobile phone radiation on spatial learning and memory in mice *Mus musculus* Balb/c using the Morris water maze (a hippocampal-dependent spatial memory task), since there is just one other study on mice with very low SAR level (0.05W/kg) showing no effects. We have applied a 2h daily dose of pulsed GSM 900MHz radiation from commercially available mobile phone for 4 days at SAR values ranging from 0.41 to 0.98W/kg. Statistical analysis revealed that during learning, exposed animals showed a deficit in transferring the acquired spatial information across training days (increased escape latency and distance swam, compared to the sham-exposed animals, on the first trial of training days 2-4). Moreover, during the memory probe-trial sham-exposed animals showed the expected preference for the target quadrant, while the exposed animals showed no preference, indicating that the exposed mice had deficits in consolidation and/or retrieval of the learned spatial information. Our results provide a basis for more thorough investigations considering reports on non-thermal effects of electromagnetic fields (EMFs).

Bouji M, Lecomte A, Hode Y, de Seze R, Villégier AS. Effects of 900 MHz radiofrequency on corticosterone, emotional memory and neuroinflammation in middle-aged rats. Exp Gerontol. 47(6):444-451, 2012.

The widespread use of mobile phones raises the question of the effects of electromagnetic fields (EMF, 900 MHz) on the brain. Previous studies reported increased levels of the glial fibrillary acidic protein (GFAP) in the rat's brain after a single exposure to 900 MHz global system for mobile (GSM) signal, suggesting a potential inflammatory process. While this result was obtained in adult rats, no data is currently available in older animals. Since the transition from middle-age to senescence is highly dependent on environment and lifestyle, we studied the reactivity of middle-aged brains to EMF exposure. We assessed the effects of a single 15 min GSM exposure (900 MHz; specific absorption rate (SAR)=6 W/kg) on GFAP expression in young adults (6

Studies that show **Cell Phone** Health Effects

week-old) and middle-aged rats (12 month-old). Brain interleukin (IL)-1 β and IL-6, plasmatic levels of corticosterone (CORT), and emotional memory were also assessed. Our data indicated that, in contrast to previously published work, acute GSM exposure did not induce astrocyte activation. Our results showed an IL-1 β increase in the olfactory bulb and enhanced contextual emotional memory in GSM-exposed middle-aged rats, and increased plasmatic levels of CORT in GSM-exposed young adults. Altogether, our data showed an age dependency of reactivity to GSM exposure in neuro-immunity, stress and behavioral parameters. Reproducing these effects and studying their mechanisms may allow a better understanding of mobile phone EMF effects on neurobiological parameters.

Masuda H, Hirata A, Kawai H, Wake K, Watanabe S, Arima T, Poullietier de Gannes F, Lagroye I, Veyret B. Local exposure of the rat cortex to radiofrequency electromagnetic fields increases local cerebral blood flow along with temperature. J Appl Physiol. 110(1):142-148, 2011.

Few studies have shown that local exposure to radiofrequency electromagnetic fields (RF) induces intensity-dependent physiological changes, especially in the brain. The aim of the present study was to detect reproducible responses to local RF exposure in the parietal cortex of anesthetized rats and to determine their dependence on RF intensity. The target cortex tissue was locally exposed to 2-GHz RF using a figure-eight loop antenna within a range of averaged specific absorption rates (10.5, 40.3, 130, and 263 W/kg averaged over 4.04 mg) in the target area. Local cerebral blood flow (CBF) and temperatures in three regions (target area, rectum, and calf hypodermis) were measured using optical fiber blood flow meters and thermometers during RF exposure. All parameters except for the calf hypodermis temperature increased significantly in exposed animals compared with sham-exposed ones during 18-min exposures. Dependence of parameter values on exposure intensity was analyzed using linear regression models. The elevation of local CBF was correlated with temperature rise in both target and rectum at the end of RF exposure. However, the local CBF elevation seemed to be elevated by the rise in target temperature, but not by that of the rectal temperature, in the early part of RF exposure or at low-intensity RF exposure. These findings suggest that local RF exposure of the rat cortex drives a regulation of CBF accompanied by a local temperature rise, and our findings may be helpful for discussing physiological changes in the local cortex region, which is locally exposed to RF.

Lovisol GA, Giardino L, Calzaà L. Effect of radiofrequency electromagnetic field exposure on in vitro models of neurodegenerative disease. Bioelectromagnetics.30(7):564-72, 2009.

In this work we tested viability, proliferation, and vulnerability of neural cells, after continuous radiofrequency (RF) electromagnetic fields exposure (global system for mobile telecommunications (GSM) modulated 900 MHz signal at a specific absorption rate (SAR) of 1 W/kg and maximum duration 144 h) generated by transverse electromagnetic cells. We used two cellular systems, SN56 cholinergic for example, SN56 cholinergic cell line and rat primary cortical neurons, and well-known neurotoxic

Studies that show **Cell Phone** Health Effects

challenges, such as glutamate, 25-35AA beta-amyloid, and hydrogen peroxide. Exposure to RF did not change viability/proliferation rate of the SN56 cholinergic cells or viability of cortical neurons. Co-exposure to RF exacerbated neurotoxic effect of hydrogen peroxide in SN56, but not in primary cortical neurons, whereas no cooperative effects of RF with glutamate and 25-35AA beta-amyloid were found. These data suggest that only under particular circumstances exposure to GSM modulated, 900 MHz signal act as a co-stressor for oxidative damage of neural cells.

Yilmaz A, Yilmaz N, Serarslan Y, Aras M, Altas M, Ozgür T, Sefil F. The effects of mobile phones on apoptosis in cerebral tissue: an experimental study on rats. Eur Rev Med Pharmacol Sci. 18(7):992-1000, 2014.

INTRODUCTION: The concern about mobile phone effects is increasing as the number of users increasing too. Different studies have different results, so this topic is still open to discussion. Aim of this report was to investigate the effects of the mobile phones on the Bcl-2 gene and p53 proteins in rat brains. **MATERIALS AND METHODS:** In the study group of 10 rats; mobile phones that spread EMW at a frequency between 1900-2100 MHz and Specific Absorption Rate range between 0.005 W/kg and 0.288 W/kg (Dialing mode), 0.004 W/kg and 0.029 W/kg (Calling mode) were attached to rat ears for simulating usage in daily life for 7 times a day during 5 minutes (3 seconds dialing mode, 4 minutes and 47 seconds of calling mode) for a four week period. Sham group (n=10) rats were only immobilized without EMW exposure. Another group of rats (n=10) were counted as control without any application. immunohistopathological examination was performed for p53 and Bcl-2 expression. **RESULTS:** Immunohistopathological examinations revealed that the samples in the study group had more p53 and Bcl-2 positive stained cells and they were stained denser. In both evaluations, these differences between the study and control group were found statistically significant ($p < 0.003$); In Bcl-2 evaluation statistically significant difference was found between study and sham group to ($p < 0.005$); however, the p53 evaluation between the study and the sham group did not show any statistically significant difference ($p > 0.005$).

CONCLUSIONS: Our results showed that the electro-magnetic waves emitted by the mobile phones may have effect on apoptosis. Besides, obtained data revealed that more realistic application of mobile phones during experiments is more important as expected.

Del Vecchio G, Giuliani A, Fernandez M, Mesirca P, Bersani F, Pinto R, Ardoino L, Lovisolo GA, Giardino L, Calzà L. Continuous exposure to 900MHz GSM-modulated EMF alters morphological maturation of neural cells. Neurosci Lett. 455(3):173-177, 2009.

The effects of radiofrequency electromagnetic field (RF-EMF) exposure on neuronal phenotype maturation have been studied in two different in vitro models: murine SN56 cholinergic cell line and rat primary cortical neurons. The samples were exposed at a dose of 1W/kg at 900 MHz GSM modulated. The phenotype analysis was carried out at 48 and 72 h (24 and 48 h of SN56 cell line differentiation) or at 24, 72, 120 h (2, 4 and 6 days in vitro for cortical neurons) of exposure, on live and immunolabeled neurons, and

Studies that show **Cell Phone** Health Effects

included the morphological study of neurite emission, outgrowth and branching. Moreover, cortical neurons were studied to detect alterations in the expression pattern of cytoskeleton regulating factors, e.g. beta-thymosin, and of early genes, e.g. c-Fos and c-Jun through real-time PCR on mRNA extracted after 24h exposure to EMF. We found that RF-EMF exposure reduced the number of neurites generated by both cell systems, and this alteration correlates to increased expression of beta-thymosin mRNA.

Del Vecchio G, Giuliani A, Fernandez M, Mesirca P, Bersani F, Pinto R, Ardoino L, Lovisolo GA, Giardino L, Calzà L. Effect of radiofrequency electromagnetic field exposure on in vitro models of neurodegenerative disease. Bioelectromagnetics. 30(7):564-572, 2009.

In this work we tested viability, proliferation, and vulnerability of neural cells, after continuous radiofrequency (RF) electromagnetic fields exposure (global system for mobile telecommunications (GSM) modulated 900 MHz signal at a specific absorption rate (SAR) of 1 W/kg and maximum duration 144 h) generated by transverse electromagnetic cells. We used two cellular systems, SN56 cholinergic for example, SN56 cholinergic cell line and rat primary cortical neurons, and well-known neurotoxic challenges, such as glutamate, 25-35AA beta-amyloid, and hydrogen peroxide. Exposure to RF did not change viability/proliferation rate of the SN56 cholinergic cells or viability of cortical neurons. Co-exposure to RF exacerbated neurotoxic effect of hydrogen peroxide in SN56, but not in primary cortical neurons, whereas no cooperative effects of RF with glutamate and 25-35AA beta-amyloid were found. These data suggest that only under particular circumstances exposure to GSM modulated, 900 MHz signal act as a co-stressor for oxidative damage of neural cells.

Curcio G, Mazzucchi E, Marca GD, Vollono C, Rossini PM. Electromagnetic fields and EEG spiking rate in patients with focal epilepsy. Clin Neurophysiol. 2014 Aug 11. pii: S1388-2457(14)00404-0. doi: 10.1016/j.clinph.2014.07.013. [Epub ahead of print]

OBJECTIVE: Despite the increase in mobile telephone technology use and possible effects on brain excitability, no studies have investigated the impact of GSM like (Global System for Mobile Communications) signal on the ongoing spiking activity in human epileptic patients. METHODS: Brain electrical (electroencephalogram, EEG) activity of 12 patients with focal epilepsy has been recorded under both Real and Sham exposure following a double-blind, crossover, counterbalanced design: before the exposure (pre-exposure/baseline session), during the Real or Sham 45min exposure (during-exposure session), and after the exposure (post-exposure session). As dependent variables both spiking activity (spikes count) and EEG quantitative indices (spectral power and coherence data) have been considered. RESULTS: Spiking activity tended to be lower under Real than under Sham exposure. EEG spectral content analysis indicated a significant increase of Gamma band under Real exposure, mainly evident in Parieto-occipital and Temporal areas. Connectivity data indicated increased interhemispheric (left temporal to right frontal Regions of Interest, ROIs) instantaneous coherence, in the Beta frequency band during-exposure with respect to baseline session. No significant modification of lagged coherence was observed. CONCLUSIONS: Acute GSM exposure in

Studies that show **Cell Phone** Health Effects

epileptic patients slightly influences their EEG properties, without reaching any clinical relevance. SIGNIFICANCE: No signs were found of an increased risk of incoming seizures for these patients as a consequence of using mobile phones.

Curcio G, Ferrara M, Limongi T, Tempesta D, Di Sante G, De Gennaro L, Quaresima V, Ferrari M. Acute mobile phones exposure affects frontal cortex hemodynamics as evidenced by functional near-infrared spectroscopy. *J Cereb Blood Flow Metab.*29(5):903-910, 2009.

This study aimed to evaluate by functional near-infrared spectroscopy (fNIRS), the effects induced by an acute exposure (40 mins) to a GSM (Global System for Mobile Communications) signal emitted by a mobile phone (MP) on the oxygenation of the frontal cortex. Eleven healthy volunteers underwent two sessions (Real and Sham exposure) after a crossover, randomized, double-blind paradigm. The whole procedure lasted 60 mins: 10-mins baseline (Bsl), 40-mins (Exposure), and 10-mins recovery (Post-Exp). Together with frontal hemodynamics, heart rate, objective and subjective vigilance, and self-evaluation of subjective symptoms were also assessed. The fNIRS results showed a slight influence of the GSM signal on frontal cortex, with a linear increase in [HHb] as a function of time in the Real exposure condition ($F(4,40)=2.67$; $P=0.04$). No other measure showed any GSM exposure-dependent changes. These results suggest that fNIRS is a convenient tool for safely and noninvasively investigating the cortical activation in MP exposure experimental settings. Given the short-term effects observed in this study, the results should be confirmed on a larger sample size and using a multichannel instrument that allows the investigation of a wider portion of the frontal cortex.

Curcio G, Valentini E, Moroni F, Ferrara M, De Gennaro L, Bertini M Psychomotor performance is not influenced by brief repeated exposures to mobile phones. *Bioelectromagnetics.*29(3):237-241,2008.

The present study investigated the presence of a cumulative effect of brief and repeated exposures to a GSM mobile phone (902.40 MHz, 217 Hz modulated; peak power of 2 W; average power of 0.25 W; SAR = 0.5 W/kg) on psychomotor functions. To this end, after each of 3 15-min exposures, both an acoustic simple reaction time task (SRTT) and a sequential finger tapping task (SFTT) were administered to 24 subjects. The present study was unable to detect the cumulative effects of brief and repeated EMF exposure on human psychomotor performance, although there was a non-statistical trend to shorter reaction times. In summary, these data show an absence of effects with these particular exposure conditions; however, possible cognitive effects induced by different signal characteristics cannot be excluded.

Bolshakov MA, Alekseev SI, Bursting responses of Lymnea neurons to microwave radiation. *Bioelectromagnetics* 13(2):119-129, 1992.

Microelectrode and voltage-clamp techniques were modified to record spontaneous electrical activity and ionic currents of Lymnea stagnalis neurons during exposure to a 900-MHz field in a waveguide-based apparatus. The field was pulse-modulated at

Studies that show **Cell Phone** Health Effects

repetition rates ranging from 0.5 to 110 pps, or it was applied as a continuous wave (CW). When subjected to pulsed waves (PW), rapid, burst-like changes in the firing rate of neurons occurred at SARs of a few W/kg. If the burst-like irregularity was present in the firing rate under control conditions, irradiation enhanced its probability of occurrence. The effect was dependent on modulation, but not on modulation frequency, and it had a threshold SAR near 0.5 W/kg. CW radiation had no effect on the firing rate pattern at the same SAR. Mediator-induced, current activation of acetylcholine, dopamine, serotonin, or gamma-aminobutyric-acid receptors of the neuronal soma was not altered during CW or PW exposures and, hence, could not have been responsible for the bursting effect.

Cao Z, Liu J, Li S, Zhao X. [Effects of electromagnetic radiation from handsets of cellular telephone on neurobehavioral function] Wei Sheng Yan Jiu 29(2):102-103, 2000. [Article in Chinese]

In order to study the effects of electromagnetic radiation from handsets of cellular telephone on neurobehavioral function, 81 staff with handsets of cellular telephone and 63 staff without handsets of cellular telephone from corporations were selected as the subjects. The subjects were investigated by questionnaire on their general health, lifestyle habit, suppress of spirit, handset using of cellular telephone, environmental exposure, morbidity, and the neurobehavioral core test battery(NCTB). The data was analyzed by chi-square, stepwise regression analysis and covariance statistics. The results showed that the average reaction time in user's group was longer than that in control group ($P < 0.01$). The time of using handset was negatively associated with corrected reaction number ($P < 0.01$). The fast reaction time and the slowest reaction time were positively associated with the length of handset using ($P < 0.01$, $P < 0.05$). The results suggested that the handset using could cause adverse health effects in neurobehavioral function.

D'Costa H, Trueman G, Tang L, Abdel-rahman U, Abdel-rahman W, Ong K, Cosic I. Human brain wave activity during exposure to radiofrequency field emissions from mobile phones. Australas Phys Eng Sci Med. 26(4):162-167, 2003.

The aim of this study was to determine whether there is an effect of mobile phone electromagnetic field emissions on the human electroencephalograph (EEG). EEG recordings from ten awake subjects were taken during exposure to radiofrequency (RF) emissions from a mobile phone positioned behind the head. Two experimental trials were conducted. In the first trial, RF exposures were generated by a GSM mobile phone with the speaker disabled and configured to transmit at full-radiated power. During the second trial, exposures were generated by a non-modified GSM mobile phone in active standby mode. For each trial, subjects were exposed in five minute intervals to a randomized, interrupted sequence of five active and five sham exposures. The experiment was conducted under single-blind conditions. The average EEG band power in active exposure recordings was compared to corresponding sham recordings. Statistical tests indicated significant difference in the full-power mode trial within the EEG alpha (8-13 Hz) and beta (13-32 Hz) bands. A subsequent statistical analysis of

Studies that show **Cell Phone** Health Effects

median spectral power in discrete EEG rhythms revealed significant differences in 7 of the 32 distinct frequencies overall. In conclusion, the results of this study lend support to EEG effects from mobile phones activated in talk-mode.

Ammari M, Brillaud E, Gamez C, Lecomte A, Sakly M, Abdelmelek H, de Seze R. Effect of a chronic GSM 900MHz exposure on glia in the rat brain. *Biomed Pharmacother.* 62(4):273-281, 2008.

Extension of the mobile phone technology raises concern about the health effects of 900MHz microwaves on the central nervous system (CNS). In this study we measured GFAP expression using immunocytochemistry method, to evaluate glial evolution 10 days after a chronic exposure (5 days a week for 24 weeks) to GSM signal for 45min/day at a brain-averaged specific absorption rate (SAR)=1.5W/kg and for 15min/day at a SAR=6W/kg in the following rat brain areas: prefrontal cortex (PfCx), caudate putamen (Cpu), lateral globus pallidus of striatum (LGP), dentate gyrus of hippocampus (DG) and cerebellum cortex (CCx). In comparison to sham or cage control animals, rats exposed to chronic GSM signal at 6W/kg have increased GFAP stained surface areas in the brain ($p<0.05$). But the chronic exposure to GSM at 1.5W/kg did not increase GFAP expression. Our results indicated that chronic exposure to GSM 900MHz microwaves (SAR=6W/kg) may induce persistent astroglia activation in the rat brain (sign of a potential gliosis).

Ammari M, Lecomte A, Sakly M, Abdelmelek H, de-Seze R. Exposure to GSM 900 MHz electromagnetic fields affects cerebral cytochrome c oxidase activity. *Toxicology.* 250(1):70-74, 2008.

The world-wide and rapidly growing use of mobile phones has raised serious concerns about the biological and health-related effects of radio frequency (RF) radiation, particularly concerns about the effects of RFs upon the nervous system. The goal of this study was conducted to measure cytochrome oxidase (CO) levels using histochemical methods in order to evaluate regional brain metabolic activity in rat brain after exposure to a GSM 900 MHz signal for 45 min/day at a brain-averaged specific absorption rate (SAR) of 1.5 W/Kg or for 15 min/day at a SAR of 6 W/Kg over seven days. Compared to the sham and control cage groups, rats exposed to a GSM signal at 6 W/Kg showed decreased CO activity in some areas of the prefrontal and frontal cortex (infralimbic cortex, prelimbic cortex, primary motor cortex, secondary motor cortex, anterior cingulate cortex areas 1 and 2 (Cg1 and Cg2)), the septum (dorsal and ventral parts of the lateral septal nucleus), the hippocampus (dorsal field CA1, CA2 and CA3 of the hippocampus and dental gyrus) and the posterior cortex (retrosplenial agranular cortex, primary and secondary visual cortex, perirhinal cortex and lateral entorhinal cortex). However, the exposure to GSM at 1.5 W/Kg did not affect brain activity. Our results indicate that 6 W/Kg GSM 900 MHz microwaves may affect brain metabolism and neuronal activity in rats.

Ammari M, Gamez C, Lecomte A, Sakly M, Abdelmelek H, De Seze R. GFAP expression in the rat brain following sub-chronic exposure to a 900 MHz electromagnetic field

Studies that show **Cell Phone** Health Effects

signal. Int J Radiat Biol. 86(5):367-375, 2010.

PURPOSE: The rapid development and expansion of mobile communications contributes to the general debate on the effects of electromagnetic fields emitted by mobile phones on the nervous system. This study aims at measuring the glial fibrillary acidic protein (GFAP) expression in 48 rat brains to evaluate reactive astrogliosis, three and 10 days after long-term head-only sub-chronic exposure to a 900 MHz electromagnetic field (EMF) signal, in male rats. **METHODS:** Sprague-Dawley rats were exposed for 45 min/day at a brain-averaged specific absorption rate (SAR) = 1.5 W/kg or 15 min/day at a SAR = 6 W/kg for five days per week during an eight-week period. GFAP expression was measured by the immunocytochemistry method in the following rat brain areas: Prefrontal cortex, cerebellar cortex, dentate gyrus of the hippocampus, lateral globus pallidus of the striatum, and the caudate putamen. **RESULTS:** Compared to the sham-treated rats, those exposed to the sub-chronic GSM (Global System for mobile communications) signal at 1.5 or 6 W/kg showed an increase in GFAP levels in the different brain areas, three and ten days after treatment. **CONCLUSION:** Our results show that sub-chronic exposures to a 900 MHz EMF signal for two months could adversely affect rat brain (sign of a potential gliosis).

Ammari M, Gamez C, Lecomte A, Sakly M, Abdelmelek H, De Seze R. GFAP expression in the rat brain following sub-chronic exposure to a 900 MHz electromagnetic field signal. Int J Radiat Biol. 86(5):367-375, 2010.

PURPOSE: The rapid development and expansion of mobile communications contributes to the general debate on the effects of electromagnetic fields emitted by mobile phones on the nervous system. This study aims at measuring the glial fibrillary acidic protein (GFAP) expression in 48 rat brains to evaluate reactive astrogliosis, three and 10 days after long-term head-only sub-chronic exposure to a 900 MHz electromagnetic field (EMF) signal, in male rats. **METHODS:** Sprague-Dawley rats were exposed for 45 min/day at a brain-averaged specific absorption rate (SAR) = 1.5 W/kg or 15 min/day at a SAR = 6 W/kg for five days per week during an eight-week period. GFAP expression was measured by the immunocytochemistry method in the following rat brain areas: Prefrontal cortex, cerebellar cortex, dentate gyrus of the hippocampus, lateral globus pallidus of the striatum, and the caudate putamen. **RESULTS:** Compared to the sham-treated rats, those exposed to the sub-chronic GSM (Global System for mobile communications) signal at 1.5 or 6 W/kg showed an increase in GFAP levels in the different brain areas, three and ten days after treatment. **CONCLUSION:** Our results show that sub-chronic exposures to a 900 MHz EMF signal for two months could adversely affect rat brain (sign of a potential gliosis).

Barth A, Winker R, Ponocny-Seliger E, Mayrhofer W, Ponocny I, Sauter C, Vana N. A meta-analysis for neurobehavioral effects due to electromagnetic field exposure emitted by GSM mobile phones. Occup Environ Med. 65(5):342-6, 2008.

BACKGROUND AND OBJECTIVE: Numerous studies have investigated the potential effects of electromagnetic fields (EMFs) emitted by GSM mobile phones (~900 MHz to ~1800 MHz) on cognitive functioning, but results have been equivocal. In order to try

Studies that show **Cell Phone** Health Effects

and clarify this issue, the current study carried out a meta-analysis on nineteen experimental studies. DESIGN: meta-analysis METHODS: Nineteen studies were taken into consideration. Ten of them were included in the meta-analysis as they fulfil several minimum requirements; for example, single-blind or double-blind experimental study design and documentation of means and standard deviation of the dependent variables. The meta-analysis aimed at comparing exposed with non-exposed subjects assuming that there is a common population effect so that one single effect size could be calculated. When homogeneity for single effect sizes was not given, an own population effect for each study and a distribution of population effects was assumed. RESULTS: Attention measured by the subtraction task seems to be affected in regard of decreased reaction time. Working memory measured by the N-back test seems to be affected too: Under condition 0-back target response time is lower under exposure, while under condition 2-back target response time increases. The number of errors under condition 2-back non-targets appears to be higher under exposure. CONCLUSION: Results of the meta-analysis suggest that EMFs may have a small impact on human attention and working memory.

Papageorgiou CC, Nanou ED, Tsiafakis VG, Kapareliotis E, Kontoangelos KA, Capsalis CN, Rabavilas AD, Soldatos CR. Acute mobile phone effects on pre-attentive operation. *Neurosci Lett*397(1-2):99-103, 2006.

There is a debate whether electromagnetic field (EMF) emitted by mobile phones (MP) have an effect on cognitive functions. Since the auditory P50 component of event-related potentials (ERPs) reflects pre-attentive processing and working memory (WM) operation, the present study was designed to investigate whether the exposure to MP-EMF affects the patterns of the P50 component of ERPs elicited during a WM test. The P50 elicited during a WM task and evoked by two warning stimuli low and high frequency (500 and 3000Hz) has been assessed in 19 normal subjects (10 women and 9 men) both without and with exposure to a 900MHz signal, emitted by a dipole antenna placed near the subjects. Results showed that the presence of MP-EMFs induced statistically significant increase in the amplitude of P50 evoked by the low frequency stimuli, at Fp1 and O1 electrode leads as compared to themselves without MP-EMF exposure. In contrast the exposure to MP-EMFs revealed statistically significant decrease of the amplitude of P50 evoked by the high frequency stimuli, at Fp1 electrode lead as compared to themselves without MP-EMF exposure. These findings provide evidence that the MP-EMF emitted by mobile phone affect pre-attentive information processing as it is reflected in P50 evoked potential. The basis of such an effect is unclear, although several possibilities exist and call for potential directions of future research.

Papageorgiou CC, Nanou ED, Tsiafakis VG, Capsalis CN, Rabavilas AD. Gender related differences on the EEG during a simulated mobile phone signal. *Neuroreport*. 15(16):2557-2560, 2004.

The present study investigated the gender-related influence of electromagnetic fields (EMF), similar to that emitted by mobile phones, on brain activity. Ten women and nine

Studies that show **Cell Phone** Health Effects

men performed a short memory task (Wechsler test), both without (baseline) and with exposure to a 900 MHz signal. The EEG energy of the total waveform and the alpha, beta, delta and theta; rhythms were calculated from the recordings of 15 scalp electrodes. Baseline EEG energy of males was greater than that of females, while exposure to EMF decreased EEG energy of males and increased that of females. Memory performance was invariant to EMF exposure and gender influences. These findings indicate that EMF may exert a gender-related influence on brain activity.

Effects on Brain

Aalto S, Haarala C, Bruck A, Sipila H, Hamalainen H, Rinne JO. Mobile phone affects cerebral blood flow in humans. *J Cereb Blood Flow Metab.* 26(7):885-890, 2006.

Mobile phones create a radio-frequency electromagnetic field (EMF) around them when in use, the effects of which on brain physiology in humans are not well known. We studied the effects of a commercial mobile phone on regional cerebral blood flow (rCBF) in healthy humans using positron emission tomography (PET) imaging. Positron emission tomography data was acquired using a double-blind, counterbalanced study design with 12 male subjects performing a computer-controlled verbal working memory task (letter 1-back). Explorative and objective voxel-based statistical analysis revealed that a mobile phone in operation induces a local decrease in rCBF beneath the antenna in the inferior temporal cortex and an increase more distantly in the prefrontal cortex. Our results provide the first evidence, suggesting that the EMF emitted by a commercial mobile phone affects rCBF in humans. These results are consistent with the postulation that EMF induces changes in neuronal activity.

Megha K, Deshmukh PS, Banerjee BD, Tripathi AK, Abegaonkar MP. Microwave radiation induced oxidative stress, cognitive impairment and inflammation in brain of Fischer rats. *Indian J Exp Biol.* 50(12):889-896, 2012.

Public concerns over possible adverse effects of microwave radiation emitted by mobile phones on health are increasing. To evaluate the intensity of oxidative stress, cognitive impairment and inflammation in brain of Fischer rats exposed to microwave radiation, male Fischer-344 rats were exposed to 900 MHz microwave radiation ($SAR = 5.953 \times 10^{-4}$ W/kg) and 1800 MHz microwave radiation ($SAR = 5.835 \times 10^{-4}$ W/kg) for 30 days (2 h/day). Significant impairment in cognitive function and induction of oxidative stress in brain tissues of microwave exposed rats were observed in comparison with sham exposed groups. Further, significant increase in level of cytokines (IL-6 and TNF-alpha) was also observed following microwave exposure. Results of the present study indicated that increased oxidative stress due to microwave exposure may contribute to cognitive impairment and inflammation in brain.

Maskey D, Kim M, Aryal B, Pradhan J, Choi IY, Park KS, Son T, Hong SY, Kim SB, Kim HG, Kim MJ. Effect of 835 MHz radiofrequency radiation exposure on calcium binding

Studies that show **Cell Phone** Health Effects

proteins in the hippocampus of the mouse brain. Brain Res.1313:232-241, 2010.

Worldwide expansion of mobile phones and electromagnetic field (EMF) exposure has raised question of their possible biological effects on the brain and nervous system. Radiofrequency (RF) radiation might alter intracellular signaling pathways through changes in calcium (Ca(2+)) permeability across cell membranes. Changes in the expression of calcium binding proteins (CaBP) like calbindin D28-k (CB) and calretinin (CR) could indicate impaired Ca(2+)homeostasis due to EMF exposure. CB and CR expression were measured with immunohistochemistry in the hippocampus of mice after EMF exposure at 835 MHz for different exposure times and absorption rates, 1 h/day for 5 days at a specific absorption rate (SAR)=1.6 W/kg, 1 h/day for 5 days at SAR=4.0 W/kg, 5 h/day for 1 day at SAR=1.6 W/kg, 5 h/day for 1 day at SAR=4.0 W/kg, daily exposure for 1 month at SAR=1.6 W/kg. Body weights did not change significantly. CB immunoreactivity (IR) displayed moderate staining of cells in the cornu ammonis (CA) areas and prominently stained granule cells. CR IR revealed prominently stained pyramidal cells with dendrites running perpendicularly in the CA area. Exposure for 1 month produced almost complete loss of pyramidal cells in the CA1 area. CaBP differences could cause changes in cellular Ca(2+)levels, which could have deleterious effect on normal hippocampal functions concerned with neuronal connectivity and integration.

Tong J, Chen S, Liu XM, Hao DM. [Effect of electromagnetic radiation on discharge activity of neurons in the hippocampus CA1 in rats]. Zhongguo Ying Yong Sheng Li Xue Za Zhi. 29(5):423-427, 2013. [Article in Chinese]

OBJECTIVE: In order to explore effect of electromagnetic radiation on learning and memory ability of hippocampus neuron in rats, the changes in discharge patterns and overall electrical activity of hippocampus neuron after electromagnetic radiation were observed. METHODS: Rat neurons discharge was recorded with glass electrode extracellular recording technology and a polygraph respectively. Radiation frequency of electromagnetic wave was 900 MHz and the power was 10 W/m². In glass electrode extracellular recording, the rats were separately irradiated for 10, 20, 30, 40, 50 and 60 min, every points repeated 10 times and updated interval of 1h, observing the changes in neuron discharge and spontaneous discharge patterns after electromagnetic radiation. In polygraph recording experiments, irradiation group rats for five days a week, 6 hours per day, repeatedly for 10 weeks, memory electrical changes in control group and irradiation group rats when they were feeding were repeatedly monitored by the implanted electrodes, observing the changes in peak electric digits and the largest amplitude in hippocampal CA1 area, and taking some electromagnetic radiation sampling sequence for correlation analysis. RESULTS: (1) Electromagnetic radiation had an inhibitory role on discharge frequency of the hippocampus CA1 region neurons. After electromagnetic radiation, discharge frequency of the hippocampus CA1 region neurons was reduced, but the changes in scale was not obvious. (2) Electromagnetic radiation might change the spontaneous discharge patterns of hippocampus CA1 region neurons, which made the explosive discharge pattern increased obviously. (3) Peak potential total number within 5 min in irradiation group was significantly reduced, the largest

Studies that show **Cell Phone** Health Effects

amplitude was less than that of control group. (4) Using mathematical method to make the correlation analysis of the electromagnetic radiation sampling sequence, that of irradiation group was less than that of control group, indicating that there was a tending to be inhibitory connection between neurons in irradiation group after electromagnetic radiation. CONCLUSION: Electromagnetic radiation may cause structure and function changes of transfer synaptic in global, make hippocampal CA1 area neurons change in the overall discharge characteristic and discharge patterns, thus lead to decrease in the ability of learning and memory.

Aboul Ezz HS, Khadrawy YA, Ahmed NA, Radwan NM, El Bakry MM. The effect of pulsed electromagnetic radiation from mobile phone on the levels of monoamine neurotransmitters in four different areas of rat brain. Eur Rev Med Pharmacol Sci. 17(13):1782-1788, 2013.

BACKGROUND: The use of mobile phones is rapidly increasing all over the world. Few studies deal with the effect of electromagnetic radiation (EMR) on monoamine neurotransmitters in the different brain areas of adult rat. **AIM:** The aim of the present study was to investigate the effect of EMR on the concentrations of dopamine (DA), norepinephrine (NE) and serotonin (5-HT) in the hippocampus, hypothalamus, midbrain and medulla oblongata of adult rats. **MATERIALS AND METHODS:** Adult rats were exposed daily to EMR (frequency 1800 MHz, specific absorption rate 0.843 W/kg, power density 0.02 mW/cm², modulated at 217 Hz) and sacrificed after 1, 2 and 4 months of daily EMR exposure as well as after stopping EMR for 1 month (after 4 months of daily EMR exposure). Monoamines were determined by high performance liquid chromatography coupled with fluorescence detection (HPLC-FD) using their native properties. **RESULTS:** The exposure to EMR resulted in significant changes in DA, NE and 5-HT in the four selected areas of adult rat brain. **CONCLUSIONS:** The exposure of adult rats to EMR may cause disturbances in monoamine neurotransmitters and this may underlie many of the adverse effects reported after EMR including memory, learning, and stress.

Ning W, Xu SJ, Chiang H, Xu ZP, Zhou SY, Yang W, Luo JH. Effects of GSM 1800 MHz on dendritic development of cultured hippocampal neurons. Acta Pharmacol Sin. 28(12):1873-1880, 2007.

AIM: To evaluate the effects of global system for mobile communications (GSM) 1800 MHz microwaves on dendritic filopodia, dendritic arborization, and spine maturation during development in cultured hippocampal neurons in rats. **METHODS:** The cultured hippocampal neurons were exposed to GSM 1800 MHz microwaves with 2.4 and 0.8 W/kg, respectively, for 15 min each day from 6 days in vitro (DIV6) to DIV14. The subtle structures of dendrites were displayed by transfection with farnesylated enhanced green fluorescent protein (F-GFP) and GFP-actin on DIV5 into the hippocampal neurons. **RESULTS:** There was a significant decrease in the density and mobility of dendritic filopodia at DIV8 and in the density of mature spines at DIV14 in the neurons exposed to GSM 1800 MHz microwaves with 2.4 W/kg. In addition, the average length of dendrites per neuron at DIV10 and DIV14 was decreased, while the dendritic

Studies that show **Cell Phone** Health Effects

arborization was unaltered in these neurons. However, there were no significant changes found in the neurons exposed to the GSM 1800 MHz microwaves with 0.8 W/kg. **CONCLUSION:** These data indicate that the chronic exposure to 2.4 W/kg GSM 1800 MHz microwaves during the early developmental stage may affect dendritic development and the formation of excitatory synapses of hippocampal neurons in culture.

Eser O, Songur A, Aktas C, Karavelioglu E, Caglar V, Aylak F, Ozguner F, Kanter M. The effect of electromagnetic radiation on the rat brain: an experimental study. Turk Neurosurg. 23(6):707-715, 2013.

AIM: The aim of this study is to determine the structural changes of electromagnetic waves in the frontal cortex, brain stem and cerebellum. MATERIAL and METHODS: 24 Wistar Albino adult male rats were randomly divided into four groups: group I consisted of control rats, and groups II-IV comprised electromagnetically irradiated (EMR) with 900, 1800 and 2450 MHz. The heads of the rats were exposed to 900, 1800 and 2450 MHz microwaves irradiation for 1h per day for 2 months. RESULTS: While the histopathological changes in the frontal cortex and brain stem were normal in the control group, there were severe degenerative changes, shrunken cytoplasm and extensively dark pyknotic nuclei in the EMR groups. Biochemical analysis demonstrated that the Total Antioxidative Capacity level was significantly decreased in the EMR groups and also Total Oxidative Capacity and Oxidative Stress Index levels were significantly increased in the frontal cortex, brain stem and cerebellum. IL-1 β level was significantly increased in the EMR groups in the brain stem. **CONCLUSION:** EMR causes to structural changes in the frontal cortex, brain stem and cerebellum and impair the oxidative stress and inflammatory cytokine system. This deterioration can cause to disease including loss of these areas function and cancer development.

Motawi TK, Darwish HA, Moustafa YM, Labib MM. Biochemical Modifications and Neuronal Damage in Brain of Young and Adult Rats After Long-Term Exposure to Mobile Phone Radiations. Cell Biochem Biophys. 2014 May 7. [Epub ahead of print]

This study investigated the effect of exposure to mobile phone radiations on oxidative stress and apoptosis in brain of rats. Rats were allocated into six groups (three young and three adult). Groups 1 and 4 were not subjected to the radiation source and served as control groups. In groups 2 and 5, the mobile phones were only connected to the global system for mobile communication, while in groups 3 and 6, the option of calling was in use. Microwaves were generated by a mobile test phone (SAR = 1.13 W/kg) during 60 days (2 h/day). Significant increments in conjugated dienes, protein carbonyls, total oxidant status, and oxidative stress index along with a significant reduction of total antioxidant capacity levels were evident after exposure. Bax/Bcl-2 ratio, caspase-3 activity, and tumor necrosis factor- α level were enhanced, whereas no DNA fragmentation was detected. The relative brain weight of young rats was greatly affected, and histopathological examination reinforced the neuronal damage. The study highlights the detrimental effects of mobile phone radiations on brain during young and adult ages. The interaction of these radiations with brain is via dissipating its antioxidant status and/or triggering apoptotic cell death.

Studies that show **Cell Phone** Health Effects

Lv B, Su C, Yang L, Xie Y, Wu T. Whole brain EEG synchronization likelihood modulated by long term evolution electromagnetic fields exposure. Conf Proc IEEE Eng Med Biol Soc. 2014:986-989, 2014.

In this paper, we aimed to investigate the possible interactions between human brain and radiofrequency electromagnetic fields (EMF) with electroencephalogram (EEG) technique. Unlike the previous studies which mainly focused on EMF effect on local brain activities, we attempted to evaluate whether the EMF emitted from Long Term Evolution (LTE) devices can modulate the functional connectivity of brain electrical activities. Ten subjects were recruited to participate in a crossover, double-blind exposure experiment which included two sessions (real and sham exposure). In each session, LTE EMF exposure (power on or off) lasted for 30 min and the EEG signals were collected with 32 channels throughout the experiment. Then we applied the synchronization likelihood method to quantify the neural synchronization over the whole brain in different frequency bands and in different EEG record periods. Our results illustrated that the short-term LTE EMF exposure would modulate the synchronization patterns of EEG activation across the whole brain.

Lv B, Chen Z, Wu T, Shao Q, Yan D, Ma L, Lu K, Xie Y. The alteration of spontaneous low frequency oscillations caused by acute electromagnetic fields exposure. Clin Neurophysiol. 2013 Sep 4. pii: S1388-2457(13)00976-0. doi: 10.1016/j.clinph.2013.07.018. [Epub ahead of print]

OBJECTIVE: The motivation of this study is to evaluate the possible alteration of regional resting state brain activity induced by the acute radiofrequency electromagnetic field (RF-EMF) exposure (30min) of Long Term Evolution (LTE) signal. **METHODS:** We designed a controllable near-field LTE RF-EMF exposure environment. Eighteen subjects participated in a double-blind, crossover, randomized and counterbalanced experiment including two sessions (real and sham exposure). The radiation source was close to the right ear. Then the resting state fMRI signals of human brain were collected before and after the exposure in both sessions. We measured the amplitude of low frequency fluctuation (ALFF) and fractional ALFF (fALFF) to characterize the spontaneous brain activity. **RESULTS:** We found the decreased ALFF value around in left superior temporal gyrus, left middle temporal gyrus, right superior temporal gyrus, right medial frontal gyrus and right paracentral lobule after the real exposure. And the decreased fALFF value was also detected in right medial frontal gyrus and right paracentral lobule. **CONCLUSIONS:** The study provided the evidences that 30min LTE RF-EMF exposure modulated the spontaneous low frequency fluctuations in some brain regions. **SIGNIFICANCE:** With resting state fMRI, we found the alteration of spontaneous low frequency fluctuations induced by the acute LTE RF-EMF exposure.

Salford LG, Brun AR, Eberhardt JL, Malmgren L, Persson BRR, Nerve cell damage in mammalian brain after exposure to microwaves from GSM mobile phones. Environ Health Persp 111:881-883, 2003.

The possible risks of radio-frequency electromagnetic fields for the human body is a growing concern for the society. We have earlier shown that weak pulsed microwaves

Studies that show **Cell Phone** Health Effects

give rise to a significant leakage of albumin through the blood-brain barrier (BBB). Now we have investigated whether a pathological leakage over the BBB might be combined with damage to the neurons. Three groups of each 8 rats were exposed for 2 hours to GSM mobile phone electromagnetic fields of different strengths. We found, and present here for the first time, highly significant ($p < 0.002$) evidence for neuronal damage in both the cortex, the hippocampus and the basal ganglia in the brains of exposed rats.

Noor NA, Mohammed HS, Ahmed NA, Radwan NM. Variations in amino acid neurotransmitters in some brain areas of adult and young male albino rats due to exposure to mobile phone radiation. Eur Rev Med Pharmacol Sci. 15(7):729-742, 2011.

BACKGROUND AND OBJECTIVES: Mobile phone radiation and health concerns have been raised, especially following the enormous increase in the use of wireless mobile telephony throughout the world. The present study aims to investigate the effect of one hour daily exposure to electromagnetic radiation (EMR) with frequency of 900 Mz (SAR 1.165 w/kg, power density 0.02 mW/cm²) on the levels of amino acid neurotransmitters in the midbrain, cerebellum and medulla of adult and young male albino rats.

MATERIALS AND METHODS: Adult and young rats were divided into two main groups (treated and control). The treated group of both adult and young rats was exposed to EMR for 1 hour daily. The other group of both adult and young animals was served as control. The determination of amino acid levels was carried out after 1 hour, 1 month, 2 months and 4 months of EMR exposure as well as after stopping radiation. **RESULTS:** Data of the present study showed a significant increase in both excitatory and inhibitory amino acids in the cerebellum of adult and young rats and midbrain of adult animals after 1 hour of EMR exposure. In the midbrain of adult animals, there was a significant increase in glycine level after 1 month followed by significant increase in GABA after 4 months. Young rats showed significant decreases in the midbrain excitatory amino acids. In the medulla, the equilibrium ratio percent (ER%) calculations showed a state of neurochemical inhibition after 4 months in case of adult animals, whereas in young animals, the neurochemical inhibitory state was observed after 1 month of exposure due to significant decrease in glutamate and aspartate levels. This state was converted to excitation after 4 months due to the increase in glutamate level. **CONCLUSION:** The present changes in amino acid concentrations may underlie the reported adverse effects of using mobile phones.

Deshmukh PS, Megha K, Banerjee BD, Ahmed RS, Chandna S, Abegaonkar MP, Tripathi AK. Detection of Low Level Microwave Radiation Induced Deoxyribonucleic Acid Damage Vis-à-vis Genotoxicity in Brain of Fischer Rats. Toxicol Int. 20(1):19-24, 2013.

BACKGROUND: Non-ionizing radiofrequency radiation has been increasingly used in industry, commerce, medicine and especially in mobile phone technology and has become a matter of serious concern in present time. **OBJECTIVE:** The present study was designed to investigate the possible deoxyribonucleic acid (DNA) damaging effects of low-level microwave radiation in brain of Fischer rats. **MATERIALS AND METHODS:** Experiments were performed on male Fischer rats exposed to microwave radiation for 30 days at three different frequencies: 900, 1800 and 2450 MHz. Animals were divided

Studies that show **Cell Phone** Health Effects

into 4 groups: Group I (Sham exposed): Animals not exposed to microwave radiation but kept under same conditions as that of other groups, Group II: Animals exposed to microwave radiation at frequency 900 MHz at specific absorption rate (SAR) 5.953×10^{-4} W/kg, Group III: Animals exposed to 1800 MHz at SAR 5.835×10^{-4} W/kg and Group IV: Animals exposed to 2450 MHz at SAR 6.672×10^{-4} W/kg. At the end of the exposure period animals were sacrificed immediately and DNA damage in brain tissue was assessed using alkaline comet assay. RESULTS: In the present study, we demonstrated DNA damaging effects of low level microwave radiation in brain. CONCLUSION: We concluded that low SAR microwave radiation exposure at these frequencies may induce DNA strand breaks in brain tissue.

Maby E, Le Bouquin Jeannes R, Faucon G. Short-term effects of GSM mobiles phones on spectral components of the human electroencephalogram. Conf Proc IEEE Eng Med Biol Soc. 1:3751-3754, 2006.

The aim of the study was to investigate whether the GSM (global system for mobile) signals affect the electrical activity of the human brain. Nine healthy subjects and six temporal epileptic patients were exposed to radiofrequencies emitted by a GSM mobile phone signals. Electroencephalographic (EEG) signals were recorded using surface electrodes with and without radiofrequency. In order to obtain a reference, a control session was also carried out. The spectral attributes of the EEG signals recorded by surface electrodes were analyzed. The significant decrease of spectral correlation coefficients under radiofrequency influence showed that the GSM signal altered the spectral arrangement of the EEG activity for healthy subjects as well as epileptic patients. For the healthy subjects, the EEG spectral energy decreased on the studied frequency band [0-40 Hz] and more precisely on occipital electrodes for the alpha-band. For the epileptic patients, these modifications were demonstrated by an increase of the power spectral density of the EEG signal. Nevertheless, these biological effects on the EEG are not sufficient to put forward some electrophysiological hypothesis.

Saikhedkar N, Bhatnagar M, Jain A, Sukhwai P, Sharma C, Jaiswal N. Effects of mobile phone radiation (900 MHz radiofrequency) on structure and functions of rat brain. Neurol Res. 2014 May 26:1743132814Y0000000392. [Epub ahead of print]

Objectives: The goals of this study were: (1) to obtain basic information about the effects of long-term use of mobile phone on cytological makeup of the hippocampus in rat brain (2) to evaluate the effects on antioxidant status, and (3) to evaluate the effects on cognitive behavior particularly on learning and memory. Methods: Rats (age 30 days, 120 ± 5 g) were exposed to 900 MHz radio waves by means of a mobile hand set for 4 hours per day for 15 days. Effects on anxiety, spatial learning, and memory were studied using open field test, elevated plus maze, Morris water maze (MWM), and classic maze test. Effects on brain antioxidant status were also studied. Cresyl violet staining was done to access the neuronal damage. Result: A significant change in behavior, i.e., more anxiety and poor learning was shown by test animals as compared to controls and sham group. A significant change in level of antioxidant enzymes and non-enzymatic antioxidants, and increase in lipid peroxidation were observed in test rats. Histological

Studies that show **Cell Phone** Health Effects

examination showed neurodegenerative cells in hippocampal sub regions and cerebral cortex. Discussion: Thus our findings indicate extensive neurodegeneration on exposure to radio waves. Increased production of reactive oxygen species due to exhaustion of enzymatic and non-enzymatic antioxidants and increased lipid peroxidation are indicating extensive neurodegeneration in selective areas of CA1, CA3, DG, and cerebral cortex. This extensive neuronal damage results in alterations in behavior related to memory and learning.

Sirav B, Seyhan N. Effects of radiofrequency radiation exposure on blood-brain barrier permeability in male and female rats. *Electromagn Biol Med.* 30(4):253-260, 2011.

During the last several decades, numerous studies have been performed aiming at the question of whether or not exposure to radiofrequency radiation (RFR) influences the permeability of the blood-brain barrier (BBB). The objective of this study was to investigate the effect of RFR on the permeability of BBB in male and female Wistar albino rats. Right brain, left brain, cerebellum, and total brain were analyzed separately in the study. Rats were exposed to 0.9 and 1.8 GHz continuous-wave (CW) RFR for 20 min (at SARs of 4.26 mW/kg and 1.46 mW/kg, respectively) while under anesthesia. Control rats were sham-exposed. Disruption of BBB integrity was detected spectrophotometrically using the Evans-blue dye, which has been used as a BBB tracer and is known to be bound to serum albumin. Right brain, left brain, cerebellum, and total brain were evaluated for BBB permeability. In female rats, no albumin extravasation was found in the brain after RFR exposure. A significant increase in albumin was found in the brains of the RF-exposed male rats when compared to sham-exposed male brains. These results suggest that exposure to 0.9 and 1.8 GHz CW RFR at levels below the international limits can affect the vascular permeability in the brain of male rats. The possible risk of RFR exposure in humans is a major concern for the society. Thus, this topic should be investigated more thoroughly in the future.

Soderqvist F, Carlberg M, Hardell L. Mobile and cordless telephones, serum transthyretin and the blood-cerebrospinal fluid barrier: a cross-sectional study. *Environ Health.* 8(1):19, 2009.

ABSTRACT: BACKGROUND: Whether low-intensity radiofrequency radiation damages the blood-brain barrier has long been debated, but little or no consideration has been given to the blood-cerebrospinal fluid barrier. In this cross-sectional study we tested whether long-term and/or short-term use of wireless telephones was associated **with changes in the serum transthyretin level, indicating altered transthyretin concentration in the cerebrospinal fluid, possibly reflecting an effect of radiation.**

METHODS: One thousand subjects, 500 of each sex aged 18-65 years, were randomly recruited using the population registry. Data on wireless telephone use were assessed by a postal questionnaire and blood samples were analyzed for serum transthyretin concentrations determined by standard immunonephelometric techniques on a BN Prospec(R) instrument. RESULTS: The response rate was 31.4%. Logistic regression of dichotomized TTR serum levels with a cut-point of 0.31 g/l on wireless telephone use yielded increased odds ratios that were statistically not significant. Linear regression of

Studies that show **Cell Phone** Health Effects

time since first use overall and on the day that blood was withdrawn gave different results for males and females: for men significantly higher serum concentrations of TTR were seen the longer an analogue telephone or a mobile and cordless desktop telephone combined had been used, and in contrast, significantly lower serum levels were seen the longer an UMTS telephone had been used. Adjustment for fractions of use of the different telephone types did not modify the effect for cumulative use or years since first use for mobile telephone and DECT, combined. For women, linear regression gave a significant association for short-term use of mobile and cordless telephones combined, indicating that the sooner blood was withdrawn after the most recent telephone call, the higher the expected transthyretin concentration.

CONCLUSIONS: In this hypothesis-generating descriptive study time since first use of mobile telephones and DECT combined was significantly associated with higher TTR levels regardless of how much each telephone type had been used. Regarding short-term use, significantly higher TTR concentrations were seen in women the sooner blood was withdrawn after the most recent telephone call on that day.

Lopez-Martin E, Relova-Quinteiro JL, Gallego-Gomez R, Peleteiro-Fernandez M, Jorge-Barreiro FJ, Ares-Pena FJ. GSM radiation triggers seizures and increases cerebral c-Fos positivity in rats pretreated with subconvulsive doses of picrotoxin. *Neurosci Lett.*398(1-2):139-144,2006.

This study investigated the effects of mobile-phone-type radiation on the cerebral activity of seizure-prone animals. When rats transformed into an experimental model of seizure-proneness by acute subconvulsive doses of picrotoxin were exposed to 2h GSM-modulated 900MHz radiation at an intensity similar to that emitted by mobile phones, they suffered seizures and the levels of the neuronal activity marker c-Fos in neocortex, paleocortex, hippocampus and thalamus increased markedly. Non-irradiated picrotoxin-treated rats did not suffer seizures, and their cerebral c-Fos counts were significantly lower. Radiation caused no such differences in rats that had not been pretreated with picrotoxin. We conclude that GSM-type radiation can induce seizures in rats following their facilitation by subconvulsive doses of picrotoxin, and that research should be pursued into the possibility that this kind of radiation may similarly affect brain function in human subjects with epileptic disorders.

Kwon MS, Vorobyev V, Kännälä S, Laine M, Rinne JO, Toivonen T, Johansson J, Teräs M, Lindholm H, Alanko T, Hämäläinen H. GSM mobile phone radiation suppresses brain glucosemetabolism. *J Cereb Blood Flow Metab.* 31(12):2293-2301, 2011.

Studies that show **Cell Phone** Health Effects

We investigated the effects of mobile phone radiation on cerebral glucose metabolism using high-resolution positron emission tomography (PET) with the (18)F-deoxyglucose (FDG) tracer. A long half-life (109 minutes) of the (18)F isotope allowed a long, natural exposure condition outside the PET scanner. Thirteen young right-handed male subjects were exposed to a pulse-modulated 902.4 MHz Global System for Mobile Communications signal for 33 minutes, while performing a simple visual vigilance task. Temperature was also measured in the head region (forehead, eyes, cheeks, ear canals) during exposure. (18)F-deoxyglucose PET images acquired after the exposure showed that relative cerebral metabolic rate of glucose was significantly reduced in the temporoparietal junction and anterior temporal lobe of the right hemisphere ipsilateral to the exposure. Temperature rise was also observed on the exposed side of the head, but the magnitude was very small. The exposure did not affect task performance (reaction time, error rate). Our results show that short-term mobile phone exposure can locally suppress brain energy metabolism in humans.

Nittby H, Brun A, Eberhardt J, Malmgren L, Persson BR, Salford LG. Increased blood-brain barrier permeability in mammalian brain 7 days after exposure to the radiation from a GSM-900 mobile phone. *Pathophysiology*. 16(2-3):103-112, 2009.

Microwaves were for the first time produced by humans in 1886 when radio waves were broadcasted and received. Until then microwaves had only existed as a part of the cosmic background radiation since the birth of universe. By the following utilization of microwaves in telegraph communication, radars, television and above all, in the modern mobile phone technology, mankind is today exposed to microwaves at a level up to 10(20) times the original background radiation since the birth of universe. Our group has earlier shown that the electromagnetic radiation emitted by mobile phones alters the permeability of the blood-brain barrier (BBB), resulting in albumin extravasation immediately and 14 days after 2h of exposure. In the background section of this report, we present a thorough review of the literature on the demonstrated effects (or lack of effects) of microwave exposure upon the BBB. Furthermore, we have continued our own studies by investigating the effects of GSM mobile phone radiation upon the blood-brain barrier permeability of rats 7 days after one occasion of 2h of exposure. Forty-eight rats were exposed in TEM-cells for 2h at non-thermal specific absorption rates (SARs) of 0mW/kg, 0.12mW/kg, 1.2mW/kg, 12mW/kg and 120mW/kg. Albumin extravasation over the BBB, neuronal albumin uptake and neuronal damage were assessed. Albumin extravasation was enhanced in the mobile phone exposed rats as compared to sham controls after this 7-day recovery period (Fisher's exact probability test, $p=0.04$ and Kruskal-Wallis, $p=0.012$), at the SAR-value of 12mW/kg (Mann-Whitney, $p=0.007$) and with a trend of increased albumin extravasation also at the SAR-values of 0.12mW/kg and 120mW/kg. There was a low, but significant correlation between the exposure level (SAR-value) and occurrence of focal albumin extravasation ($r(s)=0.33$; $p=0.04$). The present findings are in agreement with our earlier studies where we have seen increased BBB permeability immediately and 14 days after exposure. We here discuss the present findings as well as the previous results of altered BBB permeability from our and other laboratories.

Studies that show **Cell Phone** Health Effects

Volkow ND, Tomasi D, Wang GJ, Vaska P, Fowler JS, Telang F, Alexoff D, Logan J, Wong C. Effects of cell phone radiofrequency signal exposure on brain glucose metabolism. JAMA. 305(8):808-813, 2011.

CONTEXT: The dramatic increase in use of cellular telephones has generated concern about possible negative effects of radiofrequency signals delivered to the brain.

However, whether acute cell phone exposure affects the human brain is unclear.

OBJECTIVE: To evaluate if acute cell phone exposure affects brain glucose metabolism, a marker of brain activity.

DESIGN, SETTING, AND PARTICIPANTS:

Randomized crossover study conducted between January 1 and December 31, 2009, at a single US laboratory among 47 healthy participants recruited from the community. Cell phones were placed on the left and right ears and positron emission tomography with ((18)F)fluorodeoxyglucose injection was used to measure brain glucose metabolism twice, once with the right cell phone activated (sound muted) for 50 minutes ("on" condition) and once with both cell phones deactivated ("off" condition). Statistical parametric mapping was used to compare metabolism between on and off conditions using paired t tests, and Pearson linear correlations were used to verify the association of metabolism and estimated amplitude of radiofrequency-modulated electromagnetic waves emitted by the cell phone. Clusters with at least 1000 voxels (volume >8 cm³) and P < .05 (corrected for multiple comparisons) were considered significant.

MAIN OUTCOME MEASURE: Brain glucose metabolism computed as absolute metabolism ($\mu\text{mol}/100\text{ g per minute}$) and as normalized metabolism (region/whole brain).

RESULTS: Whole-brain metabolism did not differ between on and off conditions. In contrast, metabolism in the region closest to the antenna (orbitofrontal cortex and temporal pole) was significantly higher for on than off conditions (35.7 vs 33.3 $\mu\text{mol}/100\text{ g per minute}$; mean difference, 2.4 [95% confidence interval, 0.67-4.2]; P = .004). The increases were significantly correlated with the estimated electromagnetic field amplitudes both for absolute metabolism (R = 0.95, P < .001) and normalized metabolism (R = 0.89; P < .001). **CONCLUSIONS:** In healthy participants and compared with no exposure, 50-minute cell phone exposure was associated with increased brain glucose metabolism in the region closest to the antenna. This finding is of unknown clinical significance.

Ragy MM. Effect of exposure and withdrawal of 900-MHz-electromagnetic waves on brain, kidney and liver oxidative stress and some biochemical parameters in male rats. Electromagn Biol Med. 2014 Apr 8. [Epub ahead of print]

Increasing use of mobile phones in daily life with increasing adverse effects of electromagnetic radiation (EMR), emitted from mobile on some physiological processes, cause many concerns about their effects on human health. Therefore, this work was designed to study the effects of exposure to mobile phone emits 900-MHz EMR on the brain, liver and kidney of male albino rats. Thirty male adult rats were randomly divided into four groups (10 each) as follows: control group (rats without exposure to EMR), exposure group (exposed to 900-MHz EMR for 1 h/d for 60 d) and withdrawal group (exposed to 900-MHz electromagnetic wave for 1 h/d for 60 d then left for 30 d without exposure). EMR emitted from mobile phone led to a significant increase in

Studies that show **Cell Phone Health Effects**

malondialdehyde (MDA) levels and significant decrease total antioxidant capacity (TAC) levels in brain, liver and kidneys tissues. The sera activity of alanine transaminase (ALT), aspartate aminotransferase (AST), urea, creatinine and corticosterone were significantly increased ($p < 0.05$), while serum catecholamines were insignificantly higher in the exposed rats. These alterations were corrected by withdrawal. In conclusion, electromagnetic field emitting from mobile phone might produce impairments in some biochemicals changes and oxidative stress in brain, liver and renal tissue of albino rats.

Dasdag S, Akdag MZ, Kizil G, Kizil M, Cakir DU, Yokus B. Effect of 900 MHz radio frequency radiation on beta amyloid protein, protein carbonyl, and malondialdehyde in the brain. *Electromagn Biol Med.* 31(1):67-74, 2012.

Recently, many studies have been carried out in relation to 900 MHz radiofrequency radiation (RF) emitted from a mobile phone on the brain. However, there is little data concerning possible mechanisms between long-term exposure of RF radiation and biomolecules in brain. Therefore, we aimed to investigate long-term effects of 900 MHz radiofrequency radiation on beta amyloid protein, protein carbonyl, and malondialdehyde in the rat brain. The study was carried out on 17 Wistar Albino adult male rats. The rat heads in a carousel were exposed to 900 MHz radiofrequency radiation emitted from a generator, simulating mobile phones. For the study group (n: 10), rats were exposed to the radiation 2 h per day (7 days a week) for 10 months. For the sham group (n: 7), rats were placed into the carousel and the same procedure was applied except that the generator was turned off. In this study, rats were euthanized after 10 months of exposure and their brains were removed. Beta amyloid protein, protein carbonyl, and malondialdehyde levels were found to be higher in the brain of rats exposed to 900 MHz radiofrequency radiation. However, only the increase of protein carbonyl in the brain of rats exposed to 900 MHz radiofrequency radiation was found to be statistically significant ($p < 0.001$). In conclusion, 900 MHz radiation emitted from mobile/cellular phones can be an agent to alter some biomolecules such as protein. However, further studies are necessary.

Spichtig S, Scholkmann F, Chin L, Lehmann H, Wolf M. Assessment of intermittent UMTS electromagnetic field effects on blood circulation in the human auditory region using a near-infrared system. *Bioelectromagnetics.* 33(1):40-54, 2012.

The aim of the present study was to assess the potential effects of intermittent Universal Mobile Telecommunications System electromagnetic fields (UMTS-EMF) on blood circulation in the human head (auditory region) using near-infrared spectroscopy (NIRS) on two different timescales: short-term (effects occurring within 80 s) and medium-term (effects occurring within 80 s to 30 min). For the first time, we measured potential immediate effects of UMTS-EMF in real-time without any interference during exposure. Three different exposures (sham, 0.18 W/kg, and 1.8 W/kg) were applied in a controlled, randomized, crossover, and double-blind paradigm on 16 healthy volunteers. In addition to oxy-, deoxy-, and total haemoglobin concentrations ([O(2) Hb], [HHb], and [tHb], respectively), the heart rate (HR), subjective well-being, tiredness, and counting speed were recorded. During exposure to 0.18 W/kg, we found a significant short-term

Studies that show **Cell Phone** Health Effects

increase in $\Delta[\text{O}(2) \text{ Hb}]$ and $\Delta[\text{tHb}]$, which is small ($\approx 17\%$) compared to a functional brain activation. A significant decrease in the medium-term response of $\Delta[\text{HHb}]$ at 0.18 and 1.8 W/kg exposures was detected, which is in the range of physiological fluctuations. The medium-term ΔHR was significantly higher ($+1.84 \text{ bpm}$) at 1.8 W/kg than for sham exposure. The other parameters showed no significant effects. Our results suggest that intermittent exposure to UMTS-EMF has small short- and medium-term effects on cerebral blood circulation and HR.

Persson BRR, Salford LG, Brun A, Blood-brain barrier permeability in rats exposed to electromagnetic fields used in wireless communication. *Wireless Network* 3:455-461, 1997.

Biological effects of radio frequency electromagnetic fields (EMF) on the blood-brain barrier (BBB) have been studied in Fischer 344 rats of both sexes. The rats were not anesthetised during the exposure. The brains were perfused with saline for 3-4 minutes, and thereafter perfusion fixed with 4% formaldehyde for 5-6 minutes. Whole coronal sections of the brains were dehydrated and embedded in paraffin and sectioned at 5 micrometers. Albumin and fibrinogen were demonstrated immunochemically and classified as normal versus pathological leakage. In the present investigation we exposed male and female Fischer 344 rats in a Transverse Electromagnetic Transmission line chamber to microwaves of 915 MHz as continuous wave (CW) and pulse-modulated with different pulse power and at various time intervals. The CW-pulse power varied from 0.001 W to 10 W and the exposure time from 2 min to 960 min. In each experiment we exposed 4-6 rats with 2-4 controls randomly placed in excited and non-excited TEM cells, respectively. We have in total investigated 630 exposed rats at various modulation frequencies and 372 controls. The frequency of pathological rats is significantly increased ($P < 0.0001$) from 62/372 (ratio 0.17 ± 0.02) for control rats to 244/630 (ratio: 0.39 ± 0.043) in all exposed rats. Grouping the exposed animals according to the level or specific absorption energy (J/kg) give significant difference in all levels above 1.5 J/kg. The exposure was 915 MHz microwaves either pulse modulated (PW) at 217 Hz with 0.57 ms pulse width, at 50 Hz with 6.6 ms pulse width or continuous wave (CW). The frequency of pathological rats (0.17) among controls in the various groups is not significantly different. The frequency of pathological rats was 170/480 (0.35 ± 0.03) among rats exposed to pulse modulated (PW) and 74/149 (0.50 ± 0.07) among rats exposed to continuous wave exposure (CW). These results are both highly significantly different to their corresponding controls ($p < 0.0001$) and the frequency of pathological rats after exposure to pulsed radiation (PW) is significantly less ($p < 0.002$) than after exposure to continuous wave radiation (CW).

Rağbetli MC, Aydinlioğlu A, Koyun N, Rağbetli C, Bektas S, Ozdemir S. The effect of mobile phone on the number of Purkinje cells: A stereological study. *Int J Radiat Biol.* 86(7):548-54, 2010

Purpose: The World Health Organisation proposed an investigation concerning the exposure of animals to radiofrequency fields because of the possible risk factor for health. At power frequencies there is evidence to associate both childhood leukaemia

Studies that show **Cell Phone** Health Effects

and brain tumours with magnetic field exposures. There is also evidence of the effect of mobile phone exposure on both cognitive functions and the cerebellum. Purkinje cells of the cerebellum are also sensitive to high dose microwave exposure in rats. The present study investigated the effect of exposure to mobile phone on the number of Purkinje and granule neurons in the developing cerebellum. Material and methods: Male and female Swiss albino mice were housed as control and mobile phone-exposed groups. Pregnant animals in the experimental group were exposed to Global System for Mobile Communication (GSM) mobile phone radiation at 890-915 MHz at 0.95 W/Kg specific absorption rate (SAR). The cerebella were processed by frozen microtome. The sections obtained were stained with Haematoxylin-eosin and cresyl violet. For cell counting by the optical fractionator method, a pilot study was firstly performed. Cerebellar areas were analysed by using Axiovision software running on a personal computer. The optical dissectors were systematically spaced at random, and focused to the widest profile of the neuron cell nucleus. Results: A significant decrease in the number of Purkinje cells and a tendency for granule cells to increase in cerebellum was found. Conclusion: Further studies in this area are needed due to the popular use of mobile telephones and relatively high exposure on developing brain.

Sonmez OF, Odaci E, Bas O, Kaplan S. Purkinje cell number decreases in the adult female rat cerebellum following exposure to 900 MHz electromagnetic field. Brain Res. 1356:95-101, 2010.

The biological effects of electromagnetic field (EMF) exposure from mobile phones have growing concern among scientists since there are some reports showing increased risk for human health, especially in the use of mobile phones for a long duration. In the presented study, the effects on the number of Purkinje cells in the cerebellum of 16-week (16 weeks) old female rats were investigated following exposure to 900 MHz EMF. Three groups of rats, a control group (CG), sham exposed group (SG) and an electromagnetic field exposed group (EMFG) were used in this study. While EMFG group rats were exposed to 900 MHz EMF (1h/day for 28 days) in an exposure tube, SG was placed in the exposure tube but not exposed to EMF (1h/day for 28 days). The specific energy absorption rate (SAR) varied between 0.016 (whole body) and 2 W/kg (locally in the head). The CG was not placed into the exposure tube nor was it exposed to EMF during the study period. At the end of the experiment, all of the female rats were sacrificed and the number of Purkinje cells was estimated using a stereological counting technique. Histopathological evaluations were also done on sections of the cerebellum. Results showed that the total number of Purkinje cells in the cerebellum of the EMFG was significantly lower than those of CG ($p < 0.004$) and SG ($p < 0.002$). In addition, there was no significant difference at the 0.05 level between the rats' body and brain weights in the EMFG and CG or SG. Therefore, it is suggested that long duration exposure to 900 MHz EMF leads to decreases of Purkinje cell numbers in the female rat cerebellum.

Tombini M, Pellegrino G, Pasqualetti P, Assenza G, Benvenga A, Fabrizio E, Rossini PM Mobile phone emissions modulate brain excitability in patients with focal epilepsy. *Brain Stimul.* 2012 Aug 9. [Epub ahead of print]

Studies that show **Cell Phone** Health Effects

BACKGROUND: Electromagnetic fields (EMFs) emitted by mobile phones had been shown to increase cortical excitability in healthy subjects following 45 min of continuous exposure on the ipsilateral hemisphere. **OBJECTIVE:** Using Transcranial Magnetic Stimulation (TMS), the current study assessed the effects of acute exposure to mobile phone EMFs on the cortical excitability in patients with focal epilepsy. **METHODS:** Ten patients with cryptogenic focal epilepsy originating outside the primary motor area (M1) were studied. Paired-pulse TMS were applied to the M1 of both the hemisphere ipsilateral (IH) and contralateral (CH) to the epileptic focus before and immediately after real/sham exposure to the GSM-EMFs (45 min). The TMS study was carried out in all subjects in three different experimental sessions (IH and CH exposure, sham), 1 week apart, according to a crossover, double-blind and counter-balanced paradigm. **RESULTS:** The present study clearly demonstrated that an acute and relatively prolonged exposure to GSM-EMFs modulates cortical excitability in patients affected by focal epilepsy; however, in contrast to healthy subjects, these effects were evident only after EMFs exposure over the hemisphere contralateral to the epileptic focus (CH). They were characterized by a significant cortical excitability increase in the exposed hemisphere paired with slight excitability decrease in the other one (IH). Both sham and real EMFs exposure of the IH did not affect brain excitability. **CONCLUSION:** Present results suggest a significant interaction between the brain excitability changes induced by EMFs and the epileptic focus, which eliminated the excitability enhancing effects of EMFs evident only in the CH.

Perentos N, Croft RJ, McKenzie RJ, Cvetkovic D, Cosic I. The effect of GSM-like ELF radiation on the alpha band of the human resting EEG. Conf Proc IEEE Eng Med Biol Soc. 1:5680-5683, 2008.

Mobile phone handsets such as those operating in the GSM network emit extremely low frequency electromagnetic fields ranging from DC to at least 40 kHz. As a subpart of an extended protocol, the influence of these fields on the human resting EEG has been investigated in a fully counter balanced, double blind, cross-over design study that recruited 72 healthy volunteers. A decrease in the alpha frequency band was observed during the 20 minutes of ELF exposure in the exposed hemisphere only. This result suggests that ELF fields as emitted from GSM handsets during the DTX mode may have an effect on the resting alpha band of the human EEG.

Sokolovic D, Djindjic B, Nikolic J, Bjelakovic G, Pavlovic D, Kocic G, Krstic D, Cvetkovic T, Pavlovic V. Melatonin reduces oxidative stress induced by chronic exposure of microwave radiation from mobile phones in rat brain. J Radiat Res (Tokyo). 49(6):579-586, 2008.

PURPOSE: The aim of the study was to evaluate the intensity of oxidative stress in the brain of animals chronically exposed to mobile phones and potential protective effects of melatonin in reducing oxidative stress and brain injury. **MATERIALS AND METHODS:** Experiments were performed on Wistar rats exposed to microwave radiation during 20, 40 and 60 days. Four groups were formed: I group (control)- animals treated by saline, intraperitoneally (i.p.) applied daily during follow up, II group (Mel)- rats treated daily

Studies that show **Cell Phone** Health Effects

with melatonin (2 mg kg⁻¹) body weight i.p.), III group (MWs)- microwave exposed rats, IV group (MWs + Mel)- MWs exposed rats treated with melatonin (2 mg kg⁻¹) body weight i.p.). The microwave radiation was produced by a mobile test phone (SAR = 0.043-0.135 W/kg). **RESULTS:** A significant increase in the brain tissue malondialdehyde (MDA) and carbonyl group concentration was registered during exposure. Decreased activity of catalase (CAT) and increased activity of xanthine oxidase (XO) remained after 40 and 60 days of exposure to mobile phones. Melatonin treatment significantly prevented the increase in the MDA content and XO activity in the brain tissue after 40 days of exposure while it was unable to prevent the decrease of CAT activity and increase of carbonyl group contents. **CONCLUSION:** We demonstrated two important findings; that mobile phones caused oxidative damage biochemically by increasing the levels of MDA, carbonyl groups, XO activity and decreasing CAT activity; and that treatment with the melatonin significantly prevented oxidative damage in the brain.

Kesari KK, Meena R, Nirala J, Kumar J, Verma HN. Effect of 3G cell phone exposure with computer controlled 2-D stepper motor on non-thermal activation of the hsp27/p38MAPK stress pathway in rat brain. Cell Biochem Biophys. 68(2):347-358, 2014.

Cell phone radiation exposure and its biological interaction is the present concern of debate. Present study aimed to investigate the effect of 3G cell phone exposure with computer controlled 2-D stepper motor on 45-day-old male Wistar rat brain. Animals were exposed for 2 h a day for 60 days by using mobile phone with angular movement up to zero to 30°. The variation of the motor is restricted to 90° with respect to the horizontal plane, moving at a pre-determined rate of 2° per minute. Immediately after 60 days of exposure, animals were sacrificed and numbers of parameters (DNA double-strand break, micronuclei, caspase 3, apoptosis, DNA fragmentation, expression of stress-responsive genes) were performed. Result shows that microwave radiation emitted from 3G mobile phone significantly induced DNA strand breaks in brain. Meanwhile a significant increase in micronuclei, caspase 3 and apoptosis were also observed in exposed group (P < 0.05). Western blotting result shows that 3G mobile phone exposure causes a transient increase in phosphorylation of hsp27, hsp70, and p38 mitogen-activated protein kinase (p38MAPK), which leads to mitochondrial dysfunction-mediated cytochrome c release and subsequent activation of caspases, involved in the process of radiation-induced apoptotic cell death. Study shows that the oxidative stress is the main factor which activates a variety of cellular signal transduction pathways, among them the hsp27/p38MAPK is the pathway of principle stress response. Results conclude that 3G mobile phone radiations affect the brain function and cause several neurological disorders.

Kesari KK, Kumar S, Behari J. 900-MHz microwave radiation promotes oxidation in rat brain. Electromagn Biol Med. 30(4):219-234, 2011.

Recently, there have been several reports referring to detrimental effects due to radio frequency electromagnetic fields (RF-EMF) exposure. Special attention was given to investigate the effect of mobile phone exposure on the rat brain. Since the integrative

Studies that show **Cell Phone** Health Effects

mechanism of the entire body lies in the brain, it is suggestive to analyze its biochemical aspects. For this, 35-day old Wistar rats were exposed to a mobile phone for 2 h per day for a duration of 45 days where specific absorption rate (SAR) was 0.9 W/Kg. Animals were divided in two groups: sham exposed (n = 6) and exposed group (n = 6). Our observations indicate a significant decrease ($P < 0.05$) in the level of glutathione peroxidase, superoxide dismutase, and an increase in catalase activity. Moreover, protein kinase shows a significant decrease in exposed group ($P < 0.05$) of hippocampus and whole brain. Also, a significant decrease ($P < 0.05$) in the level of pineal melatonin and a significant increase ($P < 0.05$) in creatine kinase and caspase 3 was observed in exposed group of whole brain as compared with sham exposed. Finally, a significant increase in the level of ROS (reactive oxygen species) ($P < 0.05$) was also recorded. The study concludes that a reduction or an increase in antioxidative enzyme activities, protein kinase C, melatonin, caspase 3, and creatine kinase are related to overproduction of reactive oxygen species (ROS) in animals under mobile phone radiation exposure. Our findings on these biomarkers are clear indications of possible health implications.

Fragopoulou AF, Samara A, Antonelou MH, Xanthopoulou A, Papadopoulou A, Vougas K, Koutsogiannopoulou E, Anastasiadou E, Stravopodis DJ, Tsangaris GT, Margaritis LH. Brain proteome response following whole body exposure of mice to mobile phone or wireless DECT base radiation. Electromagn Biol Med.31(4):250-274, 2012.

The objective of this study was to investigate the effects of two sources of electromagnetic fields (EMFs) on the proteome of cerebellum, hippocampus, and frontal lobe in Balb/c mice following long-term whole body irradiation. Three equally divided groups of animals (6 animals/group) were used; the first group was exposed to a typical mobile phone, at a SAR level range of 0.17-0.37 W/kg for 3 h daily for 8 months, the second group was exposed to a wireless DECT base (Digital Enhanced Cordless Telecommunications/Telephone) at a SAR level range of 0.012-0.028 W/kg for 8 h/day also for 8 months and the third group comprised the sham-exposed animals. Comparative proteomics analysis revealed that long-term irradiation from both EMF sources altered significantly ($p < 0.05$) the expression of 143 proteins in total (as low as 0.003 fold downregulation up to 114 fold overexpression). Several neural function related proteins (i.e., Glial Fibrillary Acidic Protein (GFAP), Alpha-synuclein, Glia Maturation Factor beta (GMF), and apolipoprotein E (apoE)), heat shock proteins, and cytoskeletal proteins (i.e., Neurofilaments and tropomodulin) are included in this list as well as proteins of the brain metabolism (i.e., Aspartate aminotransferase, Glutamate dehydrogenase) to nearly all brain regions studied. Western blot analysis on selected proteins confirmed the proteomics data. The observed protein expression changes may be related to brain plasticity alterations, indicative of oxidative stress in the nervous system or involved in apoptosis and might potentially explain human health hazards reported so far, such as headaches, sleep disturbance, fatigue, memory deficits, and brain tumor long-term induction under similar exposure conditions.

Beason RC, Semm P. Responses of neurons to an amplitude-modulated microwave

Studies that show **Cell Phone** Health Effects

stimulus. Neurosci Lett 333(3):175-178, 2002.

In this study we investigated the effects of a pulsed radio frequency signal similar to the signal produced by global system for mobile communication telephones (900 MHz carrier, modulated at 217 Hz) on neurons of the avian brain. We found that such stimulation resulted in changes in the amount of neural activity by more than half of the brain cells. Most (76%) of the responding cells increased their rates of firing by an average 3.5-fold. The other responding cells exhibited a decrease in their rates of spontaneous activity Such responses indicate potential effects on humans using hand-held cellular phones.

Ragy MM. Effect of exposure and withdrawal of 900-MHz-electromagnetic waves on brain, kidney and liver oxidative stress and some biochemical parameters in male rats. Electromagn Biol Med. 2014 Apr 8. [Epub ahead of print]

Increasing use of mobile phones in daily life with increasing adverse effects of electromagnetic radiation (EMR), emitted from mobile on some physiological processes, cause many concerns about their effects on human health. Therefore, this work was designed to study the effects of exposure to mobile phone emits 900-MHz EMR on the brain, liver and kidney of male albino rats. Thirty male adult rats were randomly divided into four groups (10 each) as follows: control group (rats without exposure to EMR), exposure group (exposed to 900-MHz EMR for 1 h/d for 60 d) and withdrawal group (exposed to 900-MHz electromagnetic wave for 1 h/d for 60 d then left for 30 d without exposure). EMR emitted from mobile phone led to a significant increase in malondialdehyde (MDA) levels and significant decrease total antioxidant capacity (TAC) levels in brain, liver and kidneys tissues. The sera activity of alanine transaminase (ALT), aspartate aminotransferase (AST), urea, creatinine and corticosterone were significantly increased ($p < 0.05$), while serum catecholamines were insignificantly higher in the exposed rats. These alterations were corrected by withdrawal. In conclusion, electromagnetic field emitting from mobile phone might produce impairments in some biochemicals changes and oxidative stress in brain, liver and renal tissue of albino rats.

Maby E, Le Bouquin Jeannes R, Liegeois-Chauvel C, Gourevitch B, Faucon G. Analysis of auditory evoked potential parameters in the presence of radiofrequency fields using a support vector machines method. Med Biol Eng Comput. 42(4):562-568, 2004.

The paper presents a study of global system for mobile (GSM) phone radiofrequency effects on human cerebral activity. The work was based on the study of auditory evoked potentials (AEPs) recorded from healthy humans and epileptic patients. The protocol allowed the comparison of AEPs recorded with or without exposure to electrical fields. Ten variables measured from AEPs were employed in the design of a supervised support vector machines classifier. The classification performance measured the classifier's ability to discriminate features performed with or without radiofrequency exposure. Most significant features were chosen by a backward sequential selection that ranked the variables according to their pertinence for the discrimination. Finally, the most discriminating features were analysed statistically by a Wilcoxon signed rank test. For both populations, the N100 amplitudes were reduced under the influence of GSM

Studies that show **Cell Phone** Health Effects

radiofrequency (mean attenuation of -0.36 microV for healthy subjects and -0.60 microV for epileptic patients). Healthy subjects showed a N100 latency decrease (-5.23 ms in mean), which could be consistent with mild, localised heating. The auditory cortical activity in humans was modified by GSM phone radiofrequencies, but an effect on brain functionality has not been proven.

Karaca E, Durmaz B, Aktug H, Yildiz T, Guducu C, Irgi M, Koksai MG, Ozkinay F, Gunduz C, Cogulu O. The genotoxic effect of radiofrequency waves on mouse brain. J Neurooncol. 106(1):53-58, 2012.

Concerns about the health effects of radiofrequency (RF) waves have been raised because of the gradual increase in usage of cell phones, and there are scientific questions and debates about the safety of those instruments in daily life. The aim of this study is to evaluate the genotoxic effects of RF waves in an experimental brain cell culture model. Brain cell cultures of the mice were exposed to 10.715 GHz with specific absorption rate (SAR) 0.725 W/kg signals for 6 h in 3 days at 25°C to check for the changes in the micronucleus (MNI) assay and in the expression of 11 proapoptotic and antiapoptotic genes. It was found that MNI rate increased 11-fold and STAT3 expression decreased 7-fold in the cell cultures which were exposed to RF. Cell phones which spread RF may damage DNA and change gene expression in brain cells.

Chen C, Ma Q, Liu C, Deng P, Zhu G, Zhang L, He M, Lu Y, Duan W, Pei L, Li M, Yu Z, Zhou Z. Exposure to 1800 MHz radiofrequency radiation impairs neurite outgrowth of embryonic neural stem cells. Sci Rep. 2014 May 29;4:5103. doi: 10.1038/srep05103.

A radiofrequency electromagnetic field (RF-EMF) of 1800 MHz is widely used in mobile communications. However, the effects of RF-EMFs on cell biology are unclear. Embryonic neural stem cells (eNSCs) play a critical role in brain development. Thus, detecting the effects of RF-EMF on eNSCs is important for exploring the effects of RF-EMF on brain development. Here, we exposed eNSCs to 1800 MHz RF-EMF at specific absorption rate (SAR) values of 1, 2, and 4 W/kg for 1, 2, and 3 days. We found that 1800 MHz RF-EMF exposure did not influence eNSC apoptosis, proliferation, cell cycle or the mRNA expressions of related genes. RF-EMF exposure also did not alter the ratio of eNSC differentiated neurons and astrocytes. However, neurite outgrowth of eNSC differentiated neurons was inhibited after 4 W/kg RF-EMF exposure for 3 days. Additionally, the mRNA and protein expression of the proneural genes Ngn1 and NeuroD, which are crucial for neurite outgrowth, were decreased after RF-EMF exposure. The expression of their inhibitor Hes1 was upregulated by RF-EMF exposure. These results together suggested that 1800 MHz RF-EMF exposure impairs neurite outgrowth of eNSCs. More attention should be given to the potential adverse effects of RF-EMF exposure on brain development.

Bachmann M, Lass J, Kalda J, Säkki M, Tomson R, Tuulik V, Hinrikus H. Integration of differences in EEG analysis reveals changes in human EEG caused by microwave. Conf

Studies that show **Cell Phone** Health Effects

Proc IEEE Eng Med Biol Soc. 1:1597-1600, 2006.

Three different methods in combination with integration of differences in signals were applied for EEG analysis to distinguish changes in EEG caused by microwave: S-parameter, power spectral density and length distribution of low variability periods. The experiments on the effect of modulated low-level microwaves on human EEG were carried out on four different groups of healthy volunteers exposed to 450 MHz microwave radiation modulated with 7 Hz, 14 Hz, 21 Hz, 40 Hz, 70 Hz, 217 or 1000 Hz frequencies. The field power density at the scalp was 0.16 mW/cm². The EEG analysis performed for individuals with three different methods showed that statistically significant changes occur in the EEG rhythms energy and dynamics between 12% and 30% of subjects.

Wang Q, Cao ZJ, Bai XT. [Effect of 900 MHz electromagnetic fields on the expression of GABA receptor of cerebral cortical neurons in postnatal rats] Wei Sheng Yan Jiu. 34(5):546-548, 2005.

OBJECTIVE: To investigate the effects of 900 MHz microwave electromagnetic fields (EMF) on the expression of neurotransmitter GABA receptor of cerebral cortical neurons in postnatal rats. **METHODS:** Neurons were exposed to 900 MHz continuous microwave EMF (SAR = 1.15 - 3.22mW/g) for 2 hours per day in 6 consecutive days and for 12 hours at one time. GABA receptor was chosen to be the biological end. **RESULTS:** Significant changes had been observed in exposed neurons in the expression of GABA receptor. ($P < 0.01$). **CONCLUSION:** The expression of GABA receptor of neurons were significantly regulated by 900 MHz microwave, and a power "window" effect was observed in the exposed neurons.

Leung S, Croft RJ, McKenzie RJ, Iskra S, Silber B, Cooper NR, O'Neill B, Cropley V, Diaz-Trujillo A, Hamblin D, Simpson D. Effects of 2G and 3G mobile phones on performance and electrophysiology in adolescents, young adults and older adults. Clin Neurophysiol. 122(11):2203-2216, 2011.

OBJECTIVE: This study examined sensory and cognitive processing in adolescents, young adults and older adults, when exposed to 2nd (2G) and 3rd (3G) generation mobile phone signals. **METHODS:** Tests employed were the auditory 3-stimulus oddball and the N-back. Forty-one 13-15 year olds, forty-two 19-40 year olds and twenty 55-70 year olds were tested using a double-blind cross-over design, where each participant received Sham, 2G and 3G exposures, separated by at least 4 days. **RESULTS:** 3-Stimulus oddball task: Behavioural: accuracy and reaction time of responses to targets were not affected by exposure. Electrophysiological: augmented N1 was found in the 2G condition (independent of age group). N-back task: Behavioural: the combined groups performed less accurately during the 3G exposure (compared to Sham), with post hoc tests finding this effect separately in the adolescents only. Electrophysiological: delayed ERD/ERS responses of the alpha power were found in both 3G and 2G conditions (compared to Sham; independent of age group). **CONCLUSION:** Employing tasks tailored to each individual's ability level, this study provides support for an effect of acute 2G and 3G exposure on human cognitive function. **SIGNIFICANCE:** The subtlety of mobile phone

Studies that show **Cell Phone** Health Effects

effect on cognition in our study suggests that it is important to account for individual differences in future mobile phone research.

Söderqvist F, Hardell L, Carlberg M, Mild KH. Radiofrequency fields, transthyretin, and Alzheimer's disease. J Alzheimers Dis. 20(2):599-606, 2010.

Radiofrequency field (RF) exposure provided cognitive benefits in an animal study. In Alzheimer's disease (AD) mice, exposure reduced brain amyloid-beta (Abeta) deposition through decreased aggregation of Abeta and increase in soluble Abeta levels. Based on our studies on humans on RF from wireless phones, we propose that transthyretin (TTR) might explain the findings. In a cross-sectional study on 313 subjects, we used serum TTR as a marker of cerebrospinal fluid TTR. We found a statistically significantly positive beta coefficient for TTR for time since first use of mobile phones and desktop cordless phones combined ($P=0.03$). The electromagnetic field parameters were similar for the phone types. In a provocation study on 41 persons exposed for 30 min to an 890-MHz GSM signal with specific absorption rate of 1.0 Watt/kg to the temporal area of the brain, we found statistically significantly increased serum TTR 60 min after exposure. In our cross-sectional study, use of oral snuff also yielded statistically significantly increased serum TTR concentrations and nicotine has been associated with decreased risk for AD and to upregulate the TTR gene in choroid plexus but not in the liver, another source of serum TTR. TTR sequesters Abeta, thereby preventing the formation of Abeta plaques in the brain. Studies have shown that patients with AD have lowered TTR concentrations in the cerebrospinal fluid and have attributed the onset of AD to insufficient sequestering of Abeta by TTR. We propose that TTR might be involved in the findings of RF exposure benefit in AD mice.

Maby E, Jeannes Rle B, Faucon G. Scalp localization of human auditory cortical activity modified by GSM electromagnetic fields. Int J Radiat Biol. 82(7):465-472, 2006.

Purpose: This study attempted to determine whether there is a localized effect of GSM (Global System for Mobile communications) microwaves by studying the Auditory Evoked Potentials (AEP) recorded at the scalp of nine healthy subjects and six epileptic patients. Materials and methods: We determined the influence of GSM RadioFrequency (RF) on parameters characterizing the AEP in time or/and frequency domains. A parameter selection method using SVM (Support Vector Machines)-based criteria allowed us to estimate those most altered by the radiofrequencies. The topography of the parameter modifications was computed to determine the localization of the radiofrequency influence. A statistical test was conducted for selected scalp areas, in order to determine whether there were significant localized alterations due to the RF. Results: The epileptic patients showed a lengthening of the scalp component N100 (100 ms latency) in the frontal area contralateral to the radiation, which may be due to an afferent tract alteration. For the healthy subjects, an amplitude increase of the P200 wave (200 ms latency) was identified in the frontal area. Conclusions: The present study suggests that radiofrequency fields emitted by mobile phones modify the AEP. Nevertheless, no direct link between

Studies that show **Cell Phone** Health Effects

Maskey D, Kim HJ, Kim HG, Kim MJ. Calcium-binding proteins and GFAP immunoreactivity alterations in murine hippocampus after 1 month of exposure to 835MHz radiofrequency at SAR values of 1.6 and 4.0W/kg. Neurosci Lett. 506(2):292-296, 2012.

Abstract. Widespread use of wireless mobile communication has raised concerns of adverse effect to the brain owing to the proximity during use due to the electromagnetic field emitted by mobile phones. Changes in calcium ion concentrations via binding proteins can disturb calcium homeostasis; however, the correlation between calcium-binding protein (CaBP) immunoreactivity (IR) and glial cells has not been determined with different SAR values. Different SAR values [1.6 (E1.6 group) and 4.0 (E4 group) W/kg] were applied to determine the distribution of calbindin D28-k (CB), calretinin (CR), and glial fibrillary acidic protein (GFAP) IR in murine hippocampus. Compared with sham control group, decreased CB and CR IRs, loss of CB and CR immunoreactive cells and increased GFAP IR exhibiting hypertrophic cytoplasmic processes were noted in both experimental groups. E4 group showed a prominent decrement in CB and CR IR than the E1.6 group due to down-regulation of CaBP proteins and neuronal loss. GFAP IR was more prominent in the E4 group than the E1.6 group. Decrement in the CaBPs can affect the calcium-buffering capacity leading to cell death, while increased GFAP IR and changes in astrocyte morphology, may mediate brain injury due to radiofrequency exposure.

Meral I, Mert H, Mert N, Deger Y, Yoruk I, Yetkin A, Keskin S. Effects of 900-MHz electromagnetic field emitted from cellular phone on brain oxidative stress and some vitamin levels of guinea pigs. Brain Res1169:120-124, 2007.

This study was designed to demonstrate the effects of 900-MHz electromagnetic field (EMF) emitted from cellular phone on brain tissue and also blood malondialdehyde (MDA), glutathione (GSH), retinol (vitamin A), vitamin D(3) and tocopherol (vitamin E) levels, and catalase (CAT) enzyme activity of guinea pigs. Fourteen male guinea pigs, weighing 500-800 g were randomly divided into one of two experimental groups: control and treatment (EMF-exposed), each containing seven animals. Animals in treatment group were exposed to 890- to 915-MHz EMF (217-Hz pulse rate, 2-W maximum peak power, SAR 0.95 w/kg) of a cellular phone for 12 h/day (11-h 45-min stand-by and 15-min spiking mode) for 30 days. Control guinea pigs were housed in a separate room without exposing EMF of a cellular phone. Blood samples were collected through a cardiac puncture and brains were removed after decapitation for the biochemical analysis at the end of the 30 days of experimental period. It was found that the MDA level increased ($P<0.05$), GSH level and CAT enzyme activity decreased ($P<0.05$), and vitamins A, E and D(3) levels did not change ($P>0.05$) in the brain tissues of EMF-exposed guinea pigs. In addition, MDA, vitamins A, D(3) and E levels, and CAT enzyme activity increased ($P<0.05$), and GSH level decreased ($P<0.05$) in the blood of EMF-exposed guinea pigs. It was concluded that electromagnetic field emitted from cellular phone might produce oxidative stress in brain tissue of guinea pigs. However, more studies are needed to demonstrate whether these effects are harmful or/and affect the neural functions.

Studies that show **Cell Phone** Health Effects

Mausset-Bonnefont AL, Hirbec H, Bonnefont X, Privat A, Vignon J, de Seze R. Acute exposure to GSM 900-MHz electromagnetic fields induces glial reactivity and biochemical modifications in the rat brain. Neurobiol Dis. 17(3):445-454, 2004.

The worldwide proliferation of mobile phones raises the question of the effects of 900-MHz electromagnetic fields (EMF) on the brain. Using a head-only exposure device in the rat, we showed that a 15-min exposure to 900-MHz pulsed microwaves at a high brain-averaged power of 6 W/kg induced a strong glial reaction in the brain. This effect, which suggests neuronal damage, was particularly pronounced in the striatum. Moreover, we observed significant and immediate effects on the K(d) and B(max) values of N-methyl-d-aspartate (NMDA) and GABA(A) receptors as well as on dopamine transporters. Decrease of the amount of NMDA receptors at the postsynaptic membrane is also reported. Although we showed that the rat general locomotor behavior was not significantly altered on the short term, our results provide the first evidence for rapid cellular and molecular alterations in the rat brain after an acute exposure to high power GSM (Global System for Mobile communication) 900-MHz microwaves.

Nittby H, Widegren B, Krogh M, Grafström G, Berlin H, Rehn G, Eberhardt JL, Malmgren L, Persson BRR, Salford L. Exposure to radiation from global system for mobile communications at 1,800 MHz significantly changes gene expression in rat hippocampus and cortex. Environmentalist 28(4), 458-465, 2008.

We have earlier shown that radio frequency electromagnetic fields can cause significant leakage of albumin through the blood–brain barrier of exposed rats as compared to non-exposed rats, and also significant neuronal damage in rat brains several weeks after a 2 h exposure to a mobile phone, at 915 MHz with a global system for mobile communications (GSM) frequency modulation, at whole-body specific absorption rate values (SAR) of 200, 20, 2, and 0.2 mW/kg. We have now studied whether 6 h of exposure to the radiation from a GSM mobile test phone at 1,800 MHz (at a whole-body SAR-value of 13 mW/kg, corresponding to a brain SAR-value of 30 mW/kg) has an effect upon the gene expression pattern in rat brain cortex and hippocampus—areas where we have observed albumin leakage from capillaries into neurons and neuronal damage. Microarray analysis of 31,099 rat genes, including splicing variants, was performed in cortex and hippocampus of 8 Fischer 344 rats, 4 animals exposed to global system for mobile communications electromagnetic fields for 6 h in an anechoic chamber, one rat at a time, and 4 controls kept as long in the same anechoic chamber without exposure, also in this case one rat at a time. Gene ontology analysis (using the gene ontology categories biological processes, molecular functions, and cell components) of the differentially expressed genes of the exposed animals versus the control group revealed the following highly significant altered gene categories in both cortex and hippocampus: extracellular region, signal transducer activity, intrinsic to membrane, and integral to membrane. The fact that most of these categories are connected with membrane functions may have a relation to our earlier observation of albumin transport through brain capillaries.

Studies that show **Cell Phone** Health Effects

Söderqvist F, Carlberg M, Hansson Mild K, Hardell L. Exposure to an 890-MHz mobile phone-like signal and serum levels of S100B and transthyretin in volunteers. Toxicol Lett. 189(1):63-66, 2009.

Whether low-intensity non-thermal microwave radiation alters the integrity of the blood-brain barrier has been debated since the late 1970s, yet no experimental study has been carried out on humans. The aim of this study was to test, using peripheral markers, whether exposure to a mobile phone-like signal alters the integrity of the human blood-brain and blood-cerebrospinal fluid barriers. A provocation study was carried out that exposed 41 volunteers to a 30 min GSM 890 MHz signal with an average specific energy absorption rate distribution of 1.0 W/kg in the temporal area of the head as measured over any 1g of contiguous tissue. The outcome was assessed by changes in serum concentrations of two putative markers of brain barrier integrity, S100B and transthyretin. Repeated blood sampling before and after the provocation showed no statistically significant increase in the serum levels of S100B, while for transthyretin a statistically significant increase was seen in the final blood sample 60 min after the end of the provocation as compared to the prior sample taken immediately after provocation (p=0.02). The clinical significance of this finding, if any, is unknown. Further randomized studies with use of additional more brain specific markers are needed.

Schirmacher A, Winters S, Fischer S, Goeke J, Galla H, Kullnick U, Ringelstein EB, Stogbauer F, Electromagnetic fields (1.8 GHz) increase the permeability to sucrose of the blood-brain barrier in vitro. Bioelectromagnetics 21(5):338-345, 2000.

We report an investigation on the influence of high frequency electromagnetic fields (EMF) on the permeability of an in vitro model of the blood-brain barrier (BBB). Our model was a co-culture consisting of rat astrocytes and porcine brain capillary endothelial cells (BCEC). Samples were characterized morphologically by scanning electron microscopy and immunocytochemistry. The BBB phenotype of the BCEC was shown by the presence of zona occludens protein (ZO-1) as a marker for tight junctions and the close contact of the cells together with the absence of intercellular clefts. Permeability measurements using (14)C-sucrose indicated a physiological tightness which correlated with the morphological findings and verified the usefulness of our in vitro model. Samples were exposed to EMF conforming to the GSM1800-standard used in mobile telephones (1.8 GHz). The permeability of the samples was monitored over four days and compared with results of samples that were cultured identically but not exposed to EMF. Exposure to EMF increased permeability for (14)C-sucrose significantly compared to unexposed samples. The underlying pathophysiological mechanism remains to be investigated.

Narayanan SN, Kumar RS, Kedage V, Nalini K, Nayak S, Bhat PG. Evaluation of oxidant stress and antioxidant defense in discrete brain regions of rats exposed to 900 MHz radiation. Bratisl Lek Listy. 115(5):260-266, 2014.

AIM: In the current study, the effects of 900 MHz radio-frequency electromagnetic radiation (RF-EMR) on levels of thiobarbituric acid-reactive substances (TBARS), total

Studies that show **Cell Phone** Health Effects

antioxidants (TA), and glutathione S-transferase (GST) activity in discrete brain regions were studied in adolescent rats. **MATERIALS AND METHODS:** Thirty-six male Wistar rats (6-8 weeks old) were allotted into three groups (n = 12 in each group). Control group (1) remained undisturbed in their home cage; sham group (2) was exposed to mobile phone in switch off mode for four weeks; RF-EMR-exposed group (3) was exposed to 900 MHz of RF-EMR (1 hr/day with peak power density of 146.60 $\mu\text{W}/\text{cm}^2$) from an activated Global System for Mobile communication (GSM) mobile phone (kept in silent mode; no ring tone and no vibration) for four weeks. On 29th day, behavioral analysis was done. Followed by this, six animals from each group were sacrificed and biochemical parameters were studied in amygdala, hippocampus, frontal cortex, and cerebellum. **RESULTS:** Altered behavioral performances were found in RF-EMR-exposed rats. Additionally, elevated TBARS level was found with all brain regions studied. RF-EMR exposure significantly decreased TA in the amygdala and cerebellum but its level was not significantly changed in other brain regions. GST activity was significantly decreased in the hippocampus but, its activity was unaltered in other brain regions studied. **CONCLUSION:** RF-EMR exposure for a month induced oxidative stress in rat brain, but its magnitude was different in different regions studied. RF-EMR-induced oxidative stress could be one of the underlying causes for the behavioral deficits seen in rats after RF-EMR exposure (Fig. 5, Ref. 37).

Narayanan SN, Kumar RS, Potu BK, Nayak S, Bhat PG, Mailankot M. Effect of radio-frequency electromagnetic radiations (RF-EMR) on passive avoidance behaviour and hippocampal morphology in Wistar rats. Ups J Med Sci.115(2):91-96, 2010.

Abstract Introduction. The interaction of mobile phone radio-frequency electromagnetic radiation (RF-EMR) with the brain is a serious concern of our society. **Objective.** We evaluated the effect of RF-EMR from mobile phones on passive avoidance behaviour and hippocampal morphology in rats. **Materials and methods.** Healthy male albino Wistar rats were exposed to RF-EMR by giving 50 missed calls (within 1 hour) per day for 4 weeks, keeping a GSM (0.9 GHz/1.8 GHz) mobile phone in vibratory mode (no ring tone) in the cage. After the experimental period, passive avoidance behaviour and hippocampal morphology were studied. **Results.** Passive avoidance behaviour was significantly affected in mobile phone RF-EMR-exposed rats demonstrated as shorter entrance latency to the dark compartment when compared to the control rats. Marked morphological changes were also observed in the CA(3) region of the hippocampus of the mobile phone-exposed rats in comparison to the control rats. **Conclusion.** Mobile phone RF-EMR exposure significantly altered the passive avoidance behaviour and hippocampal morphology in rats.

Murbach, M., Neufeld, E., Christopoulou, M., Achermann, P. and Kuster, N. (2014), Modeling of EEG electrode artifacts and thermal ripples in human radiofrequency exposure studies. Bioelectromagnetics. doi: 10.1002/bem.21837.

The effects of radiofrequency (RF) exposure on wake and sleep electroencephalogram (EEG) have been in focus since mobile phone usage became pervasive. It has been hypothesized that effects may be explained by (1) enhanced induced fields due to RF

Studies that show **Cell Phone** Health Effects

coupling with the electrode assembly, (2) the subsequent temperature increase around the electrodes, or (3) RF induced thermal pulsing caused by localized exposure in the head. We evaluated these three hypotheses by means of both numerical and experimental assessments made with appropriate phantoms and anatomical human models. Typical and worst-case electrode placements were examined at 900 and 2140 MHz. Our results indicate that hypothesis 1 can be rejected, as the induced fields cause <20% increase in the 10 g-averaged specific absorption rate (SAR). Simulations with an anatomical model indicate that hypothesis 2 is also not supported, as the realistic worst-case electrode placement results in a maximum skin temperature increase of 0.31 °C while brain temperature elevations remained <0.1 °C. These local short-term temperature elevations are unlikely to change brain physiology during the time period from minutes to several hours after exposure. The maximum observed temperature ripple due to RF pulses is <0.001 °C for GSM-like signals and <0.004 °C for 20-fold higher pulse energy, and offers no support for hypothesis 3. Thus, the mechanism of interaction between RF and changes in the EEG power spectrum remains unknown.

Vorobyov VV, Galchenko AA, Kukushkin NI, Akoev IG, Effects of weak microwave fields amplitude modulated at ELF on EEG of symmetric brain areas in rats.

Bioelectromagnetics 18(4):293-298, 1997.

Averaged electroencephalogram (EEG) frequency spectra were studied in eight unanesthetized and unmyorelaxed adult male rats with chronically implanted carbon electrodes in symmetrical somesthetic areas when a weak (0.1-0.2 mW/cm²) microwave (MW, 945 MHz) field, amplitude-modulated at extremely low frequency (ELF) (4 Hz), was applied. Intermittent (1 min "On," 1 min "Off") field exposure (10-min duration) was used. Hemispheric asymmetry in frequency spectra (averaged data for 10 or 1 min) of an ongoing EEG was characterized by a power decrease in the 1.5-3 Hz range on the left hemisphere and by a power decrease in the 10-14 and 20-30 Hz ranges on the right hemisphere. No differences between control and exposure experiments were shown under these routines of data averaging. Significant elevations of EEG asymmetry in 10-14 Hz range were observed during the first 20 s after four from five onsets of the MW field, when averaged spectra were obtained for every 10 s. Under neither control nor pre- and postexposure conditions was this effect observed. These results are discussed with respect to interaction of MW fields with the EEG generators.

Maskey D, Pradhan J, Aryal B, Lee CM, Choi IY, Park KS, Kim SB, Kim HG, Kim MJ.

Chronic 835 MHz radiofrequency exposure to mice hippocampus alters the distribution of calbindin and GFAP immunoreactivity. Brain Res 1346:237-246, 2010.

Exponential interindividual handling in wireless communication system has raised possible doubts in the biological aspects of radiofrequency (RF) exposure on human brain owing to its close proximity to the mobile phone. In the nervous system, calcium (Ca²⁺) plays a critical role in releasing neurotransmitters, generating action potential and membrane integrity. Alterations in intracellular Ca²⁺ concentration trigger

Studies that show **Cell Phone** Health Effects

aberrant synaptic action or cause neuronal apoptosis, which may exert an influence on the cellular pathology for learning and memory in the hippocampus. Calcium binding proteins like calbindin D28-K (CB) is responsible for the maintaining and controlling $\text{Ca}(2+)$ homeostasis. Therefore, in the present study, we investigated the effect of RF exposure on rat hippocampus at 835MHz with low energy (Specific Absorption Rate: $\text{SAR}=1.6\text{W/kg}$) for 3months by using both CB and glial fibrillary acidic protein (GFAP) specific antibodies by immunohistochemical method. Decrease in CB immunoreactivity (IR) was noted in exposed (E1.6) group with loss of interneurons and pyramidal cells in CA1 area and loss of granule cells. Also, an overall increase in GFAP IR was observed in the hippocampus of E1.6. By TUNEL assay, apoptotic cells were detected in the CA1, CA3 areas and dentate gyrus of hippocampus, which reflects that chronic RF exposure may affect the cell viability. Additionally, the increase of GFAP IR due to RF exposure could be well suited with the feature of reactive astrogliosis, which is an abnormal increase in the number of astrocytes due to the loss of nearby neurons. Chronic RF exposure to the rat brain suggested that the decrease of CB IR accompanying apoptosis and increase of GFAP IR might be morphological parameters in the hippocampus damages.

Maby E, Jeannes RL, Faucon G, Liegeois-Chauvel C, De Seze R. Effects of GSM signals on auditory evoked responses. Bioelectromagnetics. 26(5):341-350, 2005.

The article presents a study of the influence of radio frequency (RF) fields emitted by mobile phones on human cerebral activity. Our work was based on the study of Auditory Evoked Potentials (AEPs) recorded on the scalp of healthy humans and epileptic patients. The protocol allowed us to compare AEPs recorded with or without exposure to RFs. To get a reference, a control session was also introduced. In this study, the correlation coefficients computed between AEPs, as well as the correlation coefficients between spectra of AEPs were investigated to detect a possible difference due to RFs. A difference in the correlation coefficients computed in control and experimental sessions was observed, but it was difficult to deduce the effect of RFs on human health.

Maaroufi K, Had-Aissouni L, Melon C, Sakly M, Abdelmelek H, Poucet B, Save E. Spatial learning, monoamines and oxidative stress in rats exposed to 900MHz electromagnetic field in combination with iron overload. Behav Brain Res. 2013 Oct 18. pii: S0166-4328(13)00624-4. doi: 10.1016/j.bbr.2013.10.016. [Epub ahead of print]

The increasing use of mobile phone technology over the last decade raises concerns about the impact of high frequency electromagnetic fields (EMF) on health. More recently, a link between EMF, iron overload in the brain and neurodegenerative disorders including Parkinson's and Alzheimer's diseases has been suggested. Co-exposure to EMF and brain iron overload may have a greater impact on brain tissues and cognitive processes than each treatment by itself. To examine this hypothesis, Long-Evans rats submitted to 900MHz exposure or combined 900MHz EMF and iron overload treatments were tested in various spatial learning tasks (navigation task in the Morris water maze, working memory task in the radial-arm maze, and object exploration task involving spatial and non spatial processing). Biogenic monoamines and metabolites (dopamine, serotonin) and oxidative stress were measured. Rats exposed to EMF were

Studies that show **Cell Phone** Health Effects

impaired in the object exploration task but not in the navigation and working memory tasks. They also showed alterations of monoamine content in several brain areas but mainly in the hippocampus. Rats that received combined treatment did not show greater behavioral and neurochemical deficits than EMF-exposed rats. None of the two treatments produced global oxidative stress. These results show that there is an impact of EMF on the brain and cognitive processes but this impact is revealed only in a task exploiting spontaneous exploratory activity. In contrast, there are no synergistic effects between EMF and a high content of iron in the brain.

Lv B, Su C, Yang L, Xie Y, Wu T. Whole brain EEG synchronization likelihood modulated by long term evolution electromagnetic fields exposure. Conf Proc IEEE Eng Med Biol Soc. 2014:986-989, 2014.

In this paper, we aimed to investigate the possible interactions between human brain and radiofrequency electromagnetic fields (EMF) with electroencephalogram (EEG) technique. Unlike the previous studies which mainly focused on EMF effect on local brain activities, we attempted to evaluate whether the EMF emitted from Long Term Evolution (LTE) devices can modulate the functional connectivity of brain electrical activities. Ten subjects were recruited to participate in a crossover, double-blind exposure experiment which included two sessions (real and sham exposure). In each session, LTE EMF exposure (power on or off) lasted for 30 min and the EEG signals were collected with 32 channels throughout the experiment. Then we applied the synchronization likelihood method to quantify the neural synchronization over the whole brain in different frequency bands and in different EEG record periods. Our results illustrated that the short-term LTE EMF exposure would modulate the synchronization patterns of EEG activation across the whole brain.

Lv B, Chen Z, Wu T, Shao Q, Yan D, Ma L, Lu K, Xie Y. The alteration of spontaneous low frequency oscillations caused by acute electromagnetic fields exposure. Clin Neurophysiol. 2013 Sep 4. pii: S1388-2457(13)00976-0. doi: 10.1016/j.clinph.2013.07.018. [Epub ahead of print]

OBJECTIVE: The motivation of this study is to evaluate the possible alteration of regional resting state brain activity induced by the acute radiofrequency electromagnetic field (RF-EMF) exposure (30min) of Long Term Evolution (LTE) signal. **METHODS:** We designed a controllable near-field LTE RF-EMF exposure environment. Eighteen subjects participated in a double-blind, crossover, randomized and counterbalanced experiment including two sessions (real and sham exposure). The radiation source was close to the right ear. Then the resting state fMRI signals of human brain were collected before and after the exposure in both sessions. We measured the amplitude of low frequency fluctuation (ALFF) and fractional ALFF (fALFF) to characterize the spontaneous brain activity. **RESULTS:** We found the decreased ALFF value around in left superior temporal gyrus, left middle temporal gyrus, right superior temporal gyrus, right medial frontal gyrus and right paracentral lobule after the real exposure. And the decreased fALFF value was also detected in right medial frontal gyrus and right paracentral lobule. **CONCLUSIONS:** The study provided the evidences that 30min LTE RF-EMF exposure modulated the spontaneous low frequency fluctuations in some brain

Studies that show **Cell Phone** Health Effects

regions. SIGNIFICANCE: With resting state fMRI, we found the alteration of spontaneous low frequency fluctuations induced by the acute LTE RF-EMF exposure.

López-Martín E, Bregains J, Relova-Quinteiro JL, Cadarso-Suárez C, Jorge-Barreiro FJ, Ares-Pena FJ. The action of pulse-modulated GSM radiation increases regional changes in brain activity and c-Fos expression in cortical and subcortical areas in a rat model of picrotoxin-induced seizure proneness. J Neurosci Res. 87(6):1484-1499, 2009.

The action of the pulse-modulated GSM radiofrequency of mobile phones has been suggested as a physical phenomenon that might have biological effects on the mammalian central nervous system. In the present study, GSM-exposed picrotoxin-pretreated rats showed differences in clinical and EEG signs, and in c-Fos expression in the brain, with respect to picrotoxin-treated rats exposed to an equivalent dose of unmodulated radiation. Neither radiation treatment caused tissue heating, so thermal effects can be ruled out. The most marked effects of GSM radiation on c-Fos expression in picrotoxin-treated rats were observed in limbic structures, olfactory cortex areas and subcortical areas, the dentate gyrus, and the central lateral nucleus of the thalamic intralaminar nucleus group. Nonpicrotoxin-treated animals exposed to unmodulated radiation showed the highest levels of neuronal c-Fos expression in cortical areas. These results suggest a specific effect of the pulse modulation of GSM radiation on brain activity of a picrotoxin-induced seizure-proneness rat model and indicate that this mobile-phone-type radiation might induce regional changes in previous preexcitability conditions of neuronal activation.

López-Martín E, Bregains J, Relova-Quinteiro JL, Cadarso-Suárez C, Jorge-Barreiro FJ, Ares-Pena FJ. The action of pulse-modulated GSM radiation increases regional changes in brain activity and c-Fos expression in cortical and subcortical areas in a rat model of picrotoxin-induced seizure proneness. J Neurosci Res. 87(6):1484-1499, 2009.

The action of the pulse-modulated GSM radiofrequency of mobile phones has been suggested as a physical phenomenon that might have biological effects on the mammalian central nervous system. In the present study, GSM-exposed picrotoxin-pretreated rats showed differences in clinical and EEG signs, and in c-Fos expression in the brain, with respect to picrotoxin-treated rats exposed to an equivalent dose of unmodulated radiation. Neither radiation treatment caused tissue heating, so thermal effects can be ruled out. The most marked effects of GSM radiation on c-Fos expression in picrotoxin-treated rats were observed in limbic structures, olfactory cortex areas and subcortical areas, the dentate gyrus, and the central lateral nucleus of the thalamic intralaminar nucleus group. Nonpicrotoxin-treated animals exposed to unmodulated radiation showed the highest levels of neuronal c-Fos expression in cortical areas. These results suggest a specific effect of the pulse modulation of GSM radiation on brain activity of a picrotoxin-induced seizure-proneness rat model and indicate that this mobile-phone-type radiation might induce regional changes in previous preexcitability conditions of neuronal activation.

Lebedeva NN, Sulimov AV, Sulimova OP, Kotrovskaya TI, Gailus T. Cellular phone

Studies that show **Cell Phone** Health Effects

electromagnetic field effects on bioelectric activity of human brain. Crit Rev Biomed Eng 28(1-2):323-337, 2000.

24 volunteers participated in the experiments. The investigation of EEG reactions to cellular phone (EMF frequency 902.4 MHz and intensity 0.06 mW/cm²) was conducted. Two experiments were performed with each subject--cellular phone exposure and Placebo Duration of the experiment was 60 min: 15 min--background; 15 min--EMF exposure or Placebo; 30 min--afterexposure. EEG was recorded in 16 standard leads with "eyes open" and "eyes closed". Special software with non-linear dynamics was developed for EEG analyses. One parameter, multichannel (global) correlation dimension, was calculated. The changes of these parameters can be evidence of brain functional state changes. As a result of EEG record processing, a significant increase of global correlation dimension during the exposure and afterexposure period was discovered, more pronounced in the case of "eyes closed". That can be viewed as the manifestation of cortex activation under phone EMF exposure.

Lass L, Tuulik V, Ferenets CR, Riisalo R, Hinrikus H. Effects of 7 Hz-modulated 450 MHz electromagnetic radiation on human performance in visual memory tasks. Int. J. Rad. Biol. 78: 937-944, 2002.

Abstract: Purpose: The aim was to examine low-level 7 Hz-modulated 450 MHz radiation effects on human performance in visually presented neuropsychological tasks associated with attention and short-term memory. Materials and methods: A homogeneous group of 100 subjects (37 female, 63 male) were randomly assigned to either the exposed (10-20 min, 0.158mW cm⁻²) or the sham-exposed group. A battery of three different tests measured attention and shortterm memory. Task 1 involved alternately selecting black digits from 1 to 25 in ascending order and white digits from 24 to 1 in descending order. The time spent on the task and the number of errors were recorded and analysed. Task 2 involved viewing a picture of 12 objects during 3 s, followed by a list of 24 words. The subject was required to select words representing previously presented objects. In task 3, an array of letters in 10 rows (60 in each row) was presented, and the subject was required to identify all examples of a particular two-letter combination. Results: The results of tasks 1 and 3 showed a significant increase in variances of errors ($p < 0.05$) in the exposed versus the shamexposed group. The results of task 2 indicated a significant decrease in errors ($p < 0.05$) in the exposed group. Conclusions: The data provide additional evidence that acute lowlevel exposure to microwaves modulated at 7 Hz can affect cognitive processes such as attention and short-term memory.

Kramarenko AV, Tan U. Effects of high-frequency electromagnetic fields on human eeg: A brain mapping study. Int J Neurosci. 113(7):1007-1019, 2003.

Cell phones emitting pulsed high-frequency electromagnetic fields (EMF) may affect the human brain, but there are inconsistent results concerning their effects on electroencephalogram (EEG). We used a 16-channel telemetric electroencephalograph (ExpertTM), to record EEG changes during exposure of human skull to EMF emitted by a mobile phone. Spatial distribution of EMF was especially concentrated around the

Studies that show **Cell Phone** Health Effects

ipsilateral eye adjacent to the basal surface of the brain. Traditional EEG was full of noises during operation of a cellular phone. Using a telemetric electroencephalograph (ExpertTM) in awake subjects, all the noise was eliminated, and EEG showed interesting changes: after a period of 10-15 s there was no visible change, the spectrum median frequency increased in areas close to antenna; after 20-40 s, a slow-wave activity (2.5-6.0 Hz) appeared in the contralateral frontal and temporal areas. These slow waves lasting for about one second repeated every 15-20 s at the same recording electrodes. After turning off the mobile phone, slow-wave activity progressively disappeared; local changes such as increased median frequency decreased and disappeared after 15-20 min. We observed similar changes in children, but the slow-waves with higher amplitude appeared earlier in children (10-20 s) than adults, and their frequency was lower (1.0-2.5 Hz) with longer duration and shorter intervals. The results suggested that cellular phones may reversibly influence the human brain, inducing abnormal slow waves in EEG of awake persons.

Krause CM, Sillanmaki L, Koivisto M, Haggqvist A, Saarela C, Revonsuo A, Laine M, Hamalainen H, Effects of electromagnetic field emitted by cellular phones on the EEG during a memory task. Neuroreport 11(4):761-764, 2000.

The effects of electromagnetic fields (EMF) emitted by cellular phones on the ERD/ERS of the 4-6 Hz, 6-8 Hz, 8-10 Hz and 10-12 Hz EEG frequency bands were studied in 16 normal subjects performing an auditory memory task. All subjects performed the memory task both with and without exposure to a digital 902 MHz EMF in counterbalanced order. The exposure to EMF significantly increased EEG power in the 8-10 Hz frequency band only. Nonetheless, the presence of EMF altered the ERD/ERS responses in all studied frequency bands as a function of time and memory task (encoding vs retrieval). Our results suggest that the exposure to EMF does not alter the resting EEG per se but modifies the brain responses significantly during a memory task.

Krause CM, Sillanmaki L, Koivisto M, Haggqvist A, Saarela C, Revonsuo A, Laine M, Hamalainen H, Effects of electromagnetic fields emitted by cellular phones on the electroencephalogram during a visual working memory task. Int J Radiat Biol 76(12):1659-1667, 2000.

PURPOSE: To examine the effects of electromagnetic fields (EMF) emitted by cellular phones on the event-related desynchronization/synchronization (ERD/ERS) responses of the 4-6, 6-8, 8-10 and 10-12Hz EEG frequency bands during cognitive processing. **MATERIALS AND METHODS:** Twenty-four subjects performed a visual sequential letter task (n-back task) with three different working memory load conditions: zero, one and two items. All subjects performed the memory task both with and without exposure to a digital 902 MHz EMF in counterbalanced order. **RESULTS:** The presence of EMF altered the ERD/ERS responses in the 6-8 and 8-10 Hz frequency bands but only when examined as a function of memory load and depending also on whether the presented stimulus was a target or not. **CONCLUSIONS:** The results suggest that the exposure to EMF modulates the responses of EEG oscillatory activity approximately 8 Hz specifically during cognitive processes.

Studies that show **Cell Phone** Health Effects

Krause CM, Haarala C, Sillanmaki L, Koivisto M, Alanko K, Revonsuo A, Laine M, Hamalainen H. Effects of electromagnetic field emitted by cellular phones on the EEG during an auditory memory task: a double blind replication study. *Bioelectromagnetics*. 25(1): 33-40, 2004.

The effects of electromagnetic fields (EMF) emitted by cellular phones on the event related desynchronization/synchronization (ERD/ERS) of the 4-6, 6-8, 8-10, and 10-12 Hz electroencephalogram (EEG) frequency bands were studied in 24 normal subjects performing an auditory memory task. This study was a systematic replication of our previous work. In the present double blind study, all subjects performed the memory task both with and without exposure to a digital 902 MHz field in a counterbalanced order. We were not able to replicate the findings from our earlier study. All eight of the significant changes in our earlier study were not significant in the present double blind replication. Also, the effect of EMF on the number of incorrect answers in the memory task was inconsistent. We previously reported no significant effect of EMF exposure on the number of incorrect answers in the memory task, but a significant increase in errors was observed in the present study. We conclude that EMF effects on the EEG and on the performance on memory tasks may be variable and not easily replicable for unknown reasons.

Köktürk S, Yardimoglu M, Celikozlu SD, Dolanbay EG, Cimbiz A. Effect of *Lycopersicon esculentum* extract on apoptosis in the rat cerebellum, following prenatal and postnatal exposure to an electromagnetic field. *Exp Ther Med*. 6(1):52-56, 2013.

The expansion of mobile phone technology has raised concerns regarding the effect of 900-MHz electromagnetic field (EMF) exposure on the central nervous system. At present, the developing human brain is regularly exposed to mobile telephones, pre- and postnatally. Several studies have demonstrated the acute effects of EMF exposure during pre- or postnatal periods; however, the chronic effects of EMF exposure are less understood. Thus, the aim of the present study was to determine the chronic effects of EMF on the pre- and postnatal rat cerebellum. The control group was maintained in the same conditions as the experimental groups, without the exposure to EMF. In the EMF1 group, the rats were exposed to EMF during pre- and postnatal periods (until postnatal day 80). In the EMF2 group, the rats were also exposed to EMF pre- and postnatally; in addition, however, they were provided with a daily oral supplementation of *Lycopersicon esculentum* extract (~2 g/kg). The number of caspase-3-labeled Purkinje neurons and granule cells present in the rats in the control and experimental groups were then counted. The neurodegenerative changes were studied using cresyl violet staining, and these changes were evaluated. In comparison with the control animals, the EMF1 group demonstrated a significant increase in the number of caspase-3-labeled Purkinje neurons and granule cells present in the cerebellum ($P < 0.001$). However, in comparison with the EMF1 group, the EMF2 group exhibited significantly fewer caspase-3-labeled Purkinje neurons and granule cells in the cerebellum. In the EMF1 group, the Purkinje neurons were revealed to have undergone dark neuron degenerative changes. However, the presence of dark Purkinje neurons was reduced in the EMF2 group,

Studies that show **Cell Phone** Health Effects

compared with the EMF1 group. The results indicated that apoptosis and neurodegeneration in rats exposed to EMF during pre- and postnatal periods may be reduced with *Lycopersicon esculentum* extract therapy.

Krause CM, Pesonen M, Haarala Bjornberg C, Hamalainen H. Effects of pulsed and continuous wave 902 MHz mobile phone exposure on brain oscillatory activity during cognitive processing. *Bioelectromagnetics*.28(4):296-308, 2007.

The aim of the current double-blind studies was to partially replicate the studies by Krause et al. [2000ab, 2004] and to further investigate the possible effects of electromagnetic fields (EMF) emitted by mobile phones (MP) on the event-related desynchronisation/synchronisation (ERD/ERS) EEG (electroencephalogram) responses during cognitive processing. Two groups, both consisting of 36 male participants, were recruited. One group performed an auditory memory task and the other performed a visual working memory task in six exposure conditions: SHAM (no EMF), CW (continuous wave EMF) and PM (pulse modulated EMF) during both left- and right-side exposure, while the EEG was recorded. In line with our previous studies, we observed that the exposure to EMF had modest effects on brain oscillatory responses in the alpha frequency range (approximately 8-12 Hz) and had no effects on the behavioural measures. The effects on the EEG were, however, varying, unsystematic and inconsistent with previous reports. We conclude that the effects of EMF on brain oscillatory responses may be subtle, variable and difficult to replicate for unknown reasons.

Imge EB, Kiliçoğlu B, Devrim E, Cetin R, Durak I. Effects of mobile phone use on brain tissue from the rat and a possible protective role of vitamin C - a preliminary study. *Int J Radiat Biol*.86(12):1044-1049, 2010.

Purpose: To evaluate effects of mobile phone use on brain tissue and a possible protective role of vitamin C. Materials and methods: Forty female rats were divided into four groups randomly (Control, mobile phone, mobile phone plus vitamin C and, vitamin C alone). The mobile phone group was exposed to a mobile phone signal (900 MHz), the mobile phone plus vitamin C group was exposed to a mobile phone signal (900 MHz) and treated with vitamin C administered orally (per os). The vitamin C group was also treated with vitamin C per os for four weeks. Then, the animals were sacrificed and brain tissues were dissected to be used in the analyses of malondialdehyde (MDA), antioxidant potential (AOP), superoxide dismutase, catalase (CAT), glutathione peroxidase (GSH-Px), xanthine oxidase, adenosine deaminase (ADA) and 5'-nucleotidase (5'-NT). Results: Mobile phone use caused an inhibition in 5'-NT and CAT activities as compared to the control group. GSH-Px activity and the MDA level were also found to be reduced in the mobile phone group but not significantly. Vitamin C caused a significant increase in the activity of GSH-Px and non-significant increase in the activities of 5'-NT, ADA and CAT enzymes. Conclusion: Our results suggest that vitamin C may play a protective role against detrimental effects of mobile phone radiation in brain tissue.

Studies that show **Cell Phone** Health Effects

Hountala CD, Maganioti AE, Papageorgiou CC, Nanou ED, Kyprianou MA, Tsiafakis VG, Rabavilas AD, Capsalis CN. The spectral power coherence of the EEG under different EMF conditions. Neurosci Lett. 441(2):188-192, 2008.

The present study introduces the concept of spectral power coherence (SPC), which reflects the pattern of coordination of the four basic EEG bands (delta, theta, alpha, and beta) at a specific location of the brain. The SPC was calculated for the pre-stimulus EEG signal during an auditory memory task under different electromagnetic field (EMF) conditions (900 MHz and 1800 MHz). The results showed that delta rhythm is less consequential in the overall cooperation between the bands than the higher frequency theta, alpha and beta rhythms. Additionally, it has been shown that the radiation effect on SPC is different for the two genders. In the absence of radiation males exhibit higher overall SPC than females. These differences disappear in the presence of 900 MHz and are reversed in the presence of 1800 MHz.

Ilhan A, Gurel A, Armutcu F, Kamisli S, Iraz M, Akyol O, Ozen S. Ginkgo biloba prevents mobile phone-induced oxidative stress in rat brain. Clin Chim Acta. 340(1-2): 153-162, 2004.

BACKGROUND: The widespread use of mobile phones (MP) in recent years has raised the research activities in many countries to determine the consequences of exposure to the low-intensity electromagnetic radiation (EMR) of mobile phones. Since several experimental studies suggest a role of reactive oxygen species (ROS) in EMR-induced oxidative damage in tissues, in this study, we investigated the effect of Ginkgo biloba (Gb) on MP-induced oxidative damage in brain tissue of rats. METHODS: Rats (EMR+) were exposed to 900 MHz EMR from MP for 7 days (1 h/day). In the EMR+Gb groups, rats were exposed to EMR and pretreated with Gb. Control and Gb-administrated groups were produced by turning off the mobile phone while the animals were in the same exposure conditions. Subsequently, oxidative stress markers and pathological changes in brain tissue were examined for each groups. RESULTS: Oxidative damage was evident by the: (i) increase in malondialdehyde (MDA) and nitric oxide (NO) levels in brain tissue, (ii) decrease in brain superoxide dismutase (SOD) and glutathione peroxidase (GSH-Px) activities and (iii) increase in brain xanthine oxidase (XO) and adenosine deaminase (ADA) activities. These alterations were prevented by Gb treatment. Furthermore, Gb prevented the MP-induced cellular injury in brain tissue histopathologically. CONCLUSION: Reactive oxygen species may play a role in the mechanism that has been proposed to explain the biological side effects of MP, and Gb prevents the MP-induced oxidative stress to preserve antioxidant enzymes activity in brain tissue.

Ingole IV, Ghosh SK. Effect of exposure to radio frequency radiation emitted by cell phone on the developing dorsal root ganglion of chick embryo: a light microscopic study. Nepal Med Coll J. 14(4):337-341, 2012.

With an ever increasing number of cell phone users since late twenty first century, magnitude of the problem of exposure to radiation emitted by cell phone is self evident. Extensive research had been devoted to incriminate or absolve it as a health hazard.

Studies that show **Cell Phone** Health Effects

Radiofrequency radiation emitted by cell phone had been stated to be a potent carcinogen, cytotoxic, genotoxic, mutagenic and neurobehavioral teratogen. Its effect on the brain had been a subject of extensive research evidently due to its proximity to the user's brain. While considering the biological effects of radiofrequency radiation, its intensity, frequency and the duration of exposure are important determinants. Nevertheless the results of these different studies have not been unequivocal. Considering the contradictory reports, the present work was undertaken to study the effect of such an exposure on the developing neural tissue of chick embryo. The processes of cell division and differentiation are fundamental to the development of any living being and are a sensitive index of any insult sustained at this stage. Neurons of dorsal root ganglion were selected for the present study as these ganglia were fully differentiated as early as fourth day of embryonic life. By varying duration of exposure, the embryos were exposed to different doses of radiation, sacrificed at different periods of incubation and subjected to histological processing. On light microscopic study it was observed that developing neurons of dorsal root ganglion suffered a damage which was dose dependent and persisted in spite of giving the exposure-free period between two exposures.

Haarala C, Aalto S, Hautzel H, Julkunen L, Rinne JO, Laine M, Krause B, Hamalainen H. Effects of a 902 MHz mobile phone on cerebral blood flow in humans: a PET study. Neuroreport. 14(16):2019-2023, 2003.

SUMMARY: Fourteen healthy right-handed subjects were scanned using PET with a [15O]water tracer during exposure to electromagnetic field (EMF) emitted by a mobile phone and a sham-exposure under double-blind conditions. During scanning, the subjects performed a visual working memory task. Exposure to an active mobile phone produced a relative decrease in regional cerebral blood flow (rCBF) bilaterally in the auditory cortex but no rCBF changes were observed in the area of maximum EMF. It is possible that these remote findings were caused by the EMF emitted by the active mobile phone. A more likely interpretation of the present findings were a result of an auditory signal from the active mobile phone. Therefore, it is not reasoned to attribute this finding to the EMF emitted by the phone. Further study on human rCBF during exposure to EMF of a mobile phone is needed.

Hamblin DL, Wood AW, Croft RJ, Stough C. Examining the effects of electromagnetic fields emitted by GSM mobile phones on human event-related potentials and performance during an auditory task. Clin Neurophysiol. 115(1):171-178, 2004.

OBJECTIVE: Due to the widespread use of mobile phones (MP), it is important to determine whether they affect human physiology. The aim of this study was to explore the sensitivity of auditory event-related potentials to electromagnetic emissions. METHODS: Twelve participants attended two sessions, 1 week apart. Participants performed an auditory oddball task while they were exposed to an active MP during one session and sham exposure during the other. Each condition lasted 1 h and order was counterbalanced. N100 and P200 latencies and amplitudes were analysed for non-target waveforms, and N200 and P300 latencies and amplitudes were analysed for target

Studies that show **Cell Phone** Health Effects

waveforms. RESULTS: In real relative to sham exposure N100 amplitude and latency to non-targets were reduced, with the reduction larger over midline and right hemisphere sites. P300 latency to targets was delayed in the real exposure condition, however as this difference was greatest at left frontal and left central sites the interpretation of this result is unclear. Reaction time increased in the real relative to sham condition. No difference in accuracy was found. CONCLUSIONS: The results suggest that MP exposure may affect neural activity, particularly in proximity to the phone, however caution should be applied due to the small sample size.

Gandhi OP, Lazzi G, Tinniswood A, Yu QS, Comparison of numerical and experimental methods for determination of SAR and radiation patterns of handheld wireless telephones. Bioelectromagnetics Suppl 4:93-101, 1999.

Some recent developments in both the numerical and experimental methods for determination of SARs and radiation patterns of handheld wireless telephones are described, with emphasis on comparison of results using the two methods. For numerical calculations, it was possible to use the Pro-Engineer CAD Files of cellular telephones for a realistic description of the device. Also, we used the expanding grid formulation of the finite-difference time-domain (FDTD) method for finer-resolution representation of the coupled region, including the antenna, and an increasingly coarser representation of the more-distant, less-coupled region. Together with the truncation of the model of the head, this procedure led to a saving of computer memory needed for SAR calculations by a factor of over 20. Automated SAR and radiation pattern measurement systems were used to validate both the calculated 1-g SARs and radiation patterns for several telephones, including some research test samples, using a variety of antennas. Even though widely different peak 1-g SARs were obtained, ranging from 0.13 to 5.41 W/kg, agreement between the calculated and the measured data for these telephones, five each at 835 and 1900 MHz, was excellent and generally within +/-20% (+/-1 dB). An important observation was that for a maximum radiated power of 600 mW at 800/900 MHz, which may be used for telephones using AMPS technology, the peak 1-g SARs can be higher than 1.6 W/kg unless antennas are carefully designed and placed further away from the head.

Frey AH, Headaches from cellular telephones: are they real and what are the implications? Environ Health Perspect 106(3):101-103, 1998.

There have been numerous recent reports of headaches occurring in association with the use of hand-held cellular telephones. Are these reported headaches real? Are they due to emissions from telephones? There is reason to believe that the answer is "yes" to both questions. There are several lines of evidence to support this conclusion. First, headaches as a consequence of exposure to low intensity microwaves were reported in the literature 30 years ago. These were observed during the course of microwave hearing research before there were cellular telephones. Second, the blood-brain barrier appears to be involved in headaches, and low intensity microwave energy exposure affects the barrier. Third, the dopamine-opiate systems of the brain appear to be involved in headaches, and low intensity electromagnetic energy exposure affects those

Studies that show **Cell Phone** Health Effects

systems. In all three lines of research, the microwave energy used was approximately the same—in frequencies, modulations, and incident energies—as those emitted by present day cellular telephones. Could the current reports of headaches be the canary in the coal mine, warning of biologically significant effects?

Freude, G, Ullsperger, P, Eggert, S, Ruppe, I, Microwaves emitted by cellular telephones affect human slow brain potentials. Eur J Appl Physiol 81(1-2):18-27, 2000.

The influence of electromagnetic fields (EMF) emitted by cellular telephones on preparatory slow brain potentials (SP) was studied in two experiments, about 6 months apart. In the first experiment, a significant decrease of SP was found during exposure to EMF in a complex visual monitoring task (VMT). This effect was replicated in the second experiment. In addition to the VMT, EMF effects on SP were analysed in two further, less demanding tasks: in a simple finger movement task to elicit a Bereitschaftspotential (BP) and in a two-stimulus task to elicit a contingent negative variation (CNV). In comparison to the VMT, no significant main EMF effects were found in BP and CNV tasks. The results accounted for a selective EMF effect on particular aspects of human information processing, but did not indicate any influence on human performance, well-being and health.

Freude, G, Ullsperger, P, Eggert, S, Ruppe, I, Effects of microwaves emitted by cellular phones on human slow brain potentials. Bioelectromagnetics 19(6):384-387, 1998.

The influence of electromagnetic fields (EMF) emitted by cellular phones on preparatory slow brain potentials (SP) was studied in two different experimental tasks: In the first, healthy male human subjects had to perform simple self-paced finger movements to elicit a Bereitschaftspotential; in the second, they performed a complex and cognitive demanding visual monitoring task (VMT). Both tasks were performed with and without EMF exposure in counterbalanced order. Whereas subjects' performance did not differ between the EMF exposure conditions, SP parameters were influenced by EMF in the VMT: EMF exposure effected a significant decrease of SPs at central and temporo-parieto-occipital brain regions, but not at the frontal one. In the simple finger movement task, EMF did not affect the Bereitschaftspotential.

Finnie JW, Blumbergs PC, Manavis J, Utteridge TD, Gebiski V, Davies RA, Vernon-Roberts B, Kuchel TR. Effect of long-term mobile communication microwave exposure on vascular permeability in mouse brain. Pathology 34(4):344-347, 2002.

AIMS: To study the effect of long-term exposure to global system for mobile communication (GSM) radiofrequency fields on vascular permeability in murine brains. METHODS: Using a purpose-designed exposure system at 900 MHz, mice were given a 60-minute far-field, whole body exposure on each of 5 days per week for 104 weeks at specific absorption rates (SAR) of 0.25, 1.0, 2.0 and 4.0 W/kg. Control mice were sham-exposed or permitted free movement in a cage to evaluate any stress-related effects. Albumin immunohistochemistry was used to detect increased vascular permeability and the efficacy of the vascular tracer was confirmed with a positive control group exposed to a clostridial toxin known to increase vascular permeability in the brain. RESULTS: In all

Studies that show **Cell Phone** Health Effects

exposed and control groups, albumin extravasation was minimal, often leptomeningeal, and was deemed insignificant as a maximum of three capillaries or venules in a given brain showed leakage from the very many blood vessels present in the three coronal brain sections. CONCLUSIONS: These results suggest that prolonged exposure to mobile telephone-type radiation produces negligible disruption to blood-brain barrier integrity at the light microscope level using endogenous albumin as a vascular tracer.

Ferreri F, Curcio G, Pasqualetti P, De Gennaro L, Fini R, Rossini PM. Mobile phone emissions and human brain excitability. *Ann Neurol*.60(2):188-196, 2006.

OBJECTIVE: To test-via Transcranial Magnetic Stimulation (TMS)-the excitability of each brain hemisphere after 'real' or 'sham' exposure to the electromagnetic field (EMF) generated by a mobile phone operating in the Global System for Mobile Communication (GSM). METHODS: Fifteen male volunteers attended two experimental sessions, one week apart, in a cross-over, double-blind paradigm. In one session the signal was turned ON (EMF-on, real exposure), in the other it was turned OFF (EMF-off, sham exposure), for 45 minutes. Motor Evoked Potentials (MEPs) were recorded using a paired-pulse paradigm (testing intracortical excitability with 1 to 17 ms interstimulus intervals), both before and at different times after exposure to the EMF. Short Intracortical Inhibition (SICI) and Facilitation (ICF) curves were evaluated both on the exposed and non-exposed hemispheres. Tympanic temperature was collected during each session. RESULTS: The intracortical excitability curve becomes significantly modified during real exposure, with SICI being reduced and ICF enhanced in the acutely exposed brain hemisphere as compared to the contralateral, non-exposed hemisphere or to sham exposure. Tympanic temperature showed no significant main effect or interactions. INTERPRETATION: These results demonstrate that GSM-EMFs modify brain excitability. Possible implications and applications are discussed.

Fayos-Fernandez J, Arranz-Faz C, Martinez-Gonzalez AM, Sanchez-Hernandez D. Effect of pierced metallic objects on sar distributions at 900 MHz. *Bioelectromagnetics*. 27(5):337-353, 2006.

. A study of the interaction between mobile phone antennas and a human head in the presence of different types of metallic objects, attached and pierced to the compressed ear, is presented in this article. Computed and measured results have been performed by considering a quasi-half-wavelength dipole as the radiating source and measurements with the DASY4 dosimetric assessment system. Two different human head models have been implemented: a homogeneously shaped sphere and a three-level head model with four different kinds of tissue. Antenna input impedance, reflection coefficient, radiation patterns, SAR distribution, absorbed power, and peak SAR values have been computed and measured for diverse scenarios, electromagnetic simulators, and organs. Despite the measuring accuracy limitations of the study, both simulated and measured results suggest that special attention has to be paid to peak SAR averaged values when wearing metallic objects close to the radiation source, since some increment of peak SAR averaged values is expected.

Studies that show **Cell Phone** Health Effects

Faucon G, Le Bouquin Jeannes R, Maby E. Short-term effects of GSM mobiles phones on spectral components of the human electroencephalogram. Conf Proc IEEE Eng Med Biol Soc. 1(1):3751-3754, 2006.

The aim of the study was to investigate whether the GSM (global system for mobile) signals affect the electrical activity of the human brain. Nine healthy subjects and six temporal epileptic patients were exposed to radiofrequencies emitted by a GSM mobile phone signals. Electroencephalographic (EEG) signals were recorded using surface electrodes with and without radiofrequency. In order to obtain a reference, a control session was also carried out. The spectral attributes of the EEG signals recorded by surface electrodes were analyzed. The significant decrease of spectral correlation coefficients under radiofrequency influence showed that the GSM signal altered the spectral arrangement of the EEG activity for healthy subjects as well as epileptic patients. For the healthy subjects, the EEG spectral energy decreased on the studied frequency band [0-40 Hz] and more precisely on occipital electrodes for the alpha-band. For the epileptic patients, these modifications were demonstrated by an increase of the power spectral density of the EEG signal. Nevertheless, these biological effects on the EEG are not sufficient to put forward some electrophysiological hypothesis.

Eulitz, C, Ullsperger, P, Freude, G, Elbert ,T, Mobile phones modulate response patterns of human brain activity. Neuroreport 9(14):3229-3232, 1998.

Mobile phones emit a pulsed high-frequency electromagnetic field (PEMF) which may penetrate the scalp and the skull. Increasingly, there is an interest in the interaction of this pulsed microwave radiation with the human brain. Our investigations show that these electromagnetic fields alter distinct aspects of the brain's electrical response to acoustic stimuli. More precisely, our results demonstrate that aspects of the induced but not the evoked brain activity during PEMF exposure can be different from those not influenced by PEMF radiation. This effect appears in higher frequency bands when subjects process task-relevant target stimuli but was not present for irrelevant standard stimuli. As the induced brain activity in higher frequency bands has been proposed to be a correlate of coherent high-frequency neuronal activity, PEMF exposure may provide means to systematically alter the pattern fluctuations in neural mass activity.

Eberhardt JL, Persson BR, Brun AE, Salford LG, Malmgren LO. Blood-brain barrier permeability and nerve cell damage in rat brain 14 and 28 days after exposure to microwaves from GSM mobile phones. Electromagn Biol Med. 27(3):215-229, 2008.

We investigated the effects of global system for mobile communication (GSM) microwave exposure on the permeability of the blood-brain barrier and signs of neuronal damage in rats using a real GSM programmable mobile phone in the 900 MHz band. Ninety-six non-anaesthetized rats were either exposed to microwaves or sham exposed in TEM-cells for 2 h at specific absorption rates of average whole-body Specific Absorption Rates (SAR) of 0.12, 1.2, 12, or 120 mW/kg. The rats were sacrificed after a recovery time of either 14 or 28 d, following exposure and the extravasation of albumin, its uptake into neurons, and occurrence of damaged neurons was assessed. Albumin

Studies that show **Cell Phone** Health Effects

extravazation and also its uptake into neurons was seen to be enhanced after 14 d (Kruskal Wallis test: $p = 0.02$ and 0.002 , respectively), but not after a 28 d recovery period. The occurrence of dark neurons in the rat brains, on the other hand, was enhanced later, after 28 d ($p = 0.02$). Furthermore, in the 28-d brain samples, neuronal albumin uptake was significantly correlated to occurrence of damaged neurons (Spearman $r = 0.41$; $p < 0.01$).

de Tommaso M, Rossi P, Falsaperla R, Francesco VD, Santoro R, Federici A. Mobile Phones exposure induces changes of Contingent Negative Variation in humans. *Neurosci Lett.* 464(2):79-83, 2009.

Event related potentials have been largely employed to test effects of GSM emissions on human brain. The aim of the present study, was the evaluation of initial Contingent Negative Variation (iCNV) changes, induced by 900MHz GSM exposure, in a double blind design in healthy volunteers, subjected to a threefold experimental condition, EXPOSED (A), a real GSM phone emitting electromagnetic power, SHAM (B), a real phone where the electromagnetic power was dissipated on an internal load and OFF (C), a phone completely switched off. Ten healthy right-handed volunteers were evaluated. The CNV was recorded during a 10minutes time interval in each of the three experimental conditions A, B, and C, in order to assess the iCNV amplitude and habituation. The iCNV amplitude decreased and habituation increased during both A and B conditions, compared with condition C. This effect was diffuse over the scalp, and there was no significant prevalence of iCNV amplitude reduction on the left side, where the phones were located. Mobile Phones exposures A and B seemed to act on brain electrical activity, reducing the arousal and expectation of warning stimulus. This evidence, limited by the low number of subjects investigated, could be explained in terms of an effect induced by both the GSM signal and the Extremely Low Frequency magnetic field produced by battery and internal circuits.

Dasdag S, Akdag MZ, Ulukaya E, Uzunlar AK, Ocak AR. Effect of mobile phone exposure on apoptotic glial cells and status of oxidative stress in rat brain. *Electromagn Biol Med.* 28(4):342-354, 2009.

The aim of this study was to investigate the effects of mobile phone exposure on glial cells in brain. The study carried out on 31 Wistar Albino adult male rats. The rat heads in a carousel exposed to 900 MHz microwave. For the study group (n:14), rats exposed to the radiation 2 h per day (7 days in a week) for 10 months. For the sham group (n:7), rats were placed into the carousel and the same procedure was applied except that the generator was turned off. For the cage control (n:10), nothing applied to rats in this group. In this study, rats were euthanized after 10 months of exposure periods and brains were removed. Brain tissues were immunohistochemically stained for the active (cleaved) caspase-3, which is a well-known apoptosis marker, and p53. The expression of the proteins was evaluated by a semi-quantitative scoring system. However, total antioxidative capacity (TAC), catalase, total oxidant status (TOS), and oxidative stress index were measured in rat brain. Final score for apoptosis in the exposed group was significantly lower than the sham ($p < 0.001$) and the cage control groups ($p < 0.01$). p53

Studies that show **Cell Phone** Health Effects

was not significantly changed by the exposure ($p > 0.05$). The total antioxidant capacity and catalase in the experimental group was found higher than that in the sham group ($p < 0.001$, $p < 0.05$). In terms of the TOS and oxidative stress index, there was no statistically significant difference between exposure and sham groups ($p > 0.05$). In conclusion, the final score for apoptosis, total antioxidant capacity and catalase in rat brain might be altered by 900 MHz radiation produced by a generator to represent exposure of global systems for mobile communication (GSM) cellular phones.

Croft RJ, Hamblin DL, Spong J, Wood AW, McKenzie RJ, Stough C. The effect of mobile phone electromagnetic fields on the alpha rhythm of human electroencephalogram. *Bioelectromagnetics*.29(1):1-10,2008.

Mobile phones (MP) emit low-level electromagnetic fields that have been reported to affect neural function in humans; however, demonstrations of such effects have not been conclusive. The purpose of the present study was to test one of the strongest findings in the literature; that of increased "alpha" power in response to MP-type radiation. Healthy participants ($N = 120$) were tested using a double-blind counterbalanced crossover design, with each receiving a 30-min Active and a 30-min Sham Exposure 1 week apart, while electroencephalogram (EEG) data were recorded. Resting alpha power (8-12 Hz) was then derived as a function of time, for periods both during and following exposure. Non-parametric analyses were employed as data could not be normalized. Previous reports of an overall alpha power enhancement during the MP exposure were confirmed (relative to Sham), with this effect larger at ipsilateral than contralateral sites over posterior regions. No overall change to alpha power was observed following exposure cessation; however, there was less alpha power contralateral to the exposure source during this period (relative to ipsilateral). Employing a strong methodology, the current findings support previous research that has reported an effect of MP exposure on EEG alpha power.

Chia SE, Chia HP, Tan JS, Prevalence of headache among handheld cellular telephone users in singapore: A community study. *Environ Health Perspect* 108(11):1059-1062, 2000.

We carried out a cross-sectional community study in Singapore to determine the prevalence of specific central nervous system (CNS) symptoms among hand-held cellular telephone (HP) users compared to nonusers and to study the association of risk factors and CNS symptoms among HP users. A total of 808 men and women between 12 and 70 years of age, who lived in one community, were selected using one-stage cluster random sampling and responses to a structured questionnaire. The prevalence of HP users was 44.8%. Headache was the most prevalent symptom among HP users compared to non-HP users, with an adjusted prevalence rate ratio of 1.31 [95% confidence interval, 1.00-1.70]. There is a significant increase in the prevalence of headache with increasing duration of usage (in minutes per day). Prevalence of headache was reduced by more than 20% among those who used hand-free equipment for their cellular telephones as compared to those who never use the equipment. The use of HPs is not associated with a significant increase of CNS symptoms other than

Studies that show **Cell Phone** Health Effects

headache.

Celikozlu SD, Ozyurt MS, Cimbiz A, Yardimoglu MY, Cayci MK, Ozay Y. The effects of long-term exposure of magnetic field via 900-MHz GSM radiation on some biochemical parameters and brain histology in rats. Electromagn Biol Med. 31(4):344-355, 2012.

The aim of this study is to determine the effects of magnetic field via cell phones on some blood parameters and neurons in the brain of rats. Animals have been classified into three groups: control, Magnetic Field (MF), and F2 groups. Throughout this study, cell phones were placed on the wall of the cages. Rats were exposed to the effects of cell phones during prenatal and postnatal periods until they were 80 days old. During the study, the exposure procedure of rats was that the phone was in standby mode for a whole day and in talking mode for 30 min per day. The waves of cell phones caused an increased blood glucose level from 96.52 ± 5.64 mg/dl to 132.14 ± 5.93 mg/dl and an increased serum protein level from 131.14 ± 6.19 mg/dl to 319.29 ± 6.73 mg/dl compared to control. Statistically, significant differences wasn't observed in the blood cholesterol concentration between the groups compared to the control. Weekly weight gain decreased in all groups compared to the control. MF exposure decreased pyramidal neuron numbers 51.15% and increased ischemic neuron numbers 73% at cortex region of brain. In addition, vascular dilatations have increased clearly in group F2. Whereas the procedure of MF did not have any effects on hippocampal pyramidal cell numbers, magnetic fields increased the amount of ischemic neurons three-fold compared to the control. In conclusion, MF affected some biochemical parameters, especially the cortex region of the brain.

Carrubba S, Frilot C 2nd, Chesson AL Jr, Marino AA. Mobile-phone pulse triggers evoked potentials. Neurosci Lett. 469(1):164-168, 2010.

If mobile-phone electromagnetic fields (EMFs) are hazardous, as suggested in the literature, processes or mechanisms must exist that allow the body to detect the fields. We hypothesized that the low-frequency pulses produced by mobile phones (217Hz) were detected by sensory transduction, as evidenced by the ability of the pulses to trigger evoked potentials (EPs). Electroencephalograms (EEGs) were recorded from six standard locations in 20 volunteers and analyzed to detect brain potentials triggered by a pulse of the type produced by mobile phones. Evoked potentials having the expected latency were found in 90% of the volunteers, as assessed using a nonlinear method of EEG analysis. Evoked potentials were not detected when the EEG was analyzed using time averaging. The possibility of systematic error was excluded by sham-exposure analyses. The results implied that mobile-phones trigger EP at the rate of 217Hz during ordinary phone use. Chronic production of the changes in brain activity might be pertinent

Carballo-Quintás M, Martínez-Silva I, Cadarso-Suárez C, Alvarez-Figueiras M, Ares-Pena FJ, López-Martín E. A study of neurotoxic biomarkers, c-fos and GFAP after acute

Studies that show **Cell Phone** Health Effects

exposure to GSM radiation at 900 MHz in the picrotoxin model of rat brains.

Neurotoxicology. 32(4):478-494, 2011.

The acute effects of microwave exposure from the Global System for Mobile Communication (GSM) were studied in rats, using 900MHz radiation at an intensity similar to mobile phone emissions. Acute subconvulsive doses of picrotoxin were then administered to the rats and an experimental model of seizure-proneness was created from the data. Seventy-two adult male Sprague-Dawley rats underwent immunochemical testing of relevant anatomical areas to measure induction of the c-fos neuronal marker after 90min and 24h, and of the glial fibrillary acidic protein (GFAP) 72h after acute exposure to a 900MHz electromagnetic field (EMF). The experimental set-up facilitated measurement of absorbed power, from which the average specific absorption rate was calculated using the finite-difference time-domain (FDTD) 2h after exposure to EMF radiation at 1.45W/kg in picrotoxin-treated rats and 1.38W/kg in untreated rats. Ninety minutes after radiation high levels of c-fos expression were recorded in the neocortex and paleocortex along with low hippocampus activation in picrotoxin treated animals. Most brain areas, except the limbic cortical region, showed important increases in neuronal activation 24h after picrotoxin and radiation. Three days after picrotoxin treatment, radiation effects were still apparent in the neocortex, dentate gyrus and CA3, but a significant decrease in activity was noted in the piriform and entorhinal cortex. During this time, glial reactivity increased with every seizure in irradiated, picrotoxin-treated brain regions. Our results reveal that c-fos and glial markers were triggered by the combined stress of non-thermal irradiation and the toxic effect of picrotoxin on cerebral tissues.

Brillaud E, Piotrowski A, de Seze R. Effect of an acute 900MHz GSM exposure on glia in the rat brain: A time-dependent study. Toxicology.238(1):23-33,2007.

Because of the increasing use of mobile phones, the possible risks of radio frequency electromagnetic fields adverse effects on the human brain has to be evaluated. In this work we measured GFAP expression, to evaluate glial evolution 2, 3, 6 and 10 days after a single GSM exposure (15min, brain averaged SAR=6W/kg, 900MHz signal) in the rat brain. A statistically significant increase of GFAP stained surface area was observed 2 days after exposure in the frontal cortex and the caudate putamen. A smaller statistically significant increase was noted 3 days after exposure in the same areas and in the cerebellum cortex. Our results confirm the Mausset-Bonnefont et al. study [Mausset-Bonnefont, A.L., Hirbec, H., Bonnefont, X., Privat, A., Vignon, J., de Seze, R., 2004. Acute exposure to GSM 900MHz electromagnetic fields induces glial reactivity and biochemical modifications in the rat brain. Neurobiol. Dis. 17, 445-454], showing the existence of glial reactivity after a 15min GSM acute exposure at a brain averaged SAR of 6W/kg. We conclude to a temporary effect, probably due to a hypertrophy of glial cells, with a temporal and a spatial modulation of the effect. Whether this effect could be harmful remains to be studied.

Bortkiewicz A, Gadzicka E, Szymczak W, Zmyślony M. Changes in tympanic temperature during the exposure to electromagnetic fields emitted by mobile phone.

Studies that show Cell Phone Health Effects

Int J Occup Med Environ Health. 25(2): 145-150, 2012.

OBJECTIVE: Mobile phones generate microwave radiation which is absorbed by exposed tissue and converted into heat. It may cause detrimental health effects. The aim of the experiment was to check if exposure to EMF emitted by mobile phone influenced the tympanic temperature. **MATERIAL AND METHODS:** Human volunteer study was performed on ten healthy young men, aged 22.1 ± 4.7 years, examined three times: 1. on a day with 2×60 min of no exposure (sham day), 2. on a day with continuous, 60 min exposure and 60 min of no exposure, 3. on a day with intermittent exposure (4×15 min "on" and 4×15 min "off"). Exposure was generated by mobile phone (frequency 900 MHz, SAR 1.23 W/kg). The study was double-blind, performed under controlled conditions (at 24°C and 70% humidity). The tympanic temperature ($T(\text{ty})$) was monitored every 10 sec by a thermistor probe placed close to the aural canal membrane in the ear opposite the one in contact with mobile phone (contralateral position). Multivariate repeated-measures analysis of variance was used to calculate the results. **RESULTS:** The mean $T(\text{ty})$ in the whole group during continuous exposure was significantly higher than during sham exposure ($p = 0.0001$). During intermittent exposure the temperature was lower than during sham day (difference was up to 0.11°C). Within an hour after continuous exposure, $T(\text{ty})$ was higher by 0.03°C and after intermittent exposure $T(\text{ty})$ was lower by 0.18°C in comparison with sham day. Two hours after exposure $T(\text{ty})$ was significantly lower ($p = 0.0001$) than after sham exposure (0.06°C and 0.26°C respectively). The trends in $T(\text{ty})$ during experiment differed significantly in relation to exposure conditions ($p < 0.05$). **CONCLUSIONS:** The results of this analysis indicate that the physiological response to EMF exposure from mobile phone was mostly related to type of exposure (continuous or intermittent).

Bilgici B, Akar A, Avci B, Tuncel OK. Effect of 900 MHz radiofrequency radiation on oxidative stress In rat brain and serum. Electromagn Biol Med. 32(1):20-29, 2013.

The increasing use of mobile telephones raises the question of possible adverse effects of the electromagnetic fields (EMF) that these phones produce. In this study, we examined the oxidative stress in the brain tissue and serum of rats that resulted from exposure to a 900-MHz EMF at a whole body average specific absorption rate (SAR) of 1.08 W/kg for 1 h/day for 3 weeks. We also examined the antioxidant effect of garlic powder (500 mg/kg/day) given orally to EMF-exposed rats. We found that malondialdehyde (MDA) ($p < 0.001$) and advanced oxidation protein product (AOPP) ($p < 0.05$) increased in rat brain tissue exposed to the EMF and that garlic reduced these effects ($p < 0.05$). There was no significant difference in the nitric oxide (NO) levels in the brain. Paraoxonase (PON) was not detected in the brain. There was a significant increase in the levels of NO ($p < 0.001$) detected in the serum after EMF exposure, and garlic intake did not affect this increase in NO. Our results suggest that there is a significant increase in brain lipid and protein oxidation after electromagnetic radiation (EMR) exposure and that garlic has a protective effect against this oxidative stress.

Bachmann M, Rubljova J, Lass J, Tomson R, Tuulik V, Hinrikus H. Adaptation of human brain bioelectrical activity to low-level microwave. Conf Proc IEEE Eng Med Biol Soc. 2007:4747-4750, 2007.

Studies that show **Cell Phone** Health Effects

The experiments of adaptation of the human brain bioelectrical activity were carried out on a group of 14 healthy volunteers exposed to 450 MHz microwave radiation modulated at 40 Hz frequencies. The field power density at the scalp was 0.16 mW/cm². Results of the study indicate that adaptation effect of human brain to low-level microwave exposure is evident. The initial increase of EEG power was compensated and even overcompensated. The adaptation phenomena were obvious in EEG alpha and beta rhythms.

Barcal J, Cendelín J, Vozech F, Zalud V. Effect of whole-body exposure to high-frequency electromagnetic field on the brain electrogeny in neurodefective and healthy mice. Prague Med Rep. 106(1):91-100, 2005.

A direct registration of brain cortical and hippocampal activity during a high-frequency electromagnetic field (HF EMF) exposure was performed. All experimental procedures were done under urethane anaesthesia (20%, 2 g/kg i.p.) in Lurcher mutant mice, wild type (healthy littermates) were used as controls. Experimental animals were exposed to the HF EMF with frequency corresponding to cellular phones. Our method is based on the use of gel electrodes (silicon tubes or glass microcapillaries filled with agar) where the connection with classical electrodes is located out of HF EMF space. ECoG evaluation showed a distinct shift to lower frequency components but clear effect has been observed only in wild type (healthy) mice whereas in Lurcher mutant mice only gentle differences between frequency spectra were found. Measurement of hippocampal rhythmicity showed gentle changes with increase of higher frequencies (i.e. opposite effect than in cortex) and changes in theta oscillations registered from a dentate gyrus and CA1 area in both types of animals (healthy and mutant). These findings support the idea about possible influencing the central nervous system by HF EMF exposure and support also some recent results about possible health risks resulting from cellular phones use.

Barcal J, Vozech F. Effect of whole-body exposure to high-frequency electromagnetic field on the brain cortical and hippocampal activity in mouse experimental model. NeuroQuantology 5:292-302, 2007.

Evaluation of the direct registration of brain cortical and hippocampal activity during a high-frequency electromagnetic field (HF-EMF) exposure was performed. Experimental procedures were done under general anesthesia (urethane, 20%, 2g/kg i.p.) in Lurcher mutant mice, wild type (healthy littermates) were used as controls. Animals were exposed to the HF-EMF with frequency corresponding to cellular phones (900 MHz). We used of gel electrodes (silicon tubes or glass microcapillary filled with agar) where the connection with classical electrodes was located out of HF-EMF space. ECoG evaluation showed a distinct shift to lower frequency components but clear effect has been observed only in wild type (healthy) mice whereas in Lurcher mutant mice only gentle differences between frequency spectra were found. Measurement of hippocampal rhythmicity showed gentle changes with increase of higher frequencies (i.e. opposite effect than in cortex) and changes in theta oscillations registered from a dentate gyrus and CA1 area in both types of animals (healthy and mutant). These findings support an

Studies that show **Cell Phone** Health Effects

idea about possible influencing the central nervous system by HF-EMF exposure and support also some recent results about possible health risks resulting from cellular phones use.

Bas O, Odaci E, Kaplan S, Acer N, Ucok K, Colakoglu S. 900 MHz electromagnetic field exposure affects qualitative and quantitative features of hippocampal pyramidal cells in the adult female rat. Brain Res. 1265:178-185, 2009.

The effects of electromagnetic fields (EMFs) emitted by mobile phones on humans hold special interest due to their use in close proximity to the brain. The current study investigated the number of pyramidal cells in the cornu ammonis (CA) of the 16-week-old female rat hippocampus following postnatal exposure to a 900 megahertz (MHz) EMF. In this study were three groups of 6 rats: control (Cont), sham exposed (Sham), and EMF exposed (EMF). EMF group rats were exposed to 900 MHz EMF (1 h/day for 28 days) in an exposure tube. Sham group was placed in the exposure tube but not exposed to EMF (1 h/day for 28 days). Cont group was not placed into the exposure tube nor were they exposed to EMF during the study period. In EMF group rats, the specific energy absorption rate (SAR) varied between 0.016 (whole body) and 2 W/kg (locally in the head). All of the rats were sacrificed at the end of the experiment and the number of pyramidal cells in the CA was estimated using the optical fractionator technique. Histopathological evaluations were made on sections of the CA region of the hippocampus. Results showed that postnatal EMF exposure caused a significant decrease of the pyramidal cell number in the CA of the EMF group ($P < 0.05$). Additionally, cell loss can be seen in the CA region of EMF group even at qualitative observation. These results may encourage researchers to evaluate the chronic effects of 900 MHz EMF on teenagers' brains.

Behari J, Kunjilwar KK, and Pyne S, Interaction of low level modulated RF radiation with $\text{Na}^+\text{-K}^+\text{-ATPase}$. Bioelectrochem Bioenerg 47:247-252, 1998.

The effect of low-level amplitude modulated radiofrequency radiation were studied on $\text{Na}^+\text{-K}^+\text{-ATPase}$ activity in the brain of developing male Wistar rats of age 23 days (body weight 55-60 g). They were exposed to carrier wave (CW) frequency 147 MHz and its sub-harmonic frequencies 73.5 and 36.75 MHz amplitude modulated (AM) at 16 and 76 Hz for 30-35 days (3 h day^{-1} , Power density 1.47 mW cm^{-2} , average specific absorption rate $9.65\text{-}6.11 \text{ W kg}^{-1}$). We observed a statistically significant increase in $\text{Na}^+\text{-K}^+\text{-ATPase}$ activity in chronically exposed rats compared to the control ones. The increase in $\text{Na}^+\text{-K}^+\text{-ATPase}$ activity was around 19-20% in the rats exposed to CW frequencies AM at 16 Hz compared to the controls, whereas the increase in $\text{Na}^+\text{-K}^+\text{-ATPase}$ activity was around 15-16% in rats exposed to the same set of CW frequencies but AM at 76 Hz. Though there was a difference in $\text{Na}^+\text{-K}^+\text{-ATPase}$ activities (3-4%) in the two groups but the difference was found to be statistically insignificant. Within the group of rats exposed to CW frequencies amplitude modulated at 16 and 76 Hz, respectively, the effect on $\text{Na}^+\text{-K}^+\text{-ATPase}$ activity was found to be independent of the magnitude of CW frequencies. An additional single short duration (20-60 min) exposure of membranes in vitro from different exposed group to the above field did not show any significant

Studies that show **Cell Phone** Health Effects

alteration on $\text{Na}^+\text{-K}^+$ -ATPase activity. It is concluded that a low level effect of amplitude modulated radiation produces statistically significant effect on $\text{Na}^+\text{-K}^+$ -ATPase activity but is insensitive to the carrier wave frequencies under investigation.

Zhou H, Su Z, Ning J, Wang C, Xie X, Qu D, Wu K, Zhang X, Pan J, Yang G. EFFECTS OF FREQUENCY, IRRADIATION GEOMETRY AND POLARISATION ON COMPUTATION OF SAR IN HUMAN BRAIN. Radiat Prot Dosimetry. 2014 Jan 6. [Epub ahead of print]

The power absorbed by the human brain has possible implications in the study of the central nervous system-related biological effects of electromagnetic fields. In order to determine the specific absorption rate (SAR) of radio frequency (RF) waves in the human brain, and to investigate the effects of geometry and polarisation on SAR value, the finite-difference time-domain method was applied for the SAR computation. An anatomically realistic model scaled to a height of 1.70 m and a mass of 63 kg was selected, which included 14 million voxels segmented into 39 tissue types. The results suggested that high SAR values were found in the brain, i.e. ~250 MHz for vertical polarisation and 900-1200 MHz both for vertical and horizontal polarisation, which may be the result of head resonance at these frequencies.

Mausset A, de Seze R, Montpeyroux F, Privat A. Effects of radiofrequency exposure on the GABAergic system in the rat cerebellum: clues from semi-quantitative immunohistochemistry. Brain Res 912(1):33-46, 2001.

The widespread use of cellular phones raises the problem of interaction of electromagnetic fields with the central nervous system (CNS). In order to measure these effects on neurotransmitter content in the CNS, we developed a protocol of neurotransmitter detection based on immunohistochemistry and image analysis. Gamma-vinyl-GABA (GVG), an inhibitor of the GABA-transaminase was injected in rats to increase GABA concentration in the CNS. The cellular GABA contents were then revealed by immunohistochemistry and semi-quantified by image analysis thanks to three parameters: optical density (O.D.), staining area, and number of positive cells. The increase in cerebellar GABA content induced by GVG 1200 mg/kg was reflected in these three parameters in the molecular and the granular layers. Therefore, control of immunohistochemistry parameters, together with appropriate image analysis, allowed both the location and the detection of variations in cellular neurotransmitter content. This protocol was used to investigate the effects of exposure to 900 MHz radiofrequencies on cerebellar GABA content. Both pulsed emission with a specific absorption rate (SAR) of 4 W/kg and continuous emission with high SAR (32 W/kg) were tested. We observed a selective diminution of the stained processes area in the Purkinje cell layer after exposure to pulsed radiofrequency and, in addition, a decrease in O.D. in the three cell layers after exposure to continuous waves. Whether this effect is, at least partly, due to a local heating of the tissues is not known. Overall, it appears that high energetic radiofrequency exposure induces a diminution in cellular GABA content in the cerebellum.

Studies that show **Cell Phone** Health Effects

Croft RJ, Leung S, McKenzie RJ, Loughran SP, Iskra S, Hamblin DL, Cooper NR. Effects of 2G and 3G mobile phones on human alpha rhythms: Resting EEG in adolescents, young adults, and the elderly. *Bioelectromagnetics*. 31(6):434-444, 2010.

The present study was conducted to determine whether adolescents and/or the elderly are more sensitive to mobile phone (MP)-related bioeffects than young adults, and to determine this for both 2nd generation (2G) GSM, and 3rd generation (3G) W-CDMA exposures. To test this, resting alpha activity (8-12 Hz band of the electroencephalogram) was assessed because numerous studies have now reported it to be enhanced by MP exposure. Forty-one 13-15 year olds, forty-two 19-40 year olds, and twenty 55-70 year olds were tested using a double-blind crossover design, where each participant received Sham, 2G and 3G exposures, separated by at least 4 days. Alpha activity, during exposure relative to baseline, was recorded and compared between conditions. Consistent with previous research, the young adults' alpha was greater in the 2G compared to Sham condition, however, no effect was seen in the adolescent or the elderly groups, and no effect of 3G exposures was found in any group. The results provide further support for an effect of 2G exposures on resting alpha activity in young adults, but fail to support a similar enhancement in adolescents or the elderly, or in any age group as a function of 3G exposure.

Central Nervous System Effects

Xu S, Zhou Z, Zhang L, Yu Z, Zhang W, Wang Y, Wang X, Li M, Chen Y, Chen C, He M, Zhang G, Zhong M. Exposure to 1800 MHz radiofrequency radiation induces oxidative damage to mitochondrial DNA in primary cultured neurons. *Brain Res*. 1311:189-196. 2010.

Increasing evidence indicates that oxidative stress may be involved in the adverse effects of radiofrequency (RF) radiation on the brain. Because mitochondrial DNA (mtDNA) defects are closely associated with various nervous system diseases and mtDNA is highly susceptible to oxidative stress, the purpose of this study was to determine whether radiofrequency radiation can cause oxidative damage to mtDNA. In this study, we exposed primary cultured cortical neurons to pulsed RF electromagnetic fields at a frequency of 1800 MHz modulated by 217 Hz at an average specific absorption rate (SAR) of 2 W/kg. At 24h after exposure, we found that RF radiation induced a significant increase in the levels of 8-hydroxyguanine (8-OHdG), a common biomarker of DNA oxidative damage, in the mitochondria of neurons. Consistent with this finding, the copy number of mtDNA and the levels of mitochondrial RNA (mtRNA) transcripts showed an obvious reduction after RF exposure. Each of these mtDNA disturbances could be reversed by pretreatment with melatonin, which is known to be an efficient in the brain. Together, these results suggested that 1800 MHz RF radiation could cause oxidative damage to mtDNA in primary cultured neurons. Oxidative damage to mtDNA may account for the neurotoxicity of RF radiation in the brain.

Pakhomov AG, [Non-thermal microwave effect on nerve fiber function]. *Biofizika*

Studies that show **Cell Phone** Health Effects

38(2):367-371, 1993. [Article in Russian]

Effects of microwave radiation (915 MHz, PW, peak SAR 20-30 W/g, pulse duration 1 mcs, 50.000 and 25.000 p.p.s.) were investigated in isolated frog nerve cord preparation. Nerve VHF heating didn't exceed 2.2 degrees C due to intense Ringer's solution perfusion. It was established that nerve irradiation simultaneously with its stimulation lead to significant decrease of action potential amplitude and peak latency. Since the equal conventional heating of the nerve caused the opposite changes (amplitude increase), the results obtained argue for non-thermal mechanism of microwave action.

Schüz J, Waldemar G, Olsen JH, Johansen C. Risks for central nervous system diseases among mobile phone subscribers: a Danish retrospective cohort study. PLoS One. 4(2):e4389, 2009

The aim of this study was to investigate a possible link between cellular telephone use and risks for various diseases of the central nervous system (CNS). We conducted a large nationwide cohort study of 420 095 persons whose first cellular telephone subscription was between 1982 and 1995, who were followed through 2003 for hospital contacts for a diagnosis of a CNS disorder. Standardized hospitalization ratios (SHRs) were derived by dividing the number of hospital contacts in the cohort by the number expected in the Danish population. The SHRs were increased by 10-20% for migraine and vertigo. No associations were seen for amyotrophic lateral sclerosis, multiple sclerosis or epilepsy in women. SHRs decreased by 30-40% were observed for dementia (Alzheimer disease, vascular and other dementia), Parkinson disease and epilepsy among men. In analyses restricted to subscribers of 10 years or more, the SHRs remained similarly increased for migraine and vertigo and similarly decreased for Alzheimer disease and other dementia and epilepsy (in men); the other SHRs were close to unity. In conclusion, the excesses of migraine and vertigo observed in this first study on cellular telephones and CNS disease deserve further attention. An interplay of a healthy cohort effect and reversed causation bias due to prodromal symptoms impedes detection of a possible association with dementia and Parkinson disease. Identification of the factors that result in a healthy cohort might be of interest for elucidation of the etiology of these diseases.

Hardell L, Söderqvist F, Carlberg M, Zetterberg H, Mild KH. Exposure to wireless phone emissions and serum beta-trace protein. Int J Mol Med. 26(2):301-306, 2010.

The lipocalin type of prostaglandin D synthase or beta-trace protein is synthesized in the choroid plexus, lepto-meninges and oligodendrocytes of the central nervous system and is secreted into the cerebrospinal fluid. beta-trace protein is the key enzyme in the synthesis of prostaglandin D2, an endogenous sleep-promoting neurohormone in the brain. Electromagnetic fields (EMF) in the radio frequency (RF) range have in some studies been associated with disturbed sleep. We studied the concentration of beta-trace protein in blood in relation to emissions from wireless phones. This study included 62 persons aged 18-30 years. The concentration of beta-trace protein decreased with increasing number of years of use of a wireless phone yielding a negative beta coefficient = -0.32, 95% confidence interval -0.60 to -0.04. Also cumulative use in hours

Studies that show **Cell Phone** Health Effects

gave a negative beta coefficient, although not statistically significant. Of the 62 persons, 40 participated in an experimental study with 30 min exposure to an 890-MHz GSM signal. No statistically significant change of beta-trace protein was found. In a similar study of the remaining 22 participants with no exposure, beta-trace protein increased significantly over time, probably due to a relaxed situation. EMF emissions may down-regulate the synthesis of beta-trace protein. This mechanism might be involved in sleep disturbances reported in persons exposed to RF fields. The results must be interpreted with caution since use of mobile and cordless phones were self-reported. Awareness of exposure condition in the experimental study may have influenced beta-trace protein concentrations.

Hamann W, Abou-Sherif S, Thompson S, Hall S. Pulsed radiofrequency applied to dorsal root ganglia causes a selective increase in ATF3 in small neurons. Eur J Pain. 10(2):171-176, 2006.

BACKGROUND: This is a "proof of concept study" to test the hypothesis that pulsed radiofrequency, PRF, produces cell stress at the primary afferent level without signs of overt thermal damage. We assumed that cell stress would result in impairment of normal function, and used the expression of activating transcription factor 3, ATF3, as an indicator of cellular "stress". **METHODS:** PRF (20ms of 500-kHz RF pulses, delivered at a rate of 2Hz; maximum temperature 42 degrees C) was delivered either to the sciatic nerve of adult rats in mid thigh, or to the L4 anterior primary ramus just distal to the intervertebral foramen. Controls were sham-operated or L4 axotomised. All tissues were examined 14 days after surgery. The percentage of CGRP- or ATF3-positive DRG neuronal somata was calculated using image analysis software (SigmaScan Pro 4). **RESULTS:** ATF3 expression was upregulated in L4 DRG neuronal cell bodies, irrespective of their size, after axotomy. It was also upregulated significantly ($p < 0.002$) and selectively, in small and medium calibre L4 DRG neurons, when PRF was applied close to the DRG just distal to the intervertebral foramen. PRF did not produce any obvious cellular changes in the nerve or L4 DRG neurons when applied to the sciatic nerve in mid-thigh. **CONCLUSION:** PRF has a biological effect, unlikely to be related to overt thermal damage. It appears to be selective in that it targets the group of neurons whose axons are the small diameter C and Adelta nociceptive fibres.

Bak M, Dudarewicz A, Zmyślony M, Sliwinska-Kowalska M. Effects of GSM signals during exposure to event related potentials (ERPs). Int J Occup Med Environ Health. 23(2):191-199, 2010.

OBJECTIVES: The primary aim of this work was to assess the effect of electromagnetic field (EMF) from the GSM mobile phone system on human brain function. The assessment was based on the assay of event related potentials (ERPs). **MATERIAL AND METHODS:** The study group consisted of 15 volunteers, including 7 men and 8 women. The test protocol comprised determination of P300 wave in each volunteer during exposure to the EMF. To eliminate possible effects of the applied test procedure on the final result, the test was repeated without EMF exposure. P300 latency, amplitude, and latency of the N1, N2, P2 waves were analysed. **RESULTS:** The statistical analysis

Studies that show **Cell Phone** Health Effects

revealed an effect of EMF on P300 amplitude. In the experiment with EMF exposure, lower P300 amplitudes were observed only at the time in which the volunteers were exposed to EMF; when the exposure was discontinued, the values of the amplitude were the same as those observed before EMF application. No such change was observed when the experiment was repeated with sham exposure, which may be considered as an indirect proof that lower P300 amplitude values were due to EMF exposure. No statistically significant changes were noted in the latencies of the N1, N2, P2 waves that precede the P300 wave, nor in the latency of the P300 itself.

CONCLUSIONS: The results suggest that exposure to GSM EMF exerts some effects on CNS, including effects on long latency ERPs.

Moretti D, Garenne A, Haro E, Poullétier de Gannes F, Lagroye I, Lévêque P, Veyret B, Lewis N. In-vitro exposure of neuronal networks to the GSM-1800 signal.

Bioelectromagnetics. 2013 Aug 1. doi: 10.1002/bem.21805. [Epub ahead of print]

The central nervous system is the most likely target of mobile telephony radiofrequency (RF) field exposure in terms of biological effects. Several electroencephalography (EEG) studies have reported variations in the alpha-band power spectrum during and/or after RF exposure, in resting EEG and during sleep. In this context, the observation of the spontaneous electrical activity of neuronal networks under RF exposure can be an efficient tool to detect the occurrence of low-level RF effects on the nervous system. Our research group has developed a dedicated experimental setup in the GHz range for the simultaneous exposure of neuronal networks and monitoring of electrical activity. A transverse electromagnetic (TEM) cell was used to expose the neuronal networks to GSM-1800 signals at a SAR level of 3.2 W/kg. Recording of the neuronal electrical activity and detection of the extracellular spikes and bursts under exposure were performed using microelectrode arrays (MEAs). This work provides the proof of feasibility and preliminary results of the integrated investigation regarding exposure setup, culture of the neuronal network, recording of the electrical activity, and analysis of the signals obtained under RF exposure. In this pilot study on 16 cultures, there was a 30% reversible decrease in firing rate (FR) and bursting rate (BR) during a 3 min exposure to RF. Additional experiments are needed to further characterize this effect.

Xu S, Ning W, Xu Z, Zhou S, Chiang H, Luo J. Chronic exposure to GSM 1800-MHz microwaves reduces excitatory synaptic activity in cultured hippocampal neurons. Neurosci Lett.398(3):253-257,2006.

The world wide proliferation of mobile phones raises the concern about the health effects of 1800-MHz microwaves on the brain. The present study assesses the effects of microwave exposure on the function of cultured hippocampal neurons of rats using whole cell patch-clamp analysis combined with immunocytochemistry. We showed that chronic exposure (15min per day for 8 days) to Global System for Mobile Communication (GSM) 1800-MHz microwaves at specific absorption rate (SAR) of 2.4W/kg induced a selective decrease in the amplitude of alpha-amino-3-hydroxy-5-methyl-4-soxazole propionic acid (AMPA) miniature excitatory postsynaptic currents (mEPSCs), whereas the frequency of AMPA mEPSCs and the amplitude of N-methyl-d-

Studies that show **Cell Phone** Health Effects

aspartate (NMDA) mEPSCs did not change. Furthermore, the GSM microwave treatment decreased the expression of postsynaptic density 95 (PSD95) in cultured neurons. Our results indicated that 2.4W/kg GSM 1800-MHz microwaves may reduce excitatory synaptic activity and the number of excitatory synapses in cultured rat hippocampal neurons.

Barteri M, Pala A, Rotella S. Structural and kinetic effects of mobile phone microwaves on acetylcholinesterase activity. Biophys Chem. 113(3):245-253, 2005.

The present study provides evidence that "in vitro" simple exposure of an aqueous solution of electric eel acetylcholinesterase (EeAChE; EC 3.1.1.7.) to cellular phone emission alters its enzymatic activity. This paper demonstrates, by combining different experimental techniques, that radio frequency (RF) radiations irreversibly affect the structural and biochemical characteristics of an important CNS enzyme. These results were obtained by using a commercial cellular phone to reproduce the reality of the human exposition. This experimental procedure provided surprising effects collected practically without experimental errors because they were obtained comparing native and irradiated sample of the same enzyme solution. Although these results cannot be used to conclude whether exposure to RF during the use of cellular phone can lead to any hazardous health effect, they may be a significant first step towards further verification of these effects on other "ex vivo" or "in vivo" biological systems.

Acar GO, Yener HM, Savrun FK, Kalkan T, Bayrak I, Enver O. Thermal effects of mobile phones on facial nerves and surrounding soft tissue. Laryngoscope.119(3):559-562, 2009.

OBJECTIVE: To investigate the possible thermal effects of microwaves from mobile phones on **facial nerves (FN)** and surrounding soft tissue. STUDY DESIGN:: A prospective study. METHODS: We studied FN conduction rate and compound muscle action potentials (CMAP) on 12 rabbits before exposure to radiofrequency radiation (RFR) emitted from a mobile phone. Also, the temperature change in the soft tissues around the FN was investigated by a four channel Luxtron fiber optic system. A mobile phone with 1900 MHz frequency was placed over the ipsilateral ear of the rabbit for 25 minutes, and FN and surrounding tissues were exposed to a 1.5 watts pulse modulated (217 packets/s) electromagnetic field. During exposure to RFR, immediately after turning off the mobile phone, and 25 minutes after the exposure temperature change in the surrounding tissue of the FN was recorded and compared to preexposure values. Additionally, another recording regarding the FN functions was done and the data were compared to preexposure values. RESULTS: The average temperature of the surrounding soft tissues was 0.39 K higher than the preexposure values during the exposure and immediately after turning off the mobile phone, and decreased to normal levels 25 minutes after the exposure, which was statistically significant. The amplitudes of FN CMAP after radiofrequency radiation exposure were significantly smaller than the preexposure amplitudes and the amplitudes were normal in the 25 minute measurement. CONCLUSION: The RFR emitted from a mobile phone can cause temporary FN dysfunction that can be due to temporary temperature increase in the

Studies that show **Cell Phone** Health Effects

soft tissue around the FN.

Ersan Odacı, Ayşe İkinci, Mehmet Yıldırım, Haydar Kaya, Metehan Akça, Hatice Hancı, Osman Fikret Sönmez, Ali Aslan, Mukadder Okuyan, Orhan Baş The Effects of 900 Megahertz Electromagnetic Field Applied in the Prenatal Period on Spinal Cord Morphology and Motor Behavior in Female Rat Pups. NeuroQuantology 11:573-581, 2013.

This study investigated the effect of a 900 megahertz (MHz) electromagnetic field (EMF) applied in the prenatal period on the spinal cord and motor behavior of female rat pups. Beginning of the study, female Sprague Dawley rats (180–250 g) were left to mate with male rats. Rats identified as pregnant were then divided into control (n=3) and EMF groups (n=3). The EMF group was exposed to 1-h 900 MHz EMF daily between days 13 and 21 of pregnancy. At 21 days old, rat pups were removed from their mothers and divided into two newborn rat groups, control (n=13) and EMF (n=10). The rotarod test was applied to the rat pups to assess motor functions and the open field test to evaluate locomotor activity. On day 32 of the study, the rat pups were decapitated, and the spinal cord in the upper thoracic region was removed. Following routine histological tests, they were stained with Cresyl fast violet. Rotarod test results revealed a significant increase in EMF group rat pups' motor functions ($p=0.037$). However, no difference was observed in the open field test results ($p>0.05$). In the EMF group' rat pups, we observed pathological changes in the spinal cord. On the basis of our results, 900 MHz EMF applied in the prenatal period affected spinal cord development. This effect was observed in the form of pathological changes in the spinal cord of rat pups, and it may be that these pathological changes led to an increase in rat pups' motor activities.

Lu Y, He M, Zhang Y, Xu S, Zhang L, He Y, Chen C, Liu C, Pi H, Yu Z, Zhou Z. Differential Pro-Inflammatory Responses of Astrocytes and Microglia Involve STAT3 Activation in Response to 1800 MHz Radiofrequency Fields. PLoS One. 2014 Oct 2;9(9):e108318. doi: 10.1371/journal.pone.0108318.

Microglia and astrocytes play important role in maintaining the homeostasis of central nervous system (CNS). Several CNS impacts have been postulated to be associated with radiofrequency (RF) electromagnetic fields exposure. Given the important role of inflammation in neural physiopathologic processes, we investigated the pro-inflammatory responses of microglia and astrocytes and the involved mechanism in response to RF fields. Microglial N9 and astroglial C8-D1A cells were exposed to 1800 MHz RF for different time with or without pretreatment with STAT3 inhibitor. Microglia and astrocytes were activated by RF exposure indicated by up-regulated CD11b and glial fibrillary acidic protein (GFAP). However, RF exposure induced differential pro-inflammatory responses in astrocytes and microglia, characterized by different expression and release profiles of IL-1 β , TNF- α , IL-6, PGE2, nitric oxide (NO), inducible nitric oxide synthase (iNOS) and cyclooxygenase 2 (COX2). Moreover, the RF exposure activated STAT3 in microglia but not in astrocytes. Furthermore, the STAT3 inhibitor Stattic ameliorated the RF-induced release of pro-inflammatory cytokines in microglia but not in astrocytes. Our results demonstrated that RF exposure differentially induced

Studies that show **Cell Phone** Health Effects

pro-inflammatory responses in microglia and astrocytes, which involved differential activation of STAT3 in microglia and astrocytes. Our data provide novel insights into the potential mechanisms of the reported CNS impacts associated with mobile phone use and present STAT3 as a promising target to protect humans against increasing RF exposure.

Zuo H, Lin T, Wang D, Peng R, Wang S, Gao Y, Xu X, Zhao L, Wang S, Su Z. RKIP Regulates Neural Cell Apoptosis Induced by Exposure to Microwave Radiation Partly Through the MEK/ERK/CREB Pathway. Mol Neurobiol. 2014 Aug 10. [Epub ahead of print]

In the present study, we investigated whether Raf-1 kinase inhibitory protein (RKIP) is important for neural cell apoptosis induced by microwave exposure and explored the role of MEK/ERK/CREB pathway regulated by RKIP in the apoptosis. Differentiated PC12 cells were exposed to continuous microwave radiation at 2.856 GHz for 5 min with average power density of 30 mW/cm². RKIP sense and anti-sense recombinant plasmids were constructed and transfected into PC12 cells, respectively. Terminal deoxynucleotidyl transferase (TdT)-mediated dUTP nick end labeling (TUNEL) staining and caspase-3 activity assay were used to detect cell apoptosis. The results showed that RKIP was downregulated after microwave exposure while the MEK/ERK/CREB signaling pathway was activated excessively. Moreover, the ratio of Bcl-2/Bax decreased, activity of caspase-3 increased, and thus apoptotic DNA fragmentation increased. RKIP overexpression significantly inhibited the phosphorylation of MEK, ERK, and CREB, while RKIP downregulation had the reverse effect. Furthermore, U0126 was found to antagonize the changes caused by RKIP downregulation after exposure to radiation. In conclusion, RKIP plays an important role in the neural cell apoptosis induced by microwave radiation, and the regulation of cell apoptosis by RKIP is partly through the MEK/ERK/CREB pathway. This suggests that RKIP may act as a key regulator of neuronal damage caused by microwave radiation. Extremely Low-Frequency Electromagnetic Fields Cause G1 Phase Arrest through the Activation of the ATM-Chk2-p21 Pathway

Ivanova Vlu, Martynova OV, Aleinik SV, Limarenko AV. [Effect of modified SHF and acoustic stimulation on spectral characteristics of the electroencephalograms of the cat brain] Biofizika. 45(5):935-940, 2000. [Article in Russian]

The effect of modulated electromagnetic fields on the spectral parameters of bioelectric brain activity in awake cats was studied by registering the electroencephalogram from the skin surface in the vertex area using carbon electrodes. In the normal electroencephalogram, spectral components in the range above 20 Hz predominated. It was shown that, upon irradiation with electromagnetic field (basic frequency 980 MHz, power density 30-50 microW/cm²), spectral components in the range of 12-18 Hz begin to prevail. A similarity in the redistribution of the power of spectral components upon both acoustic and modulated electromagnetic influences was revealed. The results suggest that there is a common neurophysiological mechanism by which modulated electromagnetic radiation and acoustic stimulation affect the electrical activity of the brain. This is consistent with the assumption that the effect of the electromagnetic field

Studies that show **Cell Phone** Health Effects

on the central nervous system is mediated through the acoustic sensory system.

Verma M, Dutta SK. Microwave induced alteration in the neuron specific enolase gene expression. Cancer Biochem Biophys. 13(4):239-244, 1993.

Exposure of pNGE7, a recombinant clone containing the coding and regulatory sequences for the expression of neuron specific enolase gene, cells to electromagnetic radiations (915 MHz, 16 Hz AM, SAR 0.05 mW/kg) resulted in the elevation of neuron specific enolase (NSE), a diagnostic marker for neuron and lung cancer. Using ion-exchange chromatography we separated the neuron specific enolase activity from the non-neuronal enolase (NNE) activity and observed an alteration in the expression of neuron specific enolase and non-neuronal enolase. The clinical applications of the present studies have been discussed.

Khudnitskii, SS, Moshkarev, EA, Fomenko, TV, [On the evaluation of the influence of cellular phones on their users]. [Article in Russian] Med Tr Prom Ekol (9):20-24, 1999.

The authors studied influence of ultrahigh frequency radiation caused by cellular phones on functional state of central nervous, cardiovascular systems and local temperature changes in cellular phones users. The head area near the phone antenna appeared to be under the most intensive heating. Ultrahigh frequency radiation induces significant changes in local temperature and in physiologic parameters of central nervous and cardiovascular systems.

Liu ML, Wen JQ, Fan YB. Potential protection of green tea polyphenols against 1800 MHz electromagnetic radiation-induced injury on rat cortical neurons. Neurotox Res. 20(3):270-276, 2011.

Radiofrequency electromagnetic fields (EMF) are harmful to public health, but the certain anti-irradiation mechanism is not clear yet. The present study was performed to investigate the possible protective effects of green tea polyphenols against electromagnetic radiation-induced injury in the cultured rat cortical neurons. In this study, green tea polyphenols were used in the cultured cortical neurons exposed to 1800 MHz EMFs by the mobile phone. We found that the mobile phone irradiation for 24 h induced marked neuronal cell death in the MTT (3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyl-tetrazolium bromide) and TUNEL (TdT mediated biotin-dUTP nicked-end labeling) assay, and protective effects of green tea polyphenols on the injured cortical neurons were demonstrated by testing the content of Bcl-2 Associated X protein (Bax) in the immunoprecipitation assay and Western blot assay. In our study results, the mobile phone irradiation-induced increases in the content of active Bax were inhibited significantly by green tea polyphenols, while the contents of total Bax had no marked changes after the treatment of green tea polyphenols. Our results suggested a neuroprotective effect of green tea polyphenols against the mobile phone irradiation-induced injury on the cultured rat cortical neurons.

Ivaschuk OI, Jones RA, Ishida-Jones T, Haggren W, Adey WR, Phillips JL, Exposure of nerve growth factor-treated PC12 rat pheochromocytoma cells to a modulated

Studies that show **Cell Phone** Health Effects

radiofrequency field at 836.55 MHz: effects on c-jun and c-fos expression.

Bioelectromagnetics 18(3):223-229, 1997.

Rat PC12 pheochromocytoma cells have been treated with nerve growth factor And then exposed to athermal levels of a packet-modulated radiofrequency field At 836.55 MHz. This signal was produced by a prototype time-domain multiple-access (TDMA) transmitter that conforms to the North American digital cellular telephone standard. Three slot average power densities were used: 0.09, 0.9, and 9 mW/cm². Exposures were for 20, 40, and 60 min and included an intermittent exposure regimen (20 min on/20 min off), resulting in total incubation times of 20, 60, and 100 min, respectively. Concurrent controls were sham exposed. After extracting total cellular RNA, Northern blot analysis was used to assess the expression of the immediate early genes, c-fos and c-jun, in all cell populations. No change in c-fos transcript levels were detected after 20 min exposure at each field intensity (20 min was the only time period at which c-fos message could be detected consistently). Transcript levels for c-jun were altered only after 20 min exposure to 9 mW/cm² (average 38% decrease).

Joubert V, Bourthoumieu S, Leveque P, Yardin C. Apoptosis is Induced by Radiofrequency Fields through the Caspase-Independent Mitochondrial Pathway in Cortical Neurons. Radiat Res. 169(1):38-45, 2008.

In the present study, we investigated whether continuous-wave (CW) radiofrequency (RF) fields induce neuron apoptosis in vitro. Rat primary neuronal cultures were exposed to a CW 900 MHz RF field with a specific absorption rate (SAR) of 2 W/kg for 24 h. During exposure, an increase of 2 degrees C was measured in the medium; control experiments with neurons exposed to 39 degrees C were then performed. Apoptosis was assessed by condensation of nuclei with 4',6-diamino-2-phenylindole (DAPI) staining observed with an epifluorescence microscope and fragmentation of DNA with TdT-mediated dUTP nick-end labeling (TUNEL) analyzed by flow cytometry. A statistically significant difference in the rate of apoptosis was found in the RF-field-exposed neurons compared to the sham-, 37 degrees C- and 39 degrees C-exposed neurons either 0 or 24 h after exposure using both methods. To assess whether the observed apoptosis was caspase-dependent or -independent, assays measuring caspase 3 activity and apoptosis-inducing factor (AIF) labeling were performed. No increase in the caspase 3 activity was found, whereas the percentage of AIF-positive nuclei in RF-field-exposed neurons was increased by three- to sevenfold compared to other conditions. Our results show that, under the experimental conditions used, exposure of primary rat neurons to CW RF fields may induce a caspase-independent pathway to apoptosis that involves AIF.

Seaman RL, Phelix CF. Acute effects of pulsed microwaves and 3-nitropropionic acid on neuronal ultrastructure in the rat caudate-putamen. Bioelectromagnetics. 26(2):82-101, 2005.

Ultrastructure of the medium sized "spiny" neuron in rat dorsal-lateral caudate-putamen was assessed after administration of 3-nitropropionic acid (3-NP) and exposure to pulsed microwaves. Sprague-Dawley male rats were given two daily intraperitoneal doses of 0 or 10 mg/kg 3-NP and 1.5 h after each dose were exposed to

Studies that show **Cell Phone** Health Effects

microwave radiation at a whole body averaged specific absorption rate (SAR) of 0 (sham exposure), 0.6, or 6 W/kg for 30 min. Microwave exposure consisted of 1.25 GHz radiation delivered as 5.9 μ s pulses with repetition frequency 10 Hz. Tissue samples taken 2-3 h after the second sham or microwave exposure showed no injury with light microscope methods. Blinded qualitative assessment of ultrastructure of randomly selected neurons from the same samples did reveal differences. Subsequent detailed, quantitative measurements showed that, when followed by sham exposure, administration of 3-NP significantly increased endoplasmic reticulum (ER) intracisternal width, ER area density, and nuclear envelope thickness. Microwave exposure at 6 W/kg alone also significantly increased these measures. Exposure of 3-NP treated animals at 6 W/kg significantly increased effects of 3-NP on ultrastructure. Although exposure at 0.6 W/kg alone did not affect ultrastructure measures, exposure of 3-NP treated animals at 0.6 W/kg reduced the effects of 3-NP. We concluded that 3-NP changed neuronal ultrastructure and that the microwave exposures used here changed neuronal ultrastructure in ways that depended on microwave SAR and neuron metabolic status. The apparent cancellation of 3-NP induced changes by exposure to pulsed microwaves at 0.6 W/kg indicated the possibility that such exposure can protect against the effects of mitochondrial toxins on the nervous system.

Zhou H, Su Z, Ning J, Wang C, Xie X, Qu D, Wu K, Zhang X, Pan J, Yang G. EFFECTS OF FREQUENCY, IRRADIATION GEOMETRY AND POLARISATION ON COMPUTATION OF SAR IN HUMAN BRAIN. Radiat Prot Dosimetry. 2014 Jan 6. [Epub ahead of print]

The power absorbed by the human brain has possible implications in the study of the central nervous system-related biological effects of electromagnetic fields. In order to determine the specific absorption rate (SAR) of radio frequency (RF) waves in the human brain, and to investigate the effects of geometry and polarisation on SAR value, the finite-difference time-domain method was applied for the SAR computation. An anatomically realistic model scaled to a height of 1.70 m and a mass of 63 kg was selected, which included 14 million voxels segmented into 39 tissue types. The results suggested that high SAR values were found in the brain, i.e. ~250 MHz for vertical polarisation and 900-1200 MHz both for vertical and horizontal polarisation, which may be the result of head resonance at these frequencies.

Hao D, Yang L, Chen S, Tong J, Tian Y, Su B, Wu S, Zeng Y. Effects of long-term electromagnetic field exposure on spatial learning and memory in rats. Neurol Sci. 2012 Feb 24. [Epub ahead of print]

With the development of communications industry, mobile phone plays an important role in daily life. Whether or not the electromagnetic radiation emitted by mobile phone causes any adverse effects on brain function has become of a great concern. This paper investigated the effect of electromagnetic field on spatial learning and memory in rats. 32 trained Wistar rats were divided into two groups: exposure group and control group. The exposure group was exposed to 916 MHz, 10w/m² mobile phone electromagnetic field (EMF) 6 h a day, 5 days a week, 10 weeks. The completion time, number of total errors and the neuron discharge signals were recorded while the rats were searching for

Studies that show **Cell Phone** Health Effects

food in an eight-arm radial maze at every weekend. The neuron signals of one exposed rat and one control rat in the maze were obtained by the implanted microelectrode arrays in their hippocampal regions. It can be seen that during the weeks 4-5 of the experiment, the average completion time and error rate of the exposure group were longer and larger than that of control group ($p < 0.05$). During the weeks 1-3 and 6-9, they were close to each other. The hippocampal neurons showed irregular firing patterns and more spikes with shorter interspike interval during the whole experiment period. It indicates that the 916 MHz EMF influence learning and memory in rats to some extent in a period during exposure, and the rats can adapt to long-term EMF exposure.

Esen F, Esen H **Effect of electromagnetic fields emitted by cellular phones on the latency of evoked electrodermal activity.** *Int J Neurosci.* **116(3):321-329, 2006.**

The widespread use of cellular phones raises the question of their possible adverse biological effects, especially on the central nervous system (CNS). Therefore, the authors examined the effect of electromagnetic fields emitted by cellular phones (CPEMFs) on the evoked neuronal activity of CNS relating to generation and representation of electrodermal activity (EDA), an index of sympathetic nervous system activity. EDA (skin resistance response; SRR) latency was lengthened approximately 200 ms with CPEMFs exposure irrespective of the head site next to mobile phone used. Hemispheric asymmetry of EDA-2 pathway, which is represented by shorter SRR latency in the right hand of the right hand responders, was also distorted with CPEMFs. Because the CNS regions including EDA-2 are also involved in tasks of motor timing and time estimation, delayed response in this neuronal network due to CPEMFs exposure may increase the response time of mobile phone users. Therefore, the findings point to the potential risks of mobile phones on the function of CNS and consequently, possible increase in the risk of phone-related driving hazards.

Calabrò E, Condello S, Currò M, Ferlazzo N, Caccamo D, Magazù S, Ientile R. **Modulation of heat shock protein response in SH-SY5Y by mobile phone microwaves.** *World J Biol Chem.* **3(2):34-40, 2012.**

AIM: To investigate putative biological damage caused by GSM mobile phone frequencies by assessing electromagnetic fields during mobile phone working.

METHODS: Neuron-like cells, obtained by retinoic-acid-induced differentiation of human neuroblastoma SH-SY5Y cells, were exposed for 2 h and 4 h to microwaves at 1800 MHz frequency bands. RESULTS: Cell stress response was evaluated by MTT assay as well as changes in the heat shock protein expression (Hsp20, Hsp27 and Hsp70) and caspase-3 activity levels, as biomarkers of apoptotic pathway. Under our experimental conditions, neither cell viability nor Hsp27 expression nor caspase-3 activity was significantly changed. Interestingly, a significant decrease in Hsp20 expression was observed at both times of exposure, whereas Hsp70 levels were significantly increased only after 4 h exposure. CONCLUSION: The modulation of the expression of Hsps in neuronal cells can be an early response to radiofrequency microwaves.

Studies that show **Cell Phone** Health Effects

Partsvania B, Sulaberidze T, Shoshiashvili L, Modebadze Z. Acute effect of exposure of mollusk single neuron to 900-MHz mobile phone radiation. Electromagn Biol Med. 30(3):170-179, 2011.

The goal of the present work was to explore the influence of commercially available cell phone irradiation on the single neuron excitability and memory processes. A Transverse Electromagnetic Cell (TEM Cell) was used to expose single neurons of mollusk to the electromagnetic field. Finite-Difference Time-Domain (FDTD) method was used for modeling the TEM Cell and the electromagnetic field interactions with living nerve ganglion and neurons. Neuron electrophysiology was investigated using standard microelectrode technique. The specific absorption rate (SAR) deposited into the single neuron was calculated to be 0.63 W/kg with a temperature increment of 0.1°C. After acute exposure, average firing threshold of the action potentials was not changed. However, the average latent period was significantly decreased. This indicates that together with latent period the threshold and the time of habituation might be altered during exposure. However, these alterations are transient and only latent period remains on the changed level.

Hearing Effects

Panda NK, Modi R, Munjal S, Virk RS. Auditory changes in mobile users: is evidence forthcoming? Otolaryngol Head Neck Surg. 144(4):581-585, 2011.

OBJECTIVE: Genuine concerns are being raised as to the potential health risks posed by electromagnetic frequency exposure secondary to mobile phone usage. This study was undertaken to assess and compare potential changes in hearing function at the level of the inner ear and central auditory pathway due to chronic exposure to electromagnetic waves from both global system for mobile communications (GSM) and code division multiple access (CDMA) mobile phone usage. DESIGN: Cohort study. SETTING: Tertiary referral center. SUBJECTS AND METHODS: One hundred twenty-five subjects who were long-term mobile phone users (more than 1 year; 63 GSM and 62 CDMA) and 58 controls who had never used mobile phones underwent audiological investigations including pure tone audiometry (250-12 kHz), tympanometry, distortion product otoacoustic emissions (DPOAE), auditory brain responses (ABR), and middle latency responses (MLRs). The changes in various parameters were studied in mobile-using and non-mobile-using ears of both GSM and CDMA subjects and corresponding ears of the controls to ascertain the effects of electromagnetic exposure. RESULTS: GSM and CDMA users were found to be at a significantly higher risk of having DPOAE absent as compared with controls ($P < .05$). They were found to have higher speech frequency thresholds and lower MLR wave and Na and Pa amplitudes. More than 3 years of mobile phone usage emerged as a risk factor ($P < .05$). The damage done was bilateral, with the quantum of damage being the same for both GSM and CDMA. CONCLUSION: Long-term and intensive GSM and CDMA mobile phone use may cause damage to cochlea as well as the auditory cortex.

Studies that show **Cell Phone** Health Effects

Panda NK, Jain R, Bakshi J, Munjal S. Audiologic disturbances in long-term mobile phone users. J Otolaryngol Head Neck Surg. 39(1):5-11, 2010.

Abstract. INTRODUCTION:There is general concern regarding the possible hazardous health effects of exposure to radiofrequency electromagnetic radiation emitted from mobile phones. This study aimed to assess the effects of chronic exposure to electromagnetic waves emitted from Global System for Mobile Communication (GSM) mobile phones on auditory functions.MATERIAL AND METHODS:A retrospective, cross-sectional, randomized, case control study was carried out in a tertiary care hospital. One hundred twelve subjects who were long-term mobile phone users (more than 1 year) and 50 controls who had never used a mobile phone underwent a battery of audiologic investigations including pure-tone audiometry (both speech and high frequency), tympanometry, distortion product otoacoustic emissions, auditory brain responses, and middle latency responses. Changes in the various parameters were studied in the mobile phone- and non-mobile phone-using ears of subjects and corresponding ears of the controls to ascertain the effects of electromagnetic exposure.RESULTS:There was no significant difference between users and controls for any of the audiologic parameters. However, trends for audiologic abnormalities were seen within the users. High-frequency loss and absent distortion product otoacoustic emissions were observed with an increase in the duration of mobile phone use, excessive use of mobile phones, and age more than 30 years. Additionally, users with some complaints during mobile phone use demonstrated absent distortion product otoacoustic emissions and abnormalities in auditory brainstem response.CONCLUSION:Long-term and intensive mobile phone use may cause inner ear damage. A large sample size would be required to reach definitive conclusions.

Oktay MF, Dasdag S. Effects of intensive and moderate cellular phone use on hearing function. Electromagn Biol Med. 25(1):13-21, 2006.

The purpose of this study is to investigate the effects of radiation emitted by mobile phones on the hearing of users. The study was carried out on three groups: 1) 20 men who have used a cellular phone frequently and spoken approximately 2 h per day for four years; 2) 20 men who have used a cellular phone for 10-20 min per day for four years; and 3) 20 healthy men who have never used a cellular phone (the control group). Brainstem evoked response audiometric (BERA) and pure tone audiometric (PTA) methods were used to measure the effects of exposure on hearing function of the subjects. In BERA measurements, I-III, III-V, and I-V interpeak latencies were evaluated. Interpeak latency of subjects in two experimental groups was compared to that of subjects in the control group. The BERA results showed no differences among the groups ($p > 0.05$).In PTA measurements, detection thresholds at 250 Hz, 500 Hz, 1000 Hz, 2000 Hz, 4000 Hz, and 8000 Hz frequencies were measured in all three groups. No differences were observed between moderate mobile phone users (10-20 min. per day) and control subjects. However, detection thresholds in those who talked approximately 2 h per day were found to be higher than those in either moderate users or control subjects. Differences at 4000 Hz for both bone and air conduction for right ears, and 500 Hz, and 4000 Hz bone and air conduction for left ears were significant for mean hearing

Studies that show **Cell Phone** Health Effects

threshold. This study shows that a higher degree of hearing loss is associated with long-term exposure to electromagnetic (EM) field generated by cellular phones.

Meo SA, Al-Drees AM. Mobile phone related-hazards and subjective hearing and vision symptoms in the Saudi population. Int J Occup Med Environ Health. 18(1):53-57, 2005

OBJECTIVES: Over the past decade utilization of mobile phones has dramatically increased. They are now an essential part of business, commerce, and communication, however, their use may lead to health problems. Therefore, the present study was designed to investigate a link between the use of mobile phones and hearing and vision symptoms in the Saudi population and also to contribute to the increase in social awareness of health problems associated with the use of these devices. **MATERIALS AND METHODS:** A total of 873 (57.04% of males and 39.86% of females) subjects using mobile phones were invited to participate in the presented study. A structured questionnaire was distributed among them to collect a detailed medical history. The Chi-square test was employed to observe the relationship between duration of calls and hearing and vision complaints. **RESULTS:** The present study showed an association between the use of mobile phones and hearing and vision complaints. About 34.59% of problems were related with impaired hearing, ear ache and/or warmth on the ear, and 5.04% of complaints with the decreased and/or blurred vision. **CONCLUSIONS:** It is concluded that the use of mobile phone is a health risk factor, and thus it is suggested that excessive use of mobile phones should be avoided and social awareness increased through health promotion activities, such as group discussions or public presentations and via electronic and printed media sources.

Velayutham P, Govindasamy GK, Raman R, Prepageran N, Ng KH. High-frequency hearing loss among mobile phone users. Indian J Otolaryngol Head Neck Surg. 66(Suppl 1):169-172, 2014.

The objective of this study is to assess high frequency hearing (above 8 kHz) loss among prolonged mobile phone users in a tertiary Referral Center. Prospective single blinded study. This is the first study that used high-frequency audiometry. The wide usage of mobile phone is so profound that we were unable to find enough non-users as a control group. Therefore we compared the non-dominant ear to the dominant ear using audiometric measurements. The study was a blinded study wherein the audiologist did not know which was the dominant ear. A total of 100 subjects were studied. Of the subjects studied 53% were males and 47% females. Mean age was 27. The left ear was dominant in 63%, 22% were dominant in the right ear and 15% did not have a preference. This study showed that there is significant loss in the dominant ear compared to the non-dominant ear ($P < 0.05$). Chronic usage mobile phone revealed high frequency hearing loss in the dominant ear (mobile phone used) compared to the non dominant ear.

Seckin E, Suren Basar F, Atmaca S, Kaymaz FF, Suzer A, Akar A, Sunan E, Koyuncu M. The effect of radiofrequency radiation generated by a Global System for Mobile

Studies that show **Cell Phone** Health Effects

Communications source on cochlear development in a rat model. J Laryngol Otol. 2014 May 1:1-6. [Epub ahead of print]

Objective: This study aimed to determine the effect of radiofrequency radiation generated by 900 and 1800 MHz Global System for Mobile Communications sources on cochlear development in the rat model. Methods: Eight pregnant albino Wistar rats were divided into three groups: control, 900 MHz and 1800 MHz. The latter two groups of pregnant rats were exposed to radiofrequency radiation for 1 hour per day starting on the 12th day of pregnancy until delivery. The rats in the control, 900 MHz and 1800 MHz groups gave birth to 24, 31 and 26 newborn rats respectively. Newborn rats in the 900 MHz and 1800 MHz groups were exposed to radiofrequency radiation for 1 hour per day for 21 days after delivery. Hearing evaluations of newborn rats were carried out using distortion product otoacoustic emissions testing. Eight newborn rats were randomly selected from each group for electron microscopic evaluation. Results: Distortion product otoacoustic emission tests revealed no significant difference among the groups, but electron microscopic evaluation revealed significant differences among the groups with regard to the number of normal, apoptotic and necrotic cells. Conclusion: The findings indicated cellular structural damage in the cochlea caused by radiofrequency radiation exposure during cochlear development in the rat model.

Kerekhanjanarong V, Supiyaphun P, Naratricoorn J, Laungpitackchumpon P. The effect of mobile phone to audiologic system. J Med Assoc Thai.88 Suppl 4:S231-234, 2005.

Mobile phones have come into widespread use. There are a lot of possible adverse effect to health. Use of mobile phone generate potentially harmful radiofrequency electromagnetic field (EMF) particularly for the hearing aspect. 98 subjects underwent hearing evaluations at Department of Otolaryngology, Faculty of Medicine, King Chulalongkorn Memorial Hospital, Chulalongkorn University. 31 males and 67 females, mean age was 30.48 +/- 9.51 years old, all subjects were investigated the hearing level by audiometry, tympanometry, otoacoustic emission (OAE) and auditory brain stem evoked response (ABR). The average of using time were 32.54 +/- 27.64 months, 57 subjects usually used the right side and 41 the left side. Average time of use per day was 26.31 +/- 30.91 minutes (range from 3 to 180 mins). When the authors compared the audiogram, both pure tone and speech audiometry, between the dominant and nondominant side, it indicated that there is no significant different. When the authors focused on the 8 subjects that used the mobile phone more than 60 mins per day. It indicated that the hearing threshold of the dominant ears was worse than the nondominant ears.

Khullar S1, Sood A2, Sood S3. Auditory Brainstem Responses and EMFs Generated by Mobile Phones. Indian J Otolaryngol Head Neck Surg. 65(Suppl 3):645-649, 2013.

There has been a manifold increase in the number of mobile phone users throughout the world with the current number of users exceeding 2 billion. However this advancement in technology like many others is accompanied by a progressive increase in the frequency and intensity of electromagnetic waves without consideration of the health consequences. The aim of our study was to advance our understanding of the

Studies that show Cell Phone Health Effects

potential adverse effects of GSM mobile phones on auditory brainstem responses (ABRs). 60 subjects were selected for the study and divided into three groups of 20 each based on their usage of mobile phones. Their ABRs were recorded and analysed for latency of waves I-V as well as interpeak latencies I-III, I-V and III-V (in ms). Results revealed no significant difference in the ABR parameters between group A (control group) and group B (subjects using mobile phones for maximum 30 min/day for 5 years). However the latency of waves was significantly prolonged in group C (subjects using mobile phones for 10 years for a maximum of 30 min/day) as compared to the control group. Based on our findings we concluded that long term exposure to mobile phones may affect conduction in the peripheral portion of the auditory pathway. However more research needs to be done to study the long term effects of mobile phones particularly of newer technologies like smart phones and 3G.

Kellenyi, L, Thuroczy, G, Faludy, B, Lenard, L, Effects of mobile GSM radiotelephone exposure on the auditory brainstem response (ABR). Neurobiology 7:79-81, 1999.

A 15-min exposure to GSM phone radiation caused an increase in auditory brainstem response in the exposed side of human subjects. Subjects also showed a hearing deficiency in the high frequency range (20 dB hearing deficiency from 2 KHz to 10 KHz).

Kaprana AE, Chimona TS, Papadakis CE, Velegrakis SG, Vardiambasis IO, Adamidis G, Velegrakis GA. Auditory brainstem response changes during exposure to GSM-900 radiation: An experimental study. Audiol Neurotol. 16(4):270-276, 2011.

Abstract. The objective of the present study was to investigate the possible electrophysiological time-related changes in auditory pathway during mobile phone electromagnetic field exposure. Thirty healthy rabbits were enrolled in an experimental study of exposure to GSM-900 radiation for 60 min and auditory brainstem responses (ABRs) were recorded at regular time-intervals during exposure. The study subjects were radiated via an adjustable power and frequency radio transmitter for GSM-900 mobile phone emission simulation, designed and manufactured according to the needs of the experiment. The mean absolute latency of waves III-V showed a statistically significant delay ($p < 0.05$) after 60, 45 and 15 min of exposure to electromagnetic radiation of 900 MHz, respectively. Interwave latency I-III was found to be prolonged after 60 min of radiation exposure in correspondence to wave III absolute latency delay. Interwave latencies I-V and III-V were found with a statistically significant delay ($p < 0.05$) after 30 min of radiation. No statistically significant delay was found for the same ABR parameters in recordings from the ear contralateral to the radiation source at 60 min radiation exposure compared with baseline ABR. The ABR measurements returned to baseline recordings 24 h after the exposure to electromagnetic radiation of 900 MHz. The prolongation of interval latencies I-V and III-V indicates that exposure to electromagnetic fields emitted by mobile phone can affect the normal electrophysiological activity of the auditory system, and these findings fit the pattern of general responses to a stressor.

Hutter HP, Moshhammer H, Wallner P, Cartellieri M, Denk-Linnert DM, Katzinger M, Ehrenberger K, Kundi M. Tinnitus and mobile phone use. Occup Environ Med.

Studies that show **Cell Phone** Health Effects

67(12):804-808, 2010.

Objectives The mechanisms that produce tinnitus are not fully understood. While tinnitus can be associated with diseases and disorders of the ear, retrocochlear diseases and vascular pathologies, there are few known risk factors for tinnitus apart from these conditions. There is anecdotal evidence of an link between mobile phone use and tinnitus, but so far there have been no systematic investigations into this possible association. **Methods** 100 consecutive patients presenting with tinnitus were enrolled in an individually matched case-control study. For each case a control subject was randomly selected from visiting outpatients matched for sex and age. The patient's history was obtained and clinical examinations were conducted to exclude patients with known underlying causes of tinnitus. Mobile phone use was assessed based on the Interphone Study protocol. ORs were computed by conditional logistic regression with years of education and living in an urban area as covariates. **Results** Mobile phone use up to the index date (onset of tinnitus) on the same side as the tinnitus did not have significantly elevated ORs for regular use and intensity or for cumulative hours of use. The risk estimate was significantly elevated for prolonged use (≥ 4 years) of a mobile phone (OR 1.95; CI 1.00 to 3.80). **Conclusions** Mobile phone use should be included in future investigations as a potential risk factor for developing tinnitus.

Cox RA, Luxton LM, Cerebral symptoms from mobile telephones. Occup Environ Med 57(6):431, 2000. (letter to the editor)

Mobile phones affect the inner ear in 5-8% of users leading to dizziness, disorientation, nausea, headache and transient confusion.

Alsanosi AA, Al-Momani MO, Hagr AA, Almomani FM, Shami IM, Al-Habeeb SF. The acute auditory effects of exposure for 60 minutes to mobile's electromagnetic field. Saudi Med J. 34(2):142-146, 2013.

OBJECTIVE: To assess the immediate consequences of 60 minutes exposure to mobile phones on hearing function by determining changes in distortion product otoacoustic emission (DPOAE) and hearing threshold levels (HTLs). **METHODS:** This prospective control clinical trial study was carried out at the Ear, Nose and Throat Department, King Abdulaziz University Hospital, Riyadh, Kingdom of Saudi Arabia from July 2009 to July 2011. The data collected included age, symptoms experienced after exposure, and HTLs and DPOAE were recorded before, and immediately after 60 minutes of exposure to the same model of mobile phone. **RESULTS:** Heat/pain was the most commonly reported symptom. In the test-ears, significant shift ($p < 0.05$) was noticed in HTLs at 1000 and 2000 Hz but not at other frequencies, while non test-ears did not reveal significant shift in HTLs. Additionally, test-ears revealed significant differences ($p < 0.05$) in DPOAE at 1000 Hz, 1400 Hz, 2000 Hz, and at the average of all frequencies, while non test-ears did not show significant differences. **CONCLUSION:** Sixty minutes of close exposure to electromagnetic fields emitted by a mobile phone had an immediate effect on HTL assessed by pure-tone audiogram and inner ear (assessed by DPOAE) in young human subjects. It also caused a number of other otologic symptoms.

Studies that show **Cell Phone** Health Effects

Mandalà M, Colletti V, Sacchetto L, Manganotti P, Ramat S, Marcocci A, Colletti L. Effect of Bluetooth headset and mobile phone electromagnetic fields on the human auditory nerve. Laryngoscope. 2013 Apr 25. doi: 10.1002/lary.24103. [Epub ahead of print]

OBJECTIVES/HYPOTHESIS: The possibility that long-term mobile phone use increases the incidence of astrocytoma, glioma and acoustic neuroma has been investigated in several studies. Recently, our group showed that direct exposure (in a surgical setting) to cell phone electromagnetic fields (EMFs) induces deterioration of auditory evoked cochlear nerve compound action potential (CNAP) in humans. To verify whether the use of Bluetooth devices reduces these effects, we conducted the present study with the same experimental protocol. **STUDY DESIGN:** Randomized trial. **METHODS:** Twelve patients underwent retrosigmoid vestibular neurectomy to treat definite unilateral Ménière's disease while being monitored with acoustically evoked CNAPs to assess direct mobile phone exposure or alternatively the EMF effects of Bluetooth headsets. **RESULTS:** We found no short-term effects of Bluetooth EMFs on the auditory nervous structures, whereas direct mobile phone EMF exposure confirmed a significant decrease in CNAPs amplitude and an increase in latency in all subjects. **CONCLUSIONS:** The outcomes of the present study show that, contrary to the finding that the latency and amplitude of CNAPs are very sensitive to EMFs produced by the tested mobile phone, the EMFs produced by a common Bluetooth device do not induce any significant change in cochlear nerve activity. The conditions of exposure, therefore, differ from those of everyday life, in which various biological tissues may reduce the EMF affecting the cochlear nerve. Nevertheless, these novel findings may have important safety implications.

Grisanti G, Parlapiano C, Tamburello CC, Tine G, Zanforlin L. Cellular phone effects on otoacoustic emissions. IEEE MTT-S Digest 2: 771-774, 1998.

A study on bioelectromagnetic effects induced by the use of TACS phones, evidencing a variation of the natural response of the auditory system is presented. This study was performed applying a method based on the registration of the evoked otoacoustic emissions (transient and distortion products). The experimental results show that modulated electromagnetic fields modify the distortion products in about all the examined subjects.

Garcia Callejo FJ, Garcia Callejo F, Pena Santamaria J, Alonso Castaneira I, Sebastian Gil E, Marco Algarra J. [Hearing level and intensive use of mobile phones] Acta Otorrinolaringol Esp. 56(5):187-191, 2005. [Article in Spanish]

INTRODUCTION: Wide studies and substantial controversies build on utilization of actual mobile phones and appearance of systemic disorders or even tumours, but there is no knowledge about an eventual involvement on early hearing loss. **PATIENTS AND METHODS:** In a group of three hundred and twenty-three healthy and normoacoustic volunteers who were usual costumers of mobile phones an audiometric evaluation was made at the beginning of its use and three years later, inquiring about the periods of time

Studies that show **Cell Phone** Health Effects

per day and year employed on direct contacts with phone. A healthy and normoacoustic control group of non users was studied too. RESULTS: Cases carried out 24.3 +/- 8.2 active contacts, reaching 50.4 +/- 27.8 days of mobile phone employment in three years. Audiometric curve was similar in cases and controls at the beginning of the study. After this follow-up, cases showed an increase on hearing threshold between 1 and 5 dB HL more than controls in speech tones ($p < 0.001$). Moreover, there was a trend to correlate time of phone use to hearing impairment, but this finding did not result statistically significative. CONCLUSIONS: Frequent management of mobile phones in a middle period of time allows to detect a mild hearing loss, but the cause of this disorder keeps unclear.

Fritze K, Wiessner C, Kuster N, Sommer C, Gass P, Hermann DM, Kiessling M, Hossmann KA, Effect of global system for mobile communication microwave exposure on the genomic response of the rat brain. Neuroscience 81(3):627-639, 1997.

The acute effect of global system for mobile communication (GSM) microwave exposure on the genomic response of the central nervous system was studied in rats by measuring changes in the messenger RNAs of hsp70, the transcription factor genes c-fos and c-jun and the glial structural gene GFAP using in situ hybridization histochemistry. Protein products of transcription factors, stress proteins and marker proteins of astroglial and microglial activation were assessed by immunocytochemistry. Cell proliferation was evaluated by bromodeoxyuridine incorporation. A special GSM radiofrequency test set, connected to a commercial cellular phone operating in the discontinuous transmission mode, was used to simulate GSM exposure. The study was conducted at time averaged and brain averaged specific absorption rates of 0.3 W/kg (GSM exposure), 1.5 W/kg (GSM exposure) and 7.5 W/kg (continuous wave exposure), respectively. Immediately after exposure, in situ hybridization revealed slight induction of hsp70 messenger RNA in the cerebellum and hippocampus after 7.5 W/kg exposure, but not at lower intensities. A slightly increased expression of c-fos messenger RNA was observed in the cerebellum, neocortex and piriform cortex of all groups subjected to immobilization, but no differences were found amongst different exposure conditions. C-jun and GFAP messenger RNAs did not increase in any of the experimental groups. 24 h after exposure, immunocytochemical analysis of FOS and JUN proteins (c-FOS, FOS B, c-JUN JUN B, JUN D), of HSP70 or of KROX-20 and -24 did not reveal any alterations. Seven days after exposure, neither increased cell proliferation nor altered expression of astroglial and microglial marker proteins were observed. In conclusion, acute high intensity microwave exposure of immobilized rats may induce some minor stress response but does not result in lasting adaptive or reactive changes of the brain.

Colletti V, Mandalà M, Manganotti P, Ramat S, Sacchetto L, Colletti L. Intraoperative observation of changes in cochlear nerve action potentials during exposure to electromagnetic fields generated by mobile phones. J Neurol Neurosurg Psychiatry.82(7):766-71, 2010

Background The rapid spread of devices generating electromagnetic fields (EMF) has raised concerns as to the possible effects of this technology on humans. The auditory system is the neural organ most frequently and directly exposed to electromagnetic

Studies that show **Cell Phone** Health Effects

activity owing to the daily use of mobile phones. In recent publications, a possible correlation between mobile phone usage and central nervous system tumours has been detected. Very recently a deterioration in otoacoustic emissions and in the auditory middle latency responses after intensive and long-term magnetic field exposure in humans has been demonstrated. **Methods** To determine with objective observations if exposure to mobile phone EMF affects acoustically evoked cochlear nerve compound action potentials, seven patients suffering from Ménière's disease and undergoing retrosigmoid vestibular neurectomy were exposed to the effects of mobile phone placed over the craniotomy for 5 min. **Results** All patients showed a substantial decrease in amplitude and a significant increase in latency of cochlear nerve compound action potentials during the 5 min of exposure to EMF. These changes lasted for a period of around 5 min after exposure. **Discussion** The possibility that EMF can produce relatively long-lasting effects on cochlear nerve conduction is discussed and analysed in light of contrasting previous literature obtained under non-surgical conditions. Limitations of this novel approach, including the effects of the anaesthetics, craniotomy and surgical procedure, are presented in detail.

Budak GG, Muluk NB, Budak B, Oztürk GG, Apan A, Seyhan N. Effects of intrauterine and extrauterine exposure to GSM-like radiofrequency on distortion product otoacoustic emissions in infant male rabbits. *Int J Pediatr Otorhinolaryngol.* 73(3):391-399, 2009.

OBJECTIVES: The aim of this study was to investigate the potential hazardous effects of intrauterine (IU) and/or extrauterine (EU) exposure to 1800 MHz Global System for Mobile Communications-like (GSM-like) radiofrequency (RF) on the cochlear functions of infant rabbits by measuring distortion product otoacoustic emission (DPOAE) response amplitudes. **METHODS:** Thirty-six white infant male New Zealand rabbits each 1-month-old were included in the study. The animals were randomly divided into four groups. Nine infant rabbits (Group 1) were not exposed to 1800 MHz GSM-like RF (Control-C). Nine infant rabbits (Group 2) were exposed to 1800 MHz GSM-like RF, 15 min daily for 14 days after they reached 1-month of age (extrauterine-EU). Nine infant rabbits (Group 3) were exposed to 1800 MHz GSM-like RF, 15 min daily for 7 days in the intrauterine period (between 15th and 22nd days of the gestational period) (intrauterine-IU). Nine infant rabbits (Group 4) were exposed to 1800 MHz GSM-like RF, 15 min daily for 7 days in the intrauterine period (between 15th and 22nd days of the gestational period) and 15 min daily for 14 days after they reached to 1-month of age (IU+EU). The cochlear functions were assessed by DPOAEs at 1.0-8.0 kHz. **RESULTS:** At 1.5 kHz, the mean DPOAE amplitude of Group 3 was higher than that of the controls and Group 2; and the mean DPOAE value of Group 4 was higher than that of the controls and Group 2. At 2.0 kHz, the mean DPOAE amplitude of Group 4 was higher than that of Group 2. At 3.0 kHz, the mean DPOAE amplitude of Group 4 was higher than that of the controls and Group 2. At 4.0 kHz, the mean DPOAE amplitude of Group 2 was lower than that of the controls, while the mean value of Group 4 was higher than the mean value of the controls and Group 2. At 6.0 kHz, the mean DPOAE amplitude of Group 2 was lower than that of the control group; however, the mean value of Group 4 was

Studies that show **Cell Phone** Health Effects

higher than that of Group 2. At 1.0 and 8.0 kHz, no significant differences were found among the four groups. CONCLUSION: Prolonged exposure and hyperthermia related to the power density of applied RF, increasing the temperature in the ear canal, may affect DPOAE amplitudes. Harmful effects of RF are mainly observed as a decrease in DPOAE amplitudes at 4.0-6.0 kHz during extrauterine exposure in infancy. During the intrauterine period, the water content of the middle and inner ear and amnion fluid may play a protective role. Therefore, children must be protected from RF exposure. The use of mobile phones at short distances from the ear of the infants should be avoided because of the lower thickness of the anatomical structure in infancy.

Tahvanainen K, Nino J, Halonen P, Kuusela T, Alanko T, Laitinen T, Lansimies E, Hietanen M, Lindholm H. Effects of cellular phone use on ear canal temperature measured by NTC thermistors. Clin Physiol Funct Imaging. 27(3):162-172, 2007.

The earlier studies using phantom models and human subjects concerning warming effects during cellular phone use have been controversial, partly because radiofrequency (RF) exposures have been variable. In this randomized, double-blind, placebo-controlled crossover trial, 30 healthy subjects were submitted to 900 MHz (2W) and 1800 MHz (1W) cellular phone RF exposure, and to sham exposure in separate study sessions. Temperature signals were recorded continuously in both ear canals before, during and after the 35-min RF exposure and the 35-min sham exposure sessions. Temperature was measured by using small-sized NTC thermistors placed in the ear canals through disposable ear plugs. The mean temperature changes were determined during a set cardiovascular autonomic function studies: during a 5-min controlled breathing test, during a 5-min spontaneous breathing test, during 7-min head-up tilting, 1-min before, during and after two consecutive Valsalva manoeuvres and during a deep breathing test. Temperatures in the exposed ear were significantly higher during RF exposures compared with sham exposure in both 900 and 1800 MHz studies with maximum differences of 1.2 +/- 0.5 degrees C (900 MHz exposure) and 1.3 +/- 0.7 degrees C (1800 MHz exposure). Temperatures in the RF-exposed ear were also significantly higher during the postexposure period compared with post-sham exposure period with maximum differences of 0.6 +/- 0.3 degrees C for 900 MHz and 0.5 +/- 0.5 degrees C for 1800 MHz. The results of this study suggest that RF exposure to a cellular phone, either using 900 or 1800 MHz with their maximal allowed antenna powers, increases the temperature in the ear canal. The reason for the ear canal temperature rising is a consequence of mobile phone battery warming during maximal antenna power use. The earlier published articles do not indicate that temperature rising in the ear canal has any significant contribution from the RF fields emitted from mobile phones.

Budak GG, Muluk NB, Budak B, Oztürk GG, Apan A, Seyhan N. Effects of GSM-like radiofrequency on distortion product otoacoustic emissions of rabbits: comparison of infants versus adults. Int J Pediatr Otorhinolaryngol. 73(8):1143-1147, 2009.

OBJECTIVES: The aim of this study is to investigate the potential hazardous effects of 1800 MHz Global System for Mobile Communications-like (GSM-like) Radiofrequency

Studies that show **Cell Phone** Health Effects

(RF) exposure on the cochlear functions of female infant and adult rabbits by measuring Distortion Product Otoacoustic Emission (DPOAE) response amplitudes. **METHODS:** Eighteen each one-month-old New Zealand White female rabbits and eighteen each 13-month-old adult rabbits were included into the study. They were randomly divided into four groups. Nine infant rabbits (Group 1) were not exposed to 1800 MHz GSM-like RF (Infant Control, C-In). Nine infant rabbits (Group 2) were exposed to 1800 MHz GSM-like RF, 15 min daily for 7 days after they reached one-month of age (Infant RF, RF-In). Nine adult rabbits were not exposed to 1800 MHz GSM-like RF, 15 min daily for 7 (Adult Control, C-Ad). Nine adult rabbits were exposed to 1800 MHz GSM-like RF, 15 min daily for 7 days (Adult RF, RF-Ad). Cochlear functions were assessed by DPOAEs at 1.0-8.0 kHz. **RESULTS:** At 1.0-2.0 and 6.0 kHz, the mean DPOAE values of Group 2 were significantly higher than that of Group 1. At 3.0-8.0 kHz, the mean DPOAE values of Group 4 were significantly lower than that of Group 1. At 6.0-8.0 kHz, the mean DPOAE values of Group 2 were significantly higher than that of Group 3. At 1.0-8.0 kHz, the mean DPOAE values of Group 4 were significantly lower than that of Group 2. At 1.0-8.0 kHz, the mean DPOAE values of Group 4 were significantly lower than that of Group 3. **CONCLUSION:** Harmful effects of GSM-like 1800 MHz RF exposure was detected more in the adult female rabbits than infant female rabbits by DPOAE measurement. Prolonged exposure and hyperthermia related to the power density of applied RFR, increasing the temperature in the ear canal, may decrease the DPOAE amplitudes. Water containing medium in the middle ear of infant rabbits may play the protective role from the RF damage.

Budak GG, Muluk NB, Oztürk GG, Budak B, Apan A, Seyhan N, Sanli C. Effects of GSM-like radiofrequency on distortion product otoacoustic emissions in pregnant adult rabbits. Clin Invest Med. 32(2):E112-116, 2009.

OBJECTIVES: To determine the effects of 1800 MHz GSM-like Radiofrequency (RFR) on the cochlear functions of pregnant adult rabbits by Distortion Product Otoacoustic Emissions (DPOAEs). **METHODS:** Eighteen 13-month-old pregnant and eighteen 13-month-old non-pregnant New Zealand White rabbits were studied. They were randomly divided into four groups. Nine pregnant rabbits (Group 2) and nine non-pregnant rabbits (Group 4) were exposed to 1800 MHz GSM-like RFR 15 min daily for 7 days. Nine pregnant (Group 1) and nine non-pregnant rabbits (Group 3) were not exposed to GSM like RFR. Cochlear functions were assessed by DPOAEs at 1.0-8.0 kHz. **RESULTS:** In all pregnant groups except 2.0 kHz, DPOAE amplitudes were not different in Group 2 and Group1. In Group 4, DPOAE amplitudes at 1.0-4.0 kHz (-1.68 dB SPL at 1.0 kHz, 3.05 dB SPL at 1.5 kHz, 2.96 dB SPL at 2.0 kHz, 1.30 dB SPL at 3.0 kHz and 12.22 dB SPL at 4.0 kHz) were lower than Group 3 (8.67 dB SPL at 1.0 kHz, 17.67 dB SPL at 1.5 kHz, 26.10 dB SPL at 2.0 kHz, 18.10 dB SPL at 3.0 kHz and 35.13 dB SPL at 4.0 kHz) ($P < 0.0125$). In the pregnant group, harmful effects of GSM-like RFR were less than in the non-pregnant group. **CONCLUSION:** GSM-like RFR caused decreases in DPOAE amplitudes mainly in non-pregnant adult rabbits. Prolonged exposure may affect the DPOAE amplitude. Recommendations are given to prevent the potential hazardous effects of RF in humans.

Studies that show **Cell Phone** Health Effects

Al-Dousary SH Mobile phone induced sensorineural hearing loss. Saudi Med J. 28(8):1283-1286, 2007.

The increased use of mobile phones worldwide has focused interest on the biological effects and possible health outcomes of exposure to radiofrequency fields from mobile phones, and their base stations. Various reports suggest that mobile phone use can cause health problems like fatigue, headache, dizziness, tension, and sleep disturbances; however, only limited research data is available in medical literature regarding interaction between electromagnetic fields emitted by mobile phones and auditory function; and the possible impact on hearing. We report a case of sensorineural hearing loss due to Global System for Mobile Communications mobile phone use, in a 42-year-old male.

Effects on Eyes

Yu Y, Yao K. Non-thermal cellular effects of lowpower microwave radiation on the lens and lens epithelial cells. J Int Med Res. 38(3):729-736, 2010.

Because of the increased use of modern radiofrequency devices, public concern about the possible health effects of exposure to microwave radiation has arisen in many countries. It is well established that high-power microwave radiation can induce cataracts via its thermal effects. It remains unclear whether low-power microwave radiation, especially at levels below the current exposure limits, is cataractogenic. This review summarizes studies on the biological effects of low-power microwave radiation on lens and lens epithelial cells (LECs). It has been reported that exposure affects lens transparency, alters cell proliferation and apoptosis, inhibits gap junctional intercellular communication, and induces genetic instability and stress responses in LECs. These results raise the question of whether the ambient microwave environment can induce non-thermal effects in the lens and whether such effects have potential health consequences. Further in vivo studies on the effects on the lens of exposure to low-power microwave radiation are needed.

Balik HH, Turgut-Balik D, Balikci K, Ozcan IC. Some ocular symptoms and sensations experienced by long term users of mobile phones. Pathol Biol (Paris). 53(2):88-91, 2005.

In this study, a survey was conducted to investigate the possible effects of long term usage of mobile phone (MP) on eyes. The studied symptoms are blurring of vision, redness on the eyes, vision disturbance, secretion of the eyes, inflammation in the eyes and lacrimation of the eyes. There is no effect on redness on the eyes and vision disturbance, but some statistical evidences are found that MP may cause blurring of vision, secretion of the eyes, inflammation in the eyes and lacrimation of the eyes. These results suggest an awareness of the symptoms and sensations.

Yu Y, Yao K, Wu W, Wang K, Chen G, Lu D. Effects of exposure to 1.8 GHz

Studies that show **Cell Phone** Health Effects

radiofrequency field on the expression of Hsps and phosphorylation of MAPKs in human lens epithelial cells. *Cell Res.* 18(12):1233-1235, 2008.

(No abstract available) Last sentence of discussion:

“Our results suggest that exposure to RF of wireless communications can induce expression of Hsp27 and Hsp70 and the activation of ERK1/2 and JNK1/2 in human LECs. The induction of Hsp27 and Hsp70, by a non-thermal stress, together with the activation of signal transduction pathways, provides reliable and sensitive biomarkers that could serve as the basis for improved mobile phone safety guidelines.”

Dovrat A, Berenson R, Bormusov E, Lahav A, Lustman T, Sharon N, Schachter L. Localized effects of microwave radiation on the intact eye lens in culture conditions. *Bioelectromagnetics.* 26(5):398-405, 2005.

A novel experimental system was used to investigate the localized effects of microwave radiation on bovine eye lenses in culture for over 2 weeks. Using this setup, we found clear evidence that this radiation has a significant impact on the eye lens. At the macroscopic level, it is demonstrated that exposure to a few mW at 1 GHz for over 36 h affects the optical function of the lens. Most importantly, self-recovery occurs if the exposure is interrupted. At the microscopic level, close examination of the lens indicates that the interaction mechanism is completely different from the mechanism-causing cataract via temperature increase. Contrary to the latter's effect, that is particularly pronounced in the vicinity of the sutures and it is assumed to be a result of local friction between the edges of the fibers consisting the lens. Even if macroscopically the lens has recovered from the irradiation, microscopically the indicators of radiation impact remain.

Lu L, Xu H, Wang X, Guo G. Increased nitric oxide synthase activity is essential for electromagnetic-pulse-induced blood-retinal barrier breakdown in vivo. *Brain Res.* 1264:104-10, 2009.

PURPOSE: To examine whether electromagnetic pulses (EMPs) affected the permeability of the blood-retinal barrier (BRB), gene expression of occludin and activity of nitric oxide synthase (NOS). **METHODS:** Sprague-Dawley (SD) rats were used and randomized into EMP and control groups. Retinas were removed immediately, and 2 h or 24 h after EMP radiation. BRB permeability was analyzed by transmission electron microscopy and Evans Blue staining. Retinal NOS activity and concentrations of nitrite and nitrate were measured. Occludin mRNA and protein levels were detected by RT-PCR and Western blotting. **RESULTS:** Exposure of SD rats to EMP resulted in increased BRB permeability, with the greatest decrease in occludin at 24 h. Moreover, this permeability defect was also correlated with significant increases in the formation of NO and induction of NOS activity in SD rats. Furthermore, we found that treatment with NOS inhibitor N-nitro-L-arginine methyl ester (L-NAME) blocked BRB breakdown and prevented the increase in NO formation and induction of NOS activity, as well as the decrease in occluding expression. **CONCLUSION:** Taken together, these results support the view that NOS-dependent NO production is an important factor that contributes to EMP-induced BRB dysfunction, and suggests that NOS induction may play an important

Studies that show **Cell Phone** Health Effects

role in BRB breakdown.

Teerapot Wessapan, Phadungsak Rattanadecho. Specific absorption rate and temperature increase in the human eye due to electromagnetic fields exposure at different frequencies. International Journal of Heat and Mass Transfer, 64:426-435, September 2013.

This study presents a numerical analysis of the specific absorption rate (SAR) and the heat transfer in a heterogeneous human eye model exposed to electromagnetic (EM) fields of 900 and 1800 MHz. In this study, the effect of operating frequency on the SAR and temperature distributions in the eye was systematically investigated. The SAR value and the temperature distribution in various tissues in the eye during exposure to EM fields were obtained by numerical simulation of EM wave propagation and a heat transfer model was then developed based on the natural convection and porous media theories. The study highlights two transport phenomena: heat and mass transfer in the eye during exposure to EM fields at different frequencies. This study indicated that when the eye exposed to EM fields at the frequencies of 900 and 1800 MHz, the highest SAR values at two chosen frequencies was in the cornea, and the highest temperature at the frequency of 900 MHz was in the anterior chamber while the highest for the frequency of 1800 MHz was in the vitreous. The temperature distribution in the eye induced by EM fields was not directly related to the SAR distribution due to the effect of the interaction among the dielectric properties, thermal properties, blood perfusion, and penetration depth of the EM power. Moreover, this study also showed that the exposure time had an influence on the temperature increase in the eye.

Teerapot Wessapan, Phadungsak Rattanadecho. Influence of ambient temperature on heat transfer in the human eye during exposure to electromagnetic fields at 900 MHz. International Journal of Heat and Mass Transfer 70: 378-388, 2014.

The topic of temperature increase in human tissue when exposed to EM fields, particularly those radiated to the eye, has been of interest for many years. This study presents a numerical analysis of the specific absorption rate (SAR) and the heat transfer in a heterogeneous two-dimensional human eye model exposed to TM-mode of electromagnetic (EM) fields of 900 MHz at various power densities. In this study, the effects of ambient temperature and power density on the temperature distributions and fluid flow in the eye during exposure to electromagnetic fields were systematically investigated. The electric field, SAR, temperature distribution and fluid flow in various tissues in the eye during exposure to EM fields were obtained by numerical simulation of EM wave propagation and a heat transfer model. The heat transfer model was then developed based on the porous media theories. The study highlights heat transfer and fluid flow in the eye during exposure to EM fields at different ambient temperatures. This study indicated that when the eye exposed to EM fields at the frequency of 900 MHz, the highest electric field intensity and SAR values at the chosen frequency was in the cornea. At the highest power density of 100 mW/cm^2 , the absorbed EM energy is converted to heat causes a further increase of 3°C in corneal temperature in cases of

Studies that show **Cell Phone** Health Effects

hot, moderate and cold ambient temperatures. The result shows important information related to a complex interaction between ambient temperature, fluid flow and temperature distribution in the eye during exposure to electromagnetic fields. Moreover, this study also showed that the power density had a strong influence on the temperature increase and fluid flow in the eye.

Balci M, Devrim E, Durak I. Effects of mobile phones on oxidant/antioxidant balance in cornea and lens of rats. *Curr Eye Res.* 32(1):21-25, 2007.

Purpose: To investigate the effects of mobile-phone-emitted radiation on the oxidant/antioxidant balance in corneal and lens tissues and to observe any protective effects of vitamin C in this setting. Methods: Forty female albino Wistar rats were assigned to one of four groups containing 10 rats each. One group received a standardized daily dose of mobile phone radiation for 4 weeks. The second group received this same treatment along with a daily oral dose of vitamin C (250 mg/kg). The third group received this dose of vitamin C alone, while the fourth group received standard laboratory care and served as a control. In corneal and lens tissues, malondialdehyde (MDA) levels and activities of superoxide dismutase (SOD), glutathione peroxidase (GSH-Px), and catalase (CAT) were measured with spectrophotometric methods. Results: In corneal tissue, MDA level and CAT activity significantly increased in the mobile phone group compared with the mobile phone plus vitamin C group and the control group ($p < 0.05$), whereas SOD activity was significantly decreased ($p < 0.05$). In the lens tissues, only the MDA level significantly increased in the mobile phone group relative to mobile phone plus vitamin C group and the control groups ($p < 0.05$). In lens tissue, significant differences were not found between the groups in terms of SOD, GSH-Px, or CAT ($p > 0.05$). Conclusions: The results of this study suggest that mobile telephone radiation leads to oxidative stress in corneal and lens tissues and that antioxidants such as vitamin C can help to prevent these effects

Lixia S, Yao K, Kaijun W, Deqiang L, Huajun H, Xiangwei G, Baohong W, Wei Z, Jianling L, Wei W. Effects of 1.8GHz radiofrequency field on DNA damage and expression of heat shock protein 70 in human lens epithelial cells. *Mutat Res.* 602(1-2):135-142, 2006.

To investigate the DNA damage, expression of heat shock protein 70 (Hsp70) and cell proliferation of human lens epithelial cells (hLEC) after exposure to the 1.8GHz radiofrequency field (RF) of a global system for mobile communications (GSM). An Xc-1800 RF exposure system was used to employ a GSM signal at 1.8GHz (217Hz amplitude-modulated) with the output power in the specific absorption rate (SAR) of 1, 2 and 3W/kg. After 2h exposure to RF, the DNA damage of hLEC was accessed by comet assay at five different incubation times: 0, 30, 60, 120 and 240min, respectively. Western blot and RT-PCR were used to determine the expression of Hsp70 in hLECs after RF exposure. The proliferation rate of cells was evaluated by bromodeoxyuridine incorporation on days 0, 1 and 4 after exposure. The results show that the difference of DNA-breaks between the exposed and sham-exposed (control) groups induced by 1 and 2W/kg

Studies that show **Cell Phone** Health Effects

irradiation were not significant at any incubation time point ($P>0.05$). The DNA damage caused by 3W/kg irradiation was significantly increased at the times of 0 and 30min after exposure ($P<0.05$), a phenomenon that could not be seen at the time points of 60, 120 or 240min ($P>0.05$). Detectable mRNA as well as protein expression of Hsp70 was found in all groups. Exposure at SARs of 2 and 3W/kg for 2h exhibited significantly increased Hsp70 protein expression ($P<0.05$), while no change in Hsp70 mRNA expression could be found in any of the groups ($P>0.05$). No difference of the cell proliferation rate between the sham-exposed and exposed cells was found at any exposure dose tested ($P>0.05$). The results indicate that exposure to non-thermal dosages of RF for wireless communications can induce no or repairable DNA damage and the increased Hsp70 protein expression in hLECs occurred without change in the cell proliferation rate. The non-thermal stress response of Hsp70 protein increase to RF exposure might be involved in protecting hLEC from DNA damage and maintaining the cellular capacity for proliferation.

Gasmelseed A. Electromagnetic energy absorption patterns in subjects with common visual disorders. *Electromagn Biol Med.* 30(3):136-145, 2011.

This article describes the analysis of electromagnetic energy absorption properties of models of the human eye with common visual disorders. The investigation addresses two types of visual disorders, namely hyperopia (or farsightedness) and myopia (or nearsightedness). Calculations were carried out using plane multilayered method with common wireless communication frequencies of 900, 1800, and 2450 MHz. The effect of wireless radiation on the eye is studied by calculation of the specific absorption rate (SAR) in three different eye models. The results of the simulations confirmed the anticipated and more complex relationship between absorption and structural variations of the eye at these frequencies.

Ozguner F, Bardak Y, Comlekci S. Protective effects of melatonin and caffeic acid phenethyl ester against retinal oxidative stress in long-term use of mobile phone: A comparative study. *Mol Cell Biochem.* 282(1-2):83-88, 2006.

There are numerous reports on the effects of electromagnetic radiation (EMR) in various cellular systems. Melatonin and caffeic acid phenethyl ester (CAPE), a component of honeybee propolis, were recently found to be potent free radical scavengers and antioxidants. Mechanisms of adverse effects of EMR indicate that reactive oxygen species may play a role in the biological effects of this radiation. The present study was carried out to compare the efficacy of the protective effects of melatonin and CAPE against retinal oxidative stress due to long-term exposure to 900 MHz EMR emitting mobile phones. Melatonin and CAPE were administered daily for 60 days to the rats prior to their EMR exposure during our study. Nitric oxide (NO, an oxidant product) levels and malondialdehyde (MDA, an index of lipid peroxidation), were used as markers of retinal oxidative stress in rats following to use of EMR. Superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GSH-Px) activities were studied to evaluate the changes of antioxidant status in retinal tissue. Retinal levels of NO and MDA increased in EMR exposed rats while both melatonin and

Studies that show **Cell Phone** Health Effects

CAPE caused a significant reduction in the levels of NO and MDA. Likewise, retinal SOD, GSH-Px and CAT activities decreased in EMR exposed animals while melatonin and CAPE caused a significant increase in the activities of these antioxidant enzymes. Treatment of EMR exposed rats with melatonin or CAPE increased the activities of SOD, GSH-Px and CAT to higher levels than those of control rats. In conclusion, melatonin and CAPE reduce retinal oxidative stress after long-term exposure to 900 MHz emitting mobile phone. Nevertheless, there was no statistically significant difference between the efficacies of these two antioxidants against to EMR induced oxidative stress in rat retina. The difference was in only GSH-Px activity in rat retina. Melatonin stimulated the retinal GSH-Px activity more efficiently than CAPE did.

Children Health and Cell Phone Use

Wang PW, Liu TL, Ko CH, Lin HC, Huang MF, Yeh YC, Yen CF. Association between Problematic Cellular Phone Use and Suicide: The Moderating Effect of Family Function and Depression. *Compr Psychiatry*. 2013 Oct 19. pii: S0010-440X(13)00282-4. doi: 10.1016/j.comppsy.2013.09.006. [Epub ahead of print]

BACKGROUND: Suicidal ideation and attempt among adolescents are risk factors for eventual completed suicide. Cellular phone use (CPU) has markedly changed the everyday lives of adolescents. Issues about how cellular phone use relates to adolescent mental health, such as suicidal ideation and attempts, are important because of the high rate of cellular phone usage among children in that age group. This study explored the association between problematic CPU and suicidal ideation and attempts among adolescents and investigated how family function and depression influence the association between problematic CPU and suicidal ideation and attempts. METHODS: A total of 5051 (2872 girls and 2179 boys) adolescents who owned at least one cellular phone completed the research questionnaires. We collected data on participants' CPU and suicidal behavior (ideation and attempts) during the past month as well as information on family function and history of depression. RESULTS: Five hundred thirty-two adolescents (10.54%) had problematic CPU. The rates of suicidal ideation were 23.50% and 11.76% in adolescents with problematic CPU and without problematic CPU, respectively. The rates of suicidal attempts in both groups were 13.70% and 5.45%, respectively. Family function, but not depression, had a moderating effect on the association between problematic CPU and suicidal ideation and attempt. CONCLUSION: This study highlights the association between problematic CPU and suicidal ideation as well as attempts and indicates that good family function may have a more significant role on reducing the risks of suicidal ideation and attempts in adolescents with problematic CPU than in those without problematic CPU.

Leena K, Tomi L, Arja RR. Intensity of mobile phone use and health compromising behaviours-how is information and communication technology connected to health-related lifestyle in adolescence? *J Adolesc*. 28(1):35-47, 2005.

The association of mobile phone use with health compromising behaviours (smoking, snuffing, alcohol) was studied in a survey comprising a representative sample of 14-16-

Studies that show **Cell Phone** Health Effects

year-olds ([Formula: see text]) in 2001. Mobile phone was used by 89% of respondents and by 13% for at least 1h daily. The intensity of use was positively associated with health compromising behaviours. The associations remained, although somewhat reduced, after including weekly spending money in the models. This study concludes that, at least in the present developmental level of communication technologies, intensive mobile phone use seems to be part of the same health-related lifestyle as health compromising behaviours.

Redmayne M, Smith E, Abramson MJ. Adolescent in-school cellphone habits: a census of rules, survey of their effectiveness, and fertility implications. Reprod Toxicol. 32(3):354-359, 2011.

We explored school cellphone rules and adolescent exposure to cellphone microwave emissions during school with a census and survey, respectively. The data were used to assess health and policy implications through a review of papers assessing reproductive bio-effects after exposure to cellphone emissions, this being most relevant to students' exposure. All schools banned private use of cellphones in class. However, 43% of student participants admitted breaking this rule. A high-exposure group of risk-takers was identified for whom prohibited in-school use was positively associated with high texting rates, carrying the phone switched-on >10h/day, and in-pocket use. The fertility literature is inconclusive, but increasingly points towards significant time- and dose-dependent deleterious effects from cellphone exposure on sperm. Genotoxic effects have been demonstrated from 'non-thermal' exposures, but not consistently. There is sufficient evidence and expert opinion to warrant an enforced school policy removing cellphones from students during the day.

Redmayne M. New Zealand adolescents' cellphone and cordless phone user-habits: are they at increased risk of brain tumours already? A cross-sectional study. Environ Health. 12(1):5, 2013.

BACKGROUND: Cellphone and cordless phone use is very prevalent among early adolescents, but the extent and types of use is not well documented. This paper explores how, and to what extent, New Zealand adolescents are typically using and exposed to active cellphones and cordless phones, and considers implications of this in relation to brain tumour risk, with reference to current research findings. **METHODS:** This cross-sectional study recruited 373 Year 7 and 8 school students with a mean age of 12.3 years (range 10.3-13.7 years) from the Wellington region of New Zealand. Participants completed a questionnaire and measured their normal body-to-phone texting distances. Main exposure-metrics included self-reported time spent with an active cellphone close to the body, estimated time and number of calls on both phone types, estimated and actual extent of SMS text-messaging, cellphone functions used and people texted. Statistical analyses used Pearson Chi2 tests and Pearson's correlation coefficient (r). Analyses were undertaken using SPSS version 19.0. **RESULTS:** Both cellphones and cordless phones were used by approximately 90% of students. A third of participants had already used a cordless phone for ≥ 7 years. In 4 years from the survey to mid-2013, the cordless phone use of 6% of participants would equal that of the

Studies that show **Cell Phone** Health Effects

highest Interphone decile (≥ 1640 hours), at the surveyed rate of use. High cellphone use was related to cellphone location at night, being woken regularly, and being tired at school. More than a third of parents thought cellphones carried a moderate-to-high health risk for their child. **CONCLUSIONS:** While cellphones were very popular for entertainment and social interaction via texting, cordless phones were most popular for calls. If their use continued at the reported rate, many would be at increased risk of specific brain tumours by their mid-teens, based on findings of the Interphone and Hardell-group studies.

Redmayne M, Smith E, and Abramson MJ. The relationship between adolescents' well-being and their wireless phone use: a cross-sectional study. Environmental Health 12(1):90, 2013.

Background. The exposure of young people to radiofrequency electromagnetic fields (RF-EMFs) has increased rapidly in recent years with their increased use of cellphones and use of cordless phones and WiFi. We sought to ascertain associations between New Zealand early-adolescents' subjective well-being and self-reported use of, or exposure to, wireless telephone and internet technology. **Methods.** In this cross-sectional survey, participants completed questionnaires in class about their cellphone and cordless phone use, their self-reported well-being, and possible confounding information such as whether they had had influenza recently or had a television in the bedroom. Parental questionnaires provided data on whether they had WiFi at home and cordless phone ownership and model. Data were analysed with Ordinal Logistic Regression adjusting for common confounders. Odds ratios (OR) and 95% confidence intervals were calculated. **Results.** The number and duration of cellphone and cordless phone calls were associated with increased risk of headaches (>6 cellphone calls over 10 minutes weekly, adjusted OR 2.4, CI 1.2-4.8; >15 minutes cordless use daily adjusted OR 1.74, CI 1.1-2.9)). Texting and extended use of wireless phones was related to having a painful 'texting' thumb). Using a wired cellphone headset was associated with tinnitus (adjusted OR 1.8, CI 1.0-3.3), while wireless headsets were associated with headache (adjusted OR 2.2, CI 1.1-4.5), feeling down/depressed (adjusted OR 2.0, CI 1.1-3.8), and waking in the night (adjusted OR 2.4, CI 1.2-4.8). Several cordless phone frequencies bands were related to tinnitus, feeling down/depressed and sleepiness at school, while the last of these was also related to modulation. Waking nightly was less likely for those with WiFi at home (adjusted OR 0.7, CI 0.4-0.99). Being woken at night by a cellphone was strongly related to tiredness at school (OR 4.1, CI 2.2-7.7). **Conclusions .** There were more statistically significant associations (36%) than could be expected by chance (5%). Several were dose-dependent relationships. To safeguard young people's well-being, we suggest limiting their use of cellphones and cordless phones to less than 15 minutes daily, and employing a speaker-phone device for longer daily use. We recommend parental measures are taken to prevent young people being woken by their cellphones.

Pedersen W. [Mobile phones, web chat, and sex among Norwegian adolescents]

Tidsskr Nor Laegeforen. 124(13-14):1756-1759, 2004. [article in Norwegian]

BACKGROUND: We investigated the associations between new interactive technology

Studies that show **Cell Phone** Health Effects

for communication, such as web chat or mobile phones, and sexual behaviour among Norwegian adolescents. **MATERIALS AND METHODS:** A representative sample of adolescents (age 13-18, N = 10,926) filled in a questionnaire during school hours; the response rate was 92%. **RESULTS:** Most adolescents have access to communication technology, but how much they use it varies. In particular with regard to mobile phones, a strong association to sexual behaviour was found. Among those who did not use the new technology, less than 10% reported having had intercourse while two out of three of the most active users reported intercourse. The associations remained significant when controls were made for age and a range of contextual, family, peer and individual factors. **INTERPRETATION:** Norwegian adolescents have changed their sexual behaviour over the last decade. The introduction and widespread use of new communication technology is one of the most salient changes over the same period. The findings suggest that this technology may in fact be of importance to teenagers' sexual socialisation.

Thomas S, Heinrich S, von Kries R, Radon K. Exposure to radio-frequency electromagnetic fields and behavioural problems in Bavarian children and adolescents. Eur J Epidemiol. 25(2):135-141, 2010.

Only few studies have so far investigated possible health effects of radio-frequency electromagnetic fields (RF EMF) in children and adolescents, although experts discuss a potential higher vulnerability to such fields. We aimed to investigate a possible association between measured exposure to RF EMF fields and behavioural problems in children and adolescents. 1,498 children and 1,524 adolescents were randomly selected from the population registries of four Bavarian (South of Germany) cities. During an Interview data on participants' mental health, socio-demographic characteristics and potential confounders were collected. Mental health behaviour was assessed using the German version of the Strengths and Difficulties Questionnaire (SDQ). Using a personal dosimeter, we obtained radio-frequency EMF exposure profiles over 24 h. Exposure levels over waking hours were expressed as mean percentage of the reference level. Overall, exposure to radiofrequency electromagnetic fields was far below the reference level. Seven percent of the children and 5% of the adolescents showed an abnormal mental behaviour. In the multiple logistic regression analyses measured exposure to RF fields in the highest quartile was associated to overall behavioural problems for adolescents (OR 2.2; 95% CI 1.1-4.5) but not for children (1.3; 0.7-2.6). These results are mainly driven by one subscale, as the results showed an association between exposure and conduct problems for adolescents (3.7; 1.6-8.4) and children (2.9; 1.4-5.9). As this is one of the first studies that investigated an association between exposure to mobile telecommunication networks and mental health behaviour more studies using personal dosimetry are warranted to confirm these findings.

Thomas S, Benke G, Dimitriadis C, Inyang I, Sim MR, Wolfe R, Croft RJ, Abramson MJ. Use of mobile phones and changes in cognitive function in adolescents. Occup Environ Med. 67(12):861-866, 2010.

Background Several studies have investigated the impact of mobile phone exposure on

Studies that show **Cell Phone** Health Effects

cognitive function in adults. However, children and adolescents are of special interest due to their developing nervous systems. Methods Data were derived from the Australian Mobile Radiofrequency Phone Exposed Users' Study (MoRPhEUS) which comprised a baseline examination of year 7 students during 2005/2006 and a 1-year follow-up. Sociodemographic and exposure data were collected with a questionnaire. Cognitive functions were assessed with a computerised test battery and the Stroop Color-Word test. Results 236 students participated in both examinations. The proportion of mobile phone owners and the number of voice calls and short message services (SMS) per week increased from baseline to follow-up. Participants with more voice calls and SMS at baseline showed less reductions in response times over the 1-year period in various computerised tasks. Furthermore, those with increased voice calls and SMS exposure over the 1-year period showed changes in response time in a simple reaction and a working memory task. No associations were seen between mobile phone exposure and the Stroop test. Conclusions We have observed that some changes in cognitive function, particularly in response time rather than accuracy, occurred with a latency period of 1 year and that some changes were associated with increased exposure. However, the increased exposure was mainly applied to those who had fewer voice calls and SMS at baseline, suggesting that these changes over time may relate to statistical regression to the mean, and not be the effect of mobile phone exposure.

Sudan M, Kheifets L, Arah O, Olsen J, Zeltzer L. Prenatal and Postnatal Cell Phone Exposures and Headaches in Children. Open Pediatr Med Journal. 6(2012):46-52, 2012.

OBJECTIVE: Children today are exposed to cell phones early in life, and may be at the greatest risk if exposure is harmful to health. We investigated associations between cell phone exposures and headaches in children. STUDY DESIGN: The Danish National Birth Cohort enrolled pregnant women between 1996 and 2002. When their children reached age seven years, mothers completed a questionnaire regarding the child's health, behaviors, and exposures. We used multivariable adjusted models to relate prenatal only, postnatal only, or both prenatal and postnatal cell phone exposure to whether the child had migraines and headache-related symptoms. RESULTS: Our analyses included data from 52,680 children. Children with cell phone exposure had higher odds of migraines and headache-related symptoms than children with no exposure. The odds ratio for migraines was 1.30 (95% confidence interval: 1.01-1.68) and for headache-related symptoms was 1.32 (95% confidence interval: 1.23-1.40) for children with both prenatal and postnatal exposure. CONCLUSIONS: In this study, cell phone exposures were associated with headaches in children, but the associations may not be causal given the potential for uncontrolled confounding and misclassification in observational studies such as this. However, given the widespread use of cell phones, if a causal effect exists it would have great public health impact.

Sudan M, Kheifets L, Arah OA, Olsen J. Cell phone exposures and hearing loss in children in the Danish National Birth Cohort. Paediatr Perinat Epidemiol. 27(3):247-257, 2013.

Studies that show **Cell Phone** Health Effects

BACKGROUND: Children today are exposed to cell phones early in life, and may be the most vulnerable if exposure is harmful to health. We investigated the association between cell phone use and hearing loss in children. **METHODS:** The Danish National Birth Cohort (DNBC) enrolled pregnant women between 1996 and 2002. Detailed interviews were conducted during gestation, and when the children were 6 months, 18 months and 7 years of age. We used multivariable-adjusted logistic regression, marginal structural models (MSM) with inverse-probability weighting, and doubly robust estimation (DRE) to relate hearing loss at age 18 months to cell phone use at age 7 years, and to investigate cell phone use reported at age 7 in relation to hearing loss at age 7. **RESULTS:** Our analyses included data from 52 680 children. We observed weak associations between cell phone use and hearing loss at age 7, with odds ratios and 95% confidence intervals from the traditional logistic regression, MSM and DRE models being 1.21 [95% confidence interval [CI] 0.99, 1.46], 1.23 [95% CI 1.01, 1.49] and 1.22 [95% CI 1.00, 1.49], respectively. **CONCLUSIONS:** Our findings could have been affected by various biases and are not sufficient to conclude that cell phone exposures have an effect on hearing. This is the first large-scale epidemiologic study to investigate this potentially important association among children, and replication of these findings is needed.

Krause CM, Bjornberg CH, Pesonen M, Hulten A, Liesivuori T, Koivisto M, Revonsuo A, Laine M, Hamalainen H. Mobile phone effects on children's event-related oscillatory EEG during an auditory memory task. *Int J Radiat Biol.* 82(6):443-450, 2006.

Purpose: To assess the effects of electromagnetic fields (EMF) emitted by mobile phones (MP) on the 1 - 20 Hz event-related brain oscillatory EEG (electroencephalogram) responses in children performing an auditory memory task (encoding and recognition). **Materials and methods:** EEG data were gathered while 15 subjects (age 10 - 14 years) performed an auditory memory task both with and without exposure to a digital 902 MHz MP in counterbalanced order. **Results:** During memory encoding, the active MP modulated the event-related desynchronization/synchronization (ERD/ERS) responses in the approximately 4 - 8 Hz EEG frequencies. During recognition, the active MP transformed these brain oscillatory responses in the approximately 4 - 8 Hz and approximately 15 Hz frequencies. **Conclusions:** The current findings suggest that EMF emitted by mobile phones has effects on brain oscillatory responses during cognitive processing in children.

Kamibepu K, Sugiura H. Impact of the mobile phone on junior high-school students' friendships in the Tokyo metropolitan area. *Cyberpsychol Behav.* 8(2):121-130, 2005.

The proportion of having keitai (Japanese mobile phone) has increased rapidly in young children. To research how junior high school students use their own keitai and to examine the impact of using it on their psychology, especially on their friendship, we recruited 651 students, grade 8, from five public junior high schools in the Tokyo metropolitan area. Each student participant completed a questionnaire that we had created. The response rates were 88.8% (n = 578) for participants. The proportion of having their own keitai was 49.3% (n = 285) and that of not having it was 50.7% (n = 293). We found that they used it much more frequently for e-mail than as a phone.

Studies that show **Cell Phone** Health Effects

Most of them exchanged e-mails between schoolmates, and more than a half of them exchanged e-mails more than 10 times a day. Sociable students estimated that their own keitai was useful for their friendship. But they experienced some insecurity or started staying up late at night engaged in e-mail exchanges, and they thought that they could not live without their own keitai. Our findings suggest that keitai having an e-mail function play a big part in the junior high-school students' daily life, and its impact on students' friendships, psychology, or health should be discussed among students to prevent keitai addiction.

Soderqvist F, Carlberg M, Hardell L. Use of wireless telephones and self-reported health symptoms: a population-based study among Swedish adolescents aged 15-19 years. *Environ Health*. 7(1):18, 2008.

ABSTRACT: BACKGROUND: Despite the last years of rapid increase in use of wireless phones little data on the use of these devices has been systematically assessed among young persons. The aim of this descriptive cross-sectional study was to assess use of wireless phones and to study such use in relation to explanatory factors and self-reported health symptoms. METHODS: A postal questionnaire comprising 8 pages of 27 questions with 75 items in total was sent to 2000 Swedish adolescents aged 15-19 years and selected from the population registry using a stratified sampling scheme. RESULTS: The questionnaire was answered by 63.5 % of the study subjects. Most participants reported access to a mobile phone (99.6%) and use increased with age; 55.6% of the 15-year-olds and 82.2% of the 19-year-olds were regular users. Girls generally reported more frequent use than boys. Use of wired hands-free equipment 'anytime' was reported by 17.4%. Cordless phones were used by 81.9%, and 67.3% were regular users. Watching TV increased the odds ratio for use of wireless phones, adjusted for age and gender. Some of the most frequently reported health complaints were tiredness, stress, headache, anxiety, concentration difficulties and sleep disturbances. Regular users of wireless phones had health symptoms more often and reported poorer perceived health than less frequent users. CONCLUSIONS: Almost all adolescence in this study used a wireless phone, girls more than boys. The most frequent use was seen among the older adolescents and those who watched TV extensively. The study further showed that perceived health and certain health symptoms seemed to be related to the use of wireless phones. However, this part of the investigation was explorative and should therefore be interpreted with caution since bias and chance findings due to multiple testing might have influenced the results. Potentially this study will stimulate more sophisticated studies that may also investigate directions of associations and whether, or to what degree, any mediation factors are involved.

Schuz J. Mobile phone use and exposures in children. *Bioelectromagnetics*. Suppl 7:S45-50,2005.

The main difference concerning the use of mobile phones (MPs) between today's children and adults is the longer lifetime exposure of children when they grow older, due to starting to use MPs at an early age. Additionally, recent trends lead to a higher frequency of use among children, including higher popularity of MPs and features

Studies that show **Cell Phone** Health Effects

specifically designed to attract children. The prevalence of MP users is already very high and reaches >90% among adolescents in some countries. In a German study, 6% of 9-10 years old children used a MP for making calls daily; 35% owned their own MP. For children, MPs are dominant sources of radio wave exposures and relevant sources of extremely low frequency magnetic fields. For very young children, however, environmental exposure to radio waves may be of concern. In conclusion, children will have a much higher cumulative exposure to radio waves than today's adults when they are at the same age. Radio wave exposure of children may be estimated more easily, because the variety of exposure sources is smaller than for adults. As long as adverse health effects cannot be ruled out with some degree of certainty, it appears to be appropriate to instruct children and their parents about a prudent use of MPs.

Heinrich S, Thomas S, Heumann C, von Kries R, Radon K. Association between exposure to radiofrequency electromagnetic fields assessed by dosimetry and acute symptoms in children and adolescents: a population based cross-sectional study. *Environ Health*. 9:75, 2010.

BACKGROUND: The increase in numbers of mobile phone users was accompanied by some concern that exposure to radiofrequency electromagnetic fields (RF EMF) might adversely affect acute health especially in children and adolescents. The authors investigated this potential association using personal dosimeters. METHODS: A 24-hour exposure profile of 1484 children and 1508 adolescents was generated in a population-based cross-sectional study in Germany between 2006 and 2008 (participation 52%). Personal interview data on socio-demographic characteristics, self-reported exposure and potential confounders were collected. Acute symptoms were assessed twice during the study day using a symptom diary. RESULTS: Only few of the large number of investigated associations were found to be statistically significant. At noon, adolescents with a measured exposure in the highest quartile during morning hours reported a statistically significant higher intensity of headache (Odd Ratio: 1.50; 95% confidence interval: 1.03, 2.19). At bedtime, adolescents with a measured exposure in the highest quartile during afternoon hours reported a statistically significant higher intensity of irritation in the evening (4th quartile 1.79; 1.23, 2.61), while children reported a statistically significant higher intensity of concentration problems (4th quartile 1.55; 1.02, 2.33). CONCLUSIONS: *We observed few statistically significant results which are not consistent over the two time points.* Furthermore, when the 10% of the participants with the highest exposure are taken into consideration the significant results of the main analysis could not be confirmed. Based on the pattern of these results, we assume that the few observed significant associations are not causal but rather occurred by chance.

Zheng F, Gao P, He M, Li M, Wang C, Zeng Q, Zhou Z, Yu Z, Zhang L. Association between mobile phone use and inattention in 7102 Chinese adolescents: a population-based cross-sectional study. *BMC Public Health*. 2014 Oct 1;14(1):1022. [Epub ahead of print]

Studies that show **Cell Phone** Health Effects

BACKGROUND: The dramatic growth of **mobile phone** (MP) use among young people has increased interest in its possible health hazards in this age group. The aim of this cross-sectional study was to investigate the association between MP use and inattention in adolescents. **METHODS:** A total of 7720 middle school students were involved in this cross-sectional study. Inattention was assessed as defined for the Attention Deficit component of Attention deficit/Hyperactivity disorder (ADHD) by the Diagnostic and Statistical Manual of Mental Disorders (4th ed., text rev. [DSM-IV-TR]). The demographic characteristics and information on MP use were included in the questionnaire. Chi-square tests and logistic regression models were used to analyze the data. **RESULTS:** In total, 7102 (91.99%) valid questionnaires were obtained. After adjusted for confounders, inattention in adolescents was significantly associated with MP ownership, the time spent on entertainment on MP per day, the position of the MP during the day and the mode of the MP at night. The strongest association between inattention and the time spent on the MP was among students who spent more than 60 minutes per day playing on their MP. **CONCLUSIONS:** Our study shows some associations between MP use and inattention in Chinese adolescents. Decreasing MP usage to less than 60 minutes per day may help adolescents to stay focused and centered.

Chiu CT, Chang YH, Chen CC, Ko MC, Li CY. Mobile phone use and health symptoms in children. J Formos Med Assoc. 2014 Aug 9. pii: S0929-6646(14)00207-1. doi: 10.1016/j.jfma.2014.07.002. [Epub ahead of print]

BACKGROUND/PURPOSE: To investigate the mobile phone (MP) use for talking in relation to health symptoms among 2042 children aged 11-15 years in Taiwan. **METHODS:** A nationwide, cross-sectional study, using the computer assisted telephone interview (CATI) technique, was conducted in 2009 to collect information on children's utilization of MPs and the perceived health symptoms reported by their parents. **RESULTS:** The overall prevalence of MP use in the past month was estimated at 63.2% [95% confidence interval (CI) = 61.1-65.3%]. MP use was associated with a significantly increased adjusted odds ratio (AOR) for headaches and migraine (1.42, 95% CI = 1.12-1.81) and skin itches (1.84, 95% CI = 1.47-2.29). Children who regularly used MPs were also considered to have a health status worse than it was 1 year ago ($\beta = 0.27$, 95% CI = 0.17-0.37). **CONCLUSION:** Although the cross-sectional design precludes the causal inference for the observed association, our study tended to suggest a need for more cautious use of MPs in children, because children are expected to experience a longer lifetime exposure to radiofrequency electromagnetic fields (RF-EMF) from MPs.

Peyman A, Rezazadeh AA, Gabriel C. Changes in the dielectric properties of rat tissue as a function of age at microwave frequencies. Phys Med Biol 46(6):1617-1629, 2001. The dielectric properties of ten rat tissues at six different ages were measured at 37 degrees C in the frequency range of 130 MHz to 10 GHz using an open-ended coaxial probe and a computer controlled network analyser. The results show a general decrease of the dielectric properties with age. The trend is more apparent for brain, skull and skin tissues and less noticeable for abdominal tissues. The variation in the dielectric properties with age is due to the changes in the water content and the organic

Studies that show **Cell Phone** Health Effects

composition of tissues. The percentage decrease in the dielectric properties of certain tissues in the 30 to 70 day old rats at cellular phone frequencies have been tabulated. These data provide an important input in the provision of rigorous dosimetry in lifetime-exposure animal experiments. The results provide some insight into possible differences in the assessment of exposure for children and adults.

de Salles AA, Bulla G, Rodriguez CE. Electromagnetic absorption in the head of adults and children due to mobile phone operation close to the head. Electromagn Biol Med. 2006;25(4):349-360, 2006.

The Specific Absorption Rate (SAR) produced by mobile phones in the head of adults and children is simulated using an algorithm based on the Finite Difference Time Domain (FDTD) method. Realistic models of the child and adult head are used. The electromagnetic parameters are fitted to these models. Comparison also are made with the SAR calculated in the children model when using adult human electromagnetic parameters values. Microstrip (or patch) antennas and quarter wavelength monopole antennas are used in the simulations. The frequencies used to feed the antennas are 1850 MHz and 850 MHz. The SAR results are compared with the available international recommendations. It is shown that under similar conditions, the 1g-SAR calculated for children is higher than that for the adults. When using the 10-year old child model, SAR values higher than 60% than those for adults are obtained.

Peyman A, Holden SJ, Watts S, Perrott R, Gabriel C. Dielectric properties of porcine cerebrospinal tissues at microwave frequencies: in vivo, in vitro and systematic variation with age. Phys Med Biol. 52(8):2229-2245, 2007.

The dielectric properties of pig cerebrospinal tissues were measured in vivo and in vitro, in the frequency range of 50 MHz-20 GHz. The total combined measurement uncertainty was calculated at each frequency point and is reported over representative frequency regions. Comparisons were made for each tissue between the two sets of data and with the literature of the past decade. The in vitro study was extended to include tissue from pigs weighing approximately 10, 50 and 250 kg to re-visit the question of the variation of dielectric properties with age. White matter and spinal chord showed significant variation as function of animal age, no age-related variations were recorded for grey matter.

Oxidative Stress

Stopczyk D, Gnitecki W, Buczynski A, Markuszewski L, Buczynski J. Med Pr 53(4):311-314, 2002. [Article in Polish]

The aim of the study was to assess in vitro the effect of electromagnetic field produced by mobile phones on the activity of superoxide dismutase (SOD-1) and the level of malonyldialdehyde (MDA) in human blood platelets. The suspension of blood platelets was exposed to the electromagnetic field with the frequency of 900 MHz for 1, 3, 5, and 7 min. Our studies demonstrated that microwaves produced by mobiles significantly

Studies that show **Cell Phone** Health Effects

depleted SOD-1 activity after 1, 5, and 7 min of exposure and increased after 3 min in comparison with the control test. There was a significant increase in the concentration of MDA after 1, 5, and 7 min and decrease after 3 min of exposure as compared with the control test. On the grounds of our results we conclude that oxidative stress after exposure to microwaves may be the reason for many adverse changes in cells and may cause a number of systemic disturbances in the human body.

Dasdag S, Akdag MZ. The link between radiofrequencies emitted from wireless technologies and oxidative stress. J Chem Neuroanat. 2015 Sep 12. pii: S0891-0618(15)00069-1. doi: 10.1016/j.jchemneu.2015.09.001. [Epub ahead of print]

Wireless communication such as cellular telephones and other types of handheld phones working with frequencies of 900MHz, 1800MHz, 2100MHz, 2450MHz have been increasing rapidly. Therefore, public opinion concern about the potential human health hazards of short and long-term effect of exposure to radiofrequency (RF) radiation. Oxidative stress is a biochemical condition, which is defined by the imbalance between reactive oxygen species (ROS) and the anti-oxidative defense. In this review, we evaluated available in vitro and in vivo studies carried out on the relation between RF emitted from mobile phones and oxidative stress. The results of the studies we reviewed here indicated that mobile phones and similar equipment or radars can be thought as a factor, which cause oxidative stress. Even some of them claimed that oxidative stress originated from radiofrequencies can be resulted with DNA damage. For this reason one of the points to think on is relation between mobile phones and oxidative stress. However, more performance is necessary especially on human exposure studies.

Esmekaya MA, Ozer C, Seyhan N. 900 MHz pulse-modulated radiofrequency radiation induces oxidative stress on heart, lung, testis and liver tissues. Gen Physiol Biophys. 30(1):84-89, 2011.

Oxidative stress may affect many cellular and physiological processes including gene expression, cell growth, and cell death. In the recent study, we aimed to investigate whether 900 MHz pulse-modulated radiofrequency (RF) fields induce oxidative damage on lung, heart and liver tissues. We assessed oxidative damage by investigating lipid peroxidation (malondialdehyde, MDA), nitric oxide (NOx) and glutathione (GSH) levels which are the indicators of tissue toxicity. A total of 30 male Wistar albino rats were used in this study. Rats were divided randomly into three groups; control group (n = 10), sham group (device off, n = 10) and 900 MHz pulsed-modulated RF radiation group (n = 10). The RF rats were exposed to 900 MHz pulsed modulated RF radiation at a specific absorption rate (SAR) level of 1.20 W/kg 20 min/day for three weeks. MDA and NOx levels were increased significantly in liver, lung, testis and heart tissues of the exposed group compared to sham and control groups ($p < 0.05$). Conversely GSH levels were significantly lower in exposed rat tissues ($p < 0.05$). No significantly difference was observed between sham and control groups. Results of our study showed that pulse-modulated RF radiation causes oxidative injury in liver, lung, testis and heart tissues mediated by lipid peroxidation, increased level of NOx and suppression of antioxidant

Studies that show **Cell Phone** Health Effects

defense mechanism.

Bodera P, Stankiewicz W, Zawada K, Antkowiak B, Paluch M, Kieliszek J, Kalicki B, Bartosiński A, Wawer I. Changes in antioxidant capacity of blood due to mutual action of electromagnetic field (1800 MHz) and opioid drug (tramadol) in animal model of persistent inflammatory state. Pharmacol Rep. 65(2):421-428, 2013.

Background: The biological effects and health implications of electromagnetic field (EMF) associated with cellular mobile telephones and related wireless systems and devices have become a focus of international scientific interest and world-wide public concern. It has also been proved that EMF influences the production of reactive oxygen species (ROS) in different tissues. Methods: Experiments were performed in healthy rats and in rats with persistent inflammatory state induced by Complete Freund's Adjuvant (CFA) injection, which was given 24 h before EMF exposure and drug application. Rats were injected with CFA or the same volume of paraffin oil into the plantar surface of the left hind paw. Animals were exposed to the far-field range of an antenna at 1800 MHz with the additional modulation which was identical to that generated by mobile phone GSM 1800. Rats were given 15 min exposure, or were sham-exposed with no voltage applied to the field generator in control groups. Immediately before EMF exposure, rats were injected intraperitoneally with tramadol in the 20 mg/kg dose or vehicle in the 1 ml/kg volume. Results: Our study revealed that single EMF exposure in 1800 MHz frequency significantly reduced antioxidant capacity both in healthy animals and those with paw inflammation. A certain synergic mode of action between applied electromagnetic fields and administered tramadol in rats treated with CFA was observed. Conclusions: The aim of the study was to examine the possible, parallel/combined effects of electromagnetic radiation, artificially induced inflammation and a centrally-acting synthetic opioid analgesic drug, tramadol, (used in the treatment of severe pain) on the antioxidant capacity of blood of rats. The antioxidant capacity of blood of healthy rats was higher than that of rats which received only tramadol and were exposed to electromagnetic fields.

Moustafa YM, Moustafa RM, Belacy A, Abou-El-Ela SH, Ali FM. Effects of acute exposure to the radiofrequency fields of cellular phones on plasma lipid peroxide and antioxidase activities in human erythrocytes. J Pharm Biomed Anal 26(4):605-608, 2001.

Radiofrequency fields of cellular phones may affect biological systems by increasing free radicals, which appear mainly to enhance lipid peroxidation, and by changing the antioxidase activities of human blood thus leading to oxidative stress. To test this, we have investigated the effect of acute exposure to radiofrequency fields of commercially available cellular phones on some parameters indicative of oxidative stress in 12 healthy adult male volunteers. Each volunteer put the phone in his pocket in standby position with the keypad facing the body. The parameters measured were lipid peroxide and the activities of superoxide dismutase (SOD), total glutathione peroxidase (GSH-Px) and catalase. The results obtained showed that the plasma level of lipid peroxide was

Studies that show **Cell Phone** Health Effects

significantly increased after 1, 2 and 4 h of exposure to radiofrequency fields of the cellular phone in standby position. Moreover, the activities of SOD and GSH-Px in human erythrocytes showed significant reduction while the activity of catalase in human erythrocytes did not decrease significantly. These results indicate that acute exposure to radiofrequency fields of commercially available cellular phones may modulate the oxidative stress of free radicals by enhancing lipid peroxidation and reducing the activation of SOD and GSH-Px, which are free radical scavengers. Therefore, these results support the interaction of radiofrequency fields of cellular phones with biological systems.

Oral B, Guney M, Ozguner F, Karahan N, Mungan T, Comlekci S, Cesur G. Endometrial Apoptosis Induced by a 900-MHz Mobile Phone: Preventive Effects of Vitamins E and C. *Adv Ther.* 23(6):957-973, 2006.

Numerous reports have described the effects induced by an electromagnetic field (EMF) in various cellular systems. The purposes of this study were to examine oxidative stress that promotes production of reactive oxygen species induced by a 900-megahertz (MHz) mobile phone and the possible ameliorating effects of vitamins E and C on endometrial tissue against EMF-induced endometrial impairment and apoptosis in rats. Animals were randomly grouped as follows: (1) sham-operated control group (n=8), (2) 900 MHz EMF-exposed group (n=8; 30 min/d for 30 d), and (3) 900 MHz EMF-exposed group, treated with vitamins E and C (n=8; 50 mg/kg intramuscularly and 20 mg/kg body weight intraperitoneally before daily EMF exposure). Malondialdehyde (an index of lipid peroxidation) was used as a marker of oxidative stress-induced endometrial impairment; Bcl-2, Bax, caspase-3, and caspase-8 were assessed immunohistochemically. In this study, increased malondialdehyde levels in endometrial tissue and apoptosis illustrated the role of the oxidative mechanism induced by exposure to a 900-MHz mobile phone-like device and vitamins E and C; via free radical scavenging and antioxidant properties, oxidative tissue injury and apoptosis were ameliorated in rat endometrium. In conclusion, exposure to 900-MHz radiation emitted by mobile phones may cause endometrial apoptosis and oxidative stress, but treatment with vitamins E and C can diminish these changes and may have a beneficial effect in preventing endometrial changes in rats.

Marjanovic AM, Pavicic I, Trosic I, Cell oxidation–reduction imbalance after modulated radiofrequency radiation. *Electromagnetic Biology and Medicine.* Posted online on August 13, 2014.

Aim of this study was to evaluate an influence of modulated radiofrequency field (RF) of 1800 MHz, strength of 30 V/m on oxidation–reduction processes within the cell. The assigned RF field was generated within Gigahertz Transversal Electromagnetic Mode cell equipped by signal generator, modulator, and amplifier. Cell line V79, was irradiated for 10, 30, and 60 min, specific absorption rate was calculated to be 1.6 W/kg. Cell metabolic activity and viability was determined by MTT assay. In order to define total protein content, colorimetric method was used. Concentration of oxidised proteins was evaluated by enzyme-linked immunosorbent assay. Reactive oxygen species (ROS)

Studies that show **Cell Phone** Health Effects

marked with fluorescent probe 2',7'-dichlorofluorescein diacetate were measured by means of plate reader device. In comparison with control cell samples, metabolic activity and total protein content in exposed cells did not differ significantly. Concentrations of carbonyl derivatives, a product of protein oxidation, insignificantly but continuously increase with duration of exposure. In exposed samples, ROS level significantly ($p < 0.05$) increased after 10 min of exposure. Decrease in ROS level was observed after 30-min treatment indicating antioxidant defence mechanism activation. In conclusion, under the given laboratory conditions, modulated RF radiation might cause impairment in cell oxidation–reduction equilibrium within the growing cells.

Kahya MC, Nazıroğlu M, Cığ B. Selenium Reduces Mobile Phone (900 MHz)-Induced Oxidative Stress, Mitochondrial Function, and Apoptosis in Breast Cancer Cells. Biol Trace Elem Res. 2014 Jun 27. [Epub ahead of print]

Exposure to mobile phone-induced electromagnetic radiation (EMR) may affect biological systems by increasing free oxygen radicals, apoptosis, and mitochondrial depolarization levels although selenium may modulate the values in cancer. The present study was designed to investigate the effects of 900 MHz radiation on the antioxidant redox system, apoptosis, and mitochondrial depolarization levels in MDA-MB-231 breast cancer cell line. Cultures of the cancer cells were divided into four main groups as controls, selenium, EMR, and EMR + selenium. In EMR groups, the cells were exposed to 900 MHz EMR for 1 h (SAR value of the EMR was 0.36 ± 0.02 W/kg). In selenium groups, the cells were also incubated with sodium selenite for 1 h before EMR exposure. Then, the following values were analyzed: (a) cell viability, (b) intracellular ROS production, (c) mitochondrial membrane depolarization, (d) cell apoptosis, and (e) caspase-3 and caspase-9 values. Selenium suppressed EMR-induced oxidative cell damage and cell viability (MTT) through a reduction of oxidative stress and restoring mitochondrial membrane potential. Additionally, selenium indicated anti-apoptotic effects, as demonstrated by plate reader analyses of apoptosis levels and caspase-3 and caspase-9 values. In conclusion, 900 MHz EMR appears to induce apoptosis effects through oxidative stress and mitochondrial depolarization although incubation of selenium seems to counteract the effects on apoptosis and oxidative stress.

Elhag MA, Nabil GM, Attia AM. Effects of electromagnetic field produced by mobile phones on the oxidant and antioxidant status of rats. Pak J Biol Sci. 10(23):4271-4274, 2007.

This study was designed to investigate the effect of EMR produced by GSM Mobile Phones (MP) on the oxidant and antioxidant status in rats. Rats were divided into three groups: (1) controls, (2) rats exposed to a fractionated dose of EMR (15 min day⁻¹) for four days (EMR-F) and (3) rats exposed to an acute dose of EMR (EMR-A). A net drop in the plasma concentration of vitamin C (-47 and -59.8%) was observed in EMR-F and EMR-A groups, respectively, when compared to controls. While, a significant decrease in the levels of lipophilic antioxidant vitamins: vitamin E (-33 and -65.8%), vitamin A (-44.4 and -46.8%) was observed in EMR-F and EMR-A groups, respectively, when compared to controls. A net drop in plasma level of reduced glutathione (GSH) (-19.8 and -35.3%) was

Studies that show **Cell Phone** Health Effects

observed in EMR-F and EMR-A groups, respectively. EMR exposure of rats produced a significant decrease in catalase (CAT) and superoxide dismutase (SOD) activities, with the values of these activities for EMR-A group is significantly lower than those of EMR-F. These results indicate that the effects of acute doses of EMR produced by mobile phones on the rat's antioxidant status is significantly higher than those of fractionated doses of the same type of radiation. On the basis of present results, it can be concluded that exposure to acute doses of EMR produced by mobile phones is more hazardous than that produced by fractionated doses of the same type of radiation.

Fattahi-Asl J, Baradaran-Ghahfarokhi M, Karbalaee M, Baradaran-Ghahfarokhi M, Baradaran-Ghahfarokhi HR. Effects of radiofrequency radiation on human ferritin: an in vitro enzymun assay. J Med Signals Sens. 2(4):235-340, 2012.

Ferritin is a macromolecule and is responsible for the long term iron storage function in human serum and plasma. Recent studies have highlighted the role of cell phone exposure on central nervous system, immune function and reproduction. The aim of this study was to investigate whether the human serum ferritin level could be interfered by the exposure to the 900 MHz GSM cell phones. Fifty human serum wells from 25 normal healthy donors were labeled with ruthenium to form a sandwich complex based on an immunoassay technique. All of them were placed into two batches, and the well heads in the first batch were exposed to 900 MHz exposure emitted from a speech mode cell phone (Nokia, Model 1202, India) for 30 min. Unexposed batch was served as the control sample under identical conditions and was compared with the exposed one in quantitative determination of ferritin using the Wilcoxon test with criterion level of $P = 0.050$. Human serum wells in the exposed batch showed a significant decrease in serum ferritin relative to the control batch ($P = 0.029$). The average \pm SD ferritin level in the exposed batch was $84.94 \pm 1.04 \mu\text{g/L}$ while it was $87.25 \pm 0.83 \mu\text{g/L}$ for the unexposed batch. Radiofrequency electromagnetic waves emitted from cell phones may lead to oxidative stress and rapid diffusion of the human ferritin level in an in vitro enzymun assay. Also, the enzyme activity can be affected. Effects of exposure from mobile phones must be considered further.

Irmak MK, Fadillioglu E, Gulec M, Erdogan H, Yagmurca M, Akyol O. Effects of electromagnetic radiation from a cellular telephone on the oxidant and antioxidant levels in rabbits. Cell Biochem Funct. 20(4):279-283, 2002.

The number of reports on the effects induced by electromagnetic radiation (EMR) in various cellular systems is still increasing. Until now no satisfactory mechanism has been proposed to explain the biological effects of this radiation. Oxygen free radicals may play a role in mechanisms of adverse effects of EMR. This study was undertaken to investigate the influence of electromagnetic radiation of a digital GSM mobile telephone (900 MHz) on oxidant and antioxidant levels in rabbits. Adenosine deaminase, xanthine oxidase, catalase, myeloperoxidase, superoxide dismutase (SOD) and glutathione peroxidase activities as well as nitric oxide (NO) and malondialdehyde levels were measured in sera and brains of EMR-exposed and sham-exposed rabbits. Serum SOD activity increased, and serum NO levels decreased in EMR-exposed animals compared to

Studies that show **Cell Phone** Health Effects

the sham group. Other parameters were not changed in either group. This finding may indicate the possible role of increased oxidative stress in the pathophysiology of adverse effect of EMR. Decreased NO levels may also suggest a probable role of NO in the adverse effect.

Guler G, Tomruk A, Ozgur E, Seyhan N. The effect of radiofrequency radiation on DNA and lipid damage in non-pregnant and pregnant rabbits and their newborns. *Gen Physiol Biophys.* 29(1):59-66, 2010.

The concerns of people on possible adverse health effects of radiofrequency radiation (RFR) generated from mobile phones as well as their supporting transmitters (base stations) have increased markedly. RFR effect on oversensitive people, such as pregnant women and their developing fetuses, and older people is another source of concern that should be considered. In this study, oxidative DNA damage and lipid peroxidation levels in the brain tissue of pregnant and non-pregnant New Zealand White rabbits and their newborns exposed to RFR were investigated. Thirteen-month-old rabbits were studied in four groups as non-pregnant-control, non-pregnant-RFR exposed, pregnant-control and pregnant-RFR exposed. They were exposed to RFR (1800 MHz GSM; 14 V/m as reference level) for 15 min/day during 7 days. Malondialdehyde (MDA) and 8-hydroxy-2'-deoxyguanosine (8-OHdG) levels were analyzed. MDA and 8-OHdG levels of non-pregnant and pregnant-RFR exposed animals significantly increased with respect to controls ($p < 0.001$, Mann-Whitney test). No difference was found in the newborns ($p > 0.05$, Mann-Whitney). There exist very few experimental studies on the effects of RFR during pregnancy. It would be beneficial to increase the number of these studies in order to establish international standards for the protection of pregnant women from RFR.

Burlaka A, Tsybulin O, Sidorik E, Lukin S, Polishuk V, Tsehmistrenko S, Yakymenko I. Overproduction of free radical species in embryonal cells exposed to low intensity radiofrequency radiation. *Exp Oncol.* 2013 Sep;35(3):219-225.

Aim: Long-term exposure of humans to low intensity radiofrequency electromagnetic radiation (RF-EMR) leads to a statistically significant increase in tumor incidence. Mechanisms of such the effects are unclear, but features of oxidative stress in living cells under RF-EMR exposure were previously reported. Our study aims to assess a production of initial free radical species, which lead to oxidative stress in the cell. Materials and Methods: Embryos of Japanese quails were exposed in ovo to extremely low intensity RF-EMR of GSM 900 MHz ($0.25 \mu\text{W}/\text{cm}^2$) during 158-360 h discontinuously (48 c - ON, 12 c - OFF) before and in the initial stages of development. The levels of superoxide ($\text{O}_2^{\cdot-}$), nitrogen oxide (NO^{\cdot}), thiobarbituric acid reactive substances (TBARS), 8-oxo-2'-deoxyguanosine (8-oxo-dG) and antioxidant enzymes' activities were assessed in cells/tissues of 38-h, 5- and 10-day RF-EMR exposed and unexposed embryos. Results: The exposure resulted in a significant persistent overproduction of superoxide and nitrogen oxide in embryo cells during all period of analyses. As a result, significantly increased levels of TBARS and 8-oxo-dG followed by significantly decreased levels of superoxide dismutase and catalase activities were developed in the exposed embryo

Studies that show **Cell Phone** Health Effects

cells. Conclusion: Exposure of developing quail embryos to extremely low intensity RF-EMR of GSM 900 MHz during at least one hundred and fifty-eight hours leads to a significant overproduction of free radicals/reactive oxygen species and oxidative damage of DNA in embryo cells. These oxidative changes may lead to pathologies up to oncogenic transformation of cells.

Tkalec M, Malarić K, Pevalek-Kozlina B. Exposure to radiofrequency radiation induces oxidative stress in duckweed *Lemna minor* L. *Sci Total Environ.* 388(1-3):78-89, 2007.

Widespread use of radiofrequency radiation emitting devices increased the exposure to electromagnetic fields (EMFs) from 300 MHz to 300 GHz. Various biological effects of exposure to these fields have been documented so far, but very little work has been carried out on plants. The aim of the present work was to investigate the physiological responses of the plant *Lemna minor* after exposure to radiofrequency EMFs, and in particular, to clarify the possible role of oxidative stress in the observed effects. Duckweed was exposed for 2 h to EMFs of 400 and 900 MHz at field strengths of 10, 23, 41 and 120 V m⁻¹. The effect of a longer exposure time (4 h) and modulation was also investigated. After exposure, parameters of oxidative stress, such as lipid peroxidation, H₂O₂ content, activities and isoenzyme pattern of antioxidative enzymes as well as HSP70 expression were evaluated. At 400 MHz, lipid peroxidation and H₂O₂ content were significantly enhanced in duckweed exposed to EMFs of 23 and 120 V m⁻¹ while other exposure treatments did not have an effect. Compared to the controls, the activities of antioxidative enzymes showed different behaviour: catalase (CAT) activity increased after most exposure treatments while pyrogallol (PPX) and ascorbate peroxidase (APX) activities were not changed. Exceptions were reduced PPX and APX activity after longer exposure at 23 V m⁻¹ and increased PPX activity after exposures at 10 and 120 V m⁻¹. By contrast, at 900 MHz almost all exposure treatments significantly increased level of lipid peroxidation and H₂O₂ content but mostly decreased PPX activity and did not affect CAT activity. Exceptions were exposures to a modulated field and to the field of 120 V m⁻¹ which increased PPX and CAT activity. At this frequency APX activity was significantly decreased after exposure at 10 V m⁻¹ and longer exposure at 23 V m⁻¹ but it increased after a shorter exposure at 23 V m⁻¹. At both frequencies no differences in isoenzyme patterns of antioxidative enzymes or HSP70 level were found between control and exposed plants. Our results showed that non-thermal exposure to investigated radiofrequency fields induced oxidative stress in duckweed as well as unspecific stress responses, especially of antioxidative enzymes. However, the observed effects markedly depended on the field frequencies applied as well as on other exposure parameters (strength, modulation and exposure time). Enhanced lipid peroxidation and H₂O₂ content accompanied by diminished antioxidative enzymes activity caused by exposure to investigated EMFs, especially at 900 MHz, indicate that oxidative stress could partly be due to changed activities of antioxidative enzymes.

Studies that show **Cell Phone** Health Effects

Cell Function Impairment

Palumbo R, Brescia F, Capasso D, Sannino A, Sarti M, Capri M, Grassilli E, Scarfi MR. Exposure to 900 MHz radiofrequency radiation induces caspase 3 activation in proliferating human lymphocytes. *Radiat Res.* 170(3):327-334, 2008.

In this study, the induction of apoptosis after exposure to 900 MHz radiofrequency radiation (GSM signal) was investigated by assessing caspase 3 activation in exponentially growing Jurkat cells and in quiescent and proliferating human peripheral blood lymphocytes (PBLs). The exposure was carried out at an average specific absorption rate of 1.35 W/kg in a dual wire patch cell exposure system where the temperature of cell cultures was accurately controlled. After 1 h exposure to the radiofrequency field, a slight but statistically significant increase in caspase 3 activity, measured 6 h after exposure, was observed in Jurkat cells (32.4%) and in proliferating human PBLs (22%). In contrast, no effect was detected in quiescent human PBLs. In the same experimental conditions, apoptosis was also evaluated in Jurkat cells by Western blot analysis and in both cell types by flow cytometry. To evaluate late effects due to caspase 3 activity, flow cytometry was also employed to assess apoptosis and viability 24 h after radiofrequency-radiation exposure in both cell types. Neither the former nor the latter was affected. Since in recent years it has been reported that caspases are also involved in processes other than apoptosis, additional cell cycle studies were carried out on proliferating T cells exposed to radiofrequency radiation; however, we found no differences between sham-exposed and exposed cultures. Further studies are warranted to investigate the biological significance of our findings of a dose-response increase in caspase 3 activity after exposure to radiofrequency radiation.

Obukhan KI, [The effect of ultrahigh-frequency radiation on adaptation thresholds and the damages to blood system cells]. *Lik Sprava* (7):71-73, 1998. [Article in Ukrainian]

Cytologic investigations designed to study bone marrow, peripheral blood, spleen, and thymus of albino rats irradiated by an electromagnetic field, 2375, 2450, and 3000 MEGS, revealed structural and functional changes in populations of megakaryocytes, immunocompetent cells as well as of undifferentiated cells, and of other types of cells that are dependent on the intensity of irradiation and permit establishing the probability-threshold levels of exposure taking account of reactions of perception and physiologic adaptation together with compensatory and regenerative processes and the injury sustained. It is shown that changes in bone marrow cells differentiation and reproduction rather than integral shifts in the peripheral blood that acquire the utmost significance. Subjected to a particular scrutiny in the paper are blast cells, which cells' repopulation was noted to be getting increased in low-intensity exposure as were disturbances in their mitosis pattern.

Nylund R, Leszczynski D. Mobile phone radiation causes changes in gene and protein

Studies that show **Cell Phone** Health Effects

expression in human endothelial cell lines and the response seems to be genome- and proteome-dependent. *Proteomics*.6(17):4769-4780, 2006.

We have examined in vitro cell response to mobile phone radiation (900 MHz GSM signal) using two variants of human endothelial cell line: EA.hy926 and EA.hy926v1. Gene expression changes were examined in three experiments using cDNA Expression Arrays and protein expression changes were examined in ten experiments using 2-DE and PDQuest software. Obtained results show that gene and protein expression were altered, in both examined cell lines, in response to one hour mobile phone radiation exposure at an average specific absorption rate of 2.8 W/kg. However, the same genes and proteins were differently affected by the exposure in each of the cell lines. This suggests that the cell response to mobile phone radiation might be genome- and proteome-dependent. Therefore, it is likely that different types of cells and from different species might respond differently to mobile phone radiation or might have different sensitivity to this weak stimulus. Our findings might also explain, at least in part, the origin of discrepancies in replication studies between different laboratories.

Zhijian C, Xiaoxue L, Wei Z, Yezhen L, Jianlin L, Deqiang L, Shijie C, Lifan J, Jiliang H. Studying the protein expression in human B lymphoblastoid cells exposed to 1.8-GHz (GSM) radiofrequency radiation (RFR) with protein microarray. *Biochem Biophys Res Commun*. 433(1):36-39, 2013.

In the present study, the protein microarray was used to investigate the protein expression in human B-cell lymphoblastoid cells intermittently exposed to 1.8-GHz GSM radiofrequency radiation (RFR) at the specific absorption rate (SAR) of 2.0W/kg for 24h. The differential expression of 27 proteins was found, which were related to DNA damage repair, apoptosis, oncogenesis, cell cycle and proliferation (ratio >1.5-fold, $P < 0.05$). The results validated with Western blot assay indicated that the expression of RPA32 was significantly down-regulated ($P < 0.05$) while the expression of p73 was significantly up-regulated in RFR exposure group ($P < 0.05$). Because of the crucial roles of those proteins in DNA repair and cell apoptosis, the results of present investigation may explain the biological effects of RFR on DNA damage/repair and cell apoptosis.

Yadav AS, Sharma MK. Increased frequency of micronucleated exfoliated cells among humans exposed in vivo to mobile telephone radiations. *Mutat Res*. 650(2):175-180, 2008.

The health concerns have been raised following the enormous increase in the use of wireless mobile telephones throughout the world. This investigation had been taken, with the motive to find out whether mobile phone radiations cause any in vivo effects on the frequency of micronucleated exfoliated cells in the exposed subjects. A total of 109 subjects including 85 regular mobile phone users (exposed) and 24 non-users (controls) had participated in this study. Exfoliated cells were obtained by swabbing the buccal-mucosa from exposed as well as sex-age-matched controls. One thousand exfoliated cells were screened from each individual for nuclear anomalies including micronuclei (MN), karyolysis (KL), karyorrhexis (KH), broken egg (BE) and binucleated (BN) cells. The average daily duration of exposure to mobile phone radiations is

Studies that show **Cell Phone** Health Effects

61.26min with an overall average duration of exposure in term of years is 2.35 years in exposed subjects along with the 9.84 ± 0.745 micronucleated cells (MNCs) and 10.72 ± 0.889 total micronuclei (TMN) as compared to zero duration of exposure along with average 3.75 ± 0.774 MNC and 4.00 ± 0.808 TMN in controls. The means are significantly different in case of MNC and TMN at 0.01% level of significance. The mean of KL in controls is 13.17 ± 2.750 and in exposed subjects is 13.06 ± 1.793 . The value of means of KH in exposed subjects (1.84 ± 0.432) is slightly higher than in controls (1.42 ± 0.737). Mean frequency of broken egg is found to be more in exposed subjects (0.65 ± 0.276) as compared to controls (0.50 ± 0.217). Frequency of presence of more than one nucleus in a cell (binucleated) is also higher in exposed (2.72 ± 0.374) in comparison to controls (0.67 ± 0.231). Although there is a slight increase in mean frequency of KH, BE and BN in exposed subjects but the difference is not found statistically significant. Correlation between 0-1, 1-2, 2-3 and 3-4 years of exposure and the frequency of MNC and TMN has been calculated and found to be positively correlated.

Nylund R, Leszczynski D. Proteomics analysis of human endothelial cell line EA.hy926 after exposure to GSM 900 radiation. *Proteomics* 4:1359-1365, 2004.

The human endothelial cell line EA.hy926 was exposed to mobile phone radiation and the effect on protein expression was examined using two-dimensional electrophoresis (2-DE). Up to 38 various proteins have statistically significantly altered their expression levels following the irradiation. Four proteins were identified with matrix-assisted laser desorption/ionization-mass spectrometry (MALDI-MS). Two of the affected proteins were determined to be isoforms of cytoskeletal vimentin. This finding supports our earlier presented working hypothesis which indicated that the mobile phone radiation might affect the cytoskeleton and might have an effect on the physiological functions that are regulated by the cytoskeleton.

Munoz S, Sebastian JL, Sancho M, Miranda JM. Transmembrane voltage induced on altered erythrocyte shapes exposed to RF fields. *Bioelectromagnetics*. 25(8):631-633, 2004.

In this article, the transmembrane voltage induced on erythrocyte, codocyte, ovalocyte and spherocyte cell models exposed to a linearly polarised electromagnetic plane wave of frequency 1800 MHz is calculated. For this purpose, a finite element (FE) numerical technique with adaptive meshing is used. The results show that the value of the induced voltage on the original erythrocyte shape is higher than the one observed on the rest of the altered cell geometries studied. The erythrocyte shape and the membrane electric permittivity are shown to play a fundamental role on the values of the induced transmembrane voltage.

Moisesescu MG, Leveque P, Verjus MA, Kovacs E, Mir LM. 900 MHz modulated electromagnetic fields accelerate the clathrin-mediated endocytosis pathway. *Bioelectromagnetics* 30(3):222-230, 2009.

We report new data regarding the molecular mechanisms of GSM-induced increase of

Studies that show **Cell Phone** Health Effects

cell endocytosis rate. Even though endocytosis represents an important physical and biological event for cell physiology, studies on modulated electromagnetic fields (EMF) effects on this process are scarce. In a previous article, we showed that fluid phase endocytosis rate increases when cultured cells are exposed to 900 MHz EMF similar to mobile phones' modulated GSM signals (217 Hz repetition frequency, 576 micro pulse width) and to electric pulses similar to the GSM electrical component. Trying to distinguish the mechanisms sustaining this endocytosis stimulation, we exposed murine melanoma cells to Lucifer Yellow (LY) or to GSM-EMF/electric pulses in the presence of drugs inhibiting the clathrin- or the caveolin-dependent endocytosis. Experiments were performed at a specific absorption rate (SAR) of 3.2 W/kg in a wire patch cell under homogeneously distributed EMF field and controlled temperature (in the range of 28.5-29.5 degrees C). Thus, the observed increase in LY uptake was not a thermal effect. Chlorpromazine and ethanol, but not Filipin, inhibited this increase. Therefore, the clathrin-dependent endocytosis is stimulated by the GSM-EMF, suggesting that the cellular mechanism affected by the modulated EMF involves vesicles that detach from the cell membrane, mainly clathrin-coated vesicles.

Velizarov, S, Raskmark, P, Kwee, S, The effects of radiofrequency fields on cell proliferation are non-thermal. *Bioelectrochem Bioenerg* 48(1):177-180, 1999.

The number of reports on the effects induced by radiofrequency (RF) electromagnetic fields and microwave (MW) radiation in various cellular systems is still increasing. Until now no satisfactory mechanism has been proposed to explain the biological effects of these fields. One of the current theories is that heat generation by RF/MW is the cause, in spite of the fact that a great number of studies under isothermal conditions have reported significant cellular changes after exposure to RF/MW. Therefore, this study was undertaken to investigate which effect MW radiation from these fields in combination with a significant change of temperature could have on cell proliferation. The experiments were performed on the same cell line, and with the same exposure system as in a previous work [S. Kwee, P. Raskmark, Changes in cell proliferation due to environmental non-ionizing radiation: 2. Microwave radiation, *Bioelectrochem. Bioenerg.*, 44 (1998), pp. 251-255]. The field was generated by signal simulation of the Global System for Mobile communications (GSM) of 960 MHz. Cell cultures, growing in microtiter plates, were exposed in a specially constructed chamber, a Transverse Electromagnetic (TEM) cell. The Specific Absorption Rate (SAR) value for each cell well was calculated for this exposure system. However, in this study the cells were exposed to the field at a higher or lower temperature than the temperature in the field-free incubator i.e., the temperature in the TEM cell was either 39 or 35 +/- 0.1 degrees C. The corresponding sham experiments were performed under exactly the same experimental conditions. The results showed that there was a significant change in cell proliferation in the exposed cells in comparison to the non-exposed (control) cells at both temperatures. On the other hand, no significant change in proliferation rate was found in the sham-exposed cells at both temperatures. This shows that biological effects due to RF/MW cannot be attributed only to a change of temperature. Since the RF/MW induced changes were of the same order of magnitude at both temperatures and also

Studies that show **Cell Phone** Health Effects

comparable to our previous results under isothermal conditions at 37 degrees C, cellular stress caused by electromagnetic fields could initiate the changes in cell cycle reaction rates. It is widely accepted that certain classes of heat-shock proteins are involved in these stress reactions.

Valbonesi P, Franzellitti S, Bersani F, Contin A, Fabbri E. Effects of the exposure to intermittent 1.8 GHz radio frequency electromagnetic fields on HSP70 expression and MAPK signaling pathways in PC12 cells. Int J Radiat Biol. 2014 Feb 11. [Epub ahead of print]

Purpose: We previously reported effects on heat shock protein 70 (HSP70) mRNA expression, a cytoprotective protein induced under stressful condition, in human trophoblast cells exposed to amplitude-modulated Global System for Mobile Communication (GSM) signals. In the present work the same experimental conditions were applied to the rat PC12 cells, in order to assess the stress responses mediated by HSP70 and by the Mitogen Activated Protein Kinases (MAPK) in neuronal-like cells, an interesting model to study possible effects of mobile phone frequencies exposure. Materials and methods: HSP70 gene expression level was evaluated by reverse transcriptase polymerase chain reaction, HSP70 protein expression and MAPK phosphorylation were assessed by Western blotting. PC12 cells were exposed for 4, 16 or 24 h to 1.8 GHz continuous wave signal (CW, carrier frequency without modulation) or to two different GSM modulation schemes, GSM-217Hz and GSM-Talk (which generates temporal changes between two different GSM signals, active during talking or listening phases respectively, thus simulating a typical conversation). Specific adsorption rate (SAR) was 2 W/kg. Results: After PC12 cells exposure to the GSM-217Hz signal for 16 or 24 h, HSP70 transcription significantly increased, whereas no effect was observed in cells exposed to the CW or GSM-Talk signals. HSP70 protein expression and three different MAPK signaling pathways were not affected by the exposure to any of the three different 1.8 GHz signals. Conclusion: The positive effect on HSP70 mRNA expression, observed only in cells exposed to the GSM-217Hz signal, is a repeatable response previously reported in human trophoblast cells and now confirmed in PC12 cells. Further investigations towards a possible role of 1.8 GHz signal modulation are therefore advisable.

Xu S, Chen G, Chen C, Sun C, Zhang D, Murbach M, Kuster N, Zeng Q, Xu Z. Cell Type-Dependent Induction of DNA Damage by 1800 MHz Radiofrequency Electromagnetic Fields Does Not Result in Significant Cellular Dysfunctions. PLoS One. 2013;8(1):e54906.

BACKGROUND: Although IARC clarifies radiofrequency electromagnetic fields (RF-EMF) as possible human carcinogen, the debate on its health impact continues due to the inconsistent results. Genotoxic effect has been considered as a golden standard to determine if an environmental factor is a carcinogen, but the currently available data for RF-EMF remain controversial. As an environmental stimulus, the effect of RF-EMF on cellular DNA may be subtle. Therefore, more sensitive method and systematic research strategy are warranted to evaluate its genotoxicity. **OBJECTIVES:** To determine whether

Studies that show **Cell Phone** Health Effects

RF-EMF does induce DNA damage and if the effect is cell-type dependent by adopting a more sensitive method γ H2AX foci formation; and to investigate the biological consequences if RF-EMF does increase γ H2AX foci formation. **METHODS:** Six different types of cells were intermittently exposed to GSM 1800 MHz RF-EMF at a specific absorption rate of 3.0 W/kg for 1 h or 24 h, then subjected to immunostaining with anti- γ H2AX antibody. The biological consequences in γ H2AX-elevated cell type were further explored with comet and TUNEL assays, flow cytometry, and cell growth assay.

RESULTS: Exposure to RF-EMF for 24 h significantly induced γ H2AX foci formation in Chinese hamster lung cells and Human skin fibroblasts (HSFs), but not the other cells. However, RF-EMF-elevated γ H2AX foci formation in HSF cells did not result in detectable DNA fragmentation, sustainable cell cycle arrest, cell proliferation or viability change. RF-EMF exposure slightly but not significantly increased the cellular ROS level.

CONCLUSIONS: RF-EMF induces DNA damage in a cell type-dependent manner, but the elevated γ H2AX foci formation in HSF cells does not result in significant cellular dysfunctions.

Miyakoshi J, Takemasa K, Takashima Y, Ding GR, Hirose H, Koyama S. Effects of exposure to a 1950 MHz radio frequency field on expression of Hsp70 and Hsp27 in human glioma cells. *Bioelectromagnetics*. 26(4):251-257, 2005.

Human glioma MO54 cells were used to investigate whether radio frequency (RF) field exposure could activate stress response genes. Cells were exposed to continuous wave 1950 MHz or sham conditions for up to 2 h. Specific absorption rates (SARs) were 1, 2, and 10 W/kg. For the cell growth experiment, cell numbers were counted at 0-4 days after exposure. Expression of Hsp27 and Hsp70, as well as the level of phosphorylated Hsp27 (78Ser) protein, was determined by Western blotting. It was found that sham exposed and RF exposed cells demonstrated a similar growth pattern up to 4 days after RFfield exposure. RFfield exposure at both 2 and 10 W/kg did not affect the growth of MO54 cells. In addition, there were no significant differences in protein expression of Hsp27 and Hsp70 between sham exposed and RF exposed cells at a SAR of 1, 2, or 10 W/kg for 1 and 2 h. However, exposure to RFfield at a SAR of 10 W/kg for 1 and 2 h decreased the protein level of phosphorylated Hsp27 (78Ser) significantly. Our results suggest that although exposure to a 1950 MHz RFfield has no effect on cell proliferation and expression of Hsp 27 and Hsp70, it may inhibit the phosphorylation of Hsp27 at Serine 78 in MO54 cells.

Mazor R, Korenstein-Ilan A, Barbul A, Eshet Y, Shahadi A, Jerby E, Korenstein R. Increased levels of numerical chromosome aberrations after in vitro exposure of human peripheral blood lymphocytes to radiofrequency electromagnetic fields for 72 hours. *Radiat Res* 169(1):28-37, 2008.

We investigated the effects of 72 h in vitro exposure of 10 human lymphocyte samples to radiofrequency electromagnetic fields (800 MHz, continuous wave) on genomic instability. The lymphocytes were exposed in a specially designed waveguide resonator at specific absorption rates (SARs) of 2.9 and 4.1 W/kg in a temperature range of 36-37 degrees C. The induced aneuploidy of chromosomes 1, 10, 11 and 17 was determined by

Studies that show **Cell Phone** Health Effects

interphase FISH using semi-automated image analysis. We observed increased levels of aneuploidy depending on the chromosome studied as well as on the level of exposure. In chromosomes 1 and 10, there was increased aneuploidy at the higher SAR, while for chromosomes 11 and 17, the increases were observed only for the lower SAR. Multisomy (chromosomal gains) appeared to be the primary contributor to the increased aneuploidy. The effect of temperature on the level of aneuploidy was examined over the range of 33.5-40 degrees C for 72 h with no statistically significant difference in the level of aneuploidy compared to 37 degrees C. These findings suggest the possible existence of an athermal effect of RF radiation that causes increased levels of aneuploidy. These results contribute to the assessment of potential health risks after continuous chronic exposure to RF radiation at SARs close to the current levels set by ICNIRP guidelines.

Sefidbakht Y, Moosavi-Movahedi AA, Hosseinkhani S, Khodaghohi F, Torkzadeh-Mahani M, Foolad F, Faraji-Dana R. Effects of 940 MHz EMF on bioluminescence and oxidative response of stable luciferase producing HEK cells. Photochem Photobiol Sci. 2014 Jun 2. [Epub ahead of print]

The effects of mobile phone frequency electromagnetic field (RF-EMF, 940 MHz) on a stable cell line (HEK293T) harbouring the firefly luciferase gene were evaluated. A waveguide exposure system with 1 W input power provided the mean specific absorption rate of $\approx 0.09 \text{ W kg}^{-1}$ in 35 mm Petri dishes. The effects of exposure duration (15, 30, 45, 60 and 90 min) on luciferase activity and oxidative response elements were investigated. Endogenous luciferase activity was reduced after 30 and 45 min of continuous exposure, while after 60 min, the exposed cell lysate showed higher luciferase activity compared with the non-exposed control. Reactive oxygen species (ROS) generation was highest in the 30 min exposed cells as studied by 2',7'-dichlorodihydrofluorescein diacetate (DCFH-DA) fluorescence. The observed boost in ROS was then followed by a sharp rise in catalase (CAT) and superoxide dismutase (SOD) activity and elevation of glutathione (GSH) during the 45 min exposure. Decrease in lipid peroxidation (malondialdehyde, MDA) was meaningful for the 45 and 60 min exposed cells. Therefore, it appears that an increase in the activity of luciferase after 60 min of continuous exposure could be associated with a decrease in ROS level caused by activation of the oxidative response. This ability in cells to overcome oxidative stress and compensate the luciferase activity could also be responsible for the adaptive response mechanism detected in ionizing radiation studies with RF-EMF pre-treatments.

Remondini D, Nylund R, Reivinen J, Poullietier de Gannes F, Veyret B, Lagroye I, Haro E, Trillo MA, Capri M, Franceschi C, Schlatterer K, Gminski R, Fitzner R, Tauber R, Schuderer J, Kuster N, Leszczynski D, Bersani F, Maercker C. Gene expression changes in human cells after exposure to mobile phone microwaves. Proteomics. 6(17):4745-4754, 2006.

Possible biological effects of mobile phone microwaves were investigated in vitro. In this study, which was part of the 5FP EU project REFLEX (Risk Evaluation of Potential Environmental Hazards From Low-Energy Electromagnetic Field Exposure Using

Studies that show **Cell Phone** Health Effects

Sensitive in vitro Methods), six human cell types, immortalized cell lines and primary cells, were exposed to 900 and 1800 MHz. RNA was isolated from exposed and sham-exposed cells and labeled for transcriptome analysis on whole-genome cDNA arrays. The results were evaluated statistically using bioinformatics techniques and examined for biological relevance with the help of different databases. NB69 neuroblastoma cells, T lymphocytes, and CHME5 microglial cells did not show significant changes in gene expression. In EA.hy926 endothelial cells, U937 lymphoblastoma cells, and HL-60 leukemia cells we found between 12 and 34 up- or down-regulated genes. Analysis of the affected gene families does not point towards a stress response. However, following microwave exposure, some but not all human cells might react with an increase in expression of genes encoding ribosomal proteins and therefore up-regulating the cellular metabolism.

Trosić I, Pavčić I. Disturbance of cell proliferation in response to mobile phone frequency radiation. *Arh Hig Rada Toksikol.* 60(1):109-115, 2009.

The aim of study was to determine the influence of mobile phone frequency radiation on the proliferation, cytoskeleton structure, and mitotic index of V79 cells after 1 h, 2 h, and 3 h of exposure. V79 cells were cultured in standard laboratory conditions and exposed to continuous-wave (CW) RF/MW radiation of 935 MHz, electric field strength of $(8.2 \pm 0.3) \text{ V m}^{-1}$, and specific absorption rate (SAR) of 0.12 W kg^{-1} . To identify proliferation kinetics, the cells were counted for each hour of exposure 24 h, 48 h, 72 h, and 96 h after respective exposures. Microtubule proteins were determined using specific immunocytochemical methods. Cell smears were analysed under a fluorescent microscope. The study included negative and positive controls. Mitotic index was determined by estimating the number of dividing cells 24 h after exposure and dividing it with the total number of cells. In comparison to the controls, cell proliferation declined in cells exposed for three hours 72 h after irradiation ($p < 0.05$). Microtubule structure was clearly altered immediately after three hours of irradiation ($p < 0.05$). The mitotic index in RF/MW-exposed cells did not differ from negative controls. However, even if exposure did not affect the number of dividing cells, it may have slowed down cell division kinetics as a consequence of microtubule impairment immediately after exposure.

Pilla AA. Electromagnetic fields instantaneously modulate nitric oxide signaling in challenged biological systems. *Biochem Biophys Res Commun.* 426(3):330-333, 2012.

This study shows that a non-thermal pulse-modulated RF signal (PRF), configured to modulate calmodulin (CaM) activation via acceleration of Ca^{2+} binding kinetics, produced an immediate nearly 3-fold increase in nitric oxide (NO) from dopaminergic MN9D cultures ($P < 0.001$). NO was measured electrochemically in real-time using a NO selective membrane electrode, which showed the PRF effect occurred within the first seconds after lipopolysaccharide (LPS) challenge. Further support that the site of action of PRF involves CaM is provided in human fibroblast cultures challenged with low serum and exposed for 15min to the identical PRF signal. In this case a CaM antagonist W-7 could be added to the culture 3h prior to PRF exposure. Those results showed the PRF

Studies that show **Cell Phone** Health Effects

signal produced nearly a two-fold increase in NO, which could be blocked by W-7 ($P < 0.001$). To the authors' knowledge this is the first report of a real-time effect of non-thermal electromagnetic fields (EMF) on NO release from challenged cells. The results provide mechanistic support for the many reported bioeffects of EMF in which NO plays a role. Thus, in a typical clinical application for acute post operative pain, or chronic pain from, e.g., osteoarthritis, EMF therapy could be employed to modulate the dynamics of NO via Ca/CaM-dependent constitutive nitric oxide synthase (cNOS) in the target tissue. This, in turn, would modulate the dynamics of the signaling pathways the body uses in response to the various phases of healing after physical or chemical insult or injury.

Pérez-Castejón C, Pérez-Bruzón RN, Llorente M, Pes N, Lacasa C, Figols T, Lahoz M, Maestú C, Vera-Gil A, Del Moral A, Azanza MJ. Exposure to ELF-pulse modulated X band microwaves increases in vitro human astrocytoma cell proliferation. *Histol Histopathol.* 24(12):1551-1561, 2009.

Common concern about the biological effects of electromagnetic fields (EMF) is increasing with the expansion of X-band microwaves (MW). The purpose of our work was to determine whether exposure to MW pulses in this range can induce toxic effects on human astrocytoma cells. Cultured astrocytoma cells (Clonetics line 1321N1) were submitted to 9.6 GHz carrier, 90% amplitude modulated by extremely low frequency (ELF)-EMF pulses inside a Gigahertz Transversal Electromagnetic Mode cell (GTEM-cell). Astrocytoma cultures were maintained inside a GTEM-incubator in standard culture conditions at 37 ± 0.1 degrees C, 5% CO₂, in a humidified atmosphere. Two experimental conditions were applied with field parameters respectively of: PW 100-120 ns; PRF 100-800 Hz; PRI 10-1.25 ms; power 0.34-0.60 mW; electric field strength 1.25-1.64 V/m; magnetic field peak amplitude 41.4-54.6 microOe. SAR was calculated to be 4.0×10^{-4} W/Kg. Astrocytoma samples were grown in a standard incubator. Reaching 70-80% confluence, cells were transferred to a GTEM-incubator. Experimental procedure included exposed human astrocytoma cells to MW for 15, 30, 60 min and 24 h and unexposed sham-control samples. Double blind method was applied. Our results showed that cytoskeleton proteins, cell morphology and viability were not modified. Statistically significant results showed increased cell proliferation rate under 24h MW exposure. Hsp-70 and Bcl-2 antiapoptotic proteins were observed in control and treated samples, while an increased expression of connexin 43 proteins was found in exposed samples. The implication of these results on increased proliferation is the subject of our current research.

Sebastian JL, Munoz S, Sancho M, Miranda JM, Analysis of the influence of the cell geometry, orientation and cell proximity effects on the electric field distribution from direct RF exposure. *Phys Med Biol* 46(1):213-225, 2001.

This paper shows the importance of using a cell model with the proper geometry, orientation and internal structure to study possible cellular effects from direct radiofrequency exposure. For this purpose, the electric field intensity is calculated, using the finite element numerical technique, in single- and multilayer spherical, cylindrical and ellipsoidal mammalian cell models exposed to linearly polarized electromagnetic

Studies that show **Cell Phone** Health Effects

plane waves of frequencies 900 and 2450 MHz. An extensive analysis is performed on the influence that the cell geometry and orientation with respect to the external field have in the value of the electric field induced in the membrane and cytoplasm. We also show the significant role that the cytoplasmic and extracellular bound water layers play in determining the electric field intensity for the cylindrical and ellipsoidal cell models. Finally, a study of the mutual interactions between cells shows that polarizing effects between cells significantly modify the values of field intensity within the cell.

Sun W, Shen X, Lu D, Lu D, Chiang H. Superposition of an incoherent magnetic field inhibited EGF receptor clustering and phosphorylation induced by a 1.8 GHz pulse-modulated radiofrequency radiation. *Int J Radiat Biol.* 2013, 89:378-383.

Purpose: The present study was conducted to investigate the effect of a temporally incoherent ('noise') magnetic field (MF) on radiofrequency radiation (RFR)-induced epidermal growth factor (EGF) receptor clustering and phosphorylation in cultured cells. Materials and methods: Human amniotic epithelial (FL) cells were exposed for 15 min to either a 1.8 GHz RFR (modulated at 217 Hz), a 2 μ T incoherent MF, or concurrently to the RFR and incoherent MF. Epidermal growth factor treatment served as the positive control. Epidermal growth factor receptor clustering on cellular membrane surface was analyzed using confocal microscopy after indirect immunofluorescence staining, and phosphorylation of EGF receptors was measured by western blot technology. Results: Exposure of FL cells to the 1.8 GHz RFR at SAR (specific absorption rate) of 0.5, 1.0, 2.0, or 4.0 W/kg for 15 min induced EGF receptor clustering and enhanced phosphorylation on tyrosine-1173 residue, whereas exposure to RFR at SAR of 0.1 W/kg for 15 min did not significantly cause these effects. Exposure to a 2 μ T incoherent MF for 15 min did not significantly affect clustering and phosphorylation of EGF receptor in FL cells. When superimposed, the incoherent MF completely inhibited EGF receptor clustering and phosphorylation induced by RFR at SAR of 0.5, 1.0, and 2.0 W/kg, but did not inhibit the effects induced at SAR of 4.0 W/kg. Conclusion: Based on the data of the experiment, it is suggested that membrane receptors could be one of the main targets by which RFR interacts with cells. An incoherent MF could block the interaction to a certain extent

Sun W, Shen X, Lu D, Fu Y, Lu D, Chiang H. A 1.8-GHz radiofrequency radiation induces EGF receptor clustering and phosphorylation in cultured human amniotic (FL) cells. *Int J Radiat Biol.* 88(3):239-244, 2012.

PURPOSE: Many studies have shown that exposure to radiofrequency radiation (RFR) could activate cellular signal transduction pathways. In the present research, we investigated the effects of exposure to a 1.8-GHz RFR at different intensities on epidermal growth factor (EGF) receptor clustering and phosphorylation in human amniotic (FL) cells. MATERIALS AND METHODS: Receptor clustering on cellular membrane surface was analyzed using immunofluorescence assessed by confocal microscopy, and phosphorylation of EGF receptors was measured by western blot technology. EGF treatment served as a positive control. RESULTS: The results showed that, compared with sham exposure, exposure to RFR at specific absorption rate (SAR) of 0.5, 1.0, 2.0, or 4.0 W/kg for 15 min significantly induced EGF receptor clustering and enhanced phosphorylation on the tyrosine-1173 residue in FL

Studies that show **Cell Phone** Health Effects

cells, whereas exposure to a SAR 0.1 W/kg radiation for 15 min did not cause a significant effect. CONCLUSION: Based on the results of this experiment, we conclude that membrane receptors could be one of the main targets that RFR interacts with cells, and the dose-rate threshold, in the case of EGF receptors, is between SAR of 0.1 and 0.5 W/kg. The results indicate a sigmoid dependence of RFR effects on intensity.

Margaritis LH, Manta AK, Kokkaliaris CD, Schiza D, Alimisis K, Barkas G, Georgiou E, Giannakopoulou O, Kollia I, Kontogianni G, Kourouzidou A, Myari A, Roumelioti F, Skouroliakou A, Sykioti V, Varda G, Xenos K, Ziomas K. *Drosophila* oogenesis as a bio-marker responding to EMF sources. *Electromagn Biol Med*. 2013 Aug 5. [Epub ahead of print]

The model biological organisms *Drosophila melanogaster* and *Drosophila virilis* have been utilized to assess effects on apoptotic cell death of follicles during oogenesis and reproductive capacity (fecundity) decline. A total of 280 different experiments were performed using newly emerged flies exposed for short time daily for 3-7 d to various EMF sources including: GSM 900/1800 MHz mobile phone, 1880-1900 MHz DECT wireless base, DECT wireless handset, mobile phone-DECT handset combination, 2.44 GHz wireless network (Wi-Fi), 2.44 GHz blue tooth, 92.8 MHz FM generator, 27.15 MHz baby monitor, 900 MHz CW RF generator and microwave oven's 2.44 GHz RF and magnetic field components. Mobile phone was used as a reference exposure system for evaluating factors considered very important in dosimetry extending our published work with *D. melanogaster* to the insect *D. virilis*. Distance from the emitting source, the exposure duration and the repeatability were examined. All EMF sources used created statistically significant effects regarding fecundity and cell death-apoptosis induction, even at very low intensity levels (0.3 V/m blue tooth radiation), well below ICNIRP's guidelines, suggesting that *Drosophila* oogenesis system is suitable to be used as a biomarker for exploring potential EMF bioactivity. Also, there is no linear cumulative effect when increasing the duration of exposure or using one EMF source after the other (i.e. mobile phone and DECT handset) at the specific conditions used. The role of the average versus the peak E-field values as measured by spectrum analyzers on the final effects is discussed.

Mancinelli F, Caraglia M, Abbruzzese A, d'Ambrosio G, Massa R, Bismuto E. Non-thermal effects of electromagnetic fields at mobile phone frequency on the refolding of an intracellular protein: myoglobin. *J Cell Biochem*. 93(1):188-196, 2004.

Non-thermal effects induced by exposure to microwave electromagnetic field (MW-EMF) at 1.95 MHz, a frequency used in mobile communication, have been observed on the refolding kinetics of the heme binding site in an intracellular protein: tuna myoglobin, starting from acidic conditions. We have selected myoglobin because it can be considered a good model to study protein interactions with MW-EMF for its well-known high-resolution crystallographic structure. Myoglobin solutions at pH 3.0 were subjected to 3 h exposure to microwave field (with a specific absorption rate of 51 ± 1 mW/g); the heme site refolding has been followed by measuring the molecular absorption in the Soret spectral region and the data were fitted to a bi-exponential

Studies that show **Cell Phone** Health Effects

model. The kinetics of exposed samples appear to be slowed by MW-EMF action. Moreover, the tryptophanyl lifetime distribution of the exposed protein, as deduced by the analysis of the fluorescence emission decay from its single tryptophan, appears sharper if compared to non-exposed protein samples. This observation suggests that the presence of MW-EMF could affect the propensity of protein molecules to populate specific conformational substates among which myoglobin molecules fluctuate at acidic pH. Changes in the structural fluctuation caused by MW perturbation can affect differently the aggregation process that occurs competitively during the protein folding, so representing a potential risk for protein "misfolding." These data suggest that MW-EMF could have also biochemical and, consequently, biological effects on eukaryotic cells that are still under investigation.

Le Quément C, Nicolaz CN, Habauzit D, Zhadobov M, Sauleau R, Le Dréan Y. Impact of 60-GHz millimeter waves and corresponding heat effect on endoplasmic reticulum stress sensor gene expression. Bioelectromagnetics. 35(6):444-451, 2014.

Emerging high data rate wireless communication systems, currently under development, will operate at millimeter waves (MMW) and specifically in the 60 GHz band for broadband short-range communications. The aim of this study was to investigate potential effects of MMW radiation on the cellular endoplasmic reticulum (ER) stress. Human skin cell lines were exposed at 60.4 GHz, with incident power densities (IPD) ranging between 1 and 20 mW/cm². The upper IPD limits correspond to the ICNIRP local exposure limit for the general public. The expression of ER-stress sensors, namely BIP and ORP150, was then examined by real-time RT-PCR. Our experimental data demonstrated that MMW radiations do not change BIP or ORP150 mRNA basal levels, whatever the cell line, the exposure duration or the IPD level. Co-exposure to the well-known ER-stress inducer thapsigargin (TG) and MMW were then assessed. Our results show that MMW exposure at 20 mW/cm² inhibits TG-induced BIP and ORP150 over expression. Experimental controls showed that this inhibition is linked to the thermal effect resulting from the MMW exposure.

Kwee S, Raskmark P, Velizarov P. Changes in cellular proteins due to environmental non-ionizing radiation. i. Heat-shock proteins. Electro- and Magnetobiology 20: 141-152, 2001.

This paper describes the effect of weak microwave fields on the amounts of heat-shock proteins in cell cultures at various temperatures. The field was generated by signal simulation of the Global System for Mobile communications (GSM) of 960 Mhz, used in portable phones. Transformed human epithelial amnion (AMA) cells, growing on glass coverslips, were exposed in a transverse electromagnetic (TEM) cell to a microwave field, generating a specific absorption rate (SAR) of 2.1 mW.kg⁻¹ in the cells. Exposure temperatures were 35, 37, and 40 ± 0.1°C, respectively, and the exposure time was 20 min. The heat-shock proteins Hsp-70 and Hsp-27 were detected by immunofluorescence. Higher amounts of Hsp-70 were present in the cells exposed at 35 and 37°C than in the sham-exposed cells. These effects can be considered to be athermal, since the field strength was much lower than the safety standard for absence of heat

Studies that show **Cell Phone** Health Effects

generation by microwave fields. There was no significant response in the case of Hsp-27.

Liu YX, Tai JL, Li GQ, Zhang ZW, Xue JH, Liu HS, Zhu H, Cheng JD, Liu YL, Li AM, Zhang Y. Exposure to 1950-MHz TD-SCDMA Electromagnetic Fields Affects the Apoptosis of Astrocytes via Caspase-3-Dependent Pathway. PLoS One. 7(8):e42332, 2012.

The usage of mobile phone increases globally. However, there is still a paucity of data about the impact of electromagnetic fields (EMF) on human health. This study investigated whether EMF radiation would alter the biology of glial cells and act as a tumor-promoting agent. We exposed rat astrocytes and C6 glioma cells to 1950-MHz TD-SCDMA for 12, 24 and 48 h respectively, and found that EMF exposure had differential effects on rat astrocytes and C6 glioma cells. A 48 h of exposure damaged the mitochondria and induced significant apoptosis of astrocytes. Moreover, caspase-3, a hallmark of apoptosis, was highlighted in astrocytes after 48 h of EMF exposure, accompanied by a significantly increased expression of bax and reduced level of bcl-2. The tumorigenicity assays demonstrated that astrocytes did not form tumors in both control and exposure groups. In contrast, the unexposed and exposed C6 glioma cells show no significant differences in both biological feature and tumor formation ability. Therefore, our results implied that exposure to the EMF of 1950-MHz TD-SCDMA may not promote the tumor formation, but continuous exposure damaged the mitochondria of astrocytes and induce apoptosis through a caspase-3-dependent pathway with the involvement of bax and bcl-2.

Irmak MK, Oztas E, Yagmurca M, Fadillioglu E, Bakir B. Effects of electromagnetic radiation from a cellular telephone on epidermal Merkel cells. J Cutan Pathol. 30(2):135-138, 2003.

The number of reports on the effects induced by electromagnetic radiation (EMR) from cellular telephones in various cellular systems is still increasing. Until now, no satisfactory mechanism has been proposed to explain the biological effects of this radiation except a role suggested for mast cells. Merkel cells may also play a role in the mechanisms of biological effects of EMR. This study was undertaken to investigate the influence of EMR from a cellular telephone (900 MHz) on Merkel cells in rats. A group of rats was exposed to a cellular telephone in speech position for 30 min. Another group of rats was sham-exposed under the same environmental conditions for 30 min. Exposure led to significantly higher exocytotic activity in Merkel cells compared with the sham exposure group. This finding may indicate the possible role of Merkel cells in the pathophysiology of the effects of EMR.

Höytö A, Luukkonen J, Juutilainen J, Naarala J. Proliferation, oxidative stress and cell death in cells exposed to 872 MHz radiofrequency radiation and oxidants. Radiat. Res. 170(2):235-243, 2008.

Human SH-SY5Y neuroblastoma and mouse L929 fibroblast cells were exposed to 872 MHz radiofrequency (RF) radiation using continuous waves (CW) or a modulated signal similar to that emitted by GSM mobile phones at a specific absorption rate (SAR) of 5 W/kg in isothermal conditions. To investigate possible combined effects with other

Studies that show **Cell Phone** Health Effects

agents, menadione was used to induce reactive oxygen species, and tert-butylhydroperoxide (t-BOOH) was used to induce lipid peroxidation. After 1 or 24 h of exposure, reduced cellular glutathione levels, lipid peroxidation, proliferation, caspase 3 activity, DNA fragmentation and viability were measured. Two statistically significant differences related to RF radiation were observed: Lipid peroxidation induced by t-BOOH was increased in SH-SY5Y (but not in L929) cells, and menadione-induced caspase 3 activity was increased in L929 (but not in SH-SY5Y) cells. Both differences were statistically significant only for the GSM-modulated signal. The other end points were not significantly affected in any of the experimental conditions, and no effects were observed from exposure to RF radiation alone. The positive findings may be due to chance, but they may also reflect effects that occur only in cells sensitized by chemical stress. Further studies are required to investigate the reproducibility and dose response of the possible effects.

Harvey C, French PW, Effects on protein kinase C and gene expression in a human mast cell line, HMC-1, following microwave exposure. *Cell Biol Int* 23(11):739-748, 2000.

We used a resonant cavity which delivered a continuous wave exposure at 864.3 MHz at an average specific absorption rate (SAR) of 7 W/kg to determine non-thermal biological effects of microwave exposure. A human mast cell line, HMC-1, was used as the biological target. Cells were given three exposures each of 20-min duration daily for 7 days. The temperature of the cell culture medium during the exposure fell to 26.5 degrees C. Effects were seen on localization of protein kinase C, and expression of three genes of 588 screened. The affected genes included the proto-oncogene c-kit, the transcription factor Nucleoside diphosphate kinase B and the apoptosis-associated gene DAD-1. Stress response genes were variably upregulated. No significant effect on morphology or on F-actin distribution was detected. We conclude that low-power microwave exposure may act on HMC-1 cells by altering gene expression via a mechanism involving activation of protein kinase C, and at temperatures well below those known to induce a heat shock response.

Gerner C, Haudek V, Schandl U, Bayer E, Gundacker N, Hutter HP, Mosgoeller W. Increased protein synthesis by cells exposed to a 1,800-MHz radio-frequency mobile phone electromagnetic field, detected by proteome profiling. *Int Arch Occup Environ Health*.83(6):691-702, 2010.

PURPOSE: To investigate whether or not low intensity radio frequency electromagnetic field exposure (RF-EME) associated with mobile phone use can affect human cells, we used a sensitive proteome analysis method to study changes in protein synthesis in cultured human cells. METHODS: Four different cell kinds were exposed to 2 W/kg specific absorption rate in medium containing (35)S-methionine/cysteine, and autoradiography of 2D gel spots was used to measure the increased synthesis of individual proteins. RESULTS: While short-term RF-EME did not significantly alter the proteome, an 8-h exposure caused a significant increase in protein synthesis in Jurkat T-cells and human fibroblasts, and to a lesser extent in activated primary human

Studies that show **Cell Phone** Health Effects

mononuclear cells. Quiescent (metabolically inactive) mononuclear cells, did not detectably respond to RF-EME. Since RF exposure induced a temperature increase of less than 0.15 degrees C, we suggest that the observed cellular response is a so called "athermal" effect of RF-EME. CONCLUSION: Our finding of an association between metabolic activity and the observed cellular reaction to low intensity RF-EME may reconcile conflicting results of previous studies. We further postulate that the observed increased protein synthesis reflects an increased rate of protein turnover stemming from protein folding problems caused by the interference of radio-frequency electromagnetic fields with hydrogen bonds. Our observations do not directly imply a health risk. However, vis-a-vis a synopsis of reports on cells stress and DNA breaks, after short and longer exposure, on active and inactive cells, our findings may contribute to the re-evaluation of previous reports.

Friedman J, Kraus S, Hauptman Y, Schiff Y, Seger R. Mechanism of a short-term ERK activation by electromagnetic fields at mobile phone frequency. *Biochem J.* 405:559-568, 2007.

The exposure to non-thermal microwave electromagnetic field generated by mobile phones affects the expression of many proteins. This effect on transcription and protein stability can be mediated by the mitogen-activated protein kinase (MAPK) cascades, which serve as central signaling pathways, and govern essentially all stimulated cellular processes. Indeed, a long-term exposure of cells to mobile phone irradiation results in the activation of p38MAPKs as well as the ERK/MAPKs. Here we studied the immediate effect of irradiation on the MAPK cascades, and found that ERKs, but not stress related MAPKs are rapidly activated in response to various frequencies and intensities. Using signaling inhibitors we delineated the mechanism that is involved in this activation. We found that the first step is mediated in the plasma membrane by NADH oxidase, which rapidly generates reactive oxygen species (ROS). These ROS then directly stimulate matrix metalloproteinases and allow them to cleave and release heparin binding-EGF. This secreted factor, activates EGF receptor, which in turn further activates the ERK cascade. Thus, this study demonstrates for the first time a detailed molecular mechanism by which electromagnetic irradiation by mobile phones induces the activation of the ERK cascade and thereby induces transcription and other cellular processes.

French PW, Donnellan M, McKenzie DR, Electromagnetic radiation at 835 MHz changes the morphology and inhibits proliferation of a human astrocytoma cell line. *Bioelectrochem Bioenerg* 43:13-18, 1997.

A human astrocytoma cell line, U-87 MG, was exposed to 835 MHz electromagnetic radiation for 20 min, 3 times per day for 7 days, at a power density of either $40 \pm 15 \text{ mWcm}^{-2}$ or $8.1 \pm 3 \text{ mWcm}^{-2}$. At the low power density, it was observed that the rate of DNA synthesis decreased, and that the cells flattened and spread out in comparison to unexposed culture. At 40 mWcm^{-2} , there were no effects seen on cell proliferation, but alteration in cell morphology included increased cell spreading and also the appearance of actin-containing blebs at localized sites on the membrane. It is hypothesised that 835

Studies that show **Cell Phone** Health Effects

MHz radiation at low power density may be affecting a signal transduction pathway involved in cell proliferation.

Franzellitti S, Valbonesi P, Contin A, Biondi C, Fabbri E. HSP70 expression in human trophoblast cells exposed to different 1.8 Ghz mobile phone signals. Radiat Res. 170(4):488-497, 2008.

The heat-shock proteins (HSPs) are important cellular stress markers and have been proposed as candidates to infer biological effects of high-frequency electromagnetic fields (EMFs). In the current study, HSP70 gene and protein expression were evaluated in cells of the human trophoblast cell line HTR-8/SVneo after prolonged exposure (4 to 24 h) to 1.8 GHz continuous-wave (CW) and different GSM signals (GSM-217Hz and GSM-Talk) to assess the possible effects of time and modulation schemes on cell responses. Inducible HSP70 protein expression was not modified by high-frequency EMFs under any condition tested. The inducible HSP70A, HSP70B and the constitutive HSC70 transcripts did not change in cells exposed to high-frequency EMFs with the different modulation schemes. Instead, levels of the inducible HSP70C transcript were significantly enhanced after 24 h exposure to GSM-217Hz signals and reduced after 4 and 16 h exposure to GSM-Talk signals. As in other cell systems, in HTR-8/SVneo cells the response to high-frequency EMFs was detected at the mRNA level after exposure to amplitude-modulated GSM signals. The present results suggest that the expression analysis for multiple transcripts, though encoding the same or similar protein products, can be highly informative and may account for subtle changes not detected at the protein level.

Duan L, Shan Y, Yu X, [Observations of changes in neurobehavioral functions in workers exposed to high-frequency radiation]. Chung Hua Yu Fang I Hsueh Tsa Chih 32(2):109-111, 1998. [Article in Chinese]

OBJECTIVE: To study the effects of exposure to high-frequency radiation on neurobehavioral function of the exposed workers and its measurement in evaluating occupational hazards caused by it. **METHODS:** Four neurobehavioral functions were tested for the workers exposed to high-frequency radiation with Neurobehavioral Core Tests Battery recommended by WHO. **RESULTS:** Scores for various indicators in exposed workers were significantly lower than those in controls, and correlated to the detection of neurasthenia in the exposed workers, to certain extent. **CONCLUSION:** Changes in neurobehavioral function in workers exposed to high-frequency radiation can reflect its important adverse effects.

Dasdag S, Akdag MZ, Erdal ME, Erdal N, Ay OI, Ay ME, Yilmaz SG, Tasdelen B, Yegin K. Long term and excessive use of 900 MHz radiofrequency radiation alter microrna expression in brain. Int J Radiat Biol. 2014 Dec 20:1-22. [Epub ahead of print].

Purpose: We still do not have any information on the interaction between radiofrequency radiation (RF) and miRNAs, which play paramount role in growth, differentiation, proliferation and cell death by suppressing one or more target genes. The purpose of this study is to bridge this gap by investigating effects of long term 900

Studies that show **Cell Phone** Health Effects

MHz mobile phone exposure on some of the miRNAs in brain tissue.

Materials and Methods: The study was carried out on fourteen Wistar Albino adult male rats by dividing them into two groups: sham (n: 7) and exposure (n: 7). Rats in the exposure group were exposed to 900 MHz RF radiation for 3 h per day (7 d a week) for twelve months (one year). The same procedure was applied to the rats in the sham group except the generator was turned off. Immediately after the last exposure, rats were sacrificed and their brains were removed. rno-miR-9-5p, rno-miR-29a-3p, rno-miR-106b-5p, rno-miR-107 and rno-miR-125a-3p in brain were investigated in detail.

Results: Results revealed that long term exposure of 900 MHz RF radiation only decreased rno-miR107 (adjP*= 0,045) value where the whole body (rms) SAR value was 0.0369 W/kg. However, our results indicated that other micro RNAs evaluated in this study was not altered by 900 MHz RF radiation. Conclusion: 900 MHz RF radiation can alter some of the miRNAs, which, in turn, may lead to adverse effects. Therefore, further studies should be performed.

Daniells, C, Duce, I, Thomas, D, Sewell, P, Tattersall, J, de Pomerai, D, Transgenic nematodes as biomonitors of microwave-induced stress. *Mutat Res* 399:55-64, 1998.

Transgenic nematodes (*Caenorhabditis elegans* strain PC72), carrying a stress-inducible reporter gene (*Escherichia coli* beta-galactosidase) under the control of a *C. elegans* hsp16 heat-shock promoter, have been used to monitor toxicant responses both in water and soil. Because these transgenic nematodes respond both to heat and toxic chemicals by synthesising an easily detectable reporter product, they afford a useful preliminary screen for stress responses (whether thermal or non-thermal) induced by microwave radiation or other electromagnetic fields. We have used a transverse electromagnetic (TEM) cell fed from one end by a source and terminated at the other end by a matched load. Most studies were conducted using a frequency of 750 MHz, at a nominal power setting of 27 dBm. The TEM cell was held in an incubator at 25 degrees C inside a shielded room; corresponding controls were shielded and placed in the same 25 degrees C incubator; additional baseline controls were held at 15 degrees C (worm growth temperature). Stress responses were measured in terms of beta-galactosidase (reporter) induction above control levels. The time-course of response to continuous microwave radiation showed significant differences from 25 degrees C controls both at 2 and 16 h, but not at 4 or 8 h. Using a 5 x 5 multiwell plate array exposed for 2 h, the 25 microwaved samples showed highly significant responses compared with a similar control array. The wells most strongly affected were those in the rows closest to the source, whereas the most distant row did not rise above control levels, suggesting a shadow effect. These differential responses are difficult to reconcile with general heating effects, although localised power absorption affords a possible explanation. Experiments in which the frequency and/or power settings were varied suggested a greater response at 21 than at 27 dBm, both at 750 and 300 MHz, although extremely variable responses were observed at 24 dBm and 750 MHz. Thus, lower power levels tended, if anything, to induce larger responses (with the above-mentioned exception), which is opposite to the trend anticipated for any simple heating effect. These results are reproducible and data acquisition is both rapid and simple. The evidence accrued to

Studies that show **Cell Phone** Health Effects

date suggests that microwave radiation causes measurable stress to transgenic nematodes, presumably reflecting increased levels of protein damage within cells (the common signal thought to trigger hsp gene induction). The response levels observed are comparable to those observed with moderate concentrations (ppm) of metal ions such as Zn^{2+} and Cu^{2+} . We conclude that this approach deserves further and more detailed investigation, but that it has already demonstrated clear biological effects of microwave radiation in terms of the activation of cellular stress responses (hsp gene induction).

d'Ambrosio G, Massa R, Scarfi MR, Zeni O, Cytogenetic damage in human lymphocytes following GSMK phase modulated microwave exposure. Bioelectromagnetics 23:7-13, 2002.

The present study investigated, using in vitro experiments on human lymphocytes, whether exposure to a microwave frequency used for mobile communication, either unmodulated or in presence of phase only modulation, can cause modification of cell proliferation kinetics and/or genotoxic effects, by evaluating the cytokinesis block proliferation index and the micronucleus frequency. In the GSM 1800 mobile communication systems the field is both phase (Gaussian minimum shift keying, GSMK) and amplitude (time domain multiple access, TDMA) modulated. The present study investigated only the effects of phase modulation, and no amplitude modulation was applied. Human peripheral blood cultures were exposed to 1.748 GHz, either continuous wave (CW) or phase only modulated wave (GSMK), for 15 min. The maximum specific absorption rate (~ 5 W/kg) was higher than that occurring in the head of mobile phone users; however, no changes were found in cell proliferation kinetics after exposure to either CW or GSMK fields. As far as genotoxicity is concerned, the micronucleus frequency result was not affected by CW exposure; however, a statistically significant micronucleus effect was found following exposure to phase modulated field. These results would suggest a genotoxic power of the phase modulation per se.

Cranfield C, Wieser HG, Al Madan J, Dobson J. Preliminary evaluation of nanoscale biogenic magnetite-based ferromagnetic transduction mechanisms for mobile phone bioeffects. IEEE Trans Nanobioscience. 2(1):40-43, 2003.

Ferromagnetic transduction models have been proposed as a potential mechanism for mobile phone bioeffects. These models are based on the coupling of RF and pulsed electromagnetic emissions to biogenic magnetite (Fe_3O_4) present in the human brain via either ferromagnetic resonance or mechanical activation of cellular ion channels. We have tested these models experimentally for the first time using a bacterial analogue (*Magnetospirillum magnetotacticum*) which produces intracellular biogenic magnetite similar to that present in the human brain. Experimental evaluation revealed that exposure to mobile phone emissions resulted in a consistent and significantly higher proportion of cell death in exposed cultures versus sham exposure ($p = 0.037$). Though there appears to be a repeatable trend toward higher cell mortality in magnetite-producing bacteria exposed to mobile phone emissions, it is not yet clear that this would extrapolate to a deleterious health effect in humans.

Studies that show **Cell Phone** Health Effects

Chiang H, Microwave and ELF electromagnetic field effects on intercellular communication, Proceedings of the 20th Annual International Conference of the IEEE Engineering in Medicine and Biology Society 20:2798-2801, 1998.

Gap junctional intercellular communication (GJIC) plays an essential role in regulation of cell growth, differentiation and wound healing. Microwave irradiation may down-regulate GJIC and the effect is strongly influenced by modulation frequency. Many studies have demonstrated that GJIC could be suppressed by ELF magnetic field (MF) and the suppression is related to the intensity of magnetic flux density and the exposure duration. Pulsed MF is more effective than sinusoidal MF in inhibiting GJIC. Inhibiting GJIC by electromagnetic field in some cases could be beneficial or detrimental. The mechanism of GJIC inhibition by ELF MF has also been studied and found that the inhibition may be mainly due to hyperphosphorylation of gap junctional connexins by PKC rather than its transcriptional or translational disregulation.

Capri M, Salvioli S, Altiglia S, Sevini F, Remondini D, Mesirca P, Bersani F, Monti D, Franceschi CAge-Dependent Effects of in Vitro Radiofrequency Exposure (Mobile Phone) on CD95+ T Helper Human Lymphocytes. Ann N Y Acad Sci. 1067:493-499, 2006.

.Recent studies on "nonthermal" effects of mobile phone radiofrequency (RF) suggest that RF can interact with cellular functions and molecular pathways. To study the possible RF effects on human lymphocyte activation, we analyzed CD25, CD95, CD28 molecules in unstimulated and stimulated CD4+ e CD8+ T cells in vitro. Peripheral blood mononuclear cells (PBMCs) from young and elderly donors were exposed or sham-exposed to RF (1,800 MHz, Specific Absorption Rate 2 W/kg) with or without mitogenic stimulation. No significant changes in the percentage of these cell subsets were found between exposed and sham-exposed lymphocytes in both young and elderly donors. Nevertheless, after RF exposure we observed a slight, but significant, downregulation of CD95 expression in stimulated CD4+ T lymphocytes from elderly, but not from young donors. This age-related result is noteworthy given the importance of such a molecule in regulation of the immune response.

Chavdoula ED, Panagopoulos DJ, Margaritis LH. Comparison of biological effects between continuous and intermittent exposure to GSM-900-MHz mobile phone radiation: detection of apoptotic cell-death features. Mutat Res. 700(1-2):51-61, 2010.

In the present study we used a 6-min daily exposure of dipteran flies, *Drosophila melanogaster*, to GSM-900MHz (Global System for Mobile Telecommunications) mobile phone electromagnetic radiation (EMR), to compare the effects between the continuous and four different intermittent exposures of 6min total duration, and also to test whether intermittent exposure provides any cumulative effects on the insect's reproductive capacity as well as on the induction of apoptotic cell death. According to our previous experiments, a 6-min continuous exposure per day for five days to GSM-900MHz and DCS-1800MHz (Digital Cellular System) mobile phone radiation, brought about a large decrease in the insect's reproductive capacity, as defined by the number of F(1) pupae. This decrease was found to be non thermal and correlated with an

Studies that show **Cell Phone** Health Effects

increased percentage of induced fragmented DNA in the egg chambers' cells at early- and mid-oogenesis. In the present experiments we show that intermittent exposure also decreases the reproductive capacity and alters the actin cytoskeleton network of the egg chambers, another known aspect of cell death that was not investigated in previous experiments, and that the effect is also due to DNA fragmentation. Intermittent exposures with 10-min intervals between exposure sessions proved to be almost equally effective as continuous exposure of the same total duration, whereas longer intervals between the exposures seemed to allow the organism the time required to recover and partly overcome the above-mentioned effects of the GSM exposure.

Effects on Hormones

Koyu A, Cesur G, Ozguner F, Akdogan M, Mollaoglu H, Ozen S. Effects of 900MHz electromagnetic field on TSH and thyroid hormones in rats. *Toxicol Lett.* 157(3):257-262, 2005.

In this study, the effects of exposure to a 900megahertz (MHz) electromagnetic field (EMF) on serum thyroid stimulating hormone (TSH) and triiodothyronine-thyroxine (T(3)-T(4)) hormones levels of adult male Sprague-Dawley rats were studied. Thirty rats were used in three independent groups, 10 of which were control (without stress and EMF), 10 of which were exposed to 900MHz EMF and 10 of which were sham-exposed. The exposures were performed 30min/day, for 5days/week for 4 weeks to 900MHz EMF. Sham-exposed animals were kept under the same environmental conditions as the study groups except with no EMF exposure. The concentration of TSH and T(3)-T(4) hormones in the rat serum was measured by using an immunoradiometric assay (IRMA) method for TSH and a radio-immunoassay (RIA) method for T(3) and T(4) hormones. TSH values and T(3)-T(4) at the 900MHz EMF group were significantly lower than the sham-exposed group ($p<0.01$). There were no statistically significant differences in serum TSH values and T(3)-T(4) hormone concentrations between the control and the sham-exposed group ($p>0.05$). These results indicate that 900MHz EMF emitted by cellular telephones decrease serum TSH and T(3)-T(4) levels.

Pawlak K, Sechman A, Nieckarz Z. Plasma thyroid hormones and corticosterone levels in blood of chicken embryos and post hatch chickens exposed during incubation to 1800 MHz electromagnetic field. *Int J Occup Med Environ Health.* 2014 Jan 31. [Epub ahead of print]

INTRODUCTION: This study attempted to determine the effect of a 1800 MHz electromagnetic field (EMF) (only carrier frequency) on thyroxine (T4), triiodothyronine (T3) and corticosterone (CORT) concentrations in the blood plasma of chick embryos, and to investigate the effect of electromagnetic field (EMF) exposure during embryogenesis on the level of these hormones in birds that are ready for slaughter.

MATERIAL AND METHODS: Throughout the incubation period, embryos from the experimental group were exposed to a 1800 MHz EMF with power density of 0.1 W/m², 10 times during 24 h for 4 min. Blood samples were collected to determine T4, T3 and

Studies that show **Cell Phone** Health Effects

CORT concentrations on the 12th (E12) and 18th (E18) day of incubation, from newly hatched chicks (D1) and from birds ready for slaughter (D42). RESULTS: The experiment showed that T4 and T3 concentrations decreased markedly and CORT levels increased in the embryos and in the newly hatched chicks exposed to EMF during embryogenesis. However, no changes were found in the level of the analyzed hormones in the birds ready for slaughter. Differences in T4 and T3 plasma concentrations between the EMF-exposed group and the embryos incubated without additional EMF were the highest in the newly hatched chicks, which may be indicative of the cumulative effect of electromagnetic field on the hypothalamo-pituitary-thyroid axis (HPT). DISCUSSION: The obtained results suggest that additional 1800 MHz radio frequency electromagnetic field inhibits function of HPT axis, however, it stimulates hypothalamo-pituitary-adrenal axis by inducing adrenal steroidogenic cells to synthesize corticosterone. Further investigations are needed to elucidate the mechanisms by which radio EMFs affect HPT and HPA axis function in the chicken embryos.

Qin F, Zhang J, Cao H, Yi C, Li JX, Nie J, Chen LL, Wang J, Tong J. Effects of 1800-MHz radiofrequency fields on circadian rhythm of plasma melatonin and testosterone in male rats. J Toxicol Environ Health A. 75(18):1120-1128, 2012.

Radiofrequency fields (RF) at 1800 MHz are known to affect melatonin (MEL) and testosterone in male rats, but it remains to be determined whether RF affected circadian rhythm of these plasma hormones. Male Sprague-Dawley rats were exposed to 1800-MHz RF at 208 $\mu\text{W}/\text{cm}^2$ power density (SAR: 0.5762 W/kg) at different zeitgeber (ZT) periods of the day, including 0 (ZT0), 4 (ZT4), 8 (ZT8), 12 (ZT12), 16 (ZT16), and 20 (ZT20) h. RF exposure was 2 h/d for 32 d. From each rat, the concentrations of plasma MEL and testosterone were determined in plasma after RF exposure and compared with controls. The results confirmed the existence of circadian rhythms in the synthesis of MEL and testosterone, but revealed an inverse relationship in peak phase of these rhythms. These rhythms were disturbed after exposure to RF, with the effect being more pronounced on MEL than testosterone. The most pronounced effect of RF exposure on MEL and testosterone appears to be in rats exposed to RF at ZT 16 and ZT0 h, respectively. Data suggest that regulation of testosterone is controlled by MEL and that MEL is more sensitive to RF exposure.

Jarupat S, Kawabata A, Tokura H, Borkiewicz A. Effects of the 1900 MHz electromagnetic field emitted from cellular phone on nocturnal melatonin secretion. J Physiol Anthropol Appl Human Sci 22(1):61-63, 2003.

Exposure to cellular phone EMF caused a significant reduction in salivary melatonin in female human subjects.

Eskander EF, Estefan SF, Abd-Rabou AA. How does long term exposure to base stations and mobile phones affect human hormone profiles? Clin Biochem. 45(1-2):157-161, 2012

OBJECTIVES: This study is concerned with assessing the role of exposure to radio frequency radiation (RFR) emitted either from mobiles or base stations and its relations

Studies that show Cell Phone Health Effects

with human's hormone profiles. **DESIGN AND METHODS:** All volunteers' samples were collected for hormonal analysis. **RESULTS:** This study showed significant decrease in volunteers' ACTH, cortisol, thyroid hormones, prolactin for young females, and testosterone levels. **CONCLUSION:** The present study revealed that high RFR effects on pituitary-adrenal axis.

Eşmekaya MA, Seyhan N, Omeroğlu S. Pulse modulated 900 MHz radiation induces hypothyroidism and apoptosis in thyroid cells: A light, electron microscopy and immunohistochemical study. *Int J Radiat Biol.* 86(12):1106-1116, 2010.

Purpose: In the present study we investigated the possible histopathological effects of pulse modulated Radiofrequency (RF) fields on the thyroid gland using light microscopy, electron microscopy and immunohistochemical methods. Materials and methods: Two months old male Wistar rats were exposed to a 900 MHz pulse-modulated RF radiation at a specific absorption rate (SAR) of 1.35 Watt/kg for 20 min/day for three weeks. The RF signals were pulse modulated by rectangular pulses with a repetition frequency of 217 Hz and a duty cycle of 1:8 (pulse width 0.576 ms). To assess thyroid endocrine disruption and estimate the degree of the pathology of the gland, we analysed structural alterations in follicular and colloidal diameters and areas, colloid content of the follicles, and height of the follicular epithelium. Apoptosis was confirmed by Transmission Electron Microscopy and assessing the activities of an initiator (caspase-9) and an effector (caspase-3) caspases that are important markers of cells undergoing apoptosis. Results: Morphological analyses revealed hypothyrophy of the gland in the 900 MHz RF exposure group. The results indicated that thyroid hormone secretion was inhibited by the RF radiation. In addition, we also observed formation of apoptotic bodies and increased caspase-3 and caspase-9 activities in thyroid cells of the rats that were exposed to modulated RF fields. Conclusion: The overall findings indicated that whole body exposure to pulse-modulated RF radiation that is similar to that emitted by global system for mobile communications (GSM) mobile phones can cause pathological changes in the thyroid gland by altering the gland structure and enhancing caspase-dependent pathways of apoptosis.

de Seze R, Fabbro-Peray P, Miro L, GSM radiocellular telephones do not disturb the secretion of antepituitary hormones in humans. *Bioelectromagnetics* 19(5):271-278, 1998.

It is known that the endocrine system of experimental animals is susceptible to perturbation by radiofrequency (RF) radiation. Because of the recent interest in health and safety issues of cellular telephones, an experiment was designed to evaluate the effect of a 900 MHz RF radiation emitted by a Global System for Mobile radiotelephone (217 Hz impulses, one-eighth duty cycle, 2 W peak power) on human endocrine functions. Twenty healthy male volunteers aged from 19 to 40 were inducted in the present experiment. Each subject was exposed to RF radiation through the use of a cellular phone 2 h/day, 5 days/wk, for 1 month. Subjects were their own control. End points were serum adrenocorticotropin, thyrotropin, growth hormone, prolactin, luteinizing hormone, and follicle stimulating hormone concentrations. These end points

Studies that show **Cell Phone** Health Effects

were determined in nine weekly blood samples obtained starting 3 weeks before the commencement of the exposure and ending 2 weeks after exposures. All but one blood sample was drawn 48 h after each weekly session. The seventh drawing was performed the morning after the last weekly exposure. Within each individual, the preexposure hormone concentration was used as a control. Results indicated that all hormone concentrations remained within normal physiologic ranges. A difference was not noted among the nine weekly samples in five of six hormones studied. There was a significant change only in thyrotropin concentration, showing a 21% decrease on the seventh sampling. Because this change recovered fully during the postexposure period, it is concluded that 1 month of intermittent exposures to RF radiation from a cellular telephone does not induce a long-lasting or cumulative effect on the hormone secretion rate of the anterior pituitary gland in humans.

Cervellati F, Valacchi G, Lunghi L, Fabbri E, Valbonesi P, Marci R, Biondi C, Vesce F. 17- β -estradiol counteracts the effects of high frequency electromagnetic fields on trophoblastic connexins and integrins. *Oxid Med Cell Longev*. 2013;2013:280850. doi: 10.1155/2013/280850.

We investigated the effect of high-frequency electromagnetic fields (HF-EMFs) and 17- β -estradiol on connexins (Cxs), integrins (Ints), and estrogen receptor (ER) expression, as well as on ultrastructure of trophoblast-derived HTR-8/SVneo cells. HF-EMF, 17- β -estradiol, and their combination induced an increase of Cx40 and Cx43 mRNA expression. HF-EMF decreased Int α 1 and β 1 mRNA levels but enhanced Int α 5 mRNA expression. All the Ints mRNA expressions were increased by 17- β -estradiol and exposure to both stimuli. ER- β mRNA was reduced by HF-EMF but augmented by 17- β -estradiol alone or with HF-EMF. ER- β immunofluorescence showed a cytoplasmic localization in sham and HF-EMF exposed cells which became nuclear after treatment with hormone or both stimuli. Electron microscopy evidenced a loss of cellular contact in exposed cells which appeared counteracted by 17- β -estradiol. We demonstrate that 17- β -estradiol modulates Cxs and Ints as well as ER- β expression induced by HF-EMF, suggesting an influence of both stimuli on trophoblast differentiation and migration.

Wood A, Loughran S, Stough C, Does evening exposure to mobile phone radiation affect subsequent melatonin production? *Int. J. Rad. Biol* 82:69-76, 2006.

Purpose: To test whether exposure to the emissions from a digital mobile phone handset prior to sleep alters the secretion of melatonin. Materials and methods: In a double-blind cross-over design, 55 adult volunteers were both actively exposed or sham-exposed (in random order on successive Sunday nights) to mobile phone emissions for 30 min (0.25 W average power). Urine collection occurred immediately prior to retiring to bed and on rising the next morning. Melatonin output was estimated from principal metabolite concentrations (6-sulphatoxymelatonin (aMT6s) via radioimmunoassay), urine volumes and creatinine concentrations. Results: Total melatonin metabolite output (concentration \times urine volume) was unchanged between the two exposure conditions (active 14.1 \pm 1.1 μ g; sham 14.6 \pm 1.3 μ g). The pre- and post-bedtime outputs considered separately were also not significantly different, although

Studies that show **Cell Phone** Health Effects

the pre-bedtime value was less for active versus sham exposure. When melatonin metabolite output was estimated from the ratio of aMT6s to creatinine concentrations, the pre-bedtime value was significantly less ($p = 0.037$) for active compared to sham. Examination of individual responses is suggestive of a small group of 'responders'. Conclusions: Total nighttime melatonin output is unchanged by mobile phone handset emissions, but there could be an effect on melatonin onset time.

Burch JB, Reif JS, Noonan CW, Ichinose T, Bachand AM, Koleber TL, Yost MG. Melatonin metabolite excretion among cellular telephone users. Int J Rad Biol 78: 1029-1036, 2002.

Abstract: Purpose: The relationship between cellular telephone use and excretion of the melatonin metabolite 6-hydroxymelatonin sulfate (6-OHMS) was evaluated in two populations of male electric utility workers (Study 1, $n=149$; Study 2, $n=77$). Materials and methods: Participants collected urine samples and recorded cellular telephone use over 3 consecutive workdays. Personal 60-Hz magnetic field (MF) and ambient light exposures were characterized on the same days using EMDEX II meters. A repeated measures analysis was used to assess the effects of cellular telephone use, alone and combined with MF exposures, after adjustment for age, participation month and light exposure. Results: No change in 6-OHMS excretion was observed among those with daily cellular telephone use >25 min in Study 1 (5 worker-days). Study 2 workers with >25 min cellular telephone use per day (13 worker-days) had lower creatinine-adjusted mean nocturnal 6-OHMS concentrations ($p=0.05$) and overnight 6-OHMS excretion ($p=0.03$) compared with those without cellular telephone use. There was also a linear trend of decreasing mean nocturnal 6-OHMS/creatinine concentrations ($p=0.02$) and overnight 6-OHMS excretion ($p=0.08$) across categories of increasing cellular telephone use. A combined effect of cellular telephone use and occupational 60-Hz MF exposure in reducing 6-OHMS excretion was also observed in Study 2. Conclusions: Exposure-related reductions in 6-OHMS excretion were observed in Study 2, where daily cellular telephone use of >25 min was more prevalent. Prolonged use of cellular telephones may lead to reduced melatonin production, and elevated 60-Hz MF exposures may potentiate the effect.

Bergamaschi A, Magrini A, Ales G, Coppetta L, Somma G. Are thyroid dysfunctions related to stress or microwave exposure (900 MHz)? Int J Immunopathol Pharmacol. 17(2 Suppl):31-36, 2004.

In the last decade, numerous scientific evidence suggested possible adverse health effects from exposure to electromagnetic fields (EMF's) and the use of mobile phones. According to some studies EMF induced changes of trans-membrane Ca^{++} flux may lead to altered metabolism and/or secretion of neurohormones including TSH, ACTH, GH, prolactin and melatonin. The aim of this research was to analyse the effects of mobile phone use on thyroid function and to evaluate the possible role of occupational stress. 2598 employees (1355 men and 1243 women) with different duties (vendors, operators and network technicians) were included in the study. Exposure to EMF's, generated by mobile phones, was assessed both by submitting a questionnaire directly to the employees and acquiring data regarding conversation times. The workers were divided

Studies that show **Cell Phone** Health Effects

into three groups on the basis of their personal mobile phone use. Moreover, a group of 160 workers with TSH values below 0.4 UI/l was characterized. No statistically significant difference regarding TSH values below 0.4 UI/l was observed among workers with different duties but there was a greater prevalence of subjects with low TSH values among 192 employees with more than 33 hrs/month conversation time; this difference was statistically significant ($p < 0.05$). On the basis of our data, it is not possible to establish whether this result is determined by exposure to EMF's from mobile phones or by the stress of using these instruments.

Philippova TM, Novoselov VI, Alekseev SI, Influence of microwaves on different types of receptors and the role of peroxidation of lipids on receptor-protein shedding. Bioelectromagnetics 15(3):183-192, 1994.

The effects of a continuous wave or pulse-modulated, 900 MHz microwave field were studied by in vitro assays of rat chemoreceptors. The pulsed field was modulated as rectangular waves at rates of 1, 6, 16, 32, 75, or 100 pps. The pulse-period to pulse-duration ratio was 5 in all cases, and specific absorption rates (SARs) ranged from 0.5 to 18 W/kg. Binding of ligands to cell membranes was differentially affected by exposure to microwaves. For example, binding of H³-glutamic acid to hippocampal cells was not altered by a 15 min exposure to a continuous wave field at 1 W/kg, but binding of H³-dihydroalprenolol to liver-cell membranes of neonates underwent a fivefold decrease under the same field conditions. This effect was not dependent on modulation or on a change in the constant of stimulus-receptor binding but depended on a shedding of the membrane's receptor elements into solution. The magnitude of inhibition correlated with the oxygen concentration in the exposed suspension. Antioxidants (dithiothreitol and ionol) inhibited the shedding of receptor elements. The microwave exposure did not cause an accumulation of products from the peroxidation of lipids (POL). Ascorbate-dependent or non-enzymatic POL was not responsible for the inhibition, and POL was not found in other model systems. However, enzymatic POL mechanisms in localized areas of receptor binding remain a possibility.

Effects on The Heart

Pakhomov AG, Dubovick BV, Degtyariv IG, Pronkevich AN, Microwave influence on the isolated heart function: I. Effect of modulation. Bioelectromagnetics 16(4):241-249, 1995.

Dependence of the microwave effect on modulation parameters (pulse width, duty ratio, and peak intensity) was studied in an isolated frog auricle preparation. The rate and amplitude of spontaneous auricle twitches were measured during and after a 2 min exposure to 915 or 885 MHz microwaves and were compared to preexposure values. The studied ranges of modulation parameters were: pulse width, 10(-6)-10(-2) s; duty ratio, 7:100000, and peak specific absorption rate, 100-3000 W/kg. Combinations of the parameters were chosen by chance, and about 400 various exposure regimes were tested. The experiments established that no regime was effective unless the average microwave

Studies that show **Cell Phone** Health Effects

power was high enough to induce preparation heating (0.1-0.4 degree C). The twitch rate instantly increased, and the amplitude decreased, as the temperature rose; similar changes could be induced by equivalent conventional heating. The data provide evidence that the effect of short-term microwave exposure on the isolated heart pacemaker and contractile functions depends on pulse modulation just as much as modulation determines the average absorbed power. These functions demonstrated no specific dependence on exposure parameters such as frequency or power windows.

Andrzejak R, Poreba R, Poreba M, Derkacz A, Skalik R, Gac P, Beck B, Steinmetz-Beck A, Pilecki W. The influence of the call with a mobile phone on heart rate variability parameters in healthy volunteers. *Ind Health*. 46(4):409-417, 2008.

It is possible that electromagnetic field (EMF) generated by mobile phones (MP) may have an influence on the autonomic nervous system (ANS) and modulates the function of circulatory system. The aim of the study was to estimate the influence of the call with a mobile phone on heart rate variability (HRV) in young healthy people. The time and frequency domain HRV analyses were performed to assess the changes in sympathovagal balance in a group of 32 healthy students with normal electrocardiogram (ECG) and echocardiogram at rest. The frequency domain variables were computed: ultra low frequency (ULF) power, very low frequency (VLF) power, low frequency (LF) power, high frequency (HF) power and LF/HF ratio was determined. ECG Holter monitoring was recorded in standardized conditions: from 08:00 to 09:00 in the morning in a sitting position, within 20 min periods: before the telephone call (period I), during the call with use of mobile phone (period II), and after the telephone call (period III). During 20 min call with a mobile phone time domain parameters such as standard deviation of all normal sinus RR intervals (SDNN [ms]--period I: 73.94+/-25.02, period II: 91.63+/-35.99, period III: 75.06+/-27.62; I-II: p<0.05, II-III: p<0.05) and standard deviation of the averaged normal sinus RR intervals for all 5-mm segments (SDANN [ms]--period I: 47.78+/-22.69, period II: 60.72+/-27.55, period III: 47.12+/-23.21; I-II: p<0.05, II-III: p<0.05) were significantly increased. As well as very low frequency (VLF [ms²]--period I: 456.62+/-214.13, period II: 566.84+/-216.99, period III: 477.43+/-203.94; I-II: p<0.05), low frequency (LF [ms²]--period I: 607.97+/-201.33, period II: 758.28+/-307.90, period III: 627.09+/-220.33; I-II: p<0.01, II-III: p<0.05) and high frequency (HF [ms²]--period I: 538.44+/-290.63, period II: 730.31+/-445.78, period III: 590.94+/-301.64; I-II: p<0.05) components were the highest and the LF/HF ratio (period I: 1.48+/-0.38, period II: 1.16+/-0.35, period III: 1.46+/-0.40; I-II: p<0.05, II-III: p<0.05) was the lowest during a call with a mobile phone. The tone of the parasympathetic system measured indirectly by analysis of heart rate variability was increased while sympathetic tone was lowered during the call with use of a mobile phone. It was shown that the call with a mobile phone may change the autonomic balance in healthy subjects. Changes in heart rate variability during the call with a mobile phone could be affected by electromagnetic field but the influence of speaking cannot be excluded.

Ozguner F, Altinbas A, Ozaydin M, Dogan A, Vural H, Kisioglu AN, Cesur G, Yildirim NG. Mobile phone-induced myocardial oxidative stress: protection by a novel antioxidant

Studies that show **Cell Phone** Health Effects

agent caffeic acid phenethyl ester. Toxicol Ind Health. 21(9):223-230, 2005.

Electromagnetic radiation (EMR) or radiofrequency fields of cellular mobile phones may affect biological systems by increasing free radicals, which appear mainly to enhance lipid peroxidation, and by changing the antioxidant defense systems of human tissues, thus leading to oxidative stress. Mobile phones are used in close proximity to the heart, therefore 900 MHz EMR emitting mobile phones may be absorbed by the heart. Caffeic acid phenethyl ester (CAPE), one of the major components of honeybee propolis, was recently found to be a potent free radical scavenger and antioxidant, and is used in folk medicine. The aim of this study was to examine 900 MHz mobile phone-induced oxidative stress that promotes production of reactive oxygen species (ROS) and the role of CAPE on myocardial tissue against possible oxidative damage in rats. Thirty rats were used in the study. Animals were randomly grouped as follows: sham-operated control group (N: 10) and experimental groups: (a) group II: 900 MHz EMR exposed group (N: 10); and (b) group III: 900 MHz EMR exposed+CAPE-treated group (N: 10). A 900 MHz EMR radiation was applied to groups II and III 30 min/day, for 10 days using an experimental exposure device. Malondialdehyde (MDA, an index of lipid peroxidation), and nitric oxide (NO, a marker of oxidative stress) were used as markers of oxidative stress-induced heart impairment. Superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GSH-Px) activities were studied to evaluate the changes of antioxidant status. In the EMR exposed group, while tissue MDA and NO levels increased, SOD, CAT and GSH-Px activities were reduced. CAPE treatment in group III reversed these effects. In this study, the increased levels of MDA and NO and the decreased levels of myocardial SOD, CAT and GSH-Px activities demonstrate the role of oxidative mechanisms in 900 MHz mobile phone-induced heart tissue damage, and CAPE, via its free radical scavenging and antioxidant properties, ameliorates oxidative heart injury. These results show that CAPE exhibits a protective effect on mobile phone-induced and free radical mediated oxidative heart impairment in rats.

Alhusseiny A, Al-Nimer M, Majeed A. Electromagnetic energy radiated from mobile phone alters electrocardiographic records of patients with ischemic heart disease. Ann Med Health Sci Res. 2(2):146-151, 2012.

BACKGROUND: Electromagnetic energy radiated from mobile phones did not show significant effect on the blood pressure, heart rate, and electrocardiographic (ECG) parameters in animals and humans. **AIM:** This study aimed to investigate the effect of radiofrequency of mobile phone on the electrocardiographic parameters in patients with history of ischemic heart disease, taking into consideration the gender factor.

SUBJECTS AND METHODS: A total number of 356 participants (129 males and 227 females) were admitted in this study. They were grouped into: subjects without cardiac diseases (Group I), patients with ischemic heart disease (Group II), and patients with history of cardiac diseases not related to myocardial ischemia (Group III).

Electrocardiogram was obtained from each patient when the mobile phone was placed at the belt level and over precordium in turn-off mode (baseline) and turn-on mode for 40 sec ringing. The records of ECG were electronically analyzed. **RESULTS:** Prolongation of QTc interval was significantly observed in male gender of Groups I and III ($P < 0.001$).

Studies that show **Cell Phone** Health Effects

Male patients of Group II showed significant QTc interval prolongation ($P = 0.01$) and changes in the voltage criteria ($P = 0.001$). These changes were not observed in female patients with ischemic heart disease. The position of mobile at the belt level or over the precordium showed effects on the heart. **CONCLUSIONS:** The radiofrequency of cell phone prolongs the QT interval in human beings and it interferes with voltage criteria of ECG records in male patients with myocardial ischemia.

Parazzini M, Ravazzani P, Tognola G, Thuróczy G, Molnar FB, Sacchettini A, Ardesi G, Mainardi LT. Electromagnetic fields produced by GSM cellular phones and heart rate variability. Bioelectromagnetics. 28(2):122-129, 2007.

In this study, 26 healthy young volunteers were submitted to 900 MHz (2 W) GSM cellular phone exposure and to sham exposure in separate sessions. The study was designed to assess cardiac regulatory mechanism in different autonomic nervous system (ANS) states during exposure to low-intensity EMF. Rest-to-stand protocol was applied to evaluate ANS in quiet condition (rest, vagal prevalence) and after a sympathetic activation (stand). The procedure is conducted twice in a double-blind design: once with a genuine EMF exposure and once with a sham exposure (at least 24 h apart). During each session three-leads electrocardiograms were recorded and RR series extracted off-line. Time domain and frequency domain HRV parameters were calculated in every phase of the protocol and during different exposures. The analysis of the data show there was no statistically significant effect due to EMF exposure both on main (i.e., RR mean) and most of the other HRV parameters. A weak interaction between some HRV parameters (i.e., SDNN, TINN, and triangular index in time domain and LF power in frequency domain analysis) and RF exposure was observed and this effect seems to be gathered around the sympathetic response to stand.

Effects on Glucose Tolerance

Meo SA, Al Rubeaan K. Effects of exposure to electromagnetic field radiation (EMFR) generated by activated mobile phones on fasting blood glucose. Int J Occup Med Environ Health. 26(2):235-241, 2013.

OBJECTIVE: Extensive use of mobile phones has been accompanied by a common public debate about possible adverse effects on human health. No study has been published so far to establish any association between the fastest growing innovation of mobile phone and fasting blood glucose. The aim was to determine the effects of exposure to electromagnetic field radiation generated by mobile phones on fasting blood glucose in Wistar Albino rats. **MATERIALS AND METHODS:** 40 Male Albino rats (Wistar Strain) were divided into 5 equally numerous groups. Group A served as the control one, group B received mobile phone radiation for less than 15 min/day, group C: 15-30 min/day, group D: 31-45 min/day, and group E: 46-60 min/day for a total period of 3 months. Fasting blood glucose was determined by using Spectrophotometer and serum insulin by Enzyme-linked Immunosorbent Assay (ELISA). The Homeostatic Model (HOMA-B) was

Studies that show Cell Phone Health Effects

applied for the assessment of β -cell function and (HOMA-IR) for resistance to insulin.

RESULTS: Wister Albino rats exposed to mobile phone radiation for longer than 15 min a day for a total period of 3 months had significantly higher fasting blood glucose ($p < 0.015$) and serum insulin ($p < 0.01$) compared to the control group. HOMA-IR for insulin resistance was significantly increased ($p < 0.003$) in the groups that were exposed for 15-30 and 46-60 min/day compared to the control rats. **CONCLUSION:** The results of the present study show an association between long-term exposure to activated mobile phones and increase in fasting blood glucose and serum insulin in Albino rats.

Bielski J, Sikorski M, [Disturbances of glucose tolerance in workers exposed to electromagnetic radiation]. Med Pr 47(3):227-231, 1996. [Article in Polish]

The study group was composed of 50 workers exposed to electromagnetic radiation (radiowaves). Out of them 31 persons (62%), employed mostly in the risk zone, showed irregular glycaemia after oral administration of 75 g of glucose. At normal blood sugar before breakfast, the glycaemia level was high following administration of glucose and it did not return to starting values after 2 hours. After 30 min from glucose administration the level accounted for 155 mg%, after 60 min-180 mg%, after 90 min-153 mg% and after 120 min-124 mg%, on average. In 10 persons (32%) with glucose tolerance disturbances, disorders in bioelectric activity of the brain (abnormal EEG record) were observed.

Effects on the Whole Body

Khalil AM, Gagaa M, Alshamali A. 8-Oxo-7, 8-dihydro-2'-deoxyguanosine as a biomarker of DNA damage by mobile phone radiation. Hum Exp Toxicol.31(7):734-740, 2012.

We examined the effect of exposure to mobile phone 1800 MHz radio frequency radiation (RFR) upon the urinary excretion of 8-oxo-7, 8-dihydro-2'-deoxyguanosine (8-oxodG), one major form of oxidative DNA damage, in adult male Sprague-Dawley rats. Twenty-four rats were used in three independent experiments (RFR exposed and control, 12 rats, each). The animals were exposed to RFR for 2 h from Global System for Mobile Communications (GSM) signal generator with whole-body-specific absorption rate of 1.0 W/kg. Urine samples were collected from the rat while housed in a metabolic cage during the exposure period over a 4-h period at 0.5, 1.0, 2.0 and 4.0 h from the beginning of exposure. In the control group, the signal generator was left in the turn-off position. The creatinine-standardized concentrations of 8-oxodG were measured. With the exception of the urine collected in the last half an hour of exposure, significant elevations were noticed in the levels of 8-oxodG in urine samples from rats exposed to RFR when compared to control animals. Significant differences were seen overall across time points of urine collection with a maximum at 1 h after exposure, suggesting repair of the DNA lesions leading to 8-oxodG formation.

Aydin B, Akar A. Effects of a 900-MHz electromagnetic field on oxidative stress

Studies that show **Cell Phone** Health Effects

parameters in rat lymphoid organs, polymorphonuclear leukocytes and plasma. Arch Med Res. 42(4):261-267, 2011.

BACKGROUND AND AIMS: The present study investigated the effects of a 900-MHz electromagnetic field (EMF) for 2 h/day for 45 days on lymphoid organs (spleen, thymus, bone marrow), polymorphonuclear leukocytes (PMNs) and plasma of rats, focusing on changes in the enzymatic and nonenzymatic antioxidant system. We determined whether there is any difference between immature and mature rats in terms of oxidative damage caused by EMF and tested recovery groups to determine whether EMF-induced damage is reversible in immature and mature rats. **METHODS:** Twenty four immature and 24 mature rats were divided randomly and equally into six groups as follows: two control groups, immature (2 weeks old) and mature (10 weeks old); two groups were exposed to 900 MHz (28.2 ± 2.1 V/m) EMF for 2 h/day for 45 days. Two recovery groups were kept for 15 days after EMF exposure. **RESULTS:** Substantial, deleterious biochemical changes were observed in oxidative stress metabolism after EMF exposure. Antioxidant enzyme activity, glutathione levels in lymphoid organs and the antioxidant capacity of the plasma decreased, but lipid peroxidation and nitric oxide levels in PMNs and plasma and also myeloperoxidase activity in PMNs increased. Oxidative damage was tissue specific and improvements seen after the recovery period were limited, especially in immature rats. **CONCLUSIONS:** In the present study, much higher levels of irreversible oxidative damage were observed in the major lymphoid organs of immature rats than in mature rats.

El Kholi SE, El Hussein EM. Effect of 60 minutes exposure to electromagnetic field on fecundity, learning and memory, speed of movement and whole body protein of the fruit fly *Drosophila melanogaster*. J Egypt Soc Parasitol. 42(3):639-648, 2012.

This study investigated the effect of four different electrical devices as source of electromagnetic field on fecundity, learning and memory function, speed of movement, in addition to the whole body proteins of the fruit fly *Drosophila melanogaster*. The results showed that exposure to EMF has no significant effect on adult fecundity (ANOVA and Duncan's test) but alters learning and memory function in *Drosophila* larvae, especially those exposed to mobile phone. Highly significant differences occurred in the larval speed of movement after exposure to EMF, with maximal effect occurred for larvae exposed to mobile phone (their speed of movement increased 2.5 times of wild type). Some protein bands serve as characters for exposure to certain electrical devices which suggest that exposure to EMF may affect the whole body proteins.

Chiabrera A, Bianco B, Moggia E, Kaufman JJ, Zeeman-Stark modeling of the RF EMF interaction with ligand binding. Bioelectromagnetics 21(4):312-324, 2000.

The influence of radiofrequency electromagnetic exposure on ligand binding to hydrophobic receptor proteins is a plausible early event of the interaction mechanism. A comprehensive quantum Zeeman-Stark model has been developed which takes into account the energy losses of the ligand ion due to its collisions inside the receptor crevice, the attracting nonlinear endogenous force due to the potential energy of the ion in the binding site, the out of equilibrium state of the ligand-receptor system due to the basal cell metabolism, and the thermal noise. The biophysical "output" is the change

Studies that show **Cell Phone** Health Effects

of the ligand binding probability that, in some instances, may be affected by a suitable low intensity exogenous electromagnetic "input" exposure, e.g., if the depth of the potential energy well of a putative receptor protein matches the energy of the radiofrequency photon. These results point toward both the possibility of the electromagnetic control of biochemical processes and the need for a new database of safety standards.

Aksen F, Dasdag S, Akdag MZ, Askin M, Dasdag MM. The effects of whole body cell phone exposure on the t1 relaxation times and trace elements in the serum of rats Electromag Biol Med. 23:7-11, 2004.

The objective of this study was to investigate the effects of radiofrequency radiation emitted from cellular phones on: (1) trace elements such as manganese, iron, copper, zinc, (2) T1 relaxation times in serum, and (3) rectal temperature of rats exposed to microwave radiation emitted from cellular phones. Sixteen Sprague–Dawley rats were separated into two groups of eight, one sham-exposed (control) and one exposed (experimental). The rats were confined in Plexiglas cages and a cellular phone was placed 0.5 cm under the cage. For the experimental group, cellular phones were activated 20 min per day, 7 days a week, for 1 month. For the control group, a cellular phone placed beneath the cage for 20 min a day was turned off. Rectal temperatures were measured weekly. For 250-mW-radiated powers, the whole body average specified absorption rate (SAR) (rms) is 0.52 W/kg and 1-g-averaged peak SAR (rms) is 3.13 W/kg. The Mann-Whitney U test was used for statistical comparisons of groups. T1 relaxation time and the values of iron and copper in the serum of the experimental group were not changed compared to the control group ($p > 0.05$). However, manganese and zinc values in the serum of the experimental group were significantly different from the control group ($p < 0.05$). The difference in rectal temperature measured before and after exposure in the experimental groups was not statistically different from control ($p > 0.05$).

Al-Khlaiwi T, Meo SA. Association of mobile phone radiation with fatigue, headache, dizziness, tension and sleep disturbance in Saudi population. Saudi Med J. 25(6):732-736, 2004.

OBJECTIVE: The widespread use of mobile phones has been increased over the past decade; they are now an essential part of business, commerce and society. The use of mobile phones can cause health problems. Therefore, the aim of the present study is to investigate the association of using mobile phones with fatigue, headache, dizziness, tension and sleep disturbance in the Saudi population and provide health and social awareness in using these devices. **METHODS:** This study was conducted in the Department of Physiology, College of Medicine, King Saud University, Riyadh, Kingdom of Saudi Arabia during the year 2002 to 2003. In the present study, a total of 437 subjects (55.1% male and 39.9% female) were invited, they have and had been using mobile phones. A questionnaire was distributed regarding detailed history and association of mobile phones with health hazards. **RESULTS:** The results of the present study showed an association between the use of mobile phones and health hazards. The

Studies that show **Cell Phone** Health Effects

overall mean percentage for these clinical findings in all groups were headache (21.6%), sleep disturbance (4.%), tension (3.9%), fatigue (3%) and dizziness (2.4%). CONCLUSION: Based on the results of the present study, we conclude that the use of mobile phones is a risk factor for health hazards and suggest that long term or excessive use of mobile phones should be avoided by health promotion activities such as group discussions, public presentations and through electronic and print media sources.

Anghileri LJ, Mayayo E, Domingo JL, Thouvenot P. Radiofrequency-induced carcinogenesis: cellular calcium homeostasis changes as a triggering factor. *InterJ RadBiol.* 81(3):205-209, 2005.

The aim was to study the effects of radiofrequency (Rf) in a mice strain characterized by age-determined carcinogenesis of lymphatic tissues. Mice were treated with a 1 h/week Rf exposure for 4 months. A group submitted to sham exposure was used as control animals. The evolution of carcinogenesis was followed up to 18 months. The maximal life span of control mice was about 24 months. All dead animals were clinically and histologically examined to give an age-determined comparative quantification of the evolving carcinogenesis. A radiocalcium tracer method permitted the evaluation of Rf effects on transmembrane transport of extracellular calcium at 1 and 24 h after exposure. The determination of induced lipid peroxidation completed this second study. The findings show that Rf provoked an earlier general lymphocyte cell infiltration, formation of lymphoblastic ascites and extranodal tumours of different histological types, as well as an increased early mortality. The results suggest that in Rf-exposed mice, carcinogenesis may be induced earlier and with different pathological forms than in control animals. The modifications in cellular calcium homeostasis and the age-determined thymus involution appear to be important factors involved in this carcinogenesis process.

Anghileri LJ, Mayayo E, Domingo JL. Iron-radiofrequency synergism in lymphomagenesis. *Immunopharmacol Immunotoxicol.* 28(1):175-183, 2006.

The parenteral **iron** administration effects on the acceleration of lymphomagenesis by radiofrequency exposure were investigated using an animal model that develops spontaneous lymphomas with ageing. Complementary studies of the in vivo uptake of ⁵⁹Fe-labeled ferric gluconate and ferric-ATP complex showed differences of absorption and excretion between both **iron** compounds. In vitro assays of their effects on calcium cellular uptake using a cell model and tissues homogenates showed a molecular structure-dependence. The current results (mortality, clinical and histopathological examinations) demonstrated a synergism between radiofrequency and ferric gluconate, and the increased risk of radiofrequency exposure when it is simultaneous to parenteral iron administration.

Anghileri LJ, Mayayo E, Domingo JL, Thouvenot P. Evaluation of health risks caused by radio frequency accelerated carcinogenesis: the importance of processes driven by the calcium ion signal. *Eur J Cancer Prev.* 15(3):191-195, 2006.

The acceleration of carcinogenesis, which was induced either by radio frequency

Studies that show **Cell Phone** Health Effects

radiation from a cellular telephone or by the ferric-ATP complex, was similar in a mouse strain characterized by age-determined carcinogenesis of lymphoid tissues. Organ hypertrophy, the presence of lymphoid blood and ascites, the development of solid tumours, and mortality were very different to those found in control animals. These results emphasize the role of calcium ion signal influx in the activation of oncogenes and the failure of thymus-determined immune defences.

Balakrishnan K, Murali V, Rathika C, Manikandan T, Malini RP, Kumar RA, Krishnan M. Hsp70 is an independent stress marker among frequent users of mobile phones. J Environ Pathol Toxicol Oncol. 33(4):339-347, 2014.

The aim of this study was to measure the serum concentrations of heat shock protein (HSP) 70 and C-reactive protein (CRP) and the expression levels of the hsp70 gene among frequent users of mobile phones (FUMPs). We enrolled 120 employees of information technology (IT)/IT enabled service companies (FUMPs; IT professionals) and 102 infrequent users of mobile phones (IFUMPs; people from non-IT professions) as controls. The serum concentrations of HSP70 and CRP were measured by enzyme-linked immunosorbant assay and hsp70 gene expression by reverse transcription polymerase chain reaction. Significantly higher concentrations of serum HSP70 ($P < 0.00012$) and CRP ($P < 0.04$) were observed among FUMPs than IFUMPs. A higher level of hsp70 gene expression (fold induction) was observed among FUMPs than IFUMPs ($P < 7.06 \times 10^{-13}$). In contrast to the duration of exposure-dependent increase of serum concentration of CRP, the serum HSP70 concentration was found to be independent of the duration of exposure to mobile phones. Thus, the study convincingly demonstrated the role of serum HSP and CRP as systemic inflammatory biomarkers for mobile phone-induced radiation.

Survival Effects

Mortazavi S, Mosleh-Shirazi M, Tavassoli A, Taheri M, Mehdizadeh A, Namazi S, Jamali A, Ghalandari R, Bonyadi S, Haghani M, Shafie M. Increased Radioresistance to Lethal Doses of Gamma Rays in Mice and Rats after Exposure to Microwave Radiation Emitted by a GSM Mobile Phone Simulator. Dose Response. 11(2):281-292, 2012.

The aim of this study was to investigate the effect of pre-irradiation with microwaves on the induction of radioadaptive response. In the 1(st) phase of the study, 110 male mice were divided into 8 groups. The animals in these groups were exposed/sham-exposed to microwave, low dose rate gamma or both for 5 days. On day six, the animals were exposed to a lethal dose (LD). In the 2(nd) phase, 30 male rats were divided into 2 groups of 15 animals. The 1(st) group received microwave exposure. The 2(nd) group (controls) received the same LD but there was no treatment before the LD. On day 5, all animals were whole-body irradiated with the LD. Statistically significant differences between the survival rate of the mice only exposed to lethal dose of gamma radiation before irradiation with a lethal dose of gamma radiation with those of the animals pre-exposed to either microwave ($p=0.02$), low dose rate gamma ($p=0.001$) or both of these

Studies that show **Cell Phone** Health Effects

physical adapting doses ($p=0.003$) were observed. Likewise, a statistically significant difference between survival rates of the rats in control and test groups was observed. Altogether, these experiments showed that exposure to microwave radiation may induce a significant survival adaptive response.

Bartsch H, Küpper H, Scheurlen U, Deerberg F, Seebald E, Dietz K, Mecke D, Probst H, Stehle T, Bartsch C. Effect of chronic exposure to a GSM-like signal (mobile phone) on survival of female Sprague-Dawley rats: Modulatory effects by month of birth and possibly stage of the solar cycle. *Neuro Endocrinol Lett.* 31(4):457-473, 2010.

During 1997-2008 two long-term (I and II) and two life-long (III and IV) experiments were performed analyzing the effect of chronic exposure to a low-intensity GSM-like signal (900 MHz pulsed with 217 Hz, 100 microW/cm² average power flux density, 38-80 mW/kg mean specific absorption rate for whole body) on health and survival of unrestrained female Sprague-Dawley rats kept under identical conditions. Radiofrequency (RF)-exposure was started at 52-70 days of age and continued for 24 (I), 17 (II) and up to 36 and 37 months, respectively (III/IV). In the first two experiments (1997-2000) 12 exposed and 12 sham-exposed animals each were observed until they were maximally 770 or 580 days old. In experiment I no adverse health effects of chronic RF-exposure were detectable, neither by macroscopic nor detailed microscopic pathological examinations. Also in experiment II no apparent macroscopic pathological changes due to treatment were apparent. Median survival time could not be estimated since in none of the groups more than 50% of the animals had died. In the course of two complete survival experiments (2002-2005; 2005-2008) 30 RF- and 30 sham-exposed animals each were followed up until their natural end or when they became moribund and had to be euthanized. A synoptical data analysis was performed. Survival data of all four groups could be fitted well by the Weibull distribution. According to this analysis median survival was significantly shortened under RF-exposure in both experiments by 9.06% (95% CI 2.7 to 15.0%) ($p=0.0064$); i.e by 72 days in experiment III and 77 days in experiment IV as compared to the corresponding sham-treated animals (III: 799 days; IV: 852 days). Both groups of animals of experiment III showed reduced median survival times by 6.25% (95% CI -0.3 to 12.4%) ($p=0.0604$) compared to the corresponding groups of experiment IV (53 days: sham-exposed animals, 48 days: RF-exposed animals) which may be due to the fact that animals of experiment III were born in October and animals of experiment IV in May indicating that the month of birth affects life span. From the results of the last two experiments it has to be concluded that chronic exposure to a low-intensity GSM-like signal may exert negative health effects and shorten survival if treatment is applied sufficiently long and the observational period covers the full life span of the animals concerned. The current data show that survival of rats kept under controlled laboratory conditions varies within certain limits depending on the month of birth. In view of our previous observations regarding an inhibitory or no effect of RF-exposure on DMBA-induced mammary cancer during the 1997-2000 period, an additional modulatory influence on a year-to-year basis should be considered which might be related to changing solar activity during the the 11-years' sunspot cycle. These potentially complex influences of the natural environment modulating the effects of

Studies that show **Cell Phone** Health Effects

anthropogenic RF-signals on health and survival require a systematic continuation of such experiments throughout solar cycle 24 which started in 2009.

Medical Implants and Devices Effects

Wilke A, Grimm W, Funck R, Maisch B, Influence of D-net (European GSM -Standard) cellular phones on pacemaker function in 50 patients with permanent pacemakers. Pacing Clin Electrophysiol 19(10):1456-1458, 1996.

The widespread use of cellular phones in the last years has prompted some recent studies to suggest an interference of pacemaker function by cellular phone usage. To determine the risk of pacemaker patients using D-net cellular phones, we tested 50 patients with permanent pacemakers after routine pacemaker check by short phone calls using a cellular phone (Ericsson, D-net, frequency 890-915 MHz, digital information coding, equivalent to the European Groupe Systemes Mobiles standard). A six-channel surface ECG was continuously recorded from each patient to detect any interactions between pacemakers and cellular phones. Phone calls were repeated during the following pacemaker settings: (1) preexisting setting; (2) minimum ventricular rate of 90 beats/min and preexisting sensitivity; and (3) minimum ventricular rate of 90 beats/min and maximum sensitivity without T wave oversensing. Only 2 (4%) of 50 patients repeatedly showed intermittent pacemaker inhibition during calls with the cellular phone. Both pacemakers had unipolar sensing. Therefore, although interactions between cellular phone use and pacemaker function appear to be rare in our study, pacemaker dependent patients in particular should avoid the use of cellular phones.

Virtanen H, Keshvari J, Lappalainen R. The effect of authentic metallic implants on the SAR distribution of the head exposed to 900, 1800 and 2450 MHz dipole near field. Phys Med Biol. 52(5):1221-1236, 2007

As the use of radiofrequency (RF) electromagnetic (EM) fields has increased along with increased use of wireless communication, the possible related health risks have also been widely discussed. One safety aspect is the interaction of medical implants and RF devices like mobile phones. In the literature, effects on active implants like pacemakers have been discussed but the studies of passive metallic (i.e. conductive) implants are rare. However, some studies have shown that the EM power absorption in tissues may be enhanced due to metallic implants. In this study, the effect of authentic passive metallic implants in the head region was examined. A half-wave dipole antenna was used as an exposure source and the specific absorption rate (SAR, W kg⁻¹) in the near field was studied numerically. The idea was to model the presumably worst cases of most common implants in an accurate MRI-based phantom. As exposure frequencies GSM (900 and 1800 MHz) and UMTS (2450 MHz) regions were considered. The implants studied were skull plates, fixtures, bone plates and ear rings. The results indicate that some of the implants, under very rare exposure conditions, may cause a notable

Studies that show **Cell Phone** Health Effects

enhancement in peak mass averaged SAR.

Yeolekar ME, Sharma A. Use of mobile phones in ICU--why not ban? J Assoc Physicians India. 52:311-313, 2004.

Due to the rapid growth of mobile telecommunications it is predicted that by 2005 there will be 1.6 billion mobile phone users worldwide. The usage of cellphones in Intensive Care Units carries with it a high incidence of interference with a number of medical devices like implantable defibrillators, cardioverters, pacemakers, monitors and other important devices like ventilators. It is in this context that this article will throw a light on complications of cellphones use in the Intensive Care Units and various strategies that can be taken to restrict their use in the Intensive Care Units.

Vagdatli E, Konstandinidou V, Adrianakis N, Tsikopoulos I, Tsikopoulos A, Mitsopoulou K. Effects of Electromagnetic Fields on Automated Blood Cell Measurements. J Lab Autom. 2014 Jan 24. [Epub ahead of print]

The aim of this study is to investigate whether the electromagnetic fields associated with mobile phones and/or laptops interfere with blood cell counts of hematology analyzers. Random blood samples were analyzed on an Aperture Impedance hematology analyzer. The analysis was performed in four ways: (A) without the presence of any mobile phone or portable computer in use, (B) with mobile phones in use (B1: one mobile, B4: four mobiles), (C) with portable computers (laptops) in use (C1: one laptop, C3: three laptops), and (D) with four mobile phones and three laptops in use simultaneously. The results obtained demonstrated a statistically significant decrease in neutrophil, erythrocyte, and platelet count and an increase in lymphocyte count, mean corpuscular volume, and red blood cell distribution width, notably in the B4 group. Despite this statistical significance, in clinical practice, only the red blood cell reduction could be taken into account, as the mean difference between the A and B4 group was 60,000 cells/ μ L. In group D, the analyzer gave odd results after 11 measurements and finally stopped working. The combined and multiple use of mobile phones and computers affects the function of hematology analyzers, leading to false results. Consequently, the use of such electronic devices must be avoided.

Schlegel RE, Grant FH, Raman S, Reynolds D Electromagnetic compatibility study of the in-vitro interaction of wireless phones with cardiac pacemakers. Biomed Instrum Technol 32(6):645-655, 1998.

This large-scale in-vitro investigation of the interaction between hand-held wireless phones and cardiac pacemakers tested 29 pacemaker models with five different phone standards. The phones were operational and suspended on a grid above a torso simulator filled with a saline bath with the pacemaker submerged at 0.5 cm. Testing consisted of 8,296 runs, during which any interactions detected were classified by type and regularity. Only a few pacemakers were responsible for a disproportionately large number of interactions. Likewise, interactions occurred during 21% of the tests using one particular phone technology, with little or no interaction resulting from use of the other standards. Other significant factors included the relative orientation of the phone

Studies that show **Cell Phone** Health Effects

and the pacemaker case, as well as the presence or absence of an injected ECG signal. The ECG signal facilitated observation of certain forms of interaction to the extent that this study indicates the importance of including an injected ECG signal in all testing. The study also supports the recommendation to maintain a separation distance of at least 6 inches between pacemakers and wireless phones. Each pacemaker reverted to its normal operation when the phone creating an interaction was turned off. This study may be useful in ongoing efforts to define test protocols, evaluate pacemaker designs, and mitigate interactions, perhaps providing the basis for future certification and screening efforts.

Trigano A, Blandeau O, Dale C, Wong MF, Wiart J. Reliability of electromagnetic filters of cardiac pacemakers tested by cellular telephone ringing. Heart Rhythm. 2(8):837-841, 2005.

BACKGROUND: State-of-the art cardiac pacemakers are protected against radiofrequency signals. Although there have been earlier clinical and in vitro reports of cellular phone interference with implantable devices, only a few studies have been performed in recent years. The ringing phase of digital GSM or PCS cellular phones includes a brief period of peak radiated power. OBJECTIVES: This study tested the protection offered by electromagnetic filters of cardiac pacemakers against cellular phone ringing. METHODS: We performed 330 consecutive tests in 158 patients at the time of routine examination in our pacemaker follow-up clinic. The programmed parameters remained unchanged before testing. During electrocardiographic monitoring, 2 single-band digital cellular phones consecutively placed over the pacemaker pocket each received a call. The phone systems tested were 1) GSM at a maximal power output of 2 W, operating on a 900 MHz carrier frequency, and 2) PCS at a maximal output of 1 W, operating on a 1800 MHz carrier frequency. RESULTS: Interference was noted in only 5 tests, due to interaction by the GSM system with 4 unprotected pacemaker models. The GSM test was negative in 12 other tests of identical pulse generator models. The overall incidence of interference was 1.5% of tests. CONCLUSIONS: Interference by cellular phone ringing occurred only with unprotected pacemaker models. Standard programming of these unprotected models was associated with a low incidence of interference.

Trigano A, Blandeau O, Dale C, Wong MF, Wiart J. Risk of cellular phone interference with an implantable loop recorder. Int J Cardiol. 116(1):126-130, 2007.

This study examined the risk of cellular phone ringing interference with implantable loop recorders (ILR). The technical manual of ILR warns of potential interference by cellular phone in close proximity to the implanted device, corrupting the data stored in memory or causing inappropriate device operation. The ringing phase of a digital Global System for Mobile Communication (GSM) or Personal Communication Services (PCS) cellular phone includes a brief burst of peak emitted power. To obviate the risk of dysfunction in recipients of implanted ILRs, the testing was performed with externally applied devices. The ILR was positioned in the left parasternal region and the telemetry wand removed after regular programming. Digital cellular telephones were placed over

Studies that show **Cell Phone** Health Effects

the device at a 1-cm distance and calls were placed. The phone systems tested were single- or dual-band receivers. The GSM used a maximal power output of 2 W, operating on a 900 MHz carrier frequency, and the PCS a maximal output of 1 W, operating on a 1800 MHz carrier frequency. The device activator was used to store the episodes encompassing the tests. Sixty nine tests were performed in 45 patients. In 61 tests, high-frequency polymorphic artifacts were visible on manually activated recordings, beginning a few seconds before the first audible ringing tone and persisting throughout the ringing phase. Cellular phone ringing in close proximity to an externally applied ILR caused bursts of high-frequency signals during electrocardiogram monitoring, without causing permanent device dysfunction or reprogramming. Cellular telephones are a potential source of electrocardiographic artifacts on ILR recordings.

Trigano AJ, Azoulay A, Rochdi M, Campillo, A Electromagnetic interference of external pacemakers by walkie-talkies and digital cellular phones: experimental study. Pacing Clin Electrophysiol 22(4 Pt 1):588-593, 1999.

A number of experimental and clinical studies have documented the risk potential of interference with implanted pacemakers by various types of cellular phones. Radiofrequency susceptibility of external medical equipment has also been reported in experimental studies. The purpose of this experimental study was to evaluate electromagnetic interference of external pacemakers by walkie-talkies and digital cellular telephones. External bipolar pacing was monitored using a digital oscilloscope to record pacemaker pulses and electromagnetic interference separately. Tests with the walkie-talkie, Private Mobile Radio (PMR) (160 MHz, 2.5 W) were conducted during the calling phase. Tests with the cellular phones, global system for mobile communications (GSM) (900 MHz, 2 W) and Digital Cellular System (DCS) (1,800 MHz, 1 W) were conducted in the test mode. Nine widely used external pacemakers from four manufacturers were tested. Various disturbances including pacing inhibition and asynchronous pacing were observed in eight pacemakers by the PMR, in four by the GSM phone, and in two by the DCS phone. The maximum distance that interference persisted ranged from 10-200 cm. This experimental study shows a potential risk of interference of external pacemakers by walkie-talkies and cellular digital phones. Appropriate warnings should be issued against the potentially serious risks of using communication devices in the vicinity of acutely ill patients treated with temporary transvenous cardiac pacemakers.

Sakakibara Y, Mitsui T, Concerns about sources of electromagnetic interference in patients with pacemakers. Jpn Heart J 40(6):737-743, 1999.

Electromagnetic noise is rapidly increasing in our environment so electromagnetic interference (EMI) with pacemakers (PM) may become a more important problem despite technological improvements in PM. The aim of this study was to evaluate the kinds of EMI which affect the quality of life of PM patients. The participants (1,942 Japanese Association for Pacemaker Patients: Pacemaker-Tomonokai) were asked to respond to a questionnaire about their major EMI troubles, and 1,567 patients (80.7%)

Studies that show **Cell Phone** Health Effects

responded by mail. The main concerns were from mobile telephones (MT) (39%), magnetic resonance imaging (MRI) (17%), electronic kitchen appliances, automobile engines and high voltage power lines. If possible, PM implantation sites should be carefully selected not only according to the physician's convenience but also considering information on each patient's habits and physical limitations.

Tri JL, Hayes DL, Smith TT, Severson RP, Cellular phone interference with external cardiopulmonary monitoring devices. Mayo Clin Proc 76(1):11-15, 2001.

OBJECTIVES: To determine the potential effect (electromagnetic interference) of cellular telephones on external cardiopulmonary monitoring devices. METHODS: For this study, we tested 17 different medical devices with 5 portable telephones (4 digital, 1 analog) to assess the potential for electromagnetic interference. The telephones were tested in a normal operating mode to simulate a typical hospital environment with patients or their families using their cellular phones. The medical devices were connected to the appropriate simulators for proper operation while the tests were under way. The screens and alarms of the medical devices were monitored while the telephones were maneuvered in the y and z planes near the devices. Clinically important interference was defined as interference that may hinder interpretation of the data or cause the equipment to malfunction. RESULTS: Any type of interference occurred in 7 (41%) of the 17 devices tested during 54.7% of the 526 tests. The incidence of clinically important interference was 7.4%. CONCLUSIONS: Cellular telephones may interfere with the operation of external cardiopulmonary monitoring devices. However, most of the test results showed that the interference would rarely be clinically important.

Tat FH, Wah KC, Hung YH. A follow-up study of electromagnetic interference of cellular phones on electronic medical equipment in the emergency department. Emerg Med (Fremantle) 14(3):315-319, 2002.

OBJECTIVE: Considering the growing use of cellular phones and the fast appearance of new phone models, the electromagnetic interference of currently popular cellular phones on electronic medical equipment was tested. METHODS: Three Personal Communication System cellular phones were put at different distances from multiple electronic medical devices, the interference effect was observed and the electromagnetic field strength measured with a spectrum analyser. RESULTS: Only two small pieces of equipment, the CO2 airway adapter and the haemoglucostix meter were affected and then only when the phone was in very close proximity. CONCLUSION: Compared to the results of our study in 1997 testing Global System for Mobile Communication phones, the Personal Communication System phones generated less electromagnetic interference. However a much larger scaled study and an accurate international electromagnetic interference standard are recommended before any change in the current restrictive hospital policy on mobile phone usage could be recommended.

Naegeli B, Osswald S, Deola M, Burkart F, Intermittent pacemaker dysfunction caused by digital mobile telephones. J Am Coll Cardiol 27(6):1471-1477, 1996.

Studies that show **Cell Phone** Health Effects

OBJECTIVES: This study was designed to evaluate possible interactions between digital mobile telephones and implanted pacemakers. **BACKGROUND:** Electromagnetic fields may interfere with normal pacemaker function. Development of bipolar sensing leads and modern noise filtering techniques have lessened this problem. However, it remains unclear whether these features also protect from high frequency noise arising from digital cellular phones. **METHODS:** In 39 patients with an implanted pacemaker (14 dual-chamber [DDD], 8 atrial-synchronized ventricular-inhibited [VDD(R)] and 17 ventricular-inhibited [VVI(R)] pacemakers), four mobile phones with different levels of power output (2 and 8 W) were tested in the standby, dialing and operating mode. During continuous electrocardiographic monitoring, 672 tests were performed in each mode with the phones positioned over the pulse generator, the atrial and the ventricular electrode tip. The tests were carried out at different sensitivity settings and, where possible, in the unipolar and bipolar pacing modes as well. **RESULTS:** In 7 (18%) of 39 patients, a reproducible interference was induced during 26 (3.9%) of 672 tests with the operating phones in close proximity (<10 cm) to the pacemaker. In 22 dual-chamber (14 DDD, 8 VDD) pacemakers, atrial triggering occurred in 7 (2.8%) of 248 and ventricular inhibition in 5 (2.8%) of 176 tests. In 17 VVI(R) systems, pacemaker inhibition was induced in 14 (5.6%) of 248 tests. Interference was more likely to occur at higher power output of the phone and at maximal sensitivity of the pacemakers (maximal vs. nominal sensitivity, 6% vs. 1.8% positive test results, $p = 0.009$). When the bipolar and unipolar pacing modes were compared in the same patients, ventricular inhibition was induced only in the unipolar mode (12.5% positive test results, $p = 0.0003$). **CONCLUSION:** Digital mobile phones in close proximity to implanted pacemakers may cause intermittent pacemaker dysfunction with inappropriate ventricular tracking and potentially dangerous pacemaker inhibition.

Morrissey JJ, Swicord M, Balzano Q. Characterization of electromagnetic interference of medical devices in the hospital due to cell phones. Health Phys 82(1):45-51, 2002.

Concern over electromagnetic interference with medical devices due to cell phone emissions has stemmed from anecdotal reports and unpublished observations of hospital staff. In an effort to characterize electromagnetic interference concerns, representative medical devices from four large teaching hospitals were exposed to standard North American and European communication signal emissions. Of 33 medical devices tested, only 4 showed disruption of critical function due to cell phone emissions at a distance of 25 cm or greater. Although other cases of electromagnetic interference were observed, these were not critically disruptive and mainly occurred when the transmitters were at full power and placed 5 cm or closer to the medical device. Overall, no cell phone signal was exempt from producing electromagnetic interference effects. While sensitive medical devices were often affected by more than one signal type, the effects were not entirely predictable based upon the results of other signals or related medical device units or models. Because a comprehensive analysis of all medical devices in all possible electromagnetic environments was not performed, the data presented here are only intended to provide a general idea of the magnitude of electromagnetic interference effects that might be encountered in a hospital environment, as well as a

Studies that show **Cell Phone** Health Effects

standard protocol for clinical engineering groups to perform ad hoc electromagnetic interference surveys and methods to manage and/or eliminate electromagnetic interference with appropriate system engineering design including supplementary communication infrastructure, medical device shielding and positioning, and appropriate cell phone user guidelines.

Kompis M, Negri S, Hausler R. Electromagnetic interference of bone-anchored hearing aids by cellular phones. Acta Otolaryngol 120(7):855-859, 2000.

We report a case of electromagnetic interference between a bone-anchored hearing aid (BAHA) and a cellular phone. A 54-year-old women was successfully treated for severe mixed conductive and sensorineural hearing loss with a BAHA. Five years after implantation, the patient experienced a sudden feeling of dizziness, accompanied by a loud buzzing sound and by a sensation of head pressure while examining a digital mobile phone. During a subsequent experiment, the buzzing sound could be reproduced and was identified as electromagnetic interference between the BAHA and digital cellular phones. Seventeen adult BAHA users from our clinic participated in a subsequent survey. Of the 13 patients with some experience of digital cellular phones, 11 reported hearing annoying noises elicited by these devices. However, no other sensation, such as dizziness, was described. Owing to the increasing number of users of both hearing aids and cellular phones, the incidence of electromagnetic interference must be expected to increase as well. Although to date there is no evidence that such interference may be harmful or dangerous to users of conventional or bone-anchored hearing aids, unexpected interference can be a frightening experience.

Kompis M, Hausler R. Electromagnetic interference of bone-anchored hearing aids by cellular phones revisited. Acta Otolaryngol 122(5):510-512, 2002.

The electromagnetic interference of the recently introduced bone-anchored hearing aid (BAHA) model "BAHA Compact" by digital cellular phones is investigated and compared with that of the older "BAHA Classic 300" model. Measurements with two different digital cellular phones in a laboratory setting indicated that the noise level due to electromagnetic interference was at least 10 dB lower for the BAHA Compact device than for the BAHA Classic 300. To compare the experience of patients using the BAHA Compact with those using a BAHA Classic 300 in an earlier study, a survey was performed. Six users of a BAHA Compact who used digital cellular phones participated in the survey. Four patients did not hear any noise associated with the use of a digital cellular phone. Two patients reported hearing quiet sounds when they were on the telephone, but not when somebody else in the vicinity used a digital cellular phone. These findings confirm that the susceptibility to electromagnetic interference of the BAHA Compact device is low.

Kainz W, Neubauer G, Alesch F, Schmid G, Jahn O. Electromagnetic compatibility of electronic implants--review of the literature. Wien Klin Wochenschr 113(23-24):903-914, 2001.

The aim of the article was to provide an overview of published studies regarding the

Studies that show **Cell Phone** Health Effects

electromagnetic compatibility (EMC) of electronic implants. The available literature was sorted according to combinations of implant types and sources of interference. Several experiments concerning the susceptibility of pacemakers to mobile phones have been performed. The results of these experiments suggest measures that may be used to prevent the disturbance of pacemakers. For instance, instead of carrying the activated mobile phone in the breast pocket it is recommended that a distance of 30 cm be maintained between the pacemaker and the mobile phone, and that the mobile phone be used on the contralateral side of the pacemaker's location. Similar measures may be recommended for patients with implantable cardioverter defibrillators when using mobile phones. Patients with electronic implants should walk rapidly through anti theft-devices because some of these devices are liable to disturb implants. Patients with cardiac pacemakers should not be subjected to magnetic resonance imaging as far as possible. For a variety of combinations of implants and interference sources, e.g. cardiac pacemakers and base station antennas, no studies were found in the literature. It is strongly recommended that trials be carried out to evaluate the potential risk for patients in these settings.

Westermarck A, Wisten A. Miniplate osteosynthesis and cellular phone create disturbance of infraorbital nerve. J Craniofac Surg 12(5):475-478, 2001.

A 37-year-old man with a zygomatic fracture underwent surgical treatment with reduction of the fracture and osteosynthesis with a miniplate on the infraorbital rim. Postoperatively, he had numbness in the distribution area of the infraorbital nerve, but he also suffered from dysesthesia in the same area during periods when he was using his hand-held mobile phone. After surgical removal of the osteosynthesis plate, the dysesthesia associated with his mobile phone was no longer present. The plate was examined in a setup where we measured the electric current that developed on the surface of the plate under the influence of the magnetic field between the phone antenna and the metal plate. The highest currents measured on the actual plate were 141 mV in air, and 21 mV in saline. These findings indicate that there might have been a correlation between the presence of the miniplate close to the infraorbital nerve, and the dysesthesia experienced by the patient, under the influence of the energy emitted from the cellular phone.

Kainz W, Neubauer G, Alesch F, Schmid G, Jahn O. Electromagnetic compatibility of electronic implants--review of the literature. Wien Klin Wochenschr 113(23-24):903-914, 2001.

The aim of the article was to provide an overview of published studies regarding the electromagnetic compatibility (EMC) of electronic implants. The available literature was sorted according to combinations of implant types and sources of interference. Several experiments concerning the susceptibility of pacemakers to mobile phones have been performed. The results of these experiments suggest measures that may be used to prevent the disturbance of pacemakers. For instance, instead of carrying the activated mobile phone in the breast pocket it is recommended that a distance of 30 cm be maintained between the pacemaker and the mobile phone, and that the mobile phone

Studies that show **Cell Phone** Health Effects

be used on the contralateral side of the pacemaker's location. Similar measures may be recommended for patients with implantable cardioverter defibrillators when using mobile phones. Patients with electronic implants should walk rapidly through anti theft-devices because some of these devices are liable to disturb implants. Patients with cardiac pacemakers should not be subjected to magnetic resonance imaging as far as possible. For a variety of combinations of implants and interference sources, e.g. cardiac pacemakers and base station antennas, no studies were found in the literature. It is strongly recommended that trials be carried out to evaluate the potential risk for patients in these settings.

Jones RP, Conway DH. The effect of electromagnetic interference from mobile communication on the performance of intensive care ventilators. Eur J Anaesthesiol. 22(8):578-583, 2005.

Electromagnetic interference produced by wireless communication can affect medical devices and hospital policies exist to address this risk. During the transfer of ventilated patients, these policies may be compromised by essential communication between base and receiving hospitals. Local wireless networks (e.g. Bluetooth) may reduce the 'spaghetti syndrome' of wires and cables seen on intensive care units, but also generate electromagnetic interference. The aim of this study was to investigate these effects on displayed and actual ventilator performance. METHODS: Five ventilators were tested: Drager Oxylog 2000, BREAS LTV-1000, Respironics BiPAP VISION, Puritan Bennett 7200 and 840. Electromagnetic interference was generated by three devices: Simoco 8020 radio handset, Nokia 7210 and Nokia 6230 mobile phone, Nokia 6230 communicating via Bluetooth with a Palm Tungsten T Personal Digital Assistant. We followed the American National Standard Recommended Practice for On-Site, Ad Hoc Testing (ANSI C63) for electromagnetic interference. We used a ventilator tester, to simulate healthy adult lungs and measure ventilator performance. The communication device under test was moved in towards each ventilator from a distance of 1 m in six axes. Alarms or error codes on the ventilator were recorded, as was ventilator performance. RESULTS: All ventilators tested, except for the Respironics VISION, showed a display error when subjected to electromagnetic interference from the Nokia phones and Simoco radio. Ventilator performance was only affected by the radio which caused the Puritan Bennett 840 to stop functioning completely. The transfer ventilators' performance were not affected by radio or mobile phone, although the mobile phone did trigger a low-power alarm. Effects on intensive care ventilators included display reset, with the ventilator restoring normal display function within 2 s, and low-power/low-pressure alarms. Bluetooth transmission had no effect on the function of all the ventilators tested. CONCLUSION: In a clinical setting, high-power-output devices such as a two-way radio may cause significant interference in ventilator function. Medium-power-output devices such as mobile phones may cause minor alarm triggers. Low-power-output devices such as Bluetooth appear to cause no interference with ventilator function.

Irnich W, Batz L, Muller R, Tobisch R, electromagnetic interference of pacemakers by mobile phones. Pacing Clin Electrophysiol 19(10):1431-1446, 1996.

Studies that show **Cell Phone** Health Effects

The topic of interference of pacemakers by mobile phones has evoked a surprisingly strong interest, not only in pacemaker patients, but also in the public opinion. The latter is the more surprising, as in the past, the problem of interference has scarcely found the attention that it deserves in the interest of the patient. It was the intention of our investigation to test as many pacemaker models as possible to determine whether incompatibility with mobile phones of different modes may exist, using an in vitro measuring setup. We had access to 231 different models of 20 manufacturers. During the measurements, a pulse generator together with a suitable lead was situated in a 0.9 g/L saline solution, and the antenna of a mobile phone was positioned as close as possible. If the pulse generator was disturbed, the antenna was elevated until interference ceased. The gap in which interference occurred was defined as "maximum interference distance." All three nets existing in Germany, the C-net (450 MHz, analogue), the D-net (900 MHz, digital pulsed), and the E-net (1,800 MHz, digital pulsed) were tested in succession. Out of 231 pulse generator models, 103 pieces corresponding to 44.6% were influenced either by C- or D-net, if both results were totaled. However, this view is misleading as no patient will use C- and D-net phones simultaneously. Separated into C- or D-net interference, the result is 30.7% for C or 34.2% for D, respectively, of all models tested. The susceptible models represent 18.6% or 27% of today's living patients, respectively. All models were resistant to the E-net. With respect to D-net phones, all pacemakers of six manufacturers proved to be unaffected. Eleven other manufacturers possessed affected and unaffected models as well. A C-net phone only prolonged up to five pacemaker periods within 10 seconds during dialing without substantial impairment to the patient. Bipolar pacemakers are as susceptible as unipolar ones. The following advice for patients and physicians can be derived from our investigations: though 27% of all patients may have problems with D-net phones (not C- or E-net), the application should generally not be questioned. On the contrary, patients with susceptible devices should be advised that a distance of 20 cm is sufficient to guarantee integrity of the pacemaker with respect to hand held phones. Portables, on the other hand, should have a distance of about 0.5 m. Pacemaker patients really suffering from mobile phones are very rare unless the phone is just positioned in the pocket over the pulse generator. The contralateral pocket or the belt position guarantees, in 99% of all patients, undisturbed operation of the pacemaker. A risk analysis reveals that the portion of patients really suffering from mobile phones is about 1 out of 100,000. Nevertheless, it would be desirable in the future if implanting physicians would use only pacemakers with immunity against mobile phones as guaranteed by the manufacturers.

Hofgartner F, Muller T, Sigel H, [Could C- and D-network mobile phones endanger patients with pacemakers]? Dtsch Med Wochenschr 121(20):646-652, 1996. [Article in German]

OBJECTIVE: To investigate prospectively the extent of potentially harmful interference of cardiac pacemakers by mobile phones in the C (analog) and D (digital) networks in use in Germany. PATIENTS AND METHODS: 104 patients (54 men, 50 women; mean age 75.8 [40-100] years) with 58 different implanted pacemaker models (43 one-chamber and 15

Studies that show **Cell Phone** Health Effects

two-chamber systems) underwent uniform tests at various functional states with three different telephones (D1 portable 8 Watt, D1 Handy model 2 Watt, C Handy model 0.5 Watt). The distances between telephone aerial and pacemaker, as well as reception sensitivity and polarity of the pacemaker were varied. All tests were done during continuous ECG monitoring. RESULTS: 28 different pacemaker types (48.3%) in 43 patients (41.3%) showed interference in the form of pacemaker inhibition and switching to interference frequencies as well as triggering of pacemaker-mediated tachycardias in the DDD mode, as well as in the temperature-regulated frequency-adaptive function. D portables influenced pacemaker function more often and at greater distance than the D Handy model, which was little different from the c network hand phone. Reduction in pacemaker sensitivity as well as switching to bipolar reception only partly eliminated the interference. CONCLUSIONS: Patients with implanted pacemakers should if possible not use mobile phones in the C and D networks. Individual testing with suitable programming of pacemaker sensitivity and polarity can reduce the risk of interference.

Irnich W, Tobisch R, [Effect of mobile phone on life-saving and life-sustaining systems]. Biomed Tech (Berl) 43(6):164-173, 1998. [Article in German]

Since the beginning of the nineties there have been warnings not to use mobile phones in the vicinity of medical devices. Functional failures of dialysis machines, respirators and defibrillators prompted the banning of their use in many hospitals in Scandinavia, and then in other countries. Since we believe that a general ban in hospitals is problematic, we decided to investigate the influence of mobile telephone on life-saving and/or life-support systems, with the aim of establishing rules for its use in hospitals. We investigated available phones of varying power of the C-, D- and E-net, as also of a cordless phone meeting the DECT standard. The aim was to identify the devices susceptible to interference and determine the minimum distances at which interference occurred. A total of 224 devices classified into 23 types of devices were examined. Nine different sets of transmission conditions were applied, giving a total of 2016 tests. Our results permit the conclusion that the ban on mobile phones in hospitals is based not on actual events, but on theoretical considerations in the absence of any practical information on the actual susceptibility of devices and their reaction to the electromagnetic fields involved. The fact that hazardous situations are very rare is due firstly to the need for the simultaneous occurrence of four coincidences, and the fail-safe feature of medical devices. We would therefore recommend that all life-saving and life-support systems that can also be used outside the hospital should be made mobile phone-proof. When apnoea monitors and respirators are protected from such interference, hazardous situations could be avoided by establishing the rule: "No portables, and mobile phones only at a distance of at least 1 metre from medical devices". With regard to emergency telephones, the minimum distance to medical devices should be at least 1.5 metres.

Jimenez A, Hernandez Madrid A, Pascual J, Gonzalez Rebollo JM, Fernandez E, Sanchez A, Ortega J, Lozano F, Munoz R, Moro C, [Electromagnetic interference between automatic defibrillators and digital and analog cellular telephones]. Rev Esp Cardiol

Studies that show **Cell Phone** Health Effects

51(5):375-382, 1998. [Article in Spanish]

BACKGROUND AND OBJECTIVES: Functional pacemaker interference by mobile telephones has been described with analogical systems and with possible greater influence, digital systems, including inhibition and inadequate pacing. The influence of both system has not been extensively studied in patients with implantable cardioverter defibrillators (ICD). **PATIENTS AND METHODS:** We studied the influence of mobile phones, both digital and analogic network, on the performance of several models of defibrillators, in a standardised test set up designed to provide high sensitivity. The purpose of our study was to establish whether there are any influences on ICD functions, both in in vivo and in in vitro models. Several mobile phones, with different transmission powers, were moved towards the defibrillator and the electrode, under continuous documentation of defibrillator sensing and interrogation afterwards. The experimental model was performed with the aid of an arrhythmia simulator (Intersim) and demo-defibrillators. The tests were repeated both in and out of a solution of saline water with an impedance within normal human limits. **RESULTS:** Partial loss of telemetry was found in 14 patients, 8 with analogical phones and 6 with digital phones. Fourteen patients showed alterations only on the surface electrocardiogram channel and five on the intracavitary channel. The same results were reproduced in the in vitro model. However, the in vitro test allowed us to simulate multiple ventricular arrhythmias, and demonstrate the normal sensing and functioning of the defibrillator during a "spontaneous" arrhythmia. After testing, we demonstrate that no real oversensing/undersensing was documented in any device. There was no evidence of ICD reprogramming or pacing inhibition. In particular, no inadequate therapies were delivered. **CONCLUSIONS:** a) in our series, we have not demonstrated clinically significant electromagnetic interferences with mobile phones of digital or analogical networks; b) the in vitro model allowed us to conclude that even if a spontaneous arrhythmia appears, the function of the defibrillator is not altered; c) the use of mobile phones seems to be safe for defibrillator patients, and d) however, some basic rules, such as to maintain the phone at least 15 cm away from the defibrillator, are advised.

Hayes DL, Wang PJ, Reynolds DW, Estes M 3rd, Griffith JL, Steffens RA, Carlo GL, Findlay GK, Johnson CM. Interference with cardiac pacemakers by cellular telephones. N Engl J Med 336(21):1473-1479, 1997.

BACKGROUND: A growing body of evidence suggests that electromagnetic interference may occur between cardiac pacemakers and wireless hand-held (cellular) telephones, posing a potential public health problem. Electromagnetic interference may occur when the pacemaker is exposed to an electromagnetic field generated by the cellular telephone. **METHODS:** In this multicenter, prospective, crossover study, we tested 980 patients with cardiac pacemakers with five types of telephones (one analogue and four digital) to assess the potential for interference. Telephones were tested in a test mode and were programmed to transmit at the maximal power, simulating the worst-case scenario; in addition, one telephone was tested during actual transmission to simulate actual use. Patients were electrocardiographically monitored while the telephones were tested at the ipsilateral ear and in a series of maneuvers directly over the pacemaker.

Studies that show **Cell Phone** Health Effects

Interference was classified according to the type and clinical significance of the effect. RESULTS: The incidence of any type of interference was 20 percent in the 5533 tests, and the incidence of symptoms was 7.2 percent. The incidence of clinically significant interference was 6.6 percent. There was no clinically significant interference when the telephone was placed in the normal position over the ear. Interference that was definitely clinically significant occurred in only 1.7 percent of tests, and only when the telephone was held over the pacemaker. Interference was more frequent with dual-chamber pacemakers (25.3 percent) than with single-chamber pacemakers (6.8 percent, $P<0.001$) and more frequent with pacemakers without feed-through filters (28.9 to 55.8 percent) than with those with such filters (0.4 to 0.8 percent, $P=0.01$). CONCLUSIONS: Cellular telephones can interfere with the function of implanted cardiac pacemakers. However, when telephones are placed over the ear, the normal position, this interference does not pose a health risk.

Geller L, Thuroczy G, Merkely B. Orv Hetil 142(36):1963-1970, 2001. [Article in Hungarian]

Electromagnetic compatibility (EMC) of cellular phones and pacemakers (PM) was examined in four different cellular phone system (NMT, GSM, RLL, DCS 1800 MHz) and in fifteen different PM type in-vitro and in-vivo in humans. After more than 1100 in-vitro and 130 in-vivo tests we concluded, that the electromagnetic immunity of the PMs which are implanted in Hungary is suitable with only few exceptions. The highest rate of EMC problems was observed with NMT 450 MHz cellular phones (10.5%-63%). There was no EMC disturbance observed with GSM and DCS 1800 MHz cellular phones. There was only one case when clinically significant symptom was noticed with only one PM type and with NMT system cellular phone when the distance of cellular phone was 3-4 cms, and the power was maximal. There was not any EMC disturbance observed with none of the cellular phone systems during normal talking and when the distance of the PM and cellular phone was more than 20 cms. Our study supports guidelines which suggest that PM patients should contact their physicians when using cellular phones and cellular phones and PMs should not get closer than 20 cms.

Glenister H, How do mobile phones affect electromedical devices? Nurs Times 94(15):44-45, 1998.

Mobile telephones and other electronic communication devices can interfere with medical equipment when used in close proximity. A study of different devices by the Medical Devices Agency showed that emergency services' radio handsets were the most likely to cause interference. It recommends that cell telephones be switched off in theatres and treatment areas and at a patients' bedsides where sensitive medical devices are in use.

Fetter JG, Ivans V, Benditt DG, Collins J, Digital cellular telephone interaction with implantable cardioverter-defibrillators. J Am Coll Cardiol 31(3):623-628, 1998.

OBJECTIVES: This study sought to determine, in vivo, whether electromagnetic interference (EMI), generated by North American Digital Communications (NADC)/Time

Studies that show **Cell Phone** Health Effects

Division Multiple Access-50-Hz (TDMA-50) mobile cellular digital telephone model AT&T 6650, disturbs normal implantable cardioverter-defibrillator (ICD) operation and to verify these observations in vitro by testing a selection of telephones representing worldwide systems. METHODS: The effects of cellular phone interference on the operation of various models of market-released ICDs from a single manufacturer, Medtronic, Inc., were tested. The in vivo clinical test was undertaken in 41 patients using the AT&T 6650 digital telephone with the NADC/TDMA-50 technology. The in vitro component of the study was examined twofold: 1) antenna generated far field; and 2) analog/digital cellular telephone near field. RESULTS: None of the ICDs tested in 41 patients were affected by oversensing of the EMI field of the cellular telephones during the in vivo study. Therefore, the binomial upper 95% confidence limit for the failure rate of 0% is 7%. The in vitro antenna-generated field testing showed that telephone modulation frequencies used in the international Global System Mobile and TDMA-50 cellular telephone technologies did not result in ICD sensing interference at the predicted electric field intensity. The in vitro near field tests were performed using both analog and digital cellular telephones in service, or in the test mode, and indicated no interaction with normal operation. However, the static magnetic field generated by the cellular telephone placed over the ICD at a distance ≤ 0.5 cm will activate the internal reed switch, resulting in temporary suspension of ventricular tachycardia and fibrillation detection. CONCLUSIONS: We conclude that TDMA-50 cellular telephones did not interfere with these types of ICDs. However, we recommend that the patient not carry or place the digital cellular telephone within 15 cm (6 in.) of the ICD.

Altamura G, Toscano S, Gentilucci G, Ammirati F, Castro A, Pandozi C, Santini M, Influence of digital and analogue cellular telephones on implanted pacemakers. Eur Heart J 18(10):1632-4161, 1997.

The aim of this study was to find out whether digital and analogue cellular 'phones affect patients with pacemakers. The study comprised continuous ECG monitoring of 200 pacemaker patients. During the monitoring certain conditions caused by interference created by the telephone were looked for: temporary or prolonged pacemaker inhibition; a shift to asynchronous mode caused by electromagnetic interference; an increase in ventricular pacing in dual chamber pacemakers, up to the programmed upper rate. The Global System for Mobile Communications system interfered with pacing 97 times in 43 patients (21.5%). During tests on Total Access of Communication System telephones, there were 60 cases of pacing interference in 35 patients (17.5%). There were 131 interference episodes during ringing vs 26 during the on/off phase; ($P < 0.0001$); 106 at maximum sensitivity level vs 51 at the 'base' value; $P < 0.0001$). Prolonged pacing inhibition (> 4 s) was seen at the pacemaker 'base' sensing value in six patients using the Global system but in only one patient using Total Access. CONCLUSION: Cellular 'phones may be dangerous for pacemaker patients. However, they can be used safely if patients do not carry the 'phone close to the pacemaker, which is the only place where high risk interference has been observed.

Chen WH, Lau CP, Leung SK, Ho DS, Lee IS, Interference of cellular phones with

Studies that show **Cell Phone** Health Effects

implanted permanent pacemakers. Clin Cardiol 19(11):881-886, 1996.

BACKGROUND AND HYPOTHESIS: Occasional reports have suggested that cellular phones may interfere with permanent pacemakers. Our investigation sought to determine systematically the effects of commercially available cellular phones on the performances of different pacing modes and sensing lead configurations of permanent implanted pacemakers. **METHODS:** We conducted the study in 29 patients implanted with single- or dual-chamber bipolar rate-adaptive permanent pacemakers (a total of nine different models and six different sensors: minute ventilation, activity sensing using either accelerometer or piezoelectric crystal, QT and oxygen saturation sensing) from four different manufacturers. Three different cellular phones with analog or digital coding with maximum power from 0.6 to 2 W were used to assess the effect of pacemaker interference. Each cellular phone was positioned at (1) above the pacemaker pocket, (2) the ear level ipsilateral to the pacemaker pocket, and (3) the contralateral ear level. Surface electrocardiograms, intracardiac electrograms, and marker channels were recorded where possible during the following maneuvers at each position: (1) calls made by a stationary phone to cellular phone, and (2) calls made from the cellular phone to a stationary phone. A total of eight different pacing modes [DDD(R), VDD(R), AAI(R) and VVI(R)] in both unipolar and bipolar sensing configurations was tested. **RESULTS:** Interference was demonstrated during cellular phone operation in 74 of 2,418 (3.1%) episodes in eight patients. Three types of interference were observed: inhibition of pacing output, rapid ventricular tracking in DDD(R) or VDD(R) mode, and asynchronous pacing. All were observed only with the cellular phone positioned above the pacemaker pocket. Interference occurred prior to and after the termination of the ringing tone of the cellular phone in 57% of cases. Cellular phones with either digital or analog technology could cause interference. Unipolar atrial lead was most susceptible to interference (relative frequency of interference: unipolar 1.8%, bipolar 0.4%, $p < 0.05$; atrial 2.9%, ventricular 1%, $p < 0.05$). There was no sensor-driven rate acceleration during all tests. In all patients, reprogramming of the sensitivity level successfully prevented cellular phone interference. **CONCLUSIONS:** Commercially available cellular phones can cause reversible interference to implanted single- or dual-chamber permanent pacemakers. The effect is maximal with high atrial unipolar sensitivity, especially in single pass VDD(R) systems. Both digital and analog cellular phones can lead to interference. Pacemaker interference can occur prior to a warning sign (ringing tone) of the phone and may have significant implications in patient safety.

Barbaro V, Bartolini P, Donato A, Militello C, Electromagnetic interference of analog cellular telephones with pacemakers. Pacing Clin Electrophysiol 19(10):1410-1418, 1996.

The aim of this study was to verify whether there is a public health risk from the interference of analog cellular telephones with pacemakers. We used a human trunk simulator to reproduce an actual implant, and two cellular telephones working with the TACS (Total Access Communication System) standard. Results showed that the electromagnetic field radiated from the analog cellular telephones interfered with a large number of the pacemakers tested (10/25). When the telephone antenna was in

Studies that show **Cell Phone** Health Effects

close proximity to the pacemaker head, pacemaker desensitizing and sensitizing and pulse inhibition was detected at the moment of an incoming call and throughout ringing. In the worst case of pulse inhibition, the pacemaker skipped three nonconsecutive beats and then resumed its normal pacing, while the desensitizing and sensitizing phenomena persisted as long as the interfering signal was on. Pulse inhibition was also observed when the connection did not succeed. Maximum sensing threshold variation was about 186% (increase) and 62% (decrease) for desensitizing and sensitizing phenomena, respectively. It was also demonstrated that the signal emitted by analog cellular telephones during the crossing of contiguous cells could induce pacemaker pulse inhibition, but under our experimental conditions this event did not seem to pose a risk for the pacemaker patient.

Bassen HI, Moore HJ, Ruggera PS, Cellular phone interference testing of implantable cardiac defibrillators in vitro. Pacing Clin Electrophysiol 21(9):1709-1715, 1998.

An in vitro study was undertaken to investigate the potential for cellular telephones to interfere with representative models of presently used ICDs. Digital cellular phones (DCPs) generate strong, amplitude modulated fields with pulse repetition rates near the physiological range sensed by the ICD as an arrhythmia. DCPs with Time Division Multiple Access (TDMA) pulsed amplitude modulation caused the most pronounced effect--high voltage firing or inhibition of pacing output of the ICDs. This electromagnetic interference (EMI) occurred only when the phones were within 2.3-5.8 cm of the ICD pulse generator that was submerged 0.5 cm in 0.18% saline. ICD performance always reverted to baseline when the cellular phones were removed from the immediate proximity of the ICD. Three models of ICDs were subjected to EMI susceptibility testing using two types of digital phones and one analog cellular phone, each operating at their respective maximum output power. EMI was observed in varying degrees from all DCPs. Inhibition of pacer output occurred in one ICD, and high voltage firing occurred in the two other ICDs, when a TDMA-11 Hz DCP was placed within 2.3 cm of the ICD. For the ICD that was most sensitive to delivering unintended therapy, inhibition followed by firing occurred at distances up to 5.8 cm. When a TDMA-50 Hz phone was placed at the minimum test distance of 2.3 cm, inhibition followed by firing was observed in one of the ICDs. EMI occurred most frequently when the lower portion of the monopole antenna of the cellular phone was placed over the ICD header.

Occhetta E, Plebani L, Bortnik M, Sacchetti G, Trevi G, Implantable cardioverter defibrillators and cellular telephones: is there any interference? Pacing Clin Electrophysiol 22(7):983-989, 1999.

The aim of our study was to consider cellular telephone interference using different cellular telephones and implantable cardioverter defibrillator (ICD) models. Thirty (26 men, 4 women) patients with ICDs were considered during follow-up. The ICD models were: Telectronics (7), CPI (7), Medtronic (7), Ventritex (5), and Ela Medical (4). All patients were monitored with surface ECG; permanent telemetric endo-ECG monitoring was activated. Then, the effect of two different European telephone systems were tested: TACS system (Sony CM-R111, 2W power) and GSM system (Motorola MG1-4A11,

Studies that show **Cell Phone** Health Effects

2 W power). For both systems, the effect during call, reception, active conversation (dialogue), and passive conversation (listening) were observed. Cellular telephones were located first in contact with the programming head, then near the leads system, and lastly, in the hands of the patient. At the end of the evaluations, memories were interrogated again to check for false arrhythmia detections. In five of these patients during arrhythmia induction at device implant (first implant or ICD replacement), we also evaluated possible interference between cellular telephones in the reception phase and the ventricular fibrillation detection phase of the ICD. All evaluated models showed significant noise in the telemetric transmission when the cellular telephone (both TACS and GSM) was located near the ICD and the programming head; noise was particularly significant during call and reception, in most cases leading to loss of telemetry. No false arrhythmia detections have been observed during tests with cellular telephones located on the ICDs. During tests performed with cellular telephones located near the leads or in the hands of patients, no telemetric noises or false arrhythmia detections were observed. During induced ventricular fibrillation and cellular telephones in reception mode near the device, the arrhythmia recognition was always correct and not delayed. In conclusion, present ICD models seem to be well protected from electromagnetic interference caused by European cellular telephones (TACS and GSM), without under-/oversensing of ventricular arrhythmias. However, cellular telephones disturb telemetry when located near the programming head. ICD patients should not be advised against the use of cellular telephones, but it has to be avoided during ICD interrogation and programming.

Cell Phone Addiction

Bianchi A, Phillips JG. Psychological predictors of problem mobile phone use.

Cyberpsychol Behav. 8(1):39-51, 2005.

Mobile phone use is banned or illegal under certain circumstances and in some jurisdictions. Nevertheless, some people still use their mobile phones despite recognized safety concerns, legislation, and informal bans. Drawing potential predictors from the addiction literature, this study sought to predict usage and, specifically, problematic mobile phone use from extraversion, self-esteem, neuroticism, gender, and age. To measure problem use, the Mobile Phone Problem Use Scale was devised and validated as a reliable self-report instrument, against the Addiction Potential Scale and overall mobile phone usage levels. Problem use was a function of age, extraversion, and low self-esteem, but not neuroticism. As extraverts are more likely to take risks, and young drivers feature prominently in automobile accidents, this study supports community concerns about mobile phone use, and identifies groups that should be targeted in any intervention campaigns.

Koylu H, Mollaoglu H, Ozguner F, Nazyroglu M, Delibab N. Melatonin modulates 900 Mhz microwave-induced lipid peroxidation changes in rat brain. Toxicol Ind Health. 22(5):211-216, 2006.

Studies that show **Cell Phone** Health Effects

Microwaves (MW) from cellular phones may affect biological systems by increasing free radicals, which may enhance lipid peroxidation levels of the brain, thus leading to oxidative damage. Melatonin is synthesized in and secreted by the pineal gland at night and exhibits anti-oxidant properties. Several studies suggest that supplementation with anti-oxidant can influence MW-induced brain damage. The present study was designed to determine the effects of MW on the brain lipid peroxidation system, and the possible protective effects of melatonin on brain degeneration induced by MW. Twenty-eight Sprague-Dawley male rats were randomly divided into three groups as follows: (1) sham-operated control group (N = 8); (2) study 900-MHz MW-exposed group (N = 8); and (3) 900-MHz MW-exposed+melatonin (100 microg/kg sc before daily MW exposure treated group) (N = 10). Cortex brain and hippocampus tissues were removed to study the levels of lipid peroxidation as malonyl dialdehyde. The levels of lipid peroxidation in the brain cortex and hippocampus increased in the MW group compared with the control group, although the levels in the hippocampus were decreased by MW+melatonin administration. The brain cortex lipid peroxidation levels were unaffected by melatonin treatment. We conclude that melatonin may prevent MW-induced oxidative changes in the hippocampus by strengthening the anti-oxidant defense system, by reducing oxidative stress products.

Synergistic Effects with Other Agents

Pakhomov AG, Dubovick BV, Degtyariov IG, Pronkevich AN, Microwave influence on the isolated heart function: II. Combined effect of radiation and some drugs. Bioelectromagnetics 16(4):250-254, 1995.

The combined effects of microwave radiation and some drugs were studied in an isolated frog auricle preparation. The experiments established that exposure to pulse-modulated 915 MHz microwaves for up to 40 min had no effect on either the rate or the amplitude of spontaneous auricle twitches, unless the average absorbed power was high enough to produce preparation heating. Treatment of the preparation with saline containing (0.6-3.0) 10^{-5} M of propranolol or (0.5-1.5) 10^{-7} M of atropine altered neither its pacemaker nor its contractile functions; these drugs also had no effect when they were combined with nonthermal microwave irradiation. Caffeine (1 mM) strongly increased the average heart power, which was calculated as the product of twitch rate and amplitude. The caffeine effect appeared to be significantly augmented (by about 15%, $P < 0.02$) under exposure to burst-type pulsed microwaves (pulse width, 1.5 msec; pause, 2.5 msec; 8 pulses/burst, 16 bursts/s; average SAR, 8-10 W/kg). By itself, this modulation was not effective; the heating of the preparation and saline during exposure was approximately 0.1 degrees C, which could not account for the detected changes. The experimental results demonstrate that caffeine treatment increases the microwave sensitivity of the frog auricle preparation and reveals primarily subthreshold, nonthermal microwave effect.

Nelson BK, Conover DL, Brightwell WS, Shaw PB, Werren D, Edwards RM, Lary JM,

Studies that show **Cell Phone** Health Effects

Marked increase in the teratogenicity of the combined administration of the industrial solvent 2-methoxyethanol and radiofrequency radiation in rats. *Teratology* 43(6):621-634, 1991.

Limited published animal research reports synergistic teratogenic effects following combined hyperthermia (induced by elevated ambient temperature) and administration of chemical teratogens. Radiofrequency (RF) radiation is widely used in occupational environments. Since RF radiation also elevates the body temperature of, and is teratogenic to, exposed animals, concurrent RF radiation and chemical agent administration may enhance teratogenicity. The present exploratory study, consisting of preliminary dose-finding studies and the primary study, was designed to investigate whether concurrent exposure of rats to RF radiation and the industrial solvent 2-methoxyethanol (2ME) can enhance the developmental toxicity of either agent acting alone. Preliminary dose-finding studies using small numbers of rats investigated the ability of various RF radiation conditions and doses of 2ME to produce external malformations (primarily of the paws) when administered on gestation day 13. Based on these preliminary studies, RF radiation exposure [sufficient to elevate rectal temperature to 42.0 degrees C (4 degrees C above normal for rats) for 30 min] and 2ME administration (150 mg/kg) were selected for the primary study. In the primary study, groups of 18 to 27 pregnant rats were administered RF radiation exposure and distilled water gavage, 2ME gavage and sham RF exposure, RF radiation exposure and 2ME gavage concurrently, or sham RF exposure and distilled water gavage. Pregnant rats were sacrificed on gestation day 20, and the offspring were examined for external malformations. Combined exposures enhanced the adverse effects produced by either experimental agent alone (no malformations were detected in the double sham group). Mean fetal malformations/litter increased from 14% after 2ME and sham RF (15/26 litters affected, with an average of 2 fetuses/litter malformed) and 30% after RF radiation and water gavage (10/18 litters affected, with an average of 4 fetuses/litter malformed), to 76% after the combined treatment (18/18 litters affected, with an average of 12 fetuses/litter malformed). In addition to a significant increase in the frequency of malformations, the severity of malformations also was enhanced by the combination treatment (on a relative severity ranking scale, the 2ME severity score was less than 1, the RF score was 3, and the combination score was 6). This study provided evidence of synergism between RF radiation and 2ME administration, but additional research will be required to characterize the extent of synergism between these two agents. Potential interactive effects between chemical and physical agents need to be investigated to determine the extent to which such interactions should impact occupational exposure standards.

Nelson BK, Conover DL, Shaw PB, Werren DM, Edwards RM, Hoberman AM, Interactive developmental toxicity of radiofrequency radiation and 2-methoxyethanol in rats. *Teratology* 50(4):275-293, 1994.

Concurrent exposures to chemical and physical agents occur in the workplace; exposed workers include those involved with the microelectronics industry, plastic sealers, and electrosurgical units. Previous animal research indicates that hyperthermia induced by

Studies that show **Cell Phone** Health Effects

an elevation in ambient temperature can potentiate the toxicity and teratogenicity of some chemical agents. We previously demonstrated that combined exposure to radiofrequency (RF; 10 MHz) radiation, which also induces hyperthermia and is teratogenic to exposed animals, and the industrial solvent, 2-methoxyethanol (2ME), produces enhanced teratogenicity in rats. The present study replicates and extends the previous research investigating the enhanced teratogenicity of combined RF radiation and 2ME exposures. The interactive dose-related teratogenicity of RF radiation (sham exposure or maintaining colonic temperatures at 42.0 degrees C for 0, 10, 20, or 30 min) and 2ME (0, 75, 100, 125, or 150 mg/kg) was investigated by administering various combinations of RF radiation and 2ME to groups of rats on gestation days 9 or 13; gestation-day 20 fetuses were examined for external, skeletal, and visceral malformations. The results are consistent with and extend our previous research findings. Synergism was observed between RF radiation and 2ME for some treatment combinations, but not for others. The study also clarified which gestational periods, RF radiation exposure durations, and 2ME doses would be most informative in future interaction studies to determine the lowest interactive effect level. Day 9 exposures generally evidenced little effect by 2ME, either by itself or in combination with RF radiation. In contrast, day 13 exposures resulted in highly significant effects from 2ME and RF radiation. The structures showing strong evidence of effects from both 2ME and RF radiation after exposure on gestation day 13 were the forepaw digits, forepaw phalanges, hindpaw digits, hindpaw phalanges, hind limbs, metacarpals, and metatarsals. Statistical analyses did not show a global synergistic effect, but did show evidence for a synergistic effect at intermediate levels of the dose ranges. Future research will address potential interactions at lower doses.

Mortazavi SMJ, Mosleh-Shirazi MA, Tavassoli AR, Taheri M, Bagheri Z, Ghalandari R, Bonyadi S, Shafie M, Haghani M. A comparative study on the increased radioresistance to lethal doses of gamma rays after exposure to microwave radiation and oral intake of flaxseed oil. Iran. J. Radiat. Res. 9(1): 9-14, 2011.

Background: Mobile phones, use electromagnetic radiation in the microwave range. On the other hand, there is only one report on radioprotective effects of flaxseed oil. The aim of this study was to investigate the effect of irradiation of rats with microwaves and/or treatment with flaxseed oil on the induction of adaptive response to a subsequent lethal dose (LD) of gamma rays. **Materials and Methods:** Eighty male rats were randomly divided into 6 groups of 13-15 animals. The animals in the 1st to 5th groups received microwave exposure, microwave+flaxseed oil (dissolved in olive oil), flaxseed (continued after LD), flaxseed, and olive oil. At day 5, all animals were whole-body irradiated with a previously reported LD 50/30 of 8 Gy gamma radiation. The 6th group (controls) received the same LD 50/30, but there was not any other treatment before or after the LD. **Results:** No death event was observed during days 1-9 after LD irradiation in either group. At day 10, death events started in the 4th group. Thirty days after irradiation of the animals, the survival fractions for the control group, as expected, was 53.3% while there was no death event in the 1st group (survival rate of 100% in

Studies that show **Cell Phone** Health Effects

microwave-pretreated animals). The survival fractions for the 2nd to 5th groups were 69.2%, 92.3%, 46.1%, and 61.5%, respectively. **Conclusion:** While these findings open new horizons in radiation protection, the radioresistance induced by microwave radiations emitted by a mobile phone may interfere with the outcome of any subsequent therapeutic application of photons or radioisotopes.

Gadhia PK, Shah T, Mistry A, Pithawala M, Tamakuwala D. A preliminary study to assess possible chromosomal damage among users of digital mobile phones.

Electromag Biol Med 22:149-159, 2003.

In a preliminary study to examine possible lymphocyte chromosomal damage, we have tested two cytogenetic endpoints, namely, chromosomal aberrations (CA) and sister chromatid exchange frequencies (SCE), in 24 mobile phone users (12 nonsmoker–nonalcoholic subjects and 12 smoker–alcoholics), who used digital mobile phones for at least 2 years, employing Gaussian Minimum Shift Keying modulations with uplink frequencies at 935–960 MHz. and downlinks at 890–915 MHz. For comparison, the control study group included another 24 individuals, matched according to their age, sex, drinking and smoking habits, as well as similar health status, working habits, and professional careers; but did not use mobile phones. Blood samples of 12 mobile users (6 smoker–alcoholic and 6 nonsmoker–nonalcoholic) and 12 controls (identical to mobile users in every respect) were further treated with a known mutagen Mitomycin-C (MMC) to find out comutagenic/synergistic effect. A complete blood picture for each individual was assessed with an automatic particle cell counter. There was a significant increase ($P < 0.05$) in dicentric chromosomes among mobile users who were smoker–alcoholic as compared to nonsmoker–nonalcoholic; the same held true for controls of both types. After MMC treatment, there was a significant increase in dicentrics ($P < 0.05$) and ring chromosomes ($P < 0.001$) in both smoker–alcoholic and nonsmoker–nonalcoholic mobile users when compared with the controls. Although SCEs showed a significant increase among mobile users, no change in cell cycle progression was noted. The hematological picture showed only minor variations between mobile users and controls.

Del Signore A, Boscolo P, Kouri S, Di Martino G, Giuliano G, Combined effects of traffic and electromagnetic fields on the immune system of fertile atopic women. Ind Health 38(3):294-300, 2000.

Object of this preliminary study was the immune response to high or low frequency electromagnetic fields (ELMF) of non-atopic and atopic fertile women with uniform exposure to toxic compounds produced by traffic. Women were divided in group A (non-atopic, non-exposed to ELMF); B (atopic, non-exposed to ELMF); C (non-atopic, exposed to ELMF); D (atopic, exposed to ELMF). "In vitro" cell proliferation of peripheral blood mononuclear cells (PBMC) of atopic women (groups B and D) stimulated by phytohaemagglutinin (PHA) was reduced. The ELMF exposed women (groups C and D) showed lower levels of blood NK CD16(+)-CD56+ lymphocyte subpopulations and of "in vitro" production of interferon-gamma (both spontaneously and in presence of PHA) by PBMC, suggesting that ELMF reduces blood cytotoxic activity. Serum IgE of the atopic

Studies that show **Cell Phone** Health Effects

women exposed to ELMF (group D) was higher than that of the other groups. Linear discriminant analysis including serum zinc and copper (essential enzymes for immune functions), blood lead and urinary transtrans muconic acid, a metabolite of benzene (markers of exposure to traffic) and key parameters of immune functions (CD16(+)-CD56+ lymphocyte subset, serum IgE, interferon-gamma produced by PBMC in presence of PHA, stimulation index of blastogenesis) showed absence of significant difference between groups A and C and a marked separation of groups B and D. This datum suggests that ELMF have a greater influence on atopic women exposed to traffic than on non-atopic ones.

Boga A, Emre M, Sertdemir Y, Akillioglu K, Binokay S, Demirhan O. The effect of 900 and 1800MHz GSM-like radiofrequency irradiation and nicotine sulfate administration on the embryonic development of *Xenopus laevis*. Ecotoxicol Environ Saf. 2014 Dec 19;113C:378-390. doi: 10.1016/j.ecoenv.2014.12.020. [Epub ahead of print]

The aim of this study was to investigate the effects of GSM-like radiofrequency electromagnetic radiation (RF EMR) and nicotine sulfate (NS) exposure on *Xenopus* embryonic development. The developmental effects of GSM-like RF-EMR (900-1800MHz, at a SAR value of 1W/kg and NS on *Xenopus laevis* embryos were investigated). Following the application of radiofrequency radiation and/or NS administration, the embryos were closely examined in order to determine their possible teratogenic effects. *Xenopus* frogs obtained from the Department of Physiology of the Cukurova University, in accordance described by the Standard Guide of the American Society for Testing and Materials (ASTM). Following the exposure of *Xenopus* embryos to RF-EMR at 900 and 1800MHz (1.0W/kg) for 4, 6 and 8h; the whole body specific energy absorption rate (SAR) of the embryos was calculated. With the exception of irradiation at 1800MHz no dramatic developmental anomalies were observed in the *Xenopus* embryos in association with RF-EMR applications. Combined RF-EMR and NS applications resulted in dramatic abnormalities and death among the *Xenopus* embryos. The study results indicated that GSM-like RF-EMR (e.g. radiation from cell phones) was not as harmful to *Xenopus* embryos as might have been expected. However, the combined effects of GSM-like RF-EMR and NS on *Xenopus* embryos were more severe than the effect of RF-EMR or NS alone. In conclusion, the study results appear to suggest that the combined use of nicotine and cell phones might result in more pronounced detrimental effects on the health of smokers.

Byun YH, Ha M, Kwon HJ, Hong YC, Leem JH, Sakong J, Kim SY, Lee CG, Kang D, Choi HD, Kim N. Mobile phone use, blood lead levels, and attention deficit hyperactivity symptoms in children: a longitudinal study. PLoS One. 2013;8(3):e59742.

BACKGROUND: Concerns have developed for the possible negative health effects of radiofrequency electromagnetic field (RF-EMF) exposure to children's brains. The purpose of this longitudinal study was to investigate the association between mobile phone use and symptoms of Attention Deficit Hyperactivity Disorder (ADHD) considering the modifying effect of lead exposure. METHODS: A total of 2,422 children at 27 elementary schools in 10 Korean cities were examined and followed up 2 years later.

Studies that show **Cell Phone** Health Effects

Parents or guardians were administered a questionnaire including the Korean version of the ADHD rating scale and questions about mobile phone use, as well as socio-demographic factors. The ADHD symptom risk for mobile phone use was estimated at two time points using logistic regression and combined over 2 years using the generalized estimating equation model with repeatedly measured variables of mobile phone use, blood lead, and ADHD symptoms, adjusted for covariates. RESULTS: The ADHD symptom risk associated with mobile phone use for voice calls but the association was limited to children exposed to relatively high lead. CONCLUSIONS: The results suggest that simultaneous exposure to lead and RF from mobile phone use was associated with increased ADHD symptom risk, although possible reverse causality could not be ruled out.

Cao Y, Xu Q, Lu MX, Jin ZD, DU HB, Li JX, Nie JH, Tong J. [Antagonistic effect of microwave on hematopoietic damage of mice induced by gamma-ray irradiation.] Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi. 27(9):525-529, 2009. [Article in Chinese]

OBJECTIVE: To investigate antagonistic effect of microwave on hematopoietic damage of mice induced by gamma-ray irradiation. METHODS: Male healthy Kunming mice were treated with low dose microwave radiation before exposure to (60)Co gamma-ray irradiation of 8.0 Gy. The 30-day survival rate and average survival time of the mice after the treatment were examined. Peripheral blood parameters and the organ indexes of thymus and spleen were also observed in the irradiated mice. After exposure to 5.0 Gy gamma irradiation, indexes of hematopoietic foci formation of bone marrow cells (CFU-GM) and the proliferation activity of BMNCs were examined. The serum concentration of hemopoietic factors (GM-CSF and IL-3) were detected by ELISA kits. RESULTS: Pre-exposure with 120 microW/cm(2) 900 MHz microwave increased the 30-day survival rate ($P < 0.05$) and the number of white blood cells of gamma-ray treated mice. The increases of the organ indexes of thymus and spleen, proliferation activity of BMNCs and CFU-GM hematopoietic foci numbers, as well as the higher serum concentration of GM-CSF and IL-3 were observed in the microwave pre-exposure group. CONCLUSION: Low dose microwave radiation may exert potential antagonistic effects on hematopoietic injuries induced by ionizing radiation. The underlying mechanisms might be related with stimulation of hematopoietic growth factors expression, promotion of HSCs/HPCs proliferation, suppression on the reduction of HSCs/HPCs caused by (60)Co gamma-ray, and enhanced construction of the hematopoietic system.

Cao Y, Zhang W, Lu MX, Xu Q, Meng QQ, Nie JH, Tong J. 900-MHz microwave radiation enhances gamma-ray adverse effects on SHG44 cells. J Toxicol Environ Health A. 72(11):727-732, 2009.

Mobile phones are widely used globally. However, the biological effects due to exposure to electromagnetic fields (EMF) produced by mobile phones are largely unknown. Environmental and occupational exposure of humans to gamma-rays is a biologically relevant phenomenon. Consequently studies were undertaken to examine the interactions between gamma-rays and EMF on human health. In this study, exposure to

Studies that show **Cell Phone** Health Effects

900-MHz EMF expanded gamma-ray damage to SHG44 cells. Preexposure EMF enhanced the decrease in cell proliferation induced by gamma-ray irradiation and the rate of apoptosis. The combination of EMF and gamma-ray exposure resulted in a synergistic effect by triggering stress response, which increased reactive oxygen species, but the expression of hsp70 at both mRNA and protein levels remained unaltered. Data indicate that the adverse effects of gamma-rays on cellular functions are strengthened by EMF.

Bodera P, Stankiewicz W, Antkowiak B, Paluch M, Kieliszek J, Sobiech J, Zdanowski R, Wojdas A, Siwicki AK, Skopińska-Różewska E. Suppressive effect of electromagnetic field on analgesic activity of tramadol in rats. Pol J Vet Sci. 15(1):95-100, 2012.

The electromagnetic fields (EMFs) have been shown to alter animal and human behavior, such as directional orientation, learning, pain perception (nociception or analgesia) and anxiety-related behaviors. The aim of this study was to evaluate the influence of electromagnetic fields of high-frequency microwaves on pain perception and anti-nociceptive activity of tramadol (TRAM) - analgetic effective in the treatment of moderate to severe acute and chronic pain states. Electromagnetic fields exposures of a) 1500 MHz frequency and b) modulated, 1800 MHz (which is identical to that generated by mobile phones) were applied. Paw withdrawal latency (PWL) to thermal stimulus was measured in vehicle or tramadol (TRAM) treated animals before and after 30, 60 and 90 minutes from injections. The differences in the level of pain (PWL) between control group and rats exposed to EMF alone in three measurements, were not observed. Tramadol alone significantly increased PWLs to thermal stimulus in comparison to vehicle results at 30 ($p < 0.001$) and 60 minutes ($p < 0.05$) after drug injection. EMF exposure of both frequencies transiently suppressed analgesic effect of tramadol, significantly reducing paw withdrawal latency in animals treated with this drug at 30 minutes from the drug injection.

Cao Y, Xu Q, Lu MX, Jin ZD, DU HB, Li JX, Nie JH, Tong J. [Antagonistic effect of microwave on hematopoietic damage of mice induced by gamma-ray irradiation.] Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi. 27(9):525-529, 2009. [Article in Chinese]

OBJECTIVE: To investigate antagonistic effect of microwave on hematopoietic damage of mice induced by gamma-ray irradiation. METHODS: Male healthy Kunming mice were treated with low dose microwave radiation before exposure to (60)Co gamma-ray irradiation of 8.0 Gy. The 30-day survival rate and average survival time of the mice after the treatment were examined. Peripheral blood parameters and the organ indexes of thymus and spleen were also observed in the irradiated mice. After exposure to 5.0 Gy gamma irradiation, indexes of hematopoietic foci formation of bone marrow cells (CFU-GM) and the proliferation activity of BMNCs were examined. The serum concentration of hemopoietic factors (GM-CSF and IL-3) were detected by ELISA kits. RESULTS: Pre-exposure with 120 microW/cm(2) 900 MHz microwave increased the 30-day survival rate ($P < 0.05$) and the number of white blood cells of gamma-ray treated mice. The increases of the organ indexes of thymus and spleen, proliferation activity of BMNCs

Studies that show **Cell Phone** Health Effects

and CFU-GM hematopoietic foci numbers, as well as the higher serum concentration of GM-CSF and IL-3 were observed in the microwave pre-exposure group. **CONCLUSION:** Low dose microwave radiation may exert potential antagonistic effects on hematopoietic injuries induced by ionizing radiation. The underlying mechanisms might be related with stimulation of hematopoietic growth factors expression, promotion of HSCs/HPCs proliferation, suppression on the reduction of HSCs/HPCs caused by (60)Co gamma-ray, and enhanced construction of the hematopoietic system.

Inflammation

Li CY, Liao MH, Lin CW, Tsai WS, Huang CC, Tang TK. Inhibitory Effects of Microwave Radiation on LPS-Induced NFκB Expression in THP-1 Monocytes. Chin J Physiol. 55(6):421-427, 2012.

Microwave radiations can be encountered regularly in daily lives. When WHO announced that microwave radiations were a kind of environmental energy which interfere with the physiological functions of the human body, great concerns have been raised over the damages microwave frequencies can do to human physiology. The immunological performance and the activities of the cellular inflammatory factor NFκB have been closely related in monocyte. Due to the effect of phorbol 12-myristate 13-acetate (PMA) on THP-1 monocytes, THP-1 monocytes would differentiate into macrophages and would then react with lipopolysaccharides (LPS), and the amount of NFκB increased in the THP-1 monocytes. Expression of cytokine is affected when cells are exposed to a frequency of 2450 MHz and at 900 W. Thus, in our experiments, an observation was made when THP-1 monocytes were stimulated with PMA and LPS to differentiate into macrophage, the amount of NFκB in cells increased exponentially, and the levels of NFκB expression were decreased by the exposure of microwave radiation. In conclusion, microwave radiations were found to inhibit the activity functions of THP-1 monocytes stimulated with PMA and LPS.

Dabrowski MP, Stankiewicz W, Kubacki R, Sobiczewska E, Szmigielski S. Immunotropic Effects in Cultured Human Blood Mononuclear Cells Pre-exposed to Low-Level 1300 MHz Pulse-Modulated Microwave Field. Electromag. Biol. Med. 22:1-13, 2003.

The samples of mononuclear cells isolated from peripheral blood of healthy donors (N = 16) were exposed to 1300 MHz pulse-modulated microwaves at 330 pps with 5 μs pulse width. The samples were exposed in an anechoic chamber at the average value of power density of $S = 10 \text{ W/m}^2$ (1 mW/cm^2). The average specific absorption rate (SAR) was measured in rectangular waveguide and the value of SAR = 0.18 W/kg was recorded. Subsequently, the exposed and control cells were assessed in the microculture system for several parameters characterizing their proliferative and immunoregulatory properties. Although the irradiation decreased the spontaneous incorporation of 3H-thymidine, the proliferative response of lymphocytes to phytohemagglutinin (PHA) and to Con A as well as the T-cell suppressive activity (SAT index) and the saturation of IL-2 receptors did not change. Nevertheless, the

Studies that show Cell Phone Health Effects

lymphocyte production of interleukin (IL)-10 increased ($P < .001$) and the concentration of IFN γ remained unchanged or slightly decreased in the culture supernatants. Concomitantly, the microwave irradiation modulated the monokine production by monocytes. The production of IL-1 β increased significantly ($P < .01$), the concentration of its antagonist (IL-1ra) dropped by half ($P < .01$) and the tumor necrosis factor (TNF- α) concentration remained unchanged. These changes of monokine proportion (IL-1 β vs. IL-1ra) resulted in significant increase of the value of LM index ($P < .01$), which reflects the activation of monocyte immunogenic function. The results indicate that pulse-modulated microwaves represent the potential of immunotropic influence, stimulating preferentially the immunogenic and proinflammatory activity of monocytes at relatively low levels of exposure.

Zhou ZD, Zeng QL, Zheng Y, Zhang JB, Chen HY, Lu DQ, Shao CS, Xia DJ. [Surface markers and functions of human dendritic cells exposed to mobile phone 1800 MHz electromagnetic fields.] *Zhejiang Da Xue Xue Bao Yi Xue Ban.* 37(1):29-33, 2008. [Article in Chinese]

OBJECTIVE: To investigate the effects of mobile phone 1800 MHz electromagnetic fields (EMF) on the surface markers and the functions of human dendritic cells (DC).

METHODS: Human DCs were exposed to intermittent 5 min on/10 min off EMF with specific absorption rates (SAR) 4 W/kg for 0 h, 1 h, 12 h or 24 h, respectively. FACS analysis was used to detect the positive percentage of DC surface markers including HLA-DR and co-stimulatory molecules such as CD80, CD86, CD40 and CD11c. CCK-8 kit was adopted to examine the function of allo-mixed lymphocyte reaction (allo-MLR) of DC, and enzyme linked immunosorbent assay (ELISA) to identify the levels of IL-12p70 and TNF-alpha secreted by DC. RESULT: Compared with the sham radiation group, after exposure to the electromagnetic fields for 1 h, 12 h, or 24 h, HLA-DR, CD80, CD86 and CD40 were all declined except CD11c. The ability of DC allo-MLR in each exposure group was decreased significantly ($P < 0.05$), especially in the 24 h exposure group. However, the secreted levels of IL-12p70 and TNF-alpha of DC in each exposure group remained no changed. Conclusion: The study showed that EMF exposure could down-regulate the surface molecules and stimulation ability of human DC.

Effects on Behavior

Shtemberg AS, Uzbekov MG, Shikhov SN, Bazian AS, Cherniakov GM, [Species specificity, age factors, and various neurochemical correlates of the animal spontaneous behavior after exposure to electromagnetic field of the ultralow intensity]. *Zh Vyssh Nerv Deiat Im I P Pavlova* 50(4):703-715, 2000. [Article in Russian] Behavioral and neurochemical reactions of small laboratory animals (mice and rats of different age) under exposure to ultralow-intensity electromagnetic fields (EMF, frequency of 4200 and 970 MHz, modulated by a quasistochastic signal in the range of 20-20,000 Hz, power density 15 microW/cm², specific body absorption rate up to 4.5

Studies that show **Cell Phone** Health Effects

mJ/kg) were studied. The EMF basically inhibited the locomotor and exploratory activity in the "open-field" test. The species- and age-specific features rather than radiation conditions dominated. However, decrease in the EMF frequency considerably intensified the observed effect. Change in animal behavior was accompanied by shifts in neurochemical processes, i.e., sharp activation of serotonergic and inhibition of morepinephrinergic system.

Narayanan SN, Kumar RS, Pavai J, Kedage V, Bhat MS, Nayak S, Bhat PG. Analysis of emotionality and locomotion in radio-frequency electromagnetic radiation exposed rats. *Neurol Sci.* 34(7):1117-1124, 2013.

In the current study the modulatory role of mobile phone radio-frequency electromagnetic radiation (RF-EMR) on emotionality and locomotion was evaluated in adolescent rats. Male albino Wistar rats (6-8 weeks old) were randomly assigned into the following groups having 12 animals in each group. Group I (Control): they remained in the home cage throughout the experimental period. Group II (Sham exposed): they were exposed to mobile phone in switch-off mode for 28 days, and Group III (RF-EMR exposed): they were exposed to RF-EMR (900 MHz) from an active GSM (Global system for mobile communications) mobile phone with a peak power density of 146.60 $\mu\text{W}/\text{cm}^2$ for 28 days. On 29th day, the animals were tested for emotionality and locomotion. Elevated plus maze (EPM) test revealed that, percentage of entries into the open arm, percentage of time spent on the open arm and distance travelled on the open arm were significantly reduced in the RF-EMR exposed rats. Rearing frequency and grooming frequency were also decreased in the RF-EMR exposed rats. Defecation boli count during the EPM test was more with the RF-EMR group. No statistically significant difference was found in total distance travelled, total arm entries, percentage of closed arm entries and parallelism index in the RF-EMR exposed rats compared to controls. Results indicate that mobile phone radiation could affect the emotionality of rats without affecting the general locomotion.

Kumar RS, Sareesh NN, Nayak S, Mailankot M. Hypoactivity of Wistar rats exposed to mobile phone on elevated plus maze. *Indian J Physiol Pharmacol.* 53(3):283-286, 2009.

No abstract available. From discussion section: "In conclusion, our preliminary results indicate mobile phone exposure induced behavioral changes in rats, expressed as deficit in open arm exploration on elevated plus-maze."

Sokolovic D, Djordjevic B, Kocic G, Babovic P, Ristic G, Stanojkovic Z, Sokolovic DM, Veljkovic A, Jankovic A, Radovanovic Z. The effect of melatonin on body mass and behaviour of rats during an exposure to microwave radiation from mobile phone. *Bratisl Lek Listy.* 113(5):265-269, 2012.

BACKGROUND: Microwave radiation (MW) produced by wireless telecommunications and a number of electrical devices used in household or in healthcare institutions may cause various disorders in human organism. On the other hand, melatonin is a potent antioxidant, immunostimulator and neuromodulator. The aim of this research was to

Studies that show Cell Phone Health Effects

determine body mass and behaviour changes in rats after a chronic microwave exposure, as well as to determine the effects of melatonin on body mass and behaviour in irradiated rats. METHODS: Wistar rats were divided into the four experimental groups: I group (control) - rats treated with 0,9 % saline, II group (Mel) - rats treated with melatonin (2 mg/kg), III group (MW) - rats exposed to MW radiation (4 h/day), IV group (MW+Mel) - rats, which were both exposed to MW radiation and received melatonin premedication (2 mg/kg). RESULTS: A significant body mass reduction was noted in animals exposed to MW radiation when compared to controls after 20, 40 and 60 days ($p < 0.001$). Furthermore, body weight was significantly increased ($p < 0.05$) in irradiated rats, which received melatonin pretreatment (MW+Mel) in comparison to irradiated group (MW) after 20 days. Microwave radiation exposed animals showed an anxiety related behaviour (agitation, irritability) after 10 days of exposure. After the radiation source removal, changes in behaviour were less noticeable. Melatonin administration to irradiated rats caused a decrease in the stress induced behaviour. CONCLUSION: Microwave radiation causes body mass decrease and anxiety related behaviour in rats, however melatonin causes a reverse of those effects on both body weight and behaviour of irradiated animals (Fig. 2, Ref. 32).

Júnior LC, Guimarães ED, Musso CM, Stabler CT, Garcia RM, Mourão-Júnior CA, Andreazzi AE. Behavior and memory evaluation of Wistar rats exposed to 1.8 GHz radiofrequency electromagnetic radiation. Neurol Res. 2014 Jan 27;1743132813Y0000000276. [Epub ahead of print]

Background: The development of communication systems has brought great social and economic benefits to society. As mobile phone use has become widespread, concerns have emerged regarding the potential adverse effects of radiofrequency electromagnetic radiation (RF-EMR) used by these devices. Objective: To verify potential effects of mobile phone radiation on the central nervous system (CNS) in an animal model. Methods: Male Wistar rats (60 days old) were exposed to RF-EMR from a Global System for Mobile (GSM) cell phone (1.8 GHz) for 3 days. At the end of the exposure, the following behavioral tests were performed: open field and object recognition. Results: Our results showed that exposed animals did not present anxiety patterns or working memory impairment, but stress behavior actions were observe. Conclusion: Given the results of the present study, we speculate that RF-EMR does not promote CNS impairment, but suggest that it may lead to stressful behavioral patterns.

Divan HA, Kheifets L, Obel C, Olsen J. Prenatal and postnatal exposure to cell phone use and behavioral problems in children. Epidemiology.19(4):523-529, 2008.

BACKGROUND: The World Health Organization has emphasized the need for research into the possible effects of radiofrequency fields in children. We examined the association between prenatal and postnatal exposure to cell phones and behavioral problems in young children. METHODS: Mothers were recruited to the Danish National Birth Cohort early in pregnancy. When the children of those pregnancies reached 7 years of age in 2005 and 2006, mothers were asked to complete a questionnaire regarding the current health and behavioral status of children, as well as past exposure

Studies that show **Cell Phone** Health Effects

to cell phone use. Mothers evaluated the child's behavior problems using the Strength and Difficulties Questionnaire. RESULTS:: Mothers of 13,159 children completed the follow-up questionnaire reporting their use of cell phones during pregnancy as well as current cell phone use by the child. Greater odds ratios for behavioral problems were observed for children who had possible prenatal or postnatal exposure to cell phone use. After adjustment for potential confounders, the odds ratio for a higher overall behavioral problems score was 1.80 (95% confidence interval = 1.45-2.23) in children with both prenatal and postnatal exposure to cell phones. CONCLUSIONS: Exposure to cell phones prenatally-and, to a lesser degree, postnatally-was associated with behavioral difficulties such as emotional and hyperactivity problems around the age of school entry. These associations may be noncausal and may be due to unmeasured confounding. If real, they would be of public health concern given the widespread use of this technology.

Divan HA, Kheifets L, Obel C, Olsen J. Cell phone use and behavioural problems in young children. J Epidemiol Community Health.66(6):524-529, 2012.

Background: Potential health effects of cell phone use in children have not been adequately examined. As children are using cell phones at earlier ages, research among this group has been identified as the highest priority by both national and international organisations. The authors previously reported results from the Danish National Birth Cohort (DNBC), which looked at prenatal and postnatal exposure to cell phone use and behavioural problems at age 7years. Exposure to cell phones prenatally, and to a lesser degree postnatally, was associated with more behavioural difficulties. The original analysis included nearly 13 000 children who reached age 7 years by November 2006. Methods: To see if a larger, separate group of DNBC children would produce similar results after considering additional confounders, children of mothers who might better represent current users of cell phones were analysed. This 'new' dataset consisted of 28 745 children with completed Age-7 Questionnaires to December 2008. Results: The highest OR for behavioural problems were for children who had both prenatal and postnatal exposure to cell phones compared with children not exposed during either time period. The adjusted effect estimate was 1.5 (95% CI 1.4 to 1.7). Conclusions: The findings of the previous publication were replicated in this separate group of participants demonstrating that cell phone use was associated with behavioural problems at age 7years in children, and this association was not limited to early users of the technology. Although weaker in the new dataset, even with further control for an extended set of potential confounders, the associations remained.

Daniels WM, Pitout IL, Afullo TJ, Mabandla MV. The effect of electromagnetic radiation in the mobile phone range on the behaviour of the rat. Metab Brain Dis. 24(4):629-641, 2009

Electromagnetic radiation (EMR) is emitted from electromagnetic fields that surround power lines, household appliances and mobile phones. Research has shown that there are connections between EMR exposure and cancer and also that exposure to EMR may result in structural damage to neurons. In a study by Salford et al. (Environ Health

Studies that show **Cell Phone** Health Effects

Perspect 111:881-883, 2003) the authors demonstrated the presence of strongly stained areas in the brains of rats that were exposed to mobile phone EMR. These darker neurons were particularly prevalent in the hippocampal area of the brain. The aim of our study was to further investigate the effects of EMR. Since the hippocampus is involved in learning and memory and emotional states, we hypothesised that EMR will have a negative impact on the subject's mood and ability to learn. We subsequently performed behavioural, histological and biochemical tests on exposed and unexposed male and female rats to determine the effects of EMR on learning and memory, emotional states and corticosterone levels. We found no significant differences in the spatial memory test, and morphological assessment of the brain also yielded non-significant differences between the groups. However, in some exposed animals there were decreased locomotor activity, increased grooming and a tendency of increased basal corticosterone levels. These findings suggested that EMR exposure may lead to abnormal brain functioning.

Razavinasab M, Moazzami K, Shabani M. Maternal mobile phone exposure alters intrinsic electrophysiological properties of CA1 pyramidal neurons in rat offspring. Toxicol Ind Health. 2014 Mar 6. [Epub ahead of print]

Some studies have shown that exposure to electromagnetic field (EMF) may result in structural damage to neurons. In this study, we have elucidated the alteration in the hippocampal function of offspring Wistar rats (n = 8 rats in each group) that were chronically exposed to mobile phones during their gestational period by applying behavioral, histological, and electrophysiological tests. Rats in the EMF group were exposed to 900 MHz pulsed-EMF irradiation for 6 h/day. Whole cell recordings in hippocampal pyramidal cells in the mobile phone groups did show a decrease in neuronal excitability. Mobile phone exposure was mostly associated with a decrease in the number of action potentials fired in spontaneous activity and in response to current injection in both male and female groups. There was an increase in the amplitude of the afterhyperpolarization (AHP) in mobile phone rats compared with the control. The results of the passive avoidance and Morris water maze assessment of learning and memory performance showed that phone exposure significantly altered learning acquisition and memory retention in male and female rats compared with the control rats. Light microscopy study of brain sections of the control and mobile phone-exposed rats showed normal morphology. Our results suggest that exposure to mobile phones adversely affects the cognitive performance of both female and male offspring rats using behavioral and electrophysiological techniques.

Valentini E, Ferrara M, Presaghi F, De Gennaro L, Curcio G. Republished review: systematic review and meta-analysis of psychomotor effects of mobile phone electromagnetic fields. Postgrad Med J. 87(1031):643-651, 2011.

OBJECTIVES Over the past 10 years there has been increasing concern about the possible behavioural effects of mobile phone use. This systematic review and meta-analysis focuses on studies published since 1999 on the human cognitive and performance effects of mobile phone-related electromagnetic fields (EMF). **METHODS**

Studies that show **Cell Phone** Health Effects

PubMed, Biomed, Medline, Biological Sciences, PsychInfo, PsycARTICLES, Environmental Sciences and Pollution Management, Neurosciences Abstracts and Web of Science professional databases were searched and 24 studies selected for meta-analysis. Each study had to have at least one psychomotor measurement result as a main outcome. Data were analysed using standardised mean difference (SMD) as the effect size measure. RESULTS Only three tasks (2-back, 3-back and simple reaction time (SRT)) displayed significant heterogeneity, but after studies with extreme SMD were excluded using sensitivity analysis, the statistical significance disappeared ($\chi^2(7)=1.63$, $p=0.20$; $\chi^2(6)=1.00$, $p=0.32$; $\chi^2(10)=14.04$, $p=0.17$, respectively). Following sensitivity analysis, the effect of sponsorship and publication bias were assessed. Meta-regression indicated a significant effect ($b1/40.12$, $p<0.05$) only for the 2-back task with mixed funding (industry and public/charity). Funnel plot inspection revealed a significant publication bias only for two cognitive tasks: SRT (Begg's rank correlation $r=0.443$; Egger's test $b=-0.652$) and the subtraction task (Egger's test $b=-0.687$). CONCLUSIONS Mobile phone-like EMF do not seem to induce cognitive and psychomotor effects. Nonetheless, the existence of sponsorship and publication biases should encourage WHO intervention to develop official research standards and guidelines. In addition, future research should address critical and neglected issues such as investigation of repeated, intensive and chronic exposures, especially in highly sensitive populations such as children.

Electromagnetic Hypersensitivity

Nordin S, Neely G, Olsson D, Sandström M. Odor and Noise Intolerance in Persons with Self-Reported Electromagnetic Hypersensitivity. Int J Environ Res Public Health. 11(9):8794-8805, 2014.

Lack of confirmation of symptoms attributed to electromagnetic fields (EMF) and triggered by EMF exposure has highlighted the role of individual factors. Prior observations indicate intolerance to other types of environmental exposures among persons with electromagnetic hypersensitivity (EHS). This study assessed differences in odor and noise intolerance between persons with EHS and healthy controls by use of subscales and global measures of the Chemical Sensitivity Scale (CSS) and the Noise Sensitivity Scale (NSS). The EHS group scored significantly higher than the controls on all CSS and NSS scales. Correlation coefficients between CSS and NSS scores ranged from 0.60 to 0.65 across measures. The findings suggest an association between EHS and odor and noise intolerance, encouraging further investigation of individual factors for understanding EMF-related symptoms.

Wilen J, Johansson A, Kalezic N, Lyskov E, Sandstrom M. Psychophysiological tests and provocation of subjects with mobile phone related symptoms. Bioelectromagnetics. 27(3):204-214, 2006.

The aim of the present study was to investigate the effect of exposure to a mobile phone-like radiofrequency (RF) electromagnetic field on persons experiencing subjective symptoms when using mobile phones (MP). Twenty subjects with MP-related symptoms

Studies that show **Cell Phone** Health Effects

were recruited and matched with 20 controls without MP-related symptoms. Each subject participated in two experimental sessions, one with true exposure and one with sham exposure, in random order. In the true exposure condition, the test subjects were exposed for 30 min to an RF field generating a maximum SAR(1g) in the head of 1 W/kg through an indoor base station antenna attached to a 900 MHz GSM MP. The following physiological and cognitive parameters were measured during the experiment: heart rate and heart rate variability (HRV), respiration, local blood flow, electrodermal activity, critical flicker fusion threshold (CFFT), short-term memory, and reaction time. No significant differences related to RF exposure conditions were detected. Also no differences in baseline data were found between subject groups, except for the reaction time, which was significantly longer among the cases than among the controls the first time the test was performed. This difference disappeared when the test was repeated. However, the cases differed significantly from the controls with respect to HRV as measured in the frequency domain. The cases displayed a shift in low/high frequency ratio towards a sympathetic dominance in the autonomous nervous system during the CFFT and memory tests, regardless of exposure condition. This might be interpreted as a sign of differences in the autonomous nervous system regulation between persons with MP related subjective symptoms and persons with no such symptoms.

Kato Y, Johansson O. Reported functional impairments of electrohypersensitive Japanese: A questionnaire survey. *Pathophysiology*.19(2) 95-100, 2012.

An increasing number of people worldwide complain that they have become electromagnetic hypersensitive (EHS). We conducted a questionnaire survey of EHS persons in Japan. The aim was to identify electromagnetic fields (EMF) and plausible EMF sources that caused their symptoms. Postal questionnaires were distributed via a self-help group, and 75 participants (95% women) responded. Reported major complaints were "fatigue/tiredness" (85%), "headache", "concentration, memory, and thinking" difficulty (81%, respectively). Seventy-two per cent used some form of complementary/alternative therapy. The most plausible trigger of EHS onset was a mobile phone base station or personal handy-phone system (37%). Sixty-five percent experienced health problems to be due to the radiation from other passengers' mobile phones in trains or buses, and 12% reported that they could not use public transportation at all. Fifty-three percent had a job before the onset, but most had lost their work and/or experienced a decrease in income. Moreover, 85.3% had to take measures to protect themselves from EMF, such as moving to low EMF areas, or buying low EMF electric appliances. EHS persons were suffering not only from their symptoms, but also from economical and social problems.

Johansson A, Nordin S, Heiden M, Sandström M. Symptoms, personality traits, and stress in people with mobile phone-related symptoms and electromagnetic hypersensitivity. *J Psychosom Res*. 68(1):37-45, 2010.

OBJECTIVE: Some people report symptoms that they associate with electromagnetic field (EMF) exposure. These symptoms may be related to specific EMF sources or to electrical equipment in general (perceived electromagnetic hypersensitivity, EHS).

Studies that show **Cell Phone** Health Effects

Research and clinical observations suggest a difference between mobile phone (MP)-related symptoms and EHS with respect to symptom prevalence, psychological factors, and health prognosis. This study assessed prevalence of EMF-related and EMF-nonrelated symptoms, anxiety, depression, somatization, exhaustion, and stress in people with MP-related symptoms or EHS versus a population-based sample and a control sample without EMF-related symptoms. **METHODS:** Forty-five participants with MP-related symptoms and 71 with EHS were compared with a population-based sample (n=106) and a control group (n=63) using self-report questionnaires. **RESULTS:** The EHS group reported more symptoms than the MP group, both EMF-related and EMF-nonrelated. The MP group reported a high prevalence of somatosensory symptoms, whereas the EHS group reported more neurasthenic symptoms. As to self-reported personality traits and stress, the case groups differed only on somatization and listlessness in a direct comparison. In comparison with the reference groups, the MP group showed increased levels of exhaustion and depression but not of anxiety, somatization, and stress; the EHS group showed increased levels for all of the conditions except for stress. **CONCLUSION:** The findings support the idea of a difference between people with symptoms related to specific EMF sources and people with general EHS with respect to symptoms and anxiety, depression, somatization, exhaustion, and stress. The differences are likely to be important in the management of patients.

Hagström M, Auranen J, Ekman R. Electromagnetic hypersensitive Finns: Symptoms, perceived sources and treatments, a questionnaire study. Pathophysiology. 2013 Apr 1. pii: S0928-4680(13)00002-3.

The aim was to analyze the subjective experiences of Finns who describe themselves as suffering from electromagnetic hypersensitivity (EHS), their symptoms, self-perceived sources of the health complaints and the effectiveness of medical and complementary alternative therapies. A total of 395 questionnaires were mailed to self-diagnosed EHS persons. Of the participants 345 belonged to a Finnish self-help group and 50 came from outside of the group. The return rate of the study was 52.1% (206) and 80.9% of the respondents were women. Before the onset of EHS the most common health complaints were different types of allergies (35.1%, 68). During the acute phase of EHS the most common symptoms were nervous system related: "stress" (60.3%, 117), "sleeping disorders" (59.3%, 115) and "fatigue" (57.2%, 111). The sources that were most often reported to have triggered EHS were: "personal computers" (50.8%, 94) and "mobile phones" (47.0%, 87). The same devices were also claimed to cause the most symptoms during the acute phase. After the acute phase of EHS had passed, the respondents still claimed to react to these same digital and wireless devices while their reactions to basic electrical appliances were reduced. According to 76% of 157 respondents the reduction or avoidance of electromagnetic fields (EMF) helped in their full or partial recovery. The best treatments for EHS were given as: "dietary change" (69.4%), "nutritional supplements" (67.8%) and "increased physical exercise" (61.6%). The official treatment recommendations of psychotherapy (2.6%) and medication (-4.2%) were not significantly helpful. According to the present results the official treatment protocols should take better account the EHS person's own experiences. The

Studies that show **Cell Phone** Health Effects

avoidance of electromagnetic radiation and fields effectively removed or lessened the symptoms in EHS persons.

De Luca C, Chung Sheun Thai J, Raskovic D, Cesareo E, Caccamo D, Trukhanov A, Korkina L. Metabolic and genetic screening of electromagnetic hypersensitive subjects as a feasible tool for diagnostics and intervention. Mediators Inflamm. 2014;2014:924184. doi: 10.1155/2014/924184. Epub 2014 Apr 9.

Growing numbers of "electromagnetic hypersensitive" (EHS) people worldwide self-report severely disabling, multiorgan, non-specific symptoms when exposed to low-dose electromagnetic radiations, often associated with symptoms of multiple chemical sensitivity (MCS) and/or other environmental "sensitivity-related illnesses" (SRI). This cluster of chronic inflammatory disorders still lacks validated pathogenetic mechanism, diagnostic biomarkers, and management guidelines. We hypothesized that SRI, not being merely psychogenic, may share organic determinants of impaired detoxification of common physic-chemical stressors. Based on our previous MCS studies, we tested a panel of 12 metabolic blood redox-related parameters and of selected drug-metabolizing-enzyme gene polymorphisms, on 153 EHS, 147 MCS, and 132 control Italians, confirming MCS altered ($P < 0.05$ - 0.0001) glutathione-(GSH), GSH-peroxidase/S-transferase, and catalase erythrocyte activities. We first described comparable-though milder-metabolic pro-oxidant/proinflammatory alterations in EHS with distinctively increased plasma coenzyme-Q10 oxidation ratio. Severe depletion of erythrocyte membrane polyunsaturated fatty acids with increased $\omega 6/\omega 3$ ratio was confirmed in MCS, but not in EHS. We also identified significantly ($P = 0.003$) altered distribution-versus-control of the CYP2C19*1/*2 SNP variants in EHS, and a 9.7-fold increased risk (OR: 95% C.I. = 1.3-74.5) of developing EHS for the haplotype (null)GSTT1 + (null)GSTM1 variants. Altogether, results on MCS and EHS strengthen our proposal to adopt this blood metabolic/genetic biomarkers' panel as suitable diagnostic tool for SRI.

Effects on Blood

Mousavy SJ, Riazi GH, Kamarei M, Aliakbarian H, Sattarahmady N, Sharifizadeh A, Safarian S, Ahmad F, Moosavi-Movahedi AA. Effects of mobile phone radiofrequency on the structure and function of the normal human hemoglobin. Int J Biol Macromol. 44(3):278-285, 2009

Widespread use of mobile phones has increased the human exposure to electromagnetic fields (EMFs). It is required to investigate the effect of EMFs on the biological systems. In this paper the effect of mobile phone RF (910MHz and 940 MHz) on structure and function of HbA was investigated. Oxygen affinity was measured by sodium dithionite with UV-vis spectrophotometer. Structural changes were studied by circular dichroism and fluorescence spectroscopy. The results indicated that mobile phone EMFs altered oxygen affinity and tertiary structure of HbA. Furthermore, the decrease of oxygen affinity of HbA corresponded to the EMFs intensity and time of

Studies that show **Cell Phone** Health Effects

exposure.

Shahbazi-Gahrouei D, Mortazavi SM, Nasri H, Baradaran A, Baradaran-Ghahfarokhi M, Baradaran-Ghahfarokhi HR. Mobile phone radiation interferes laboratory immunoassays: Example chorionic gonadotropin assays. Pathophysiology. 19(1):43-47, 2012.

The radiofrequency radiation is of concern in hospital laboratories as the microwaves have many health effects even on immune functions. The aim of this study was, however, to evaluate the effects of cell phone radiation on chorionic gonadotropin immunoassays of human serum. Two cell phones with 0.69 and 1.09W/kg (head SAR) emitting 900MHz radiation were used. Sixty wells with five human serum concentrations (0, 10, 100, 250, 500mIU/mL) were used in three batches. The well heads in each batch were exposed to 900MHz emitted from these phones, and the 0.69, 1.09W/kg exposed batches were compared with the unexposed controls. Radiation exposure from mobile phones altered the measured serum levels especially in the wells with 100, 250, 500mIU/mL hormone concentrations. Exposure at 1.09W/kg SAR caused a significant loss compared to 0.69W/kg SAR exposure. In conclusion, the microwave exposures may require attention in laboratories using immunoassays.

Ruan P, Yong J, Shen H, Zheng X. Monitoring dynamic reactions of red blood cells to UHF electromagnetic waves radiation using a novel micro-imaging technology. Electromagn Biol Med. 31(4):365-374, 2012.

Multiple state-of-the-art techniques, such as multi-dimensional micro-imaging, fast multi-channel micro-spectrophotometry, and dynamic micro-imaging analysis, were used to dynamically investigate various effects of cell under the 900 MHz electromagnetic radiation. Cell changes in shape, size, and parameters of Hb absorption spectrum under different power density electromagnetic waves radiation were presented in this article. Experimental results indicated that the isolated human red blood cells (RBCs) do not have obviously real-time responses to the ultra-low density (15 $\mu\text{W}/\text{cm}^2$, 31 $\mu\text{W}/\text{cm}^2$) electromagnetic wave radiation when the radiation time is not more than 30 min; however, the cells do have significant reactions in shape, size, and the like, to the electromagnetic waves radiation with power densities of 1 mW/cm^2 and 5 mW/cm^2 . The data also reveal the possible influences and statistical relationships among living human cell functions, radiation amount, and exposure time with high-frequency electromagnetic waves. The results of this study may be significant on protection of human being and other living organisms against possible radiation affections of the high-frequency electromagnetic waves.

Zotti-Martelli L, Peccatori M, Maggini V, Ballardini M, Barale R. Individual responsiveness to induction of micronuclei in human lymphocytes after exposure in vitro to 1800-MHz microwave radiation. Mutat Res. 582(1-2):42-52, 2005.

The widespread application of microwaves is of great concern in view of possible consequences for human health. Many in vitro studies have been carried out to detect possible effects on DNA and chromatin structure following exposure to microwave

Studies that show **Cell Phone** Health Effects

radiation. The aim of this study is to assess the capability of microwaves, at different power densities and exposure times, to induce genotoxic effects as evaluated by the in vitro **m micronucleus (MN)** assay on peripheral blood lymphocytes from nine different healthy donors, and to investigate also the possible inter-individual response variability. Whole blood samples were exposed for 60, 120 and 180min to continuous microwave radiation with a frequency of 1800MHz and power densities of 5, 10 and 20mW/cm². Reproducibility was tested by repeating the experiment 3 months later. Multivariate analysis showed that lymphocyte proliferation indices were significantly different among donors ($p < 0.004$) and between experiments ($p < 0.01$), whereas the applied power density and the exposure time did not have any effect on them. Both spontaneous and induced MN frequencies varied in a highly significant way among donors ($p < 0.009$) and between experiments ($p < 0.002$), and a statistically significant increase of MN, although rather low, was observed dependent on exposure time ($p = 0.0004$) and applied power density ($p = 0.0166$). A considerable decrease in spontaneous and induced MN frequencies was measured in the second experiment. The results show that microwaves are able to induce MN in short-time exposures to medium power density fields. Our data analysis highlights a wide inter-individual variability in the response, which was confirmed to be a characteristic reproducible trait by means of the second experiment.

Zmyslony M, Politanski P, Rajkowska E, Szymczak W, Jajte J. Acute exposure to 930 MHz CW electromagnetic radiation in vitro affects reactive oxygen species level in rat lymphocytes treated by iron ions. Bioelectromagnetics. 25(5):324-328, 2004.

The aim of this study was to test the hypothesis that the 930 MHz continuous wave (CW) electromagnetic field, which is the carrier of signals emitted by cellular phones, affects the reactive oxygen species (ROS) level in living cells. Rat lymphocytes were used in the experiments. A portion of the lymphocytes was treated with iron ions to induce oxidative processes. Exposures to electromagnetic radiation (power density 5 W/m², theoretical calculated SAR = 1.5 W/kg) were performed within a GTEM cell. Intracellular ROS were measured by the fluorescent probe dichlorofluorescein diacetate (DCF-DA). The results show that acute (5 and 15 min) exposure does not affect the number of produced ROS. If, however, FeCl₂ with final concentration 10 microg/ml was added to the lymphocyte suspensions to stimulate ROS production, after both durations of exposure, the magnitude of fluorescence (ROS level during the experiment) was significantly greater in the exposed lymphocytes. The character of the changes in the number of free radicals observed in our experiments was qualitatively compatible with the theoretical prediction from the model of electromagnetic radiation effect on radical pairs.

Sannino A, Sarti M, Reddy SB, Prihoda TJ, Vijayalaxmi, Scarfi MR. Induction of adaptive response in human blood lymphocytes exposed to radiofrequency radiation. Radiat Res. 171(6):735-742, 2009.

Abstract Sannino, A., Sarti, M., Reddy, S. B., Prihoda, T. J., Vijayalaxmi and Scarfi, M. R. Induction of Adaptive Response in Human Blood Lymphocytes Exposed to

Studies that show **Cell Phone** Health Effects

Radiofrequency Radiation. Radiat. Res. 171, 735-742 (2009). The incidence of micronuclei was evaluated to assess the induction of an adaptive response to non-ionizing radiofrequency (RF) radiation in peripheral blood lymphocytes collected from five different human volunteers. After stimulation with phytohemagglutinin for 24 h, the cells were exposed to an adaptive dose of 900 MHz RF radiation used for mobile communications (at a peak specific absorption rate of 10 W/kg) for 20 h and then challenged with a single genotoxic dose of mitomycin C (100 ng/ml) at 48 h. Lymphocytes were collected at 72 h to examine the frequency of micronuclei in cytokinesis-blocked binucleated cells. Cells collected from four donors exhibited the induction of adaptive response (i.e., responders). Lymphocytes that were pre-exposed to 900 MHz RF radiation had a significantly decreased incidence of micronuclei induced by the challenge dose of mitomycin C compared to those that were not pre-exposed to 900 MHz RF radiation. These preliminary results suggested that the adaptive response can be induced in cells exposed to non-ionizing radiation. A similar phenomenon has been reported in cells as well as in animals exposed to ionizing radiation in several earlier studies. However, induction of adaptive response was not observed in the remaining donor (i.e., non-responder). The incidence of micronuclei induced by the challenge dose of mitomycin C was not significantly different between the cells that were pre-exposed and unexposed to 900 MHz RF radiation. Thus the overall data indicated the existence of heterogeneity in the induction of an adaptive response between individuals exposed to RF radiation and showed that the less time-consuming micronucleus assay can be used to determine whether an individual is a responder or non-responder.

Sannino A, Zeni O, Sarti M, Romeo S, Reddy SB, Belisario MA, Prihoda TJ, Vijayalaxmi, Scarfi MR. Induction of adaptive response in human blood lymphocytes exposed to 900 MHz radiofrequency fields: influence of cell cycle. Int J Radiat Biol. 87(9):993-999, 2011.

PURPOSE: To investigate the influence of cell cycle on the adaptive response (AR) induced by the exposure of human blood lymphocytes to radiofrequency fields (RF).

MATERIALS AND METHODS: Human peripheral blood lymphocytes in G(0)-, G(1)- or S-phase of the cell cycle were exposed for 20 hours to an adaptive dose (AD) of 900 MHz RF at an average specific absorption rate of 1.25 W/kg and then treated with a challenge dose (CD) of 100 ng/ml mitomycin C (MMC). Un-exposed and sham-exposed controls as well as cells treated with MMC alone were included in the study. The incidence of micronuclei (MN) was evaluated to determine the induction of AR. **RESULTS:** The results indicated that the cells which were exposed to AD of RF in G(0)- and G(1)-phase of the cell cycle did not exhibit AR while such a response was observed when the cells were exposed to AD of RF in S-phase of the cell cycle. **CONCLUSIONS:** These results confirmed the observations reported in our previous investigation where AR was observed in human blood lymphocytes exposed to AD of RF in S-phase of the cell cycle and further suggested that the timing of AD exposure of RF is important to elicit AR.

Studies that show **Cell Phone** Health Effects

Sannino A, Zeni O, Romeo S, Massa R, Gialanella G, Grossi G, Manti L, Vijayalaxmi, Scarfi MR. Adaptive response in human blood lymphocytes exposed to non-ionizing radiofrequency fields: resistance to ionizing radiation-induced damage. *J Radiat Res.* 2013 Aug 26. [Epub ahead of print]

The aim of this preliminary investigation was to assess whether human peripheral blood lymphocytes which have been pre-exposed to non-ionizing radiofrequency fields exhibit an adaptive response (AR) by resisting the induction of genetic damage from subsequent exposure to ionizing radiation. Peripheral blood lymphocytes from four healthy donors were stimulated with phytohemagglutinin for 24 h and then exposed for 20 h to 1950 MHz radiofrequency fields (RF, adaptive dose, AD) at an average specific absorption rate of 0.3 W/kg. At 48 h, the cells were subjected to a challenge dose (CD) of 1.0 or 1.5 Gy X-irradiation (XR, challenge dose, CD). After a 72 h total culture period, cells were collected to examine the incidence of micronuclei (MN). There was a significant decrease in the number of MN in lymphocytes exposed to RF + XR (AD + CD) as compared with those subjected to XR alone (CD). These observations thus suggested a RF-induced AR and induction of resistance to subsequent damage from XR. There was variability between the donors in RF-induced AR. The data reported in our earlier investigations also indicated a similar induction of AR in human blood lymphocytes that had been pre-exposed to RF (AD) and subsequently treated with a chemical mutagen, mitomycin C (CD). Since XR and mitomycin-C induce different kinds of lesions in cellular DNA, further studies are required to understand the mechanism(s) involved in the RF-induced adaptive response.

Sarimov, R., Malmgren, L.O.G., Markova, E., Persson, B.R.R., Belyaev, I.Y. Nonthermal GSM microwaves affect chromatin conformation in human lymphocytes similar to heat shock. *IEEE Trans Plasma Sci* 32:1600-1608, 2004.

Here we investigated whether microwaves (MWs) of Global System for Mobile Communication (GSM) induce changes in chromatin conformation in human lymphocytes. Effects of MWs were studied at different frequencies in the range of 895-915 MHz in experiments with lymphocytes from seven healthy persons. Exposure was performed in transverse electromagnetic transmission line cell (TEM-cell) using a GSM test-mobile phone. All standard modulations included 2 W output power in the pulses, specific absorbed rate (SAR) being 5.4 mW/kg. Changes in chromatin conformation, which are indicative of stress response and genotoxic effects, were measured by the method of anomalous viscosity time dependencies (AVTD). Heat shock and treatment with the genotoxic agent camptothecin, were used as positive controls. 30-min exposure to MWs at 900 and 905 MHz resulted in statistically significant condensation of chromatin in lymphocytes from 1 of 3 tested donors. This condensation was similar to effects of heat shock within the temperature window of 40/spl deg/C-44/spl deg/C. Analysis of pooled data from all donors showed statistically significant effect of 30-min exposure to MWs. Stronger effects of MWs was found following 1-h exposure. In replicated experiments, cells from four out of five donors responded to 905 MHz. Responses to 915 MHz were observed in cells from 1 out of 5 donors, $p < 0.002$. Dependent on donor, condensation, 3 donors, or decondensation, 1 donor, of

Studies that show **Cell Phone** Health Effects

chromatin was found in response to 1-h exposure. Analysis of pooled data from all donors showed statistically significant effect of 1-h exposure to MWs. In cells from one donor, this effect was frequency-dependent ($p < 0.01$). Effects of MWs correlated statistically significantly with effects of heat shock and initial state of chromatin before exposure. MWs at 895 and 915 MHz affected chromatin conformation in transformed lymphocytes. The conclusion-GSM microwaves under specific conditions of exposure affected human lymphocytes similar to stress response. The data suggested that the MW effects differ at various GSM frequencies and vary between donors.

Markova E, Hillert L, Malmgren L, Persson BR, Belyaev IY. Microwaves from GSM Mobile Telephones Affect 53BP1 and gamma-H2AX Foci in Human Lymphocytes from Hypersensitive and Healthy Persons. Environ Health Perspect. 113(9):1172-1177, 2005.

The data on biologic effects of nonthermal microwaves (MWs) from mobile telephones are diverse, and these effects are presently ignored by safety standards of the International Commission for Non-Ionizing Radiation Protection (ICNIRP). In the present study, we investigated effects of MWs of Global System for Mobile Communication (GSM) at different carrier frequencies on human lymphocytes from healthy persons and from persons reporting hypersensitivity to electromagnetic fields (EMFs). We measured the changes in chromatin conformation, which are indicative of stress response and genotoxic effects, by the method of anomalous viscosity time dependence, and we analyzed tumor suppressor p53-binding protein 1 (53BP1) and phosphorylated histone H2AX (gamma-H2AX), which have been shown to colocalize in distinct foci with DNA double-strand breaks (DSBs), using immunofluorescence confocal laser microscopy. We found that MWs from GSM mobile telephones affect chromatin conformation and 53BP1/gamma-H2AX foci similar to heat shock. For the first time, we report here that effects of MWs from mobile telephones on human lymphocytes are dependent on carrier frequency. On average, the same response was observed in lymphocytes from hypersensitive and healthy subjects. Key words: 53BP1 and gamma-H2AX foci, chromatin, DNA double-strand breaks, hypersensitivity to electromagnetic fields, stress response.

Lu YS, Huang BT, Huang YX. Reactive oxygen species formation and apoptosis in human peripheral blood mononuclear cell induced by 900MHz mobile phone radiation. Oxid Med Cell Longev. 2012:740280, 2012.

We demonstrate that reactive oxygen species (ROS) plays an important role in the process of apoptosis in human peripheral blood mononuclear cell (PBMC) which is induced by the radiation of 900MHz radiofrequency electromagnetic field (RFEMF) at a specific absorption rate (SAR) of ~ 0.4 W/kg when the exposure lasts longer than two hours. The apoptosis is induced through the mitochondrial pathway and mediated by activating ROS and caspase-3, and decreasing the mitochondrial potential. The activation of ROS is triggered by the conformation disturbance of lipids, protein, and DNA induced by the exposure of GSM RFEMF. Although human PBMC was found to have a self-protection mechanism of releasing carotenoid in response to oxidative stress to lessen the further increase of ROS, the imbalance between the antioxidant defenses

Studies that show **Cell Phone** Health Effects

and ROS formation still results in an increase of cell death with the exposure time and can cause about 37% human PBMC death in eight hours.

El-Bediwi AB, Saad M, El-Kott AF, Eid E. Influence of Electromagnetic Radiation Produced by Mobile Phone on Some Biophysical Blood Properties in Rats. *Cell Biochem Biophys*. 2012 Oct 10. [Epub ahead of print]

Effects of electromagnetic radiation produced by mobile phone on blood viscosity, plasma viscosity, hemolysis, Osmotic fragility, and blood components of rats have been investigated. Experimental results show that there are significant change on blood components and its viscosity which affects on a blood circulation due to many body problems. Red blood cells, White blood cells, and Platelets are broken after exposure to electromagnetic radiation produced by mobile phone. Also blood viscosity and plasma viscosity values are increased but Osmotic fragility value decreased after exposure to electromagnetic radiation produced by mobile phone.

Stankiewicz W, Dąbrowski MP, Kubacki R, Sobiczewska E, Szmigielski S Immunotropic influence of 900 MHz microwave GSM signal on human blood immune cells activated in vitro. *Electromag Biol Med* 25(1): 45-51, 2006.

In an earlier study we reported that G₀ phase peripheral blood mononuclear cells (PBMC) exposed to low-level (SAR = 0.18 W/kg) pulse-modulated 1300 MHz microwaves and subsequently cultured, demonstrate changed immune activity (Dabrowski et al., 2003). We investigated whether cultured immune cells induced into the active phases of cell cycle (G₁, S) and then exposed to microwaves will also be sensitive to electromagnetic field. An anechoic chamber of our design containing a microplate with cultured cells and an antenna emitting microwaves (900 MHz simulated GSM signal, 27 V/m, SAR 0.024 W/kg) was placed inside the ASSAB incubator. The microcultures of PBMC exposed to microwaves demonstrated significantly higher response to mitogens and higher immunogenic activity of monocytes (LM index) than control cultures. LM index, described in detail elsewhere (Dabrowski et al., 2001), represents the monokine influence on lymphocyte mitogenic response. The results suggest that immune activity of responding lymphocytes and monocytes can be additionally intensified by 900 MHz microwaves.

Effects on Wellbeing

Szyjowska A, Gadzicka E, Szymczak W, Bortkiewicz A. The risk of subjective symptoms in mobile phone users in Poland - An epidemiological study. *Int J Occup Med Environ Health*. 2014 Apr 1. [Epub ahead of print]

OBJECTIVES: To assess the type and incidence of subjective symptoms related to the use of mobile phones in Polish users. **MATERIAL AND METHODS:** The study was conducted in 2005 using a questionnaire survey. Although it has been quite a long time, up to now, no such data have been published for Poland. The questionnaire consisted of 53 questions concerning sex, age, education, general health, characteristics of a mobile

Studies that show **Cell Phone** Health Effects

phone (hand-held, loud-speaking unit) as well as the habits associated with its use (frequency and duration of calls, text messages, etc.) and complaints associated with using a mobile phone. **RESULTS:** As many as 1800 questionnaires were sent. The response was obtained from 587 subjects aged 32.6 ± 11.3 (48.9% women, 51.1% men); the age did not differ significantly between men and women. The subjects owned a cell phone for an average of 3 years. Majority of the respondents used the phone intensively, i.e. daily (74%) or almost daily (20%). Headaches were reported significantly more often by the people who talked frequently and long in comparison with other users (63.2% of the subjects, $p = 0.0029$), just like the symptoms of fatigue (45%, $p = 0.013$). Also, the feeling of warmth around the ear and directly to the auricle was reported significantly more frequently by the intensive mobile phone users, compared with other mobile phone users (47.3%, $p = 0.00004$ vs. 44.6%, $p = 0.00063$, respectively). Most symptoms appeared during or immediately after a call and disappeared within 2 h after the call. Continuous headache, persisting for longer than 6 h since the end of a call, was reported by 26% of the subjects. **CONCLUSIONS:** Our results show that the mobile phone users may experience subjective symptoms, the intensity of which depends on the intensity of use of mobile phones.

Szykowska A, Bortkiewicz A, Szymczak W, Makowiec-Dabrowska T. [Subjective symptoms related to mobile phone use--a pilot study] *Pol Merkuriusz Lek.* 19(112):529-532, 2005. [Article in Polish]

Research findings indicate that the use of mobile phones may lead to a number of symptoms such as headache, impaired concentration and memory, fatigue. In Poland this problem has not as yet been addressed by scientific studies. **THE AIM:** The present project was undertaken to investigate whether the symptoms of ill health reported by young people may be associated with the use of mobile phone. **MATERIAL AND METHODS:** A survey using a self-reported questionnaire was conducted among randomly selected university students in Lodz, Central Poland. The questionnaire was designed specifically for this study and contained items on health condition and complaints as well as on frequency of mobile phone use. The number of questionnaires necessary for the study was assessed using the simple random sample method. Out of the 160 copies distributed among the students, 140 (87.5%) were completed. Eventually, 117 questionnaires were subject to analysis; the data from respondents who reported health problems (neck trauma in a car accident, chronic sinusitis and arterial hypertension) were excluded. The following statistical methods were used to analyse questionnaire data: t-Student test for equal and unequal variances or F-Snedecor test for comparing parameters in two study groups, Fisher exact test for comparing frequency, and single and multiple logistic regression models for quantitative risk assessment of negative health outcomes in relation to exposure level and with control for confounders. The subjects were 61 (52.1%) males and 56 females (47.9%). **RESULTS:** Most of the subjects (62%) assessed their health condition as good, 31% as very good and 7% as fair. 70% complained of headache and 20% of dizziness. Impaired concentration occurred in 56% of respondents. Facial dermatitis was reported by 11%. The most prevalent symptom related to mobile phone use was the thermal sensation

Studies that show **Cell Phone** Health Effects

within the auricle and behind/around the ear. This was reported by 33 subjects (28.2%). Out of 82 subjects who complained of headache, only 8 (6.8%) related this symptom to mobile phone use. Only 10 subjects of 65 reporting impaired concentration thought it could be associated with their using a mobile phone. The symptoms and health complaints reported by the respondents in no case were the reason for a medical check-up or taking any medication. CONCLUSIONS: The large number of young people complaining of headache and impaired concentration calls for further research to investigate the underlying reasons. It cannot be excluded that one of them may be exposure to EMF emitted by mobile phone. The explanation should be sought through further experimental and epidemiologic studies.

Santini R, Seigne M, Bonhomme-Faivre L, Bouffet S, Defrasne E, Sage M. Symptoms experienced by users of digital cellular phones: a pilot study in a French engineering school. Pathol Biol (Paris) 49(3):222-226, 2001. [Article in French]

A survey study, using questionnaire, was conducted in 161 students and workers in a French engineering school on symptoms experienced during use of digital cellular phones. A significant increase in concentration difficult ($p < 0.05$) was reported by users of 1800-MHz (DCS) cellular phones compared to 900-MHz (GSM) phone users. In users of cellular phones, women significantly ($p < 0.05$) complained more often of sleep disturbance than men. This sex difference for sleep complaint is not observed between women and men non-users of cellular phone. The use of both cellular phones and VDT significantly ($p \leq 0.05$) increased concentration difficulty. Digital cellular phone users also significantly ($p < 0.05$) more often complained of discomfort, warmth, and picking on the ear during phone conversation in relation with calling duration per day and number of calls per day. The complaint warmth on the ear might be a signal to users for stopping the call.

Sandstrom M, Wilen J, Oftedal G, Hansson Mild K, Mobile phone use and subjective symptoms. Comparison of symptoms experienced by users of analogue and digital mobile phones. Occup Med (Lond) 51(1):25-35, 2001.

In 1995 many people reported symptoms such as headaches, feelings of discomfort, warmth behind/around or on the ear and difficulties concentrating while using mobile phones. The number of complaints was higher for people using the digital (GSM) system, i.e. with pulse modulated fields, than for those using the analogue (NMT) system. Our main hypothesis was that GSM users experience more symptoms than NMT users. An epidemiological investigation was initiated including 6379 GSM users and 5613 NMT 900 users in Sweden, and 2500 from each category in Norway. The adjusted odds ratio did not indicate any increased risk for symptoms for GSM users compared with NMT 900 users. Our hypothesis was therefore disproved. However, we observed a statistically significant lower risk for sensations of warmth on the ear for GSM users compared with NMT 900 users. The same trend was seen in Norway for sensations of warmth behind/around the ear and in Sweden for headaches and fatigue. Factors distinguishing the two systems (radio frequency emission, phone temperatures and various ergonomic factors) may be responsible for these results, as well as for a

Studies that show **Cell Phone** Health Effects

secondary finding: a statistically significant association between calling time/number of calls per day and the prevalence of warmth behind/around or on the ear, headaches and fatigue.

Rubin GJ, Cleare AJ, Wessely S. Psychological factors associated with self-reported sensitivity to mobile phones. J Psychosom Res. 64(1):1-9; discussion 11-12, 2008

OBJECTIVE: Some people report symptoms associated with mobile phone use. A minority also report "electrosensitivity," experiencing symptoms following exposure to other electrical devices. Research suggests that electromagnetic fields do not trigger these symptoms. In this study, we examined the differences between these two "sensitive" groups and healthy controls. METHODS: Fifty-two people who reported sensitivity to mobile phones, 19 people who reported sensitivity to mobile phones and "electrosensitivity," and 60 nonsensitive controls completed a questionnaire assessing the following: primary reason for using a mobile phone, psychological health, symptoms of depression, modern health worries (MHW), general health status, symptom severity, and the presence of other medically unexplained syndromes. RESULTS: Perceived sensitivity was associated with an increased likelihood of using a mobile phone predominantly for work (3% of controls, 13% of those sensitive to mobile phones, and 21% of those reporting "electrosensitivity") and greater MHW concerning radiation [mean (S.D.) on a scale of 1-5: 2.0 (1.0), 2.7 (0.9), and 4.0 (0.8), respectively]. Participants who reported "electrosensitivity" also experienced greater depression, greater worries about tainted food and toxic interventions, worse general health on almost every measure, and a greater number of other medically unexplained syndromes compared to participants from the other two groups. No group differences were observed with regards to psychiatric cases. CONCLUSIONS: The data illustrate that patients reporting "electrosensitivity" experience substantially worse health than either healthy individuals or people who report sensitivity to mobile phones but who do not adopt the label "electrosensitivity." Clinicians and researchers would be wise to pay greater attention to this subdivision.

Oftedal G, Wilen J, Sandstrom M, Mild KH, Symptoms experienced in connection with mobile phone use. Occup Med (Lond) 50(4):237-245, 2000.

Many people in Norway and Sweden reported headaches, fatigue, and other symptoms experienced in connection with the use of a mobile phone (MP). Therefore, we initiated a cross-sectional epidemiological study among 17,000 people, all using an MP in their job. Thirty-one percent of the respondents in Norway and 13% of those in Sweden had experienced at least one symptom in connection with MP use. Next to the sensations of warmth on the ear and behind/around the ear, burning sensations in the facial skin and headaches were most commonly reported. Most symptoms usually began during or within half an hour after the call and lasted for up to 2 h. Relatively few had consulted a physician or been on sick leave because of the symptoms, but about 45% among those with an MP attributed symptom had taken steps to reduce the symptom. These results suggest an awareness of the symptoms, but not necessarily a serious health problem.

Khan MM. Adverse effects of excessive mobile phone use.Int J Occup Med Environ

Studies that show **Cell Phone** Health Effects

Health. 21(4):289-293, 2008.

Introduction: Research findings indicate that the use of mobile phones may lead to a number of symptoms such as headache, impaired concentration and memory, and also fatigue. Materials and Methods: The present study was designed to investigate whether the symptoms of ill health reported by young people may be associated with the use of mobile phone (MP) and to analyze its influence on health and development of medical students. The questionnaire was designed specifically for this study and contained items regarding health condition and health complaints as well as the frequency of MP use. The response rate was 86.6% (286 of 330 forms, completed by 73.77% males and 26.22% females). Results: Most of the subjects (83.57%) had some knowledge about the adverse effects of MP use. 76.92% of the students carried one mobile, and 23.08% more than one. 55.94%, of the subjects reported the average daily MP use of less than 30 min, 27.97%, of 30-60 min, 11.53%, of 60-90 min and 4.54% of more than 90 min. 16.08% of the subjects complained of headache and 24.48% of fatigue. Impaired concentration was reported by 34.27% of respondents, memory disturbances by 40.56%, sleeplessness by 38.8%, hearing problems by 23.07%, and facial dermatitis by 16.78%. The sensation of warmth within the auricle and behind/around the ear was reported by 28.32%. Out of 286 subjects who participated in this study, 44.4% related their symptoms to mobile phone use. Conclusions: The findings of the present study indicate that mobile phones play a large part in the daily life of medical students. Therefore, its impact on psychology and health should be discussed among the students to prevent the harmful effects of mobile phone use.

Thomé S, Härenstam A, Hagberg M. Mobile phone use and stress, sleep disturbances, and symptoms of depression among young adults--a prospective cohort study.BMC Public Health. 11:66, 2011.

BACKGROUND: Because of the quick development and widespread use of mobile phones, and their vast effect on communication and interactions, it is important to study possible negative health effects of mobile phone exposure. The overall aim of this study was to investigate whether there are associations between psychosocial aspects of mobile phone use and mental health symptoms in a prospective cohort of young adults. METHODS: The study group consisted of young adults 20-24 years old (n = 4156), who responded to a questionnaire at baseline and 1-year follow-up. Mobile phone exposure variables included frequency of use, but also more qualitative variables: demands on availability, perceived stressfulness of accessibility, being awakened at night by the mobile phone, and personal overuse of the mobile phone. Mental health outcomes included current stress, sleep disorders, and symptoms of depression. Prevalence ratios (PRs) were calculated for cross-sectional and prospective associations between exposure variables and mental health outcomes for men and women separately. RESULTS: There were cross-sectional associations between high compared to low mobile phone use and stress, sleep disturbances, and symptoms of depression for the men and women. When excluding respondents reporting mental health symptoms at baseline, high mobile phone use was associated with sleep disturbances and symptoms of depression for the men and symptoms of depression for the women at 1-

Studies that show **Cell Phone** Health Effects

year follow-up. All qualitative variables had cross-sectional associations with mental health outcomes. In prospective analysis, overuse was associated with stress and sleep disturbances for women, and high accessibility stress was associated with stress, sleep disturbances, and symptoms of depression for both men and women. CONCLUSIONS: High frequency of mobile phone use at baseline was a risk factor for mental health outcomes at 1-year follow-up among the young adults. The risk for reporting mental health symptoms at follow-up was greatest among those who had perceived accessibility via mobile phones to be stressful. Public health prevention strategies focusing on attitudes could include information and advice, helping young adults to set limits for their own and others' accessibility.

Ikeda K, Nakamura K. Association between mobile phone use and depressed mood in Japanese adolescents: a cross-sectional study. Environ Health Prev Med. 2013 Dec 18. [Epub ahead of print]

OBJECTIVES: Mobile phones are commonly used by adolescents. The aim of this study was to clarify associations between duration of mobile phone use and psychological mood in high school students. **METHODS:** This cross-sectional study included 2,785 high school students in Niigata, Japan. A self-administered questionnaire was used to elicit information on sex, school year, hours of mobile phone use, psychological mood status, and possible confounders. Psychological mood outcomes were evaluated with the Mood Inventory, developed and validated in 1994, which includes five subcomponents with total scores ranging from 8 to 32 (higher score indicates stronger feeling): "Tension and excitement," "Refreshing mood," "Fatigue," "Depressed mood," and "Anxious mood." Analysis of covariance with Bonferroni's multiple comparison was used to compare mean values among quartiles of hours of mobile phone use. **RESULTS:** Among the respondents, mean mobile phone use per week was 24 (median 18) h. Long-duration mobile phone use was associated with female students, no participation in sports club activities, early mobile phone use, and fewer hours spent sleeping (all $P < 0.001$). Overall associations between hours of mobile phone use and total scores were significant for "Depressed mood" (P for trend = 0.005), "Tension and excitement" (P for trend < 0.001), and "Fatigue" (P for trend < 0.001). Total scores for "Depressed mood," "Tension and excitement," and "Fatigue" of the fourth quartile (≥ 33 h/week) of mobile phone use were significantly higher than for other quartiles (all $P < 0.05$). **CONCLUSIONS:** Increased duration of mobile phone use is associated with unfavorable psychological mood, in particular, a depressed mood. Decreasing mobile phone use may help maintain appropriate mental health in very long-duration users.

Hocking, B, Preliminary report: symptoms associated with mobile phone use. Occup Med (Lond);48(6):357-360, 1998.

Mobile phone use is ubiquitous, although the alleged health effects of low level radio-frequency radiation (RFR) used in transmission are contentious. Following isolated reports of headache-like symptoms arising in some users, a survey has been conducted to characterize the symptoms sometimes associated with mobile phone usage. A notice of interest in cases was placed in a major medical journal and this was publicized by the media. Respondents were interviewed by telephone using a structured questionnaire.

Studies that show **Cell Phone** Health Effects

Forty respondents from diverse occupations described unpleasant sensations such as a burning feeling or a dull ache mainly occurring in the temporal, occipital or auricular areas. The symptoms often began minutes after beginning a call, but could come on later during the day. The symptoms usually ceased within an hour after the call, but could last until evening. Symptoms did not occur when using an ordinary handset, and were different from ordinary headaches. There were several reports suggestive of intracranial effects. Three respondents reported local symptoms associated with wearing their mobile phone on their belts. There was one cluster of cases in a workplace. Seventy-five per cent of cases were associated with digital mobile phones. Most of the respondents obtained relief by altering their patterns of telephone usage or type of phone. Cranial and other diverse symptoms may arise associated with mobile phone usage. Physicians and users alike should be alert to this. Further work is needed to determine the range of effects, their mechanism and the possible implications for safety limits of RFR.

Hocking B , Microwave sickness: a reappraisal. Occup Med (Lond) 51(1):66-69, 2001.

Microwave sickness (MWS) has been a disputed condition. The syndrome involves the nervous system and includes fatigue, headaches, dysaesthesia and various autonomic effects in radiofrequency radiation workers. This paper describes the early reports of the syndrome from Eastern Europe and notes the scepticism expressed about them in the West, before considering comprehensive recent reports by Western specialists and a possible neurological basis for the condition. It is concluded that MWS is a medical entity which should be recognized as a possible risk for radiofrequency radiation workers.

Hocking B. Management of radiofrequency radiation overexposures. Aust Fam Physician 30(4):339-342, 2001.

BACKGROUND: Radiofrequency radiation (RFR) has been in use for some time but is now proliferative with the burgeoning radiocommunications industry. OBJECTIVE: To inform the profession of the possible health effects from overexposure to radiofrequency radiation (RFR) and the clinical approach to cases. An introduction to the health effects of overexposure to RFR is given. A clinical approach to integrating the patient's symptoms and the circumstances of the exposure is given. Emergency treatment and ongoing care is outlined, and sources of expert advice given. CONCLUSION: Overexposure to RFR is a complex injury. Advice is given in this article for emergency care and planning ongoing care.

Hanson Mild, K, Oftedal, G, Sandstrom, M, Wilen, J, Tynes, T, Haugsdal, B, Hauger E, Comparison of symptoms experienced by users of analogue and digital mobile phones: a Swedish-Norwegian epidemiological study. Arbetslivsrapport 1998:23.

Study of mobile phone users showed a statistically significant association between calling time/number of calls per day and the prevalence of warmth behind/around the ear, headaches, and fatigue.

Studies that show **Cell Phone** Health Effects

Hallberg O. Adverse health indicators correlating with sparsely populated areas in Sweden. *Eur J Cancer Prev.* 16(1):71-76, 2007.

Earlier studies on health characteristics in Sweden have pointed at a sudden trend change in general health indicators around 1997. The decline was worse in areas with less estimated coverage by the mobile phone system; that is, areas where the average output power from mobile phone handsets is expected to be higher. In this study, health parameters were related to the population density, which is a well defined, rather than an estimated variable. Statistics were obtained from different authorities in Sweden. Data were correlated to the population densities in the 21 different counties of Sweden as well as to estimates of average mobile phone output power. Several health quality measures showed that people in sparsely populated counties in Sweden (as well as in Denmark and Norway) have suffered more illness, and lengthier recovery than people in more densely populated areas since 1997. This is in strong contrast to the situation 20 years ago, when the countryside was the healthiest place in which to live. The indicators strongly correlated with estimated mobile phone area coverage and estimated power output. The indicator statistics suggest that the decline in health in Sweden is not a primary consequence of low population density by itself, but that other factors related to population density are causative. The two factors having the strongest correlation with decreased health quality were the estimated average power output from mobile phones (positive correlation) and the reported coverage from the global system for mobile communication base stations (negative correlation) in each county.

Hallberg O, Johansson O. Long-term sickness and mobile phone use. *J Aust Coll Nutr & Env med* 23:11-12, 2004.

The number of people unable to work due to long-term sickness is drastically increasing in Sweden. In this paper we take a close look at the development of mobile phone communication to see how it possibly relates to the health impairment of the Swedish population. Official data was collected regarding mobile phone use and long-term absence from work. The co-variation between those data sets was used to estimate future development of long-term absence rates under the hypothesis that there is a connection between the two sets of data. It was concluded that future long-term absence rates will continue to increase as long as the annual number of ear-heating minute per year is increasing.

Gómez-Perretta C, Navarro EA, Segura J, Portolés M. Subjective symptoms related to GSM radiation from mobile phone base stations: a cross-sectional study. *BMJ Open.* 3(12):e003836, 2013. doi: 10.1136/bmjopen-2013-003836.

OBJECTIVES: We performed a re-analysis of the data from Navarro et al (2003) in which health symptoms related to microwave exposure from mobile phone base stations (BSs) were explored, including data obtained in a retrospective inquiry about fear of exposure from BSs. DESIGN: Cross-sectional study. SETTING: La Ñora (Murcia), Spain. PARTICIPANTS: Participants with known illness in 2003 were subsequently disregarded: 88 participants instead of 101 (in 2003) were analysed. Since weather circumstances can influence exposure, we restricted data to measurements made under similar weather conditions. OUTCOMES

Studies that show **Cell Phone** Health Effects

AND METHODS: A statistical method indifferent to the assumption of normality was employed: namely, binary logistic regression for modelling a binary response (eg, suffering fatigue (1) or not (0)), and so exposure was introduced as a predictor variable. This analysis was carried out on a regular basis and bootstrapping (95% percentile method) was used to provide more accurate CIs. RESULTS: The symptoms most related to exposure were lack of appetite (OR=1.58, 95% CI 1.23 to 2.03); lack of concentration (OR=1.54, 95% CI 1.25 to 1.89); irritability (OR=1.51, 95% CI 1.23 to 1.85); and trouble sleeping (OR=1.49, 95% CI 1.20 to 1.84). Changes in -2 log likelihood showed similar results. Concerns about the BSs were strongly related with trouble sleeping (OR =3.12, 95% CI 1.10 to 8.86). The exposure variable remained statistically significant in the multivariate analysis. The bootstrapped values were similar to asymptotic CIs. CONCLUSIONS: This study confirms our preliminary results. We observed that the incidence of most of the symptoms was related to exposure levels-independently of the demographic variables and some possible risk factors. Concerns about adverse effects from exposure, despite being strongly related with sleep disturbances, do not influence the direct association between exposure and sleep.

Frick U, Rehm J, Eichhammer P. Risk perception, somatization, and self report of complaints related to electromagnetic fields--a randomized survey study. *Int J Hyg Environ Health*. 205(5):353-360, 2002.

Exposure to electromagnetic fields (EMF) as well as EMF-related complaints has increased over the past decades. However, it is unclear whether these complaints are related to the electromagnetic or other physical properties of these fields per se, to salience of EMF in media, or to both. What is the prevalence of EMF-related complaints in the general population? What are the influencing factors on this prevalence? Does reporting of EMF-related symptoms depend on cognitive factors? To answer these questions, a survey with random variation of three cognitive factors was performed. As expected, EMF-related complaints were reported more by females and people with higher somatization tendency. Age had no significant linear effect on EMF-related complaints. The cognitive condition of threat produced a significant contrast effect among people with high somatization tendency on EMF-related complaints. Cognition can influence reporting of EMF-related effects. Thus, in future research of such effects, psychologically influencing factors should be included. Also risk communication should incorporate knowledge about social cognition.

Effects on Glands

Abu Khadra KM, Khalil AM, Abu Samak M, Aljaberi A. Evaluation of selected biochemical parameters in the saliva of young males using mobile phones. *Electromagn Biol Med*. 2014 Feb 5. [Epub ahead of print]

Abstract The biochemical status in the saliva of 12 males before/after using mobile phone has been evaluated. Radio frequency signals of 1800 MHz (continuous wave transmission, 217 Hz modulate and Global System for Mobile Communications [GSM - non-DTX]) with 1.09 w/kg specific absorption rate (SAR) value were used for 15 and

Studies that show **Cell Phone** Health Effects

30 min. Cell phone radiation induced a significant increase of superoxide dismutase (SOD); there was a statistically significant effect of talking time on the levels of SOD, $F(2, 33) = 8.084$, $p < 0.05$, $\omega = 0.53$. The trend analysis suggests a significant quadratic trend, $F(1, 33) = 4.891$, $p < 0.05$; indicating that after 15 min of talking the levels of SOD increased, but as talking time increased the SOD activity started to drop. In contrast to this, there was no statistically significant effect of talking time on the level of salivary albumin, cytochrome c, catalase or uric acid. Results suggest that exposure to electromagnetic radiation may exert an oxidative stress on human cells as evidenced by the increase in the concentration of the superoxide radical anion released in the saliva of cell phone users.

Hashemipour MS, Yarbakht M, Gholamhosseini A, Famori H. Effect of mobile phone use on salivary concentrations of protein, amylase, lipase, immunoglobulin A, lysozyme, lactoferrin, peroxidase and C-reactive protein of the parotid gland. J Laryngol Otol. 2014 Apr 17:1-9. [Epub ahead of print]

Background: The possibility of side effects associated with the electromagnetic waves emitted from mobile phones is a controversial issue. The present study aimed to evaluate the effect of mobile phone use on parotid gland salivary concentrations of protein, amylase, lipase, immunoglobulin A, lysozyme, lactoferrin, peroxidase and C-reactive protein. Methods: Stimulated salivary samples were collected simultaneously from both parotid glands of 86 healthy volunteers. Salivary flow rate and salivary concentrations of proteins, amylase, lipase, lysozyme, lactoferrin, peroxidase, C-reactive protein and immunoglobulin A, were measured. Data were analysed using t-tests and one-way analyses of variance. Results: Salivary flow rate and parotid gland salivary concentrations of protein were significantly higher on the right side compared to the left in those that predominantly held mobile phones on the right side. In addition, there was a decrease in concentrations of amylase, lipase, lysozyme, lactoferrin and peroxidase. Conclusion: The side of dominant mobile phone use was associated with differences in salivary flow rate and parotid gland salivary concentrations, in right-dominant users. Although mobile phone use influenced salivary composition, the relationship was not significant.

Goldwein O, Aframian DJ. The influence of handheld mobile phones on human parotid gland secretion. Oral Dis.16(2):146-150, 2010.

Abstract. BACKGROUND: Handheld mobile phones (MPHs) have become a 'cultural' accessory device, no less so than a wrist watch. Nevertheless, the use of MPHs has given rise to great concern because of possible adverse health effects from exposure to the radiofrequency radiation (RFR) emitted by the device. Previous studies suggested correlation between MPH and salivary gland tumors. OBJECTIVE: To evaluate whether MPH induces physiologic changes in the adjacent parotid gland, located on the dominant side, in terms of secretion rates and protein levels in the secreted saliva. MATERIALS AND METHOD: Stimulated parotid saliva was collected simultaneously from both glands in 50 healthy volunteers whose MPH use was on a dominant side of the head. RESULTS: A significantly higher saliva secretion rate was noticed in the dominant

Studies that show **Cell Phone** Health Effects

MPH side compared with that in the non-dominant side. Lower total protein concentration was obtained in the dominant compared with the non-dominant MPH side among the right dominant MPH users. CONCLUSIONS: Parotid glands adjacent to handheld MPH in use respond by elevated salivary rates and decreased protein secretion reflecting the continuous insult to the glands. This phenomenon should be revealed to the worldwide population and further exploration by means of large-scale longitudinal studies is warranted.

Pereira C, Edwards M, Parotid nodular fasciitis in a mobile phone user. J Laryngol Otol 114(11):886-887, 2000.

We describe the first case of nodular fasciitis affecting the deep lobe of the parotid gland in a 39-year-old male telephone engineer and its possible association with the high usage of mobile phones.

Hamzany Y, Feinmesser R, Shpitzer T, Mizrachi A, Hilly O, Hod R, Bahar G, Otradnov I, Gavish M, Nagler RM. Is human saliva an indicator of the adverse health effects of using mobile phones? Antioxid Redox Signal. 18(6):622-627, 2013.

Increasing use of mobile phones creates growing concern regarding harmful effects of radiofrequency non-ionizing electromagnetic radiation (NIER) on human tissues located close to the ear where phones are commonly held for long periods of time. We studied 20 subjects in the 'mobile phone group' who had a mean duration of mobile phone use of 12.5 years (range 8-15) and a mean time use of 29.6 hours per month (range 8-100). Deaf individuals served as controls. We compared salivary outcomes (secretion, oxidative damage indices, flow rate and composition) between mobile phone users and non-users. We report significant increase in all salivary oxidative stress indices studied in mobile phone users. Salivary flow, total protein, albumin and amylase activity were decreased in mobile phone users. These observations lead to the hypothesis that the use of mobile phones may cause oxidative stress and modify salivary function.

Effects on Sleep

Yogesh S, Abha S, Priyanka S. Mobile usage and sleep patterns among medical students. Indian J Physiol Pharmacol. 58(1):100-103, 2014.

Exposure of humans to radio frequency electromagnetic field (EMF) both during receiving and transmitting the signals has amplified public and scientific debate about possible adverse effects on human health. The study was designed with the objective of assessing the extent of mobile phone use amongst medical students and finding correlation if any between the hours of usage of mobile to sleep pattern and quality. hundred medical students grouped as cases (n = 57) (> 2 hours/day of mobile usage) and control (n = 43) (\leq 2 hours/day of mobile usage) were examined for their sleep quality & pattern by Pittsburg sleep Quality Index (PSQI). Differences between groups were

Studies that show **Cell Phone** Health Effects

examined with the Mann Whitney "U" test for proportions (Quantitative values) and with Student's 't' test for continuous variables. The association of variables was analyzed by Spearman Rank's correlation. Probability was set at < 0.05 as significant. Sleep disturbance, latency and day dysfunction was more in cases especially females. A significant association of hours of usage and sleep indices were observed in both genders (males $r = 0.25$; $p = 0.04$, females $r = 0.31$; $p = 0.009$). Evening usage of mobile phone in cases showed a statistically significant negative association (-0.606 ; $p = 0.042$) with Sleep quality (higher PSQI means sleep deprivation). Students using mobile for > 2 hours/day may cause sleep deprivation and day sleepiness affecting cognitive and learning abilities of medical students.

Mohammed HS, Fahmy HM, Radwah NM, Elsayed AA. Non-thermal continuous and modulated electromagnetic radiation fields effects on sleep EEG of rats. J Adv Res 4(2) 181-187, 2013.

In the present study, the alteration in the sleep EEG in rats due to chronic exposure to low-level non-thermal electromagnetic radiation was investigated. Two types of radiation fields were used; 900 MHz unmodulated wave and 900 MHz modulated at 8 and 16 Hz waves. Animals has exposed to radiation fields for 1 month (1 h/day). EEG power spectral analyses of exposed and control animals during slow wave sleep (SWS) and rapid eye movement sleep (REM sleep) revealed that the REM sleep is more susceptible to modulated radiofrequency radiation fields (RFR) than the SWS. The latency of REM sleep increased due to radiation exposure indicating a change in the ultradian rhythm of normal sleep cycles. The cumulative and irreversible effect of radiation exposure was proposed and the interaction of the extremely low frequency radiation with the similar EEG frequencies was suggested.

Mann, K, Roschke, J, Effects of pulsed high-frequency electromagnetic fields on human sleep. Neuropsychobiology 33(1):41-47, 1996.

In the present study we investigated the influence of pulsed high-frequency electromagnetic fields of digital mobile radio telephones on sleep in healthy humans. Besides a hypnotic effect with shortening of sleep onset latency, a REM suppressive effect with reduction of duration and percentage of REM sleep was found. Moreover, spectral analysis revealed qualitative alterations of the EEG signal during REM sleep with an increased spectral power density. Knowing the relevance of REM sleep for adequate information processing in the brain, especially concerning mnestic functions and learning processes, the results emphasize the necessity to carry out further investigations on the interaction of this type of electromagnetic fields and the human organism.

Lustenberger C, Murbach M, Dürr R, Schmid MR, Kuster N, Achermann P, Huber R. Stimulation of the brain with radiofrequency electromagnetic field pulses affects sleep-dependent performance improvement. Brain Stimul. 6(5):805-811, 2013.

BACKGROUND: Sleep-dependent performance improvements seem to be closely related to sleep spindles (12-15 Hz) and sleep slow-wave activity (SWA, 0.75-4.5 Hz). Pulse-modulated

Studies that show **Cell Phone** Health Effects

radiofrequency electromagnetic fields (RF EMF, carrier frequency 900 MHz) are capable to modulate these electroencephalographic (EEG) characteristics of sleep. **OBJECTIVE:** The aim of our study was to explore possible mechanisms how RF EMF affect cortical activity during sleep and to test whether such effects on cortical activity during sleep interact with sleep-dependent performance changes. **METHODS:** Sixteen male subjects underwent 2 experimental nights, one of them with all-night 0.25-0.8 Hz pulsed RF EMF exposure. All-night EEG was recorded. To investigate RF EMF induced changes in overnight performance improvement, subjects were trained for both nights on a motor task in the evening and the morning. **RESULTS:** We obtained good sleep quality in all subjects under both conditions (mean sleep efficiency > 90%). After pulsed RF EMF we found increased SWA during exposure to pulse-modulated RF EMF compared to sham exposure ($P < 0.05$) toward the end of the sleep period. Spindle activity was not affected. Moreover, subjects showed an increased RF EMF burst-related response in the SWA range, indicated by an increase in event-related EEG spectral power and phase changes in the SWA range. Notably, during exposure, sleep-dependent performance improvement in the motor sequence task was reduced compared to the sham condition (-20.1%, $P = 0.03$). **CONCLUSION:** The changes in the time course of SWA during the exposure night may reflect an interaction of RF EMF with the renormalization of cortical excitability during sleep, with a negative impact on sleep-dependent performance improvement.

Lowden A, Akerstedt T, Ingre M, Wiholm C, Hillert L, Kuster N, Nilsson JP, Arnetz B. Sleep after mobile phone exposure in subjects with mobile phone-related symptoms. *Bioelectromagnetics*. 32(1):4-14, 2011.

Several studies show increases in activity for certain frequency bands (10-14 Hz) and visually scored parameters during sleep after exposure to radiofrequency electromagnetic fields. A shortened REM latency has also been reported. We investigated the effects of a double-blind radiofrequency exposure (884 MHz, GSM signaling standard including non-DTX and DTX mode, time-averaged 10 g psSAR of 1.4 W/kg) on self-evaluated sleepiness and objective EEG measures during sleep. Forty-eight subjects (mean age 28 years) underwent 3 h of controlled exposure (7:30-10:30 PM; active or sham) prior to sleep, followed by a full-night polysomnographic recording in a sleep laboratory. The results demonstrated that following exposure, time in Stages 3 and 4 sleep (SWS, slow-wave sleep) decreased by 9.5 min (12%) out of a total of 78.6 min, and time in Stage 2 sleep increased by 8.3 min (4%) out of a total of 196.3 min compared to sham. The latency to Stage 3 sleep was also prolonged by 4.8 min after exposure. Power density analysis indicated an enhanced activation in the frequency ranges 0.5-1.5 and 5.75-10.5 Hz during the first 30 min of Stage 2 sleep, with 7.5-11.75 Hz being elevated within the first hour of Stage 2 sleep, and bands 4.75-8.25 Hz elevated during the second hour of Stage 2 sleep. No pronounced power changes were observed in SWS or for the third hour of scored Stage 2 sleep. No differences were found between controls and subjects with prior complaints of mobile phone-related symptoms. The results confirm previous findings that RF exposure increased the EEG alpha range in the sleep EEG, and indicated moderate impairment of SWS. Furthermore, reported differences in sensitivity to mobile phone use were not reflected in sleep

Studies that show **Cell Phone** Health Effects

parameters.

Wiholm C, Lowden A, Kuster N, Hillert L, Arnetz BB, Akerstedt T, Moffat SD. Mobile phone exposure and spatial memory. *Bioelectromagnetics*. 30(1):59-65, 2009.

Radiofrequency (RF) emission during mobile phone use has been suggested to impair cognitive functions, that is, working memory. This study investigated the effects of a 2 1/2 h RF exposure (884 MHz) on spatial memory and learning, using a double-blind repeated measures design. The exposure was designed to mimic that experienced during a real-life mobile phone conversation. The design maximized the exposure to the left hemisphere. The average exposure was peak spatial specific absorption rate (psSAR10g) of 1.4 W/kg. The primary outcome measure was a "virtual" spatial navigation task modeled after the commonly used and validated Morris Water Maze. The distance traveled on each trial and the amount of improvement across trials (i.e., learning) were used as dependent variables. The participants were daily mobile phone users, with and without symptoms attributed to regular mobile phone use. Results revealed a main effect of RF exposure and a significant RF exposure by group effect on distance traveled during the trials. The symptomatic group improved their performance during RF exposure while there was no such effect in the non-symptomatic group. Until this new finding is further investigated, we can only speculate about the cause.

Loughran SP, McKenzie RJ, Jackson ML, Howard ME, Croft RJ. Individual differences in the effects of mobile phone exposure on human sleep: rethinking the problem. *Bioelectromagnetics*. 33(1):86-93, 2012.

Mobile phone exposure-related effects on the human electroencephalogram (EEG) have been shown during both waking and sleep states, albeit with slight differences in the frequency affected. This discrepancy, combined with studies that failed to find effects, has led many to conclude that no consistent effects exist. We hypothesised that these differences might partly be due to individual variability in response, and that mobile phone emissions may in fact have large but differential effects on human brain activity. Twenty volunteers from our previous study underwent an adaptation night followed by two experimental nights in which they were randomly exposed to two conditions (Active and Sham), followed by a full-night sleep episode. The EEG spectral power was increased in the sleep spindle frequency range in the first 30 min of non-rapid eye movement (non-REM) sleep following Active exposure. This increase was more prominent in the participants that showed an increase in the original study. These results confirm previous findings of mobile phone-like emissions affecting the EEG during non-REM sleep. Importantly, this low-level effect was also shown to be sensitive to individual variability. Furthermore, this indicates that previous negative results are not strong evidence for a lack of an effect and, given the far-reaching implications of mobile phone research, we may need to rethink the interpretation of results and the manner in which research is conducted in this field.

Studies that show **Cell Phone** Health Effects

Loughran SP, Wood AW, Barton JM, Croft RJ, Thompson B, Stough C. The effect of electromagnetic fields emitted by mobile phones on human sleep. *Neuroreport*. 16(17):1973-1976, 2005.

Previous research has suggested that exposure to radiofrequency electromagnetic fields increases electroencephalogram spectral power in non-rapid eye movement sleep. Other sleep parameters have also been affected following exposure. We examined whether aspects of sleep architecture show sensitivity to electromagnetic fields emitted by digital mobile phone handsets. Fifty participants were exposed to electromagnetic fields for 30 min prior to sleep. Results showed a decrease in rapid eye movement sleep latency and increased electroencephalogram spectral power in the 11.5-12.25 Hz frequency range during the initial part of sleep following exposure. These results are evidence that mobile phone exposure prior to sleep may promote rapid eye movement sleep and modify the sleep electroencephalogram in the first non-rapid eye movement sleep period.

Liu H, Chen G, Pan Y, Chen Z, Jin W, Sun C, Chen C, Dong X, Chen K, Xu Z, Zhang S, Yu Y. (2014) Occupational Electromagnetic Field Exposures Associated with Sleep Quality: A Cross-Sectional Study. *PLoS ONE* 9(10): e110825. doi:10.1371/journal.pone.0110825.

BACKGROUND: Exposure to electromagnetic field (EMF) emitted by mobile phone and other machineries concerns half the world's population and raises the problem of their impact on human health. The present study aims to explore the effects of electromagnetic field exposures on sleep quality and sleep duration among workers from electric power plant. METHODS: A cross-sectional study was conducted in an electric power plant of Zhejiang Province, China. A total of 854 participants were included in the final analysis. The detailed information of participants was obtained by trained investigators using a structured questionnaire, which including socio-demographic characteristics, lifestyle variables, sleep variables and electromagnetic exposures. Physical examination and venous blood collection were also carried out for every study subject. RESULTS: After grouping daily occupational electromagnetic exposure into three categories, subjects with long daily exposure time had a significantly higher risk of poor sleep quality in comparison to those with short daily exposure time. The adjusted odds ratios were 1.68 (95%CI: 1.18, 2.39) and 1.57 (95%CI: 1.10, 2.24) across tertiles. Additionally, among the subjects with long-term occupational exposure, the longer daily occupational time apparently increased the risk of poor sleep quality (OR (95%CI): 2.12 (1.23~3.66) in the second tertile; 1.83 (1.07~3.15) in the third tertile). There was no significant association of long-term occupational exposure duration, monthly electric fee or years of mobile-phone use with sleep quality or sleep duration. CONCLUSIONS: The findings showed that daily occupational EMF exposure was positively associated with poor sleep quality. It implies EMF exposure may damage human sleep quality rather than sleep duration.

Lebedeva NN, Sulimov AV, Sulimova OP, Korotkovskaya TI, Gailus T, Investigation of brain potentials in sleeping humans exposed to the electromagnetic field of mobile

Studies that show **Cell Phone** Health Effects

phones. Crit Rev Biomed Eng 29(1):125-133, 2001.

An investigation was made of 8-hour EEG tracings of sleeping humans exposed to the electromagnetic field of a GSM-standard mobile phone. To analyze the EEG-patterns, manual scoring, nonlinear dynamics, and spectral analysis were employed. It was found that, when human beings were exposed to the electromagnetic field of a cellular phone, their cerebral cortex biopotentials revealed an increase in the alpha-range power density as compared to the placebo experiment. It was also found that the dimension of EEG correlation dynamics and the relation of sleep stages changed under the influence of the electromagnetic field of a mobile phone.

Huber R, Graf T, Cote KA, Wittmann L, Gallmann E, Matter D, Schuderer J, Kuster N, Borbely AA, Achermann P, Exposure to pulsed high-frequency electromagnetic field during waking affects human sleep EEG. Neuroreport 11(15):3321-3325, 2000.

The aim of the study was to investigate whether the electromagnetic field (EMF) emitted by digital radiotelephone handsets affects brain physiology. Healthy, young male subjects were exposed for 30 min to EMF (900 MHz; spatial peak specific absorption rate 1 W/kg) during the waking period preceding sleep. Compared with the control condition with sham exposure, spectral power of the EEG in non-rapid eye movement sleep was increased. The maximum rise occurred in the 9.75-11.25 Hz and 12.5-13.25 Hz band during the initial part of sleep. These changes correspond to those obtained in a previous study where EMF was intermittently applied during sleep. Unilateral exposure induced no hemispheric asymmetry of EEG power. The present results demonstrate that exposure during waking modifies the EEG during subsequent sleep. Thus the changes of brain function induced by pulsed high-frequency EMF outlast the exposure period.

Regel SJ, Tinguely G, Schuderer J, Adam M, Kuster N, Landolt HP, Achermann P. Pulsed radio-frequency electromagnetic fields: dose-dependent effects on sleep, the sleep EEG and cognitive performance. J Sleep Res. 16(3):253-258, 2007.

To establish a dose-response relationship between the strength of electromagnetic fields (EMF) and previously reported effects on the brain, we investigated the influence of EMF exposure by varying the signal intensity in three experimental sessions. The head of 15 healthy male subjects was unilaterally exposed for 30 min prior to sleep to a pulse-modulated EMF (GSM handset like signal) with a 10 g-averaged peak spatial specific absorption rate of (1) 0.2 W kg⁻¹, (2) 5 W kg⁻¹, or (3) sham exposed in a double-blind, crossover design. During exposure, subjects performed two series of three computerized cognitive tasks, each presented in a fixed order [simple reaction time task, two-choice reaction time task (CRT), 1-, 2-, 3-back task]. Immediately after exposure, night-time sleep was polysomnographically recorded for 8 h. Sleep architecture was not affected by EMF exposure. Analysis of the sleep electroencephalogram (EEG) revealed a dose-dependent increase of power in the spindle frequency range in non-REM sleep. Reaction speed decelerated with increasing field intensity in the 1-back task, while accuracy in the CRT and N-back task were not affected in a dose-dependent manner. In summary, this study reveals first indications of a dose-response relationship between

Studies that show **Cell Phone** Health Effects

EMF field intensity and its effects on brain physiology as demonstrated by changes in the sleep EEG and in cognitive performance.

Pelletier A, Delanaud S, de Seze R, Bach V, Libert JP, Loos N. Does Exposure to a Radiofrequency Electromagnetic Field Modify Thermal Preference in Juvenile Rats? PLoS One. 2014 Jun 6;9(6):e99007. doi: 10.1371/journal.pone.0099007. eCollection 2014.

Some studies have shown that people living near a mobile phone base station may report sleep disturbances and discomfort. Using a rat model, we have previously shown that chronic exposure to a low-intensity radiofrequency electromagnetic field (RF-EMF) was associated with paradoxical sleep (PS) fragmentation and greater vasomotor tone in the tail. Here, we sought to establish whether sleep disturbances might result from the disturbance of thermoregulatory processes by a RF-EMF. We recorded thermal preference and sleep stage distribution in 18 young male Wistar rats. Nine animals were exposed to a low-intensity RF-EMF (900 MHz, 1 V.m⁻¹) for five weeks and nine served as non-exposed controls. Thermal preference was assessed in an experimental chamber comprising three interconnected compartments, in which the air temperatures (Ta) were set to 24°C, 28°C and 31°C. Sleep and tail skin temperature were also recorded. Our results indicated that relative to control group, exposure to RF-EMF at 31°C was associated with a significantly lower tail skin temperature (-1.6°C) which confirmed previous data. During the light period, the exposed group preferred to sleep at Ta = 31°C and the controls preferred Ta = 28°C. The mean sleep duration in exposed group was significantly greater (by 15.5%) than in control group (due in turn to a significantly greater amount of slow wave sleep (SWS, +14.6%). Similarly, frequency of SWS was greater in exposed group (by 4.9 episodes.h⁻¹). The PS did not differ significantly between the two groups. During the dark period, there were no significant intergroup differences. We conclude that RF-EMF exposure induced a shift in thermal preference towards higher temperatures. The shift in preferred temperature might result from a cold thermal sensation. The change in sleep stage distribution may involve signals from thermoreceptors in the skin. Modulation of SWS may be a protective adaptation in response to RF-EMF exposure.

HuberR, TreyerV, BorbélyAA, SchudererJ, GottseligJM, LandoltH-P, WerthE, BertholdT, KusterN, BuckA, AchermannP, Electromagnetic fields, such as those from mobile phones, alter regional cerebral blood flow and sleep and waking EEG. J Sleep Res 11: 289-295, 2002.

Usage of mobile phones is rapidly increasing, but there is limited data on the possible effects of electromagnetic field (EMF) exposure on brain physiology. We investigated the effect of EMF vs. sham control exposure on waking regional cerebral blood flow (rCBF) and on waking and sleep electroencephalogram (EEG) in humans. In Experiment 1, positron emission tomography (PET) scans were taken after unilateral head exposure to 30-min pulse-modulated 900 MHz electromagnetic field (pm-EMF). In Experiment 2, night-time sleep was polysomnographically recorded after EMF exposure. Pulse-modulated EMF exposure increased relative rCBF in the dorsolateral prefrontal cortex

Studies that show **Cell Phone** Health Effects

ipsilateral to exposure. Also, pm-EMF exposure enhanced EEG power in the alpha frequency range prior to sleep onset and in the spindle frequency range during stage 2 sleep. Exposure to EMF without pulse modulation did not enhance power in the waking or sleep EEG. We previously observed EMF effects on the sleep EEG (A. A. Borbély, R. Huber, T. Graf, B. Fuchs, E. Gallmann and P. Achermann. *Neurosci. Lett.*, 1999, 275: 207-210; R. Huber, T. Graf, K. A. Cote, L. Wittmann, E. Gallmann, D. Matter, J. Schuderer, N. Kuster, A. A. Borbély, and P. Achermann. *Neuroreport*, 2000, 11: 3321-3325), but the basis for these effects was unknown. The present results show for the first time that (1) pm-EMF alters waking rCBF and (2) pulse modulation of EMF is necessary to induce waking and sleep EEG changes. Pulse-modulated EMF exposure may provide a new, non-invasive method for modifying brain function for experimental, diagnostic and therapeutic purposes.

Huber R, Schuderer J, Graf T, Jutz K, Borbely AA, Kuster N, Achermann P. Radio frequency electromagnetic field exposure in humans: Estimation of SAR distribution in the brain, effects on sleep and heart rate. *Bioelectromagnetics* 24(4):262-276, 2003.

In two previous studies we demonstrated that radiofrequency electromagnetic fields (RF EMF) similar to those emitted by digital radiotelephone handsets affect brain physiology of healthy young subjects exposed to RF EMF (900 MHz; spatial peak specific absorption rate [SAR] 1 W/kg) either during sleep or during the waking period preceding sleep. In the first experiment, subjects were exposed intermittently during an 8 h nighttime sleep episode and in the second experiment, unilaterally for 30 min prior to a 3 h daytime sleep episode. Here we report an extended analysis of the two studies as well as the detailed dosimetry of the brain areas, including the assessment of the exposure variability and uncertainties. The latter enabled a more in depth analysis and discussion of the findings. Compared to the control condition with sham exposure, spectral power of the non-rapid eye movement sleep electroencephalogram (EEG) was initially increased in the 9-14 Hz range in both experiments. No topographical differences with respect to the effect of RF EMF exposure were observed in the two experiments. Even unilateral exposure during waking induced a similar effect in both hemispheres. Exposure during sleep reduced waking after sleep onset and affected heart rate variability. Exposure prior to sleep reduced heart rate during waking and stage 1 sleep. The lack of asymmetries in the effects on sleep EEG, independent of bi- or unilateral exposure of the cortex, may indicate involvement of subcortical bilateral projections to the cortex in the generation of brain function changes, especially since the exposure of the thalamus was similar in both experiments (approx. 0.1 W/kg).

Huber R, Treyer V, Schuderer J, Berthold T, Buck A, Kuster N, Landolt HP, Achermann P. Exposure to pulse-modulated radio frequency electromagnetic fields affects regional cerebral blood flow. *Eur J Neurosci.* 21(4):1000-1006, 2005.

We investigated the effects of radio frequency electromagnetic fields (RF EMF) similar to those emitted by mobile phones on waking regional cerebral blood flow (rCBF) in 12 healthy young men. Two types of RF EMF exposure were applied: a 'base-station-like' and a 'handset-like' signal. Positron emission tomography scans were taken after 30 min

Studies that show **Cell Phone** Health Effects

unilateral head exposure to pulse-modulated 900 MHz RF EMF (10 g tissue-averaged spatial peak-specific absorption rate of 1 W/kg for both conditions) and sham control. We observed an increase in relative rCBF in the dorsolateral prefrontal cortex on the side of exposure. The effect depended on the spectral power in the amplitude modulation of the RF carrier such that only 'handset-like' RF EMF exposure with its stronger low-frequency components but not the 'base-station-like' RF EMF exposure affected rCBF. This finding supports our previous observation that pulse modulation of RF EMF is necessary to induce changes in the waking and sleep EEG, and substantiates the notion that pulse modulation is crucial for RF EMF-induced alterations in brain physiology.

Hung CS, Anderson C, Horne JA, McEvoy P. Mobile phone 'talk-mode' signal delays EEG-determined sleep onset. *Neurosci Lett.* 421: 82-86, 2007.

Mobile phones signals are pulse-modulated microwaves, and EEG studies suggest that the extremely low-frequency (ELF) pulse modulation has sleep effects. However, 'talk', 'listen' and 'standby' modes differ in the ELF (2, 8, and 217Hz) spectral components and specific absorption rates, but no sleep study has differentiated these modes. We used a GSM900 mobile phone controlled by a base-station simulator and a test SIM card to simulate these three specific modes, transmitted at 12.5% (23dBm) of maximum power. At weekly intervals, 10 healthy young adults, sleep restricted to 6h, were randomly and single-blind exposed to one of: talk, listen, standby and sham (nil signal) modes, for 30min, at 13:30h, whilst lying in a sound-proof, lit bedroom, with a thermally insulated silent phone beside the right ear. Bipolar EEGs were recorded continuously, and subjective ratings of sleepiness obtained every 3min (before, during and after exposure). After exposure the phone and base-station were switched off, the bedroom darkened, and a 90min sleep opportunity followed. We report on sleep onset using: (i) visually scored latency to onset of stage 2 sleep, (ii) EEG power spectral analysis. There was no condition effect for subjective sleepiness. Post-exposure, sleep latency after talk mode was markedly and significantly delayed beyond listen and sham modes. This condition effect over time was also quite evident in 1-4Hz EEG frontal power, which is a frequency range particularly sensitive to sleep onset. It is possible that 2, 8, 217Hz modulation may differentially affect sleep onset.

Rusterholz T, Bersagliere A, Kuster N, Achermann P. Sleep EEG alterations: effects of different pulse-modulated radio frequency electromagnetic fields. *J Sleep Res.* 21(1):50-58, 2012.

Previous studies have observed increases in electroencephalographic power during sleep in the spindle frequency range (approximately 11-15 Hz) after exposure to mobile phone-like radio frequency electromagnetic fields (RF EMF). Results also suggest that pulse modulation of the signal is crucial to induce these effects. Nevertheless, it remains unclear which specific elements of the field are responsible for the observed changes. We investigated whether pulse-modulation frequency components in the range of sleep spindles may be involved in mediating these effects. Thirty young healthy men were exposed, at weekly intervals, to three different conditions for 30 min

Studies that show **Cell Phone** Health Effects

directly prior to an 8-h sleep period. Exposure consisted of a 900-MHz RF EMF, pulse modulated at 14 Hz or 217 Hz, and a sham control condition. Both active conditions had a peak spatial specific absorption rate of 2 W kg^{-1} . During exposure subjects performed three different cognitive tasks (measuring attention, reaction speed and working memory), which were presented in a fixed order. Electroencephalographic power in the spindle frequency range was increased during non-rapid eye movement sleep (2nd episode) following the 14-Hz pulse-modulated condition. A similar but non-significant increase was also observed following the 217-Hz pulse-modulated condition. Importantly, this exposure-induced effect showed considerable individual variability. Regarding cognitive performance, no clear exposure-related effects were seen. Consistent with previous findings, our results provide further evidence that pulse-modulated RF EMF alter brain physiology, although the time-course of the effect remains variable across studies. Additionally, we demonstrated that modulation frequency components within a physiological range may be sufficient to induce these effects.

Schmid MR, Loughran SP, Regel SJ, Murbach M, Bratic Grunauer A, Schmid MR, Murbach M, Lustenberger C, Maire M, Kuster N, Achermann P, Loughran SP. Sleep EEG alterations: effects of pulsed magnetic fields versus pulse-modulated radio frequency electromagnetic fields. *J Sleep Res.*21(6):620-629, 2012.

Studies have repeatedly shown that electroencephalographic power during sleep is enhanced in the spindle frequency range following radio frequency electromagnetic field exposures pulse-modulated with fundamental frequency components of 2, 8, 14 or 217 Hz and combinations of these. However, signals used in previous studies also had significant harmonic components above 20 Hz. The current study aimed: (i) to determine if modulation components above 20 Hz, in combination with radio frequency, are necessary to alter the electroencephalogram; and (ii) to test the demodulation hypothesis, if the same effects occur after magnetic field exposure with the same pulse sequence used in the pulse-modulated radio frequency exposure. In a randomized double-blind crossover design, 25 young healthy men were exposed at weekly intervals to three different conditions for 30 min before sleep. Cognitive tasks were also performed during exposure. The conditions were a 2-Hz pulse-modulated radio frequency field, a 2-Hz pulsed magnetic field, and sham. Radio frequency exposure increased electroencephalogram power in the spindle frequency range. Furthermore, delta and theta activity (non-rapid eye movement sleep), and alpha and delta activity (rapid eye movement sleep) were affected following both exposure conditions. No effect on sleep architecture and no clear impact of exposure on cognition was observed. These results demonstrate that both pulse-modulated radio frequency and pulsed magnetic fields affect brain physiology, and the presence of significant frequency components above 20 Hz are not fundamental for these effects to occur. Because responses were not identical for all exposures, the study does not support the hypothesis that effects of radio frequency exposure are based on demodulation of the signal only.

Studies that show **Cell Phone** Health Effects

Effects On Skin

Pacini S, Ruggiero M, Sardi I, Aterini S, Gulisano F, Gulisano M. Exposure to global system for mobile communication (GSM) cellular phone radiofrequency alters gene expression, proliferation, and morphology of human skin fibroblasts. *Oncol Res* 13(1):19-24, 2002.

Human skin fibroblasts were exposed to global system for mobile communication (GSM) cellular phone radiofrequency for 1 h. GSM exposure induced alterations in cell morphology and increased the expression of mitogenic signal transduction genes (e.g., MAP kinase kinase 3, G2/mitotic-specific cyclin G1), cell growth inhibitors (e.g., transforming growth factor-beta), and genes controlling apoptosis (e.g., bax). A significant increase in DNA synthesis and intracellular mitogenic second messenger formation matched the high expression of MAP kinase family genes. These findings show that these electromagnetic fields have significant biological effects on human skin fibroblasts.

Strobos MA, Coenraads PJ, De Jongste MJ, Ubels FL. Dermatitis caused by radio-frequency electromagnetic radiation. *Contact Dermatitis* 44(5):309, 2001.

A case report of a woman who developed dermatitis to a transmitter placed on the abdomen that sent radio waves to an implanted neurostimulatory receiver for angina. Patch tests with the plastic, rubber and glue of the transmitter were negative, as well as those with various components of the device from the manufacturer. She had skin symptoms only after starting stimulation, with spontaneous improvement in between times.

Ozguner F, Aydin G, Mollaoglu H, Gokalp O, Koyu A, Cesur G. Prevention of mobile phone induced skin tissue changes by melatonin in rat: an experimental study. *Toxicol Ind Health*. 20(6-10):133-139, 2004.

Most of the mobile phones in Turkey emit 900 MHz radiation which is mainly absorbed by the skin and, to a lesser extent, muscle. The aim of this study was to investigate the effects the 900 MHz electromagnetic irradiation emitted by these devices on the induction of histopathologic changes in skin and the effect of melatonin (Mel) on any of these changes. Thirty male Wistar-Albino rats were used in the study. The experimental groups were composed of: a nontreated control group, an irradiated group (IR) without Mel and an irradiated with Mel treatment group (IR + Mel). 900 MHz radiation was applied to IR group for 10 days (30 min/day). The IR + Mel group received 10 mg/kg per day melatonin in tap water for 10 days before irradiation. At the end of the tenth day, the skin graft was excized from the thoraco-abdominal area. Histopathologic changes in skin were analyzed. In the IR group, increased thickness of stratum corneum, atrophy of epidermis, papillomatosis, basal cell proliferation, increased granular cell layer (hypergranulosis) in epidermis and capillary proliferation, impairment in collagen tissue distribution and separation of collagen bundles in dermis were all observed compared to the control group. Most of these changes, except hypergranulosis, were prevented

Studies that show **Cell Phone** Health Effects

with melatonin treatment. In conclusion, exposure to 900 MHz radiation emitted by mobile phones caused mild skin changes. Furthermore, melatonin treatment can reduce these changes and may have a beneficial effect to prevent 900 MHz mobile phone-induced rat skin changes.

Nam KC, Kim SW, Kim SC, Kim DW. Effects of RF exposure of teenagers and adults by CDMA cellular phones. Bioelectromagnetics. 27(7):509-514, 2006.

Many cellular phone provocation studies have been conducted since the question of increased health risk from extended usage of cellular phones became a social issue. Internationally, most studies have been conducted regarding the effects of GSM cellular phones on blood pressure and heart rate of adult volunteers. On the other hand, very few provocation studies have been conducted regarding the physiological effects of CDMA phones on teenagers. In this study, two volunteer groups consisting of 21 teenagers and 21 adults were exposed to 300 mW of radio frequency (RF) electromagnetic field emitted by a CDMA cellular phone for half an hour. Physiological parameters such as systolic and diastolic blood pressures, heart rate, respiration rate, and skin resistance were simultaneously measured. All the parameters for both groups were unaffected during the exposure except for decreased skin resistance of the teenager group ($P < .0001$). For the regrouped 23 male and 19 female subjects, all the parameters for both groups were unaffected during the exposure except for decreased skin resistance of the male subjects ($P = .0026$). Those resistances at 10 min after the terminated exposure returned to the resistances at rest regardless of the different groups of age and sex.

Monfrecola G, Moffa G, Procaccini EM. Non-ionizing electromagnetic radiations, emitted by a cellular phone, modify cutaneous blood flow. Dermatology. 207(1):10-14, 2003.

BACKGROUND: Our surroundings are full of non-ionizing electromagnetic radiation (EMR) of different frequency and power. The non-ionizing EMRs emitted by television, computer and cellular phone (CF) sets have been increasing over the past few years. OBJECTIVE: The aim of our study was to assess the effects of non-ionizing EMRs (frequency 3×10^8 to 3×10^{11} Hz), emitted by CFs, on cutaneous blood flow in healthy volunteers. METHODS: Thirty healthy volunteers (14 male and 16 female; age: 18-53 years) entered the study. Measurements of cutaneous blood flow were taken under standard conditions (temperature and humidity), using a laser Doppler He-Ne flowmeter that was applied to the ear skin by an optical fibre probe. Microflow values were recorded without CF contact with the skin (T0), with the CF turned off but in contact with the ear skin (T1), with CF contact and turned on (T2), with CF contact, turned on and receiving (T3). The microflow values were also recorded backwards: with CF contact and set turned on (T4), with CF contact and turned off (T5), without CF contact (T6). RESULTS: The mean value of basal microflow (T0), expressed as perfusion units (PU), was 51.26 ± 11.93 PU. During the T1 phase, the microflow increase was 61.38%; in T2 it was 131.74%, in T3 157.67%, in T4 139.21% and in T5 122.90%; in T6, the microflow value was 57.58 ± 10 PU (similar to the basal microflow). Statistically

Studies that show **Cell Phone** Health Effects

significant cutaneous microflow values ($p < 0.050$) were observed comparing the T1 to T5 values with basal microflow (T0). Furthermore, in comparison with T1 values (CF turned off in contact with the ear skin), the T2, T3 and T4 data were statistically significant (T2 vs. T1: $t = 7.763$ with $p < 0.050$; T3 vs. T1: $t = 9.834$ with $p < 0.050$; T4 vs. T1: $t = 8.885$ with $p < 0.050$).

Simon D, Daubos A, Pain C, Fitoussi R, Vié K, Taieb A, de Benetti L, Cario-André M. Exposure to acute electromagnetic radiation of mobile phone exposure range alters transiently skin homeostasis of a model of pigmented reconstructed epidermis. *Int J Cosmet Sci.* 35(1):27-34, 2013.

Exposure to electromagnetic radiations (EMR) produced by mobile phone concerns half the world's population and raises the problem of their impact on human health. In this study, we looked at the effects of mobile phone exposure (GSM basic, 900MHz, SAR 2 mW g⁻¹), 6 h) on a model of pigmented skin. We have analysed the expression and localization of various markers of keratinocyte and melanocyte differentiation 2, 6, 18 and 24 h after EMR exposure of reconstructed epidermis containing either only keratinocytes or a combination of keratinocytes and melanocytes grown on dead de-epidermized dermis, using histology, immunohistochemistry and Western blot. No changes were found in epidermal architecture, localization of epidermal markers, presence of apoptotic cells and the induction of p53 in both types of epidermis (with or without melanocytes) after exposure to EMR. In pigmented reconstructs, no change in the location and dendricity of melanocytes and in melanin transfer to neighbouring keratinocytes was detected after EMR exposure. Loricrin, cytokeratin 14 were significantly decreased at 6 h. The level of all markers increased at 24 h as compared to 6 h post-EMR exposure, associated with a significant decrease of the 20S proteasome activity. Our data indicate that exposure to 900MHz frequency induces a transient alteration of epidermal homeostasis, which may alter the protective capacity of the skin against external factors. Presence or absence of melanocytes did not modify the behaviour of reconstructs after EMR exposure.

Seishima M, Oyama Z, Oda M. Cellular phone dermatitis with chromate allergy. *Dermatology.* 207(1):48-50, 2003.

BACKGROUND: A patient with allergic contact dermatitis caused by hexavalent chromium plating on a cellular phone has already been reported. **OBJECTIVES:** This study described the clinical characteristics and results of patch tests in 8 patients with contact dermatitis possibly caused by handling a cellular phone. **PATIENTS:** The 8 patients were 4 males and 4 females aged from 14 to 54 years. They each noticed skin eruptions after 9-25 days of using a cellular phone. All patients had erythema, and 7 had papules on the hemilateral auricle or in the preauricular region. Three of 8 patients had a history of metal allergy. Chromate, aluminium and acrylonitrile-butadiene-styrene copolymer were used as plating on the cellular phones used by these patients. **METHODS:** Closed patch tests and photopatch tests were performed using metal standard antigens. **RESULTS:** The patch test was positive for 0.5, 0.1 and 0.05%

Studies that show **Cell Phone** Health Effects

potassium dichromate in all 8 patients. The photopatch test showed the same results. One patient was positive for 2% cobalt chloride and one for 5% nickel sulfate.

CONCLUSION: It is important to consider the possibility of contact dermatitis due to a cellular phone, possibly caused by chromate, when the patients have erythema and papules on the hemilateral auricle or in the preauricular region.

Loos N, György T, Ghosn R, Brenet-Dufour V, Liabeuf S, Selmaoui B, Jean-Pierre L, Bach V, Diouf M, de Seze R. Is the effect of mobile phone radiofrequency waves on human skin perfusion non-thermal? Microcirculation. 2013 Apr 17. doi: 10.1111/micc.12062. [Epub ahead of print]

OBJECTIVE: to establish whether skin micro blood flow can be modified by exposure to the radiofrequency waves emitted by a mobile phone when the latter is held against the jaw and ear. METHODS: Variations in skin micro blood flow and skin temperature in adult volunteers were simultaneously recorded with a thermostatic laser Doppler system during a 20-minute "radiofrequency" exposure session and a 20-minute "sham" session. The skin microvessels' vasodilatory reserve was assessed with a heat challenge at the end of the protocol. RESULTS: During the radiofrequency exposure session, skin micro blood flow increased (vs. baseline) more than during the sham exposure session. The sessions did not differ significant in terms of the skin temperature time-course response. The skin microvessels' vasodilatory ability was found to be greater during radiofrequency exposure than during sham exposure. CONCLUSIONS: Our results reveal the existence of a specific vasodilatory effect of mobile phone radiofrequency emission from mobile phones.

Karinen A, Heinavaara S, Nylund R, Leszczynski D. Mobile phone radiation might alter protein expression in human skin. BMC Genomics. 9(1):77, 2008.

ABSTRACT: BACKGROUND: Earlier we have shown that the mobile phone radiation (radiofrequency modulated electromagnetic fields; RF-EMF) alters protein expression in human endothelial cell line. This does not mean that similar response will take place in human body exposed to this radiation. Therefore, in this pilot human volunteer study, using proteomics approach, we have examined whether a local exposure of human skin to RF-EMF will cause changes in protein expression in living people. RESULTS: Small area of forearm's skin in 10 female volunteers was exposed to RF-EMF (specific absorption rate SAR=1.3W/kg) and punch biopsies were collected from exposed and non exposed areas of skin. Proteins extracted from biopsies were separated using 2-DE and protein expression changes were analyzed using PDQuest software. Analysis has identified 8 proteins that were statistically significantly affected (Anova and Wilcoxon tests). Two of the proteins were present in all 10 volunteers. This suggests that protein expression in human skin might be affected by the exposure to RF-EMF. The number of affected proteins was similar to the number of affected proteins observed in our earlier in vitro studies. CONCLUSIONS: This is the first study showing that molecular level changes might take place in human volunteers in response to exposure to RF-EMF. Our study confirms that proteomics screening approach can identify protein targets of RF-EMF in human volunteers.

Studies that show **Cell Phone** Health Effects

Sanchez S, Milochau A, Ruffie G, Poullietier de Gannes F, Lagroye I, Haro E, Surleve-Bazeille JE, Billaudel B, Lassegues M, Veyret B. Human skin cell stress response to GSM-900 mobile phone signals. *FEBS J.* 273(24):5491-5507, 2006.

In recent years, possible health hazards due to radiofrequency radiation (RFR) emitted by mobile phones have been investigated. Because several publications have suggested that RFR is stressful, we explored the potential biological effects of Global System for Mobile phone communication at 900 MHz (GSM-900) exposure on cultures of isolated human skin cells and human reconstructed epidermis (hRE) using human keratinocytes. As cell stress markers, we studied Hsc70, Hsp27 and Hsp70 heat shock protein (HSP) expression and epidermis thickness, as well as cell proliferation and apoptosis. Cells were exposed to GSM-900 under optimal culture conditions, for 48 h, using a specific absorption rate (SAR) of 2 W.kg⁻¹. This SAR level represents the recommended limit for local exposure to a mobile phone. The various biological parameters were analysed immediately after exposure. Apoptosis was not induced in isolated cells and there was no alteration in hRE thickness or proliferation. No change in HSP expression was observed in isolated keratinocytes. By contrast, a slight but significant increase in Hsp70 expression was observed in hREs after 3 and 5 weeks of culture. Moreover, fibroblasts showed a significant decrease in Hsc70, depending on the culture conditions. These results suggest that adaptive cell behaviour in response to RFR exposure, depending on the cell type and culture conditions, is unlikely to have deleterious effects at the skin level.

Allergies Effects

Kimata H. Enhancement of allergic skin wheal responses by microwave radiation from mobile phones in patients with atopic eczema/dermatitis syndrome. *Int Arch Allergy Immunol* 129(4):348-350, 2002.

Microwave radiation from mobile phones enhanced skin wheal responses induced by house dust mite and Japanese cedar pollen while it had no effect on wheal responses induced by histamine in patients with atopic eczema/dermatitis syndrome (AEDS). Microwave radiation also increased plasma levels of substance P (SP) and vasoactive intestinal peptide (VIP) in patients with AEDS. These results indicate that microwave radiation from mobile phones may enhance allergen-induced wheal responses in association with the release of SP and VIP. This finding may be useful in elucidating the pathophysiology and treatment of AEDS.

Kimata H. Enhancement of allergic skin wheal responses in patients with atopic eczema/dermatitis syndrome by playing video games or by a frequently ringing mobile phone. *Eur J Clin Invest.* 33(6):513-517, 2003.

BACKGROUND: Playing video games causes physical and psychological stress, including increased heart rate and blood pressure and aggression-related feelings. Use of mobile phones is very popular in Japan, and frequent ringing is a common and intrusive part of

Studies that show **Cell Phone** Health Effects

Japanese life. Atopic eczema/dermatitis syndrome is often exacerbated by stress. Stress increases serum IgE levels, skews cytokine pattern towards Th2 type, enhances allergen-induced skin wheal responses, and triggers mast cell degranulation via substance P, vasoactive intestinal peptide and nerve growth factor. **MATERIALS AND METHODS:** (1) In the video game study, normal subjects (n = 25), patients with allergic rhinitis (n = 25) or atopic eczema/dermatitis syndrome (n = 25) played a video game (STREET FIGHTER II) for 2 h. Before and after the study, allergen-induced wheal responses, plasma levels of substance P, vasoactive intestinal peptide and nerve growth factor, and in vitro production of total IgE, anti-house dust mite IgE and cytokines were measured. (2) In the mobile phone study, normal subjects (n = 27), patients with allergic rhinitis (n = 27) or atopic eczema/dermatitis syndrome (n = 27) were exposed to 30 incidences of ringing mobile phones during 30 min. Before and after the study, allergen-induced wheal responses, plasma levels of substance P, vasoactive intestinal peptide and nerve growth factor were measured. **RESULTS:** Playing video games had no effect on the normal subjects or the patients with allergic rhinitis. In contrast, playing video games significantly enhanced allergen-induced skin wheal responses and increased plasma levels of substance P, vasoactive intestinal peptide and nerve growth factors in the patients with atopic eczema/dermatitis syndrome. Moreover, playing video games enhanced in vitro production of total IgE and anti-house dust mite IgE with concomitant increased production of IL-4, IL-10 and IL-13 and decreased production of IFN-gamma and IL-12 in the patients with atopic eczema/dermatitis syndrome. However, exposure to frequently ringing mobile phones significantly enhanced allergen-induced skin wheal responses, plasma levels of substance P, vasoactive intestinal peptide and nerve growth factors in the patients with atopic eczema/dermatitis syndrome, but not in the normal subjects or the patients with allergic rhinitis. **CONCLUSION:** Playing video games enhanced allergic responses with a concomitant increased release of substance P, vasoactive intestinal peptide and nerve growth factor, and skewing of the cytokine pattern toward Th2 type in the patients with atopic eczema/dermatitis syndrome. In addition, exposure to frequently ringing mobile phones also enhanced allergic responses with a concomitant increased release of substance P, vasoactive intestinal peptide and nerve growth factor. Collectively, high technology causes stress, which in turn may aggravate symptoms of atopic eczema/dermatitis syndrome.

Kimata H. Laughter counteracts enhancement of plasma neurotrophin levels and allergic skin wheal responses by mobile phone-mediated stress. Behav Med. 29(4):149-152, 2004.

Laughter caused by viewing a comic video (Rowan Atkinson's The Best Bits of Mr. Bean) reduced the plasma nerve growth factor, neurotrophin-3 levels, and allergic skin wheal responses in patients with atopic dermatitis, whereas viewing a nonhumorous video (weather information) failed to do so. In contrast, stress induced by writing mail on a mobile phone enhanced the plasma nerve growth factor, neurotrophin-3 levels, and allergic skin wheal responses. However, previewing the comic video counteracted mobile phone-mediated enhancement of plasma neurotrophins or allergic skin wheal responses, whereas previewing the weather information failed to do so. Taken together,

Studies that show **Cell Phone** Health Effects

these results suggest that, in patients with atopic dermatitis, writing mail on a mobile phone causes stress and enhances allergic responses with a concomitant increase in plasma neurotrophins that are counteracted by laughter. These results may be useful in the study of pathophysiology and treatment of atopic dermatitis.

Kimata H. Microwave radiation from cellular phones increases allergen-specific IgE production. Allergy 60(6):838-839, 2005. (no abstract available).

Effects on Critical Organs

Ozgur E, Sahin D, Tomruk A, Guler G, Sepici-Dinçel A, Altan N, Seyhan N. The Effects of N-acetyl-L-cysteine and Epigallocatechin-3-gallate on Liver Tissue Protein Oxidation and Antioxidant Enzyme Levels After the Exposure to Radio Frequency Radiation. Int J Radiat Biol. 2014 Sep 24:1-19. [Epub ahead of print]

PURPOSE: The widespread and sustained use of mobile and cordless phones causes unprecedented increase of radiofrequency radiation (RFR). The aim of this experimental study was to investigate the effect of 900 MHz Global System for Mobile Communications (GSM) modulated RFR (average whole body Specific Absorption Rate (SAR) of 0,4 W/kg, 10 or 20 min daily for consecutive 7 days) to the liver tissue of guinea pigs and the protective effects of antioxidant treatments. **MATERIALS and METHODS:** Adult male guinea pigs were randomly divided into nine groups as; Group I (Sham/saline), Group II (Sham/EGCG), Group III (Sham/NAC), Group IV (10-min RF-exposure/saline), Group V (20-min RF-exposure/saline), Group VI (10-min RF-exposure/EGCG), Group VII (20-min RF-exposure/EGCG), Group VIII (10-min RF-exposure/NAC), Group IX (20-min RF-exposure/NAC). Protein oxidation (PCO), advanced oxidation protein products (AOPP) and antioxidant enzyme activities of superoxide dismutase (SOD) were evaluated after the exposure and the treatments with N-acetylcysteine (NAC) and (-)-epigallocatechin-3-gallate (EGCG). **RESULTS and CONCLUSIONS:** Significant decreases in the activities of SOD were observed in the liver of guinea pigs after RFR exposure. Protein damage did not change due to RFR exposure. On the other hand, only NAC treatment induces increase PCO levels, whereas EGCG treatment alone elevated the level of AOPP. Due to antioxidants have pro-oxidant behavior, the well decided doses and treatment time tables of NAC and EGCG is needed.

Ozgur E, Güler G, Seyhan N. Mobile phone radiation-induced free radical damage in the liver is inhibited by the antioxidants n-acetyl cysteine and epigallocatechin-gallate. Int J Radiat Biol.86(11):935-945, 2010.

Purpose: To investigate oxidative damage and antioxidant enzyme status in the liver of guinea pigs exposed to mobile phone-like radiofrequency radiation (RFR) and the potential protective effects of N-acetyl cysteine (NAC) and epigallocatechin-gallate (EGCG) on the oxidative damage. **Materials and methods:** Nine groups of guinea pigs were used to study the effects of exposure to an 1800-MHz Global System for Mobile Communications (GSM)-modulated signal (average whole body Specific Absorption Rate

Studies that show **Cell Phone** Health Effects

(SAR) of 0.38 W/kg, 10 or 20 min per day for seven days) and treatment with antioxidants. Results: Significant increases in malondialdehyde (MDA) and total nitric oxide (NO(x)) levels and decreases in activities of superoxide dismutase (SOD), myeloperoxidase (MPO) and glutathione peroxidase (GSH-Px) were observed in the liver of guinea pigs after RFR exposure. Only NAC treatment induces increase in hepatic GSH-Px activities, whereas EGCG treatment alone attenuated MDA level. Extent of oxidative damage was found to be proportional to the duration of exposure ($P < 0.05$).

Conclusion: Mobile phone-like radiation induces oxidative damage and changes the activities of antioxidant enzymes in the liver. The adverse effect of RFR may be related to the duration of mobile phone use. NAC and EGCG protect the liver tissue against the RFR-induced oxidative damage and enhance antioxidant enzyme activities.

Devrim E, Ergüder IB, Kılıçoğlu B, Yaykaşlı E, Cetin R, Durak I. Effects of electromagnetic radiation use on oxidant/antioxidant status and dna turn-over enzyme activities in erythrocytes and heart, kidney, liver, and ovary tissues from rats: possible protective role of Vitamin C. Toxicol Mech Methods.18(9):679-683, 2008.

ABSTRACT In this study, the aim was to investigate possible effects of Electromagnetic Radiation (EMR) use on oxidant and antioxidant status in erythrocytes and kidney, heart, liver, and ovary tissues from rats, and possible protective role of vitamin C. For this aim, 40 Wistar albino female rats were used throughout the study. The treatment group was exposed to EMR in a frequency of 900 MHz, the EMR plus vitamin C group was exposed to the same EMR frequency and given vitamin C (250 mg/kg/day) orally for 4 weeks. There were 10 animals in each group including control and vitamin C groups. At the end of the study period, blood samples were obtained from the animals to get erythrocyte sediments. Then the animals were sacrificed and heart, kidney, liver, and ovary tissues were removed. Malondialdehyde (MDA) levels and superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GSH-Px), xanthine oxidase (XO), and adenosine deaminase (ADA) enzyme activities were measured in the tissues and erythrocytes. It was observed that MDA level, XO, and GSH-Px activities significantly increased in the EMR group as compared with those of the control group in the erythrocytes. In the kidney tissues, it was found that MDA level and CAT activity significantly increased, whereas XO and ADA activities decreased in the cellular phone group as compared with those of the control group. However, in the heart tissues it was observed that MDA level, ADA, and XO activities significantly decreased in the cellular phone group as compared with those of the control group. The results suggest that EMR at the frequency generated by a cell phone causes oxidative stress and peroxidation in the erythrocytes and kidney tissues from rats. In the erythrocytes, vitamin C seems to make partial protection against the oxidant stress.

Ozguner F, Oktem F, Armagan A, Yilmaz R, Koyu A, Demirel R, Vural H, Uz E. Comparative analysis of the protective effects of melatonin and caffeic acid phenethyl ester (CAPE) on mobile phone-induced renal impairment in rat. Mol Cell Biochem. 276(1-2):31-37, 2005.

Melatonin and caffeic acid phenethyl ester (CAPE), a component of honeybee propolis,

Studies that show **Cell Phone** Health Effects

were recently found to be potent free radical scavengers and antioxidants. There are a number of reports on the effects induced by electromagnetic radiation (EMR) in various cellular systems. Mechanisms of adverse effects of EMR indicate that reactive oxygen species may play a role in the biological effects of this radiation. The present study was carried out to compare the protective effects of melatonin and CAPE against 900 MHz EMR emitted mobile phone-induced renal tubular injury. Melatonin was administered whereas CAPE was given for 10 days before the exposure. Urinary N-acetyl-beta-D-glucosaminidase (NAG, a marker of renal tubular injury) and malondialdehyde (MDA, an index of lipid peroxidation), were used as markers of oxidative stress-induced renal impairment in rats exposed to EMR. Superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GSH-Px) activities were studied to evaluate the changes of antioxidant status in renal tissue. Urinary NAG and renal MDA were increased in EMR exposed rats while both melatonin and CAPE caused a significant reduction in the levels of these parameters. Likewise, renal SOD and GSH-Px activities were decreased in EMR exposed animals while melatonin caused a significant increase in the activities of these antioxidant enzymes but CAPE did not. Melatonin caused a significant decrease in urinary NAG activity and MDA levels which were increased because of EMR exposure. CAPE also reduced elevated MDA levels in EMR exposed renal tissue, but the effect of melatonin was more potent than that of CAPE. Furthermore, treatment of EMR exposed rats with melatonin increased activities of SOD and GSH-Px to higher levels than those of control rats. In conclusion, melatonin and CAPE prevent renal tubular injury by reducing oxidative stress and protect the kidney from oxidative damage induced by 900 MHz mobile phone. Nevertheless, melatonin seems to be a more potent antioxidant compared with CAPE in kidney.

Ozguner F, Oktem F, Ayata A, Koyu A, Yilmaz HR. A novel antioxidant agent caffeic acid phenethyl ester prevents long-term mobile phone exposure-induced renal impairment in rat. Prognostic value of malondialdehyde, N-acetyl-beta-D-glucosaminidase and nitric oxide determination. Mol Cell Biochem. 277(1-2):73-80, 2005.

Caffeic acid phenethyl ester (CAPE), a flavonoid like compound, is one of the major components of honeybee propolis. It has been used in folk medicine for many years in Middle East countries. It was found to be a potent free radical scavenger and antioxidant recently. The aim of this study was to examine long-term applied 900 MHz emitting mobile phone-induced oxidative stress that promotes production of reactive oxygen species (ROS) and, was to investigate the role of CAPE on kidney tissue against the possible electromagnetic radiation (EMR)-induced renal impairment in rats. In particular, the ROS such as superoxide and nitric oxide (NO) may contribute to the pathophysiology of EMR-induced renal impairment. Malondialdehyde (MDA, an index of lipid peroxidation) levels, urinary N-acetyl-beta-D-glucosaminidase (NAG, a marker of renal tubular injury) and nitric oxide (NO, an oxidant product) levels were used as markers of oxidative stress-induced renal impairment and the success of CAPE treatment. The activities of superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GSH-Px) in renal tissue were determined to evaluate the changes of antioxidant status. The rats used in the study were randomly grouped (10 each) as

Studies that show **Cell Phone** Health Effects

follows: i) Control group (without stress and EMR), ii) Sham-operated rats stayed without exposure to EMR (exposure device off), iii) Rats exposed to 900 MHz EMR (EMR group), and iv) A 900 MHz EMR exposed + CAPE treated group (EMR + CAPE group). In the EMR exposed group, while tissue MDA, NO levels and urinary NAG levels increased ($p < 0.0001$), the activities of SOD, CAT, and GSH-Px in renal tissue were reduced ($p < 0.001$). CAPE treatment reversed these effects as well ($p < 0.0001$, $p < 0.001$ respectively). In conclusion, the increase in NO and MDA levels of renal tissue, and in urinary NAG with the decrease in renal SOD, CAT, GSH-Px activities demonstrate the role of oxidative mechanisms in 900 MHz mobile phone-induced renal tissue damage, and CAPE, via its free radical scavenging and antioxidant properties, ameliorates oxidative renal damage. These results strongly suggest that CAPE exhibits a protective effect on mobile phone-induced and free radical mediated oxidative renal impairment in rats.

Oktem F, Ozguner F, Mollaoglu H, Koyu A, Uz E. Oxidative damage in the kidney induced by 900-MHz-emitted mobile phone: protection by melatonin. Arch Med Res.36(4):350-355, 2005.

BACKGROUND: The mobile phones emitting 900-MHz electromagnetic radiation (EMR) may be mainly absorbed by kidneys because they are often carried in belts. Melatonin, the chief secretory product of the pineal gland, was recently found to be a potent free radical scavenger and antioxidant. The aim of this study was to examine 900-MHz mobile phone-induced oxidative stress that promotes production of reactive oxygen species (ROS) on renal tubular damage and the role of melatonin on kidney tissue against possible oxidative damage in rats. METHODS: The animals were randomly grouped as follows: 1) sham-operated control group and 2) study groups: i) 900-MHz EMR exposed (30 min/day for 10 days) group and ii) 900-MHz EMR exposed+melatonin (100 $\mu\text{g kg}^{-1}$) s.c. before the daily EMR exposure) treated group. Malondialdehyde (MDA), an index of lipid peroxidation, and urine N-acetyl-beta-d-glucosaminidase (NAG), a marker of renal tubular damage were used as markers of oxidative stress-induced renal impairment. Superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GSH-Px) activities were studied to evaluate the changes of antioxidant status. RESULTS: In the EMR-exposed group, while tissue MDA and urine NAG levels increased, SOD, CAT, and GSH-Px activities were reduced. Melatonin treatment reversed these effects as well. In this study, the increase in MDA levels of renal tissue and in urine NAG and also the decrease in renal SOD, CAT, GSH-Px activities demonstrated the role of oxidative mechanism induced by 900-MHz mobile phone exposure, and melatonin, via its free radical scavenging and antioxidant properties, ameliorated oxidative tissue injury in rat kidney. CONCLUSIONS: These results show that melatonin may exhibit a protective effect on mobile phone-induced renal impairment in rats.

Mugunthan N, Anbalagan J, Meenachi S, Samy AS. EXPOSURE OF MICE TO 900 - 1900 MHZ RADIATIONS FROM CELL PHONE RESULTING IN MICROSCOPIC CHANGES IN THE KIDNEY. IJCRR. 6(16): 44-49, 2014

Objective: The study was to evaluate possible effects of chronic exposure to 900 - 1900

Studies that show **Cell Phone** Health Effects

MHz radiations emitted from 2G cell phone on kidney of mice at the histological level. Methods: Mice were exposed to 2G ultra-high frequency radiation, 48 minutes per day for a period of 30 to 180 days. The amount of electromagnetic field (EMF) exposed was measured by radiation frequency meter. The sham control mice were subject to similar conditions without 2G exposure. Six animals each were sacrificed at the end of 30, 60, 90, 120, 150 and 180 days of exposure in the experimental group after 24 hours of last exposure. Same numbers of control animals were sacrificed on similar period. Both kidneys were harvested and processed for histomorphometric study. Kidneys size, weight and volume were measured and analysed. Kidney sections were analysed under the light microscope and structural changes were studied. Results: In 2G exposed group the kidney weight and volume was significantly reduced in the first month. Kidney weight alone was significantly increased in the fifth month. Glomerulus showed dilated capillaries and increased urinary space. Proximal convoluted tubule showed wider lumen with reduced cell size. Brush border interrupted at places and vacuolated cytoplasm and pyknotic nuclei. Wider lumen with decreased cell size and marked basal striations were found in the distal convoluted tubule. Conclusion: Chronic exposure to ultra-high frequency radiation from 2G cell phone could cause microscopic changes in glomerulus, proximal and distal convoluted tubules of the kidney.

Luo YP, Ma HR, Chen JW, Li JJ, Li CX. [Effect of American Ginseng Capsule on the liver oxidative injury and the Nrf2 protein expression in rats exposed by electromagnetic radiation of frequency of cell phone.] [Article in Chinese]. Zhongguo Zhong Xi Yi Jie He Za Zhi. 34(5):575-580, 2014. (In Chinese)

OBJECTIVE: To observe the effect of American Ginseng Capsule (AGC) on the liver oxidative injury and the Nrf2 protein expression in the liver tissue of rats exposed by 900 MHz cell phone electromagnetic radiation. METHODS: Totally 40 male SD rats were randomly divided into the normal control group, the model group, the Shuifei Jibin Capsule (SJC) group, and the AGC group, 10 in each group. Rats in the normal control group were not irradiated. Rats in the rest three groups were exposed by imitated 900 MHz cellular phone for 4 h in 12 consecutive days. Meanwhile, rats in the SJC group and the AGC group were intragastrically administrated with suspension of SJC and AGC (1 mL/200 g body weight) respectively. Normal saline was administered to rats in the normal control group and the model group. The histomorphological changes of the liver tissue were observed by HE staining. Contents of malonic dialdehyde (MDA), superoxide dismutase (SOD), glutathione (GSH), and glutathione peroxidase (GSH-PX) were detected by colorimetry. The Nrf2 protein expression of hepatocytes was detected by immunohistochemical assay and Western blot. RESULTS: Compared with the normal control group, hepatocyte nucleus was atrophied or partially disappeared, the contents of liver MDA and Nrf2 protein obviously increased ($P < 0.05$, $P < 0.01$); contents of liver SOD and GSH decreased ($P < 0.05$) in the model group. Compared with the model group, karyopyknosis was obviously attenuated and approached to the normal level in the SJC group and the AGC group. The contents of liver MDA and Nrf2 protein expression decreased ($P < 0.05$), and the contents of liver SOD, GSH, and GSH-PX

Studies that show **Cell Phone** Health Effects

obviously increased ($P < 0.05$) in the SJC group. The contents of liver MDA and the Nrf2 protein expression decreased ($P < 0.05$), and contents of SOD and GSH obviously increased in the AGC group ($P < 0.01$, $P < 0.05$). CONCLUSIONS: The electromagnetic radiation induced by 900 MHz cell phone could affect the expression of Nrf2 protein, induce oxidative injury, and induce abnormal morphology of liver cells. SJC and AGC could promote the morphological recovery of the liver cells. Its mechanism might be related to affecting the expression of Nrf2 protein and attenuating oxidative damage of liver cells.

Koca O, Gökçe AM, Oztürk MI, Ercan F, Yurdakul N, Karaman MI. Effects of intensive cell phone (philips genic 900) use on the rat kidney tissue. Urol J. 10(2):886-891. 2013.

PURPOSE: To investigate effects of electromagnetic radiation (EMR) emitted by cell phones on the rat kidney tissue. **MATERIALS AND METHODS:** Twenty-one male Albino rats were divided into 3 groups, each comprising 7 rats. Group 1 was exposed to a cell phone in speech mode for 8 hours/day for 20 days and their kidneys were removed. Group 2 was exposed to EMR for 20 days and then their kidneys were removed after an interval of 20 days. Cell phone used in the present study was Philips Genie 900, which has the highest specific absorption rate on the market. **RESULTS:** Light microscopic examination of the kidney tissues obtained from the first group of rats revealed glomerular damage, dilatation of Bowman's capsule, formation of large spaces between the tubules, tubular damage, perivascular edema, and inflammatory cell infiltration. The mean severity score was 4.64 ± 1.7 in group 1, 4.50 ± 0.8 in group 2, and 0 in group 3. While there was no significant difference between group 1 and group 2 ($P > .05$), the mean severity scores of groups 1 and 2 were significantly higher than that of the control group ($P = .001$ for each). **CONCLUSION:** Considering the damage in rat kidney tissue caused by EMR-emitting cell phones, high-risk individuals should take protective measures.

Koca O, Gokce AM, Akyuz M, Ercan F, Yurdakul N, Karaman MI. A new problem in inflammatory bladder diseases: Use of mobile phones! Int Braz J Urol. 40(4):520-525, 2014.

PURPOSE: Technological developments provide a lot of conveniences to our lives. This issue is one of the risks that arise along with these conveniences. In our study we tried to understand the impact of electromagnetic waves from mobile phones on bladder tissue. **MATERIALS AND METHODS:** Twenty-one adult male albino rats were divided into three equal groups. Group 1 was exposed to electromagnetic wave for 8 hours per day for 20 days and then their bladders were taken off immediately. Group 2 was firstly exposed to electromagnetic wave for 8 hours per day for 20 days then secondly another for 20 days without exposition to electromagnetic wave and then their bladders were taken off. Group 3 was the control group and they were not exposed to electromagnetic wave. **RESULTS:** Under microscopic examination of bladder tissue, in the first group severe inflammatory cell infiltration was seen in lamina propria and muscle layer in contrast to intact urothelium. In the second group mild inflammatory cell infiltration was seen in lamina propria and muscle layer. The mean scores for the three groups were 5.5 ± 2.5 , 0.8 ± 1.3 and 1.2 ± 1.5 respectively. Mean score of group 1 was

Studies that show Cell Phone Health Effects

statistically higher than others ($p = 0.001$). CONCLUSION: Intensive use of mobile phones has negative impact on bladder tissue as well as the other organs. Keeping a minimum level of mobile phone use makes it easy to be kept under control of diseases in which inflammation is an etiologic factor.

Mercury Release in Dental Amalgams

Mortazavi SM, Daiee E, Yazdi A, Khiabani K, Kavousi A, Vazirinejad R, Behnejad B, Ghasemi M, Mood MB. Mercury release from dental amalgam restorations after magnetic resonance imaging and following mobile phone use. Pak J Biol Sci. 11(8):1142-1146, 2008.

In the 1st phase of this study, thirty patients were investigated. Five milliliter stimulated saliva was collected just before and after MRI. The magnetic flux density was 0.23 T and the duration of exposure of patients to magnetic field was 30 minutes. In the 2nd phase, fourteen female healthy University students who had not used mobile phones before the study and did not have any previous amalgam restorations were investigated. Dental amalgam restoration was performed for all 14 students. Their urine samples were collected before amalgam restoration and at days 1, 2, 3 and 4 after restoration. The mean \pm SD saliva Hg concentrations of the patients before and after MRI were 8.6 ± 3.0 and 11.3 ± 5.3 microg L⁻¹, respectively ($p < 0.01$). A statistical significant ($p < 0.05$) higher concentration was observed in the students used mobile phone. The mean \pm SE urinary Hg concentrations of the students who used mobile phones were 2.43 ± 0.25 , 2.71 ± 0.27 , 3.79 ± 0.25 , 4.8 ± 0.27 and 4.5 ± 0.32 microg L⁻¹ before the amalgam restoration and at days 1, 2, 3 and 4, respectively. Whereas the respective Hg concentrations in the controls, were 2.07 ± 0.22 , 2.34 ± 0.30 , 2.51 ± 0.25 , 2.66 ± 0.24 and 2.76 ± 0.32 microg L⁻¹. It appears that MRI and microwave radiation emitted from mobile phones significantly release mercury from dental amalgam restoration. Further research is needed to clarify whether other common sources of electromagnetic field exposure may cause alterations in dental amalgam and accelerate the release of mercury.

Effects on Other Living Things

Panagopoulos DJ, Chavdoula ED, Karabarbounis A, Margaritis LH. Comparison of bioactivity between GSM 900 MHz and DCS 1800 MHz mobile telephony radiation. Electromagn Biol Med. 26(1):33-44, 2007.

An increasing number of studies find that pulsed Radio Frequency (RF), electromagnetic radiation of both systems of digital mobile telephony, established and commonly used in Europe during the last years, GSM 900 MHz (Global System for Mobile telecommunications) and DCS 1800 MHz (Digital Cellular System), exert intense biological action on different organisms and cells (Hardell et al., 2006; Hyland, 2000; Kundi, 2004; Panagopoulos et al., 2004, 2007). The two types of cellular telephony

Studies that show **Cell Phone** Health Effects

radiation use different carrier frequencies and give different frequency spectra, but they usually also differ in intensity, as GSM 900 MHz antennas operate at about double the power output than the corresponding DCS 1800 MHz ones. In our present experiments, we used a model biological system, the reproductive capacity of *Drosophila melanogaster*, to compare the biological activity between the two systems of cellular mobile telephony radiation. Both types of radiation were found to decrease significantly and non thermally the insect's reproductive capacity, but GSM 900 MHz seems to be even more bioactive than DCS 1800 MHz. The difference seems to be dependent mostly on field intensity and less on carrier frequency.

Zareen N, Khan MY, Minhas LA. Dose related shifts in the developmental progress of chick embryos exposed to mobile phone induced electromagnetic fields. *J Ayub Med Coll Abbottabad*.21(1):130-134, 2009.

BACKGROUND: The possible adverse effects of Electromagnetic Fields (EMFs) emitted from mobile phones present a major public concern today. Some studies indicate EMFs effects on genes, free radical production, immunological and carcinogenic effects. On the other hand there are studies which do not support the hypothesis of any biological impacts of EMFs. This study was designed to observe the effects of mobile phone induced EMFs on survival and general growth and development of chick embryo, investigating dose-response relationship if any. **METHODS:** This was an experimental study in which developing chick embryos were exposed to different doses of mobile phone induced EMFs. For this purpose a mobile phone was placed in the incubator in the centre of fertilised eggs in silent ringing mode and was 'rung' upon from any other line or cell phone. After incubation for 10 or 15 days the eggs were opened and the developmental mile-stones of the surviving embryos were compared with the non exposed subgroup. **RESULTS:** EMFs exposure significantly decreased the survivability of the chick embryos. The lower doses of EMFs caused growth retardation. However, this effect of growth retardation reallocated to partial growth enhancement on increasing the dose of EMFs and shifted over to definite growth enhancement on further raising the dose. **CONCLUSION:** There is an adverse effect of EMFs exposure on embryo survivability. Chick embryos developmental process is influenced by EMFs. However, these effects are variable depending upon the dose of EMFs exposure.

Zareen N, Khan MY, Minhas LA. Derangement of chick embryo retinal differentiation caused by radiofrequency electromagnetic fields. *Congenit Anom (Kyoto)*. 49(1):15-19, 2009.

The possible adverse effects of radiofrequency electromagnetic fields (EMF) emitted from mobile phones present a major public concern. Biological electrical activities of the human body are vulnerable to interference from oscillatory aspects of EMF, which affect fundamental cellular activities, in particular, the highly active development process of embryos. Some studies highlight the possible health hazards of EMF, while others contest the hypothesis of biological impact of EMF. The present study was designed to observe the histomorphological effects of EMF emitted by a mobile phone on the retinae of developing chicken embryos. Fertilized chicken eggs were exposed to a ringing mobile set on silent tone placed in the incubator at different ages of

Studies that show **Cell Phone Health Effects**

development. After exposure for the scheduled duration the retinae of the embryos were dissected out and processed for histological examination. The control and experimental embryos were statistically compared for retinal thickness and epithelial pigmentation grades. Contrasting effects of EMF on the retinal histomorphology were noticed, depending on the duration of exposure. The embryos exposed for 10 post-incubation days exhibited decreased retinal growth and mild pigmentation of the epithelium. Growth retardation reallocated to growth enhancement on increasing EMF exposure for 15 post-incubation days, with a shift of pigmentation grade from mild to intense. We conclude that EMF emitted by a mobile phone cause derangement of chicken embryo retinal differentiation.

Soran ML, Stan M, Niinemets U, Copolovici L. Influence of microwave frequency electromagnetic radiation on terpene emission and content in aromatic plants. J Plant Physiol. 171(15):1436-1443, 2014

Influence of environmental stress factors on both crop and wild plants of nutritional value is an important research topic. The past research has focused on rising temperatures, drought, soil salinity and toxicity, but the potential effects of increased environmental contamination by human-generated electromagnetic radiation on plants have little been studied. Here we studied the influence of microwave irradiation at bands corresponding to wireless router (WLAN) and mobile devices (GSM) on leaf anatomy, essential oil content and volatile emissions in *Petroselinum crispum*, *Apium graveolens* and *Anethum graveolens*. Microwave irradiation resulted in thinner cell walls, smaller chloroplasts and mitochondria, and enhanced emissions of volatile compounds, in particular, monoterpenes and green leaf volatiles (GLV). These effects were stronger for WLAN-frequency microwaves. Essential oil content was enhanced by GSM-frequency microwaves, but the effect of WLAN-frequency microwaves was inhibitory. There was a direct relationship between microwave-induced structural and chemical modifications of the three plant species studied. These data collectively demonstrate that human-generated microwave pollution can potentially constitute a stress to the plants.

Tkalec M, Malaric K, Pevalek-Kozlina B. Influence of 400, 900, and 1900 MHz electromagnetic fields on *Lemna minor* growth and peroxidase activity. Bioelectromagnetics. 26(3):185-193, 2005.

Increased use of radio and microwave frequencies requires investigations of their effects on living organisms. Duckweed (*Lemna minor* L.) has been commonly used as a model plant for environmental monitoring. In the present study, duckweed growth and peroxidase activity was evaluated after exposure in a Gigahertz Transversal Electromagnetic (GTEM) cell to electric fields of frequencies 400, 900, and 1900 MHz. The growth of plants exposed for 2 h to the 23 V/m electric field of 900 MHz significantly decreased in comparison with the control, while an electric field of the same strength but at 400 MHz did not have such effect. A modulated field at 900 MHz strongly inhibited the growth, while at 400 MHz modulation did not influence the growth significantly. At both frequencies a longer exposure mostly decreased the

Studies that show **Cell Phone Health Effects**

growth and the highest electric field (390 V/m) strongly inhibited the growth. Exposure of plants to lower field strength (10 V/m) for 14 h caused significant decrease at 400 and 1900 MHz while 900 MHz did not influence the growth. Peroxidase activity in exposed plants varied, depending on the exposure characteristics. Observed changes were mostly small, except in plants exposed for 2 h to 41 V/m at 900 MHz where a significant increase (41%) was found. Our results suggest that investigated electromagnetic fields (EMFs) might influence plant growth and, to some extent, peroxidase activity. However, the effects of EMFs strongly depended on the characteristics of the field exposure.

Singh HP, Sharma VP, Batish DR, Kohli RK. Cell phone electromagnetic field radiations affect rhizogenesis through impairment of biochemical processes. *Environ Monit Assess.* 184(4):1813-1821, 2012.

Indiscriminate adoption and use of cell phone technology has tremendously increased the levels of electromagnetic field radiations (EMFr) in the natural environment. It has raised the concerns among the scientists regarding the possible risks of EMFr to living organisms. However, not much has been done to assess the damage caused to plants that are continuously exposed to EMFr present in the environment. The present study investigated the biochemical mechanism of interference of 900 MHz cell phone EMFr with root formation in mung bean (*Vigna radiata* syn. *Phaseolus aureus*) hypocotyls, a model system to study rhizogenesis in plants. Cell phone EMFr enhanced the activities of proteases (by 1.52 to 2.33 times), polyphenol oxidases (by 1.5 to 4.3 times), and peroxidases (by 1.5 to 2.0 times) in mung bean hypocotyls over control. Further, EMFr enhanced malondialdehyde (an indicator of lipid peroxidation), hydrogen peroxide, and proline content, indicating a reactive oxygen species-mediated oxidative damage in hypocotyls. It was confirmed by the upregulation in the activities of antioxidant enzymes (superoxide dismutase, ascorbate peroxidase, guaiacol peroxidase, catalase, and glutathione reductase) suggesting their possible role in providing protection against EMFr-induced oxidative damage. The study concluded that cell phone radiations affect the process of rhizogenesis through biochemical alterations that manifest as oxidative damage resulting in root impairment.

Tkalec M, Malarić K, Pavlica M, Pevalek-Kozlina B, Vidaković-Cifrek Z. Effects of radiofrequency electromagnetic fields on seed germination and root meristematic cells of *Allium cepa* L. *Mutat Res.* 672(2):76-81, 2009.

The effects of exposure to radiofrequency electromagnetic fields (RF-EMFs) on seed germination, primary root growth as well as mitotic activity and mitotic aberrations in root meristematic cells were examined in *Allium cepa* L. cv. Srebrnjak Majski. Seeds were exposed for 2h to EMFs of 400 and 900MHz at field strengths of 10, 23, 41 and 120Vm(-1). The effect of longer exposure time (4h) and field modulation was investigated at 23Vm(-1) as well. Germination rate and root length did not change significantly after exposure to radiofrequency fields under any of the treatment conditions. At 900MHz, exposures to EMFs of higher field strengths (41 and 120Vm(-1)) or to modulated fields showed a significant increase of the mitotic index compared with corresponding controls, while the percentage of mitotic abnormalities increased after all

Studies that show **Cell Phone** Health Effects

exposure treatments. On the other hand, at 400MHz the mitotic index increased only after exposure to modulated EMF. At this frequency, compared with the control higher numbers of mitotic abnormalities were found after exposure to modulated EMF as well as after exposure to EMFs of higher strengths (41 and 120Vm(-1)). The types of aberration induced by the EMFs of both frequencies were quite similar, mainly consisting of lagging chromosomes, vagrants, disturbed anaphases and chromosome stickiness. Our results show that non-thermal exposure to the radiofrequency fields investigated here can induce mitotic aberrations in root meristematic cells of A. cepa. The observed effects were markedly dependent on the field frequencies applied as well as on field strength and modulation. Our findings also indicate that mitotic effects of RF-EMF could be due to impairment of the mitotic spindle.

Sharma VP, Singh HP, Batish DR, Kohli RK. Cell phone radiations affect early growth of Vigna radiata (mung bean) through biochemical alterations. Z Naturforsch C. 65(1-2):66-72, 2010.

The indiscriminate use of wireless technologies, particularly of cell phones, has increased the health risks among living organisms including plants. We investigated the impact of cell phone electromagnetic field (EMF) radiations (power density, 8.55 microW cm(-2)) on germination, early growth, proteins and carbohydrate contents, and activities of some enzymes in Vigna radiata. Cell phone EMF radiations significantly reduced the seedling length and dry weight of V radiata after exposure for 0.5, 1, 2, and 4 h. Furthermore, the contents of proteins and carbohydrates were reduced in EMF-exposed plants. However, the activities of proteases, alpha-amylases, beta-amylases, polyphenol oxidases, and peroxidases were enhanced in EMF-exposed radicles indicating their role in providing protection against EMF-induced stress. The study concludes that cell phone EMFs impair early growth of V radiata seedlings by inducing biochemical changes.

Sharma VP, Singh HP, Kohli RK, Batish DR. Mobile phone radiation inhibits Vigna radiata (mung bean) root growth by inducing oxidative stress. Sci Total Environ. 407(21):5543-7, 2009.

During the last couple of decades, there has been a tremendous increase in the use of cell phones. It has significantly added to the rapidly increasing EMF smog, an unprecedented type of pollution consisting of radiation in the environment, thereby prompting the scientists to study the effects on humans. However, not many studies have been conducted to explore the effects of cell **phone** EMFr on growth and biochemical changes in plants. We investigated whether EMFr from cell phones inhibit growth of Vigna radiata (mung bean) through induction of conventional stress responses. Effects of cell **phone** EMFr (power density: 8.55 microW cm(-2); 900 MHz band width; for 1/2, 1, 2, and 4 h) were determined by measuring the generation of reactive oxygen species (ROS) in terms of malondialdehyde and hydrogen peroxide (H₂O₂) content, root oxidizability and changes in levels of antioxidant enzymes. Our results showed that cell **phone** EMFr significantly inhibited the germination (at > or =2 h), and radicle and plumule growths (> or =1 h) in mung bean in a time-dependent

Studies that show **Cell Phone** Health Effects

manner. Further, cell **phone** EMFr enhanced MDA content (indicating lipid peroxidation), and increased H₂O₂ accumulation and root oxidizability in mung bean roots, thereby inducing oxidative stress and **cellular** damage. In response to EMFr, there was a significant upregulation in the activities of scavenging enzymes, such as superoxide dismutases, ascorbate peroxidases, guaiacol peroxidases, catalases and glutathione reductases, in mung bean roots. The study concluded that cell **phone** EMFr inhibit root growth of mung bean by inducing ROS-generated oxidative stress despite increased activities of antioxidant enzymes.

Roux D, Faure C, Bonnet P, Girard S, Ledoigt G, Davies E, Gendraud M, Paladian F, Vian A. A possible role for extra-cellular ATP in plant responses to high frequency, low amplitude electromagnetic field. to the reports of health hazards among mobile-phone users. *Plant Signal Behav.* 3(6):383-385, 2008.

In parallel to evoking the accumulation of stress-related transcripts, exposure to low level 900 MHz EMF affected the levels of ATP, the main energy molecule of the cell. Its concentration dropped rapidly (27% after 30 min) in response to EMF exposure, along with a 18% decrease in the adenylate energy charge (AEC), a good marker of cell energy status. One could interpret this decrease in ATP and AEC in a classical way, i.e., as the result of an increase in cellular energy usage, but recent work brings exciting new insights in pointing out a signalling function for ATP, especially in the stress physiology context where it could trigger both reactive oxygen species and calcium movement (this latter being involved in plant responses to EMF exposure). In this addendum, we discuss our results within this new perspective for ATP function

Roux D, Vian A, Girard S, Bonnet P, Paladian F, Davies E, Ledoigt G. High frequency (900 MHz) low amplitude (5 V m⁻¹) electromagnetic field: a genuine environmental stimulus that affects transcription, translation, calcium and energy charge in tomato. *Planta.* 227(4):883-891, 2008.

Using an especially-designed facility, the Mode Stirred Reverberation Chamber, we exposed tomato plants (*Lycopersicon esculentum* Mill. VFN8) to low level (900 MHz, 5 V m⁻¹) electromagnetic fields for a short period (10 min) and measured changes in abundance of three specific mRNA soon after exposure. Within minutes of electromagnetic stimulation, stress-related mRNA (calmodulin, calcium-dependent protein kinase and proteinase inhibitor) accumulated in a rapid, large and 3-phase manner typical of an environmental stress response. Accumulation of these transcripts into the polysomal RNA also took place (indicating that the encoded proteins were translated) but was delayed (indicating that newly-synthesized mRNA was not immediately recruited into polysomes). Transcript accumulation was maximal at normal Ca²⁺ levels and was depressed at higher Ca²⁺, especially for those encoding calcium-binding proteins. Removal of Ca²⁺ (by addition of chelating agents or Ca²⁺ channel blocker) led to total suppression of mRNA accumulation. Finally, 30 min after the electromagnetic treatment, ATP concentration and adenylate energy charge were transiently decreased, while transcript accumulation was totally prevented by application of the uncoupling reagent, CCCP. These responses occur very soon after

Studies that show **Cell Phone** Health Effects

exposure, strongly suggesting that they are the direct consequence of application of radio-frequency fields and their similarities to wound responses strongly suggests that this radiation is perceived by plants as an injurious stimulus.

Tsybulin O, Sidorik E, Brieieva O , Buchynska L, Kyrylenko S, Henshel D, IYakymenko I. GSM 900 MHz cellular phone radiation can either stimulate or depress early embryogenesis in Japanese quails depending on the duration of exposure. Int J Rad Biol. Posted online on April 11, 2013. (doi:10.3109/09553002.2013.791408)

Purpose: Our study was designed to assess the effects of low intensity radiation of a GSM (Global System for Mobile communication) 900 MHz cellular phone on early embryogenesis in dependence on the duration of exposure. *Materials and methods:* Embryos of Japanese Quails were exposed *in ovo* to GSM 900 MHz cellular phone radiation during initial 38 h of brooding or alternatively during 158 h (120 h before brooding plus initial 38 h of brooding) discontinuously with 48 sec ON (average power density $0.25 \mu\text{W}/\text{cm}^2$, specific absorption rate $3 \mu\text{W}/\text{kg}$) followed by 12 sec OFF intervals. A number of differentiated somites was assessed microscopically. Possible DNA damage evoked by irradiation was assessed by an alkaline comet assay. *Results:* Exposure to radiation from a GSM 900 MHz cellular phone led to a significantly altered number of differentiated somites. In embryos irradiated during 38 h the number of differentiated somites increased ($p < 0.001$), while in embryos irradiated during 158 h this number decreased ($p < 0.05$). The lower duration of exposure led to a significant ($p < 0.001$) decrease in a level of DNA strand breaks in cells of 38-hour embryos, while the higher duration of exposure resulted in a significant ($p < 0.001$) increase in DNA damage as compared to the control. *Conclusion:* Effects of GSM 900 MHz cellular phone radiation on early embryogenesis can be either stimulating or deleterious depending on the duration of exposure.

Tsybulin O, Sidorik E, Kyrylenko S, Henshel D, Yakymenko I. GSM 900 MHz microwave radiation affects embryo development of Japanese quails. Electromagn Biol Med. 31(1):75-86, 2012.

A wide range of non thermal biological effects of microwave radiation (MW) was revealed during the last decades. A number of reports showed evident hazardous effects of MW on embryo development in chicken. In this study, we aimed at elucidating the effects of MW emitted by a commercial model of GSM 900 MHz cell phone on embryo development in quails (*Coturnix coturnix japonica*) during both short and prolonged exposure. For that, fresh fertilized eggs were irradiated during the first 38 h or 14 days of incubation by a cell phone in "connecting" mode activated continuously through a computer system. Maximum intensity of incident radiation on the egg's surface was $0.2 \mu\text{W}/\text{cm}^2$. The irradiation led to a significant ($p < 0.001$) increase in numbers of differentiated somites in 38-hour exposed embryos and to a significant ($p < 0.05$) increase in total survival of embryos from exposed eggs after 14 days exposure. We hypothesized that observed facilitating effect was due to enhancement of metabolism in exposed embryos provoked via peroxidation mechanisms. Indeed, a level of thiobarbituric acid (TBA) reactive substances was significantly ($p < 0.05-0.001$) higher

Studies that show **Cell Phone** Health Effects

in brains and livers of hatchlings from exposed embryos. Thus, observed effects of radiation from commercial GSM 900 MHz cell phone on developing quail embryos signify a possibility for non-thermal impact of MW on embryogenesis. We suggest that the facilitating effect of low doses of irradiation on embryo development can be explained by a hormesis effect induced by reactive oxygen species (ROS). Future studies need to be done to clarify this assumption.

Tafforeau M, Verdus M-C, Norris V, White G, Demarty M, Thellier M, Ripoll C. SIMS study of the calcium-deprivation step related to epidermal meristem production induced in flax by cold shock or radiation from a GSM telephone. J Trace Microprobe Tech 20(4):611-623, 2002.

Exposing seedlings of the flax, *Linum usitatissimum* L., to a variety of weak environmental stresses plus a 2-day calcium deprivation triggers the common response of production of epidermal meristems in the hypocotyls. Here, we show that the same response was induced by a 1 min cold shock. Epidermal meristem production was also induced by a single 2-h exposure to radiation emitted at 0.9 GHz at non-thermal levels by a GSM telephone. This flax-based system is therefore well suited to studying the effects of low intensity stimuli, including those of electromagnetic radiation. To begin to determine the underlying mechanisms, in which calcium is implicated, it is desirable to analyse the changes in ions in the tissues affected. We therefore performed a Secondary Ion Mass Spectrometry (SIMS) study of the distribution of the main inorganic cations in the hypocotyl of control and calcium-deprived seedlings. This showed decreases in calcium, sodium and potassium and an increase in magnesium that did not alter substantially the overall ratio of divalent to monovalent cations.

Bastide M, Youbibier-Simoa BJ, Lebecq JC, Giaimis J. Toxicologic study of electromagnetic radiation emitted by television and video display screens and cellular telephones on chickens and mice. Indoor Built Environ 10:291-298, 2001.

The effects of continuous exposure of chick embryos and young chickens to the electromagnetic fields (EMFs) emitted by video display units (VDUs) and GSM cell phone radiation, either the whole spectrum emitted or attenuated by a copper gauze, were investigated. Permanent exposure to the EMFs radiated by a VDU was associated with significantly increased fetal loss (47-68%) and markedly depressed levels of circulating specific antibodies (IgG), corticosterone and melatonin. We have also shown that under chronic exposure conditions, GSM cell phone radiation was harmful to chick embryos, stressful for healthy mice and, in this species, synergistic with cancer insofar as it depleted stress hormones. The same pathological results were observed after substantial reduction of the microwaves radiated from the cell phone by attenuating them with a copper gauze.

Vian A, Roux D, Girard S, Bonnet P, Paladian F, Davies E, Ledoigt G. Microwave irradiation affects gene expression in plants. Plant Signal Behav. 1(2):67-70, 2006.

The physiological impact of nonionizing radiation has long been considered negligible. However, here we use a carefully calibrated stimulation system that mimics the

Studies that show **Cell Phone** Health Effects

characteristics (isotropy and homogeneity) of electromagnetic fields present in the environment to measure changes in a molecular marker (mRNA encoding the stress-related bZIP transcription factor), and show that low amplitude, short duration, 900 MHz EMF evokes the accumulation of this mRNA. Accumulation is rapid (peaking 5-15 min after stimulation) and strong (3.5-fold), and is similar to that evoked by mechanical stimulations.

Panagopoulos DJ, Chavdoula ED, Nezis IP, Margaritis LH Cell death induced by GSM 900-MHz and DCS 1800-MHz mobile telephony radiation. Mutat Res.626(1-2):69-78, 2007.

In the present study, the TUNEL (Terminal deoxynucleotide transferase dUTP Nick End Labeling) assay - a well known technique widely used for detecting fragmented DNA in various types of cells - was used to detect cell death (DNA fragmentation) in a biological model, the early and mid stages of oogenesis of the insect *Drosophila melanogaster*. The flies were exposed in vivo to either GSM 900-MHz (Global System for Mobile telecommunications) or DCS 1800-MHz (Digital Cellular System) radiation from a common digital mobile phone, for few minutes per day during the first 6 days of their adult life. The exposure conditions were similar to those to which a mobile phone user is exposed, and were determined according to previous studies of ours [D.J.

Panagopoulos, A. Karabarounis, L.H. Margaritis, Effect of GSM 900-MHz mobile phone radiation on the reproductive capacity of *D. melanogaster*, *Electromagn. Biol. Med.* 23 (1) (2004) 29-43; D.J. Panagopoulos, N. Messini, A. Karabarounis, A.L. Philippetis, L.H. Margaritis, Radio frequency electromagnetic radiation within "safety levels" alters the physiological function of insects, in: P. Kostarakis, P. Stavroulakis (Eds.), *Proceedings of the Millennium International Workshop on Biological Effects of Electromagnetic Fields*, Heraklion, Crete, Greece, October 17-20, 2000, pp. 169-175, ISBN: 960-86733-0-5; D.J. Panagopoulos, L.H. Margaritis, Effects of electromagnetic fields on the reproductive capacity of *D. melanogaster*, in: P. Stavroulakis (Ed.), *Biological Effects of Electromagnetic Fields*, Springer, 2003, pp. 545-578], which had shown a large decrease in the oviposition of the same insect caused by GSM radiation. Our present results suggest that the decrease in oviposition previously reported, is due to degeneration of large numbers of egg chambers after DNA fragmentation of their constituent cells, induced by both types of mobile telephony radiation. Induced cell death is recorded for the first time, in all types of cells constituting an egg chamber (follicle cells, nurse cells and the oocyte) and in all stages of the early and mid-oogenesis, from germarium to stage 10, during which programmed cell death does not physiologically occur. Germarium and stages 7-8 were found to be the most sensitive developmental stages also in response to electromagnetic stress induced by the GSM and DCS fields and, moreover, germarium was found to be even more sensitive than stages 7-8.

Panagopoulos DJ, Karabarounis A, Margaritis LH. Effect of gsm 900-mhz mobile phone radiation on the reproductive capacity of *drosophila melanogaster*. *Electromag Biol Med* 23:29-43, 2004.

Pulsed radio frequency, (RF), electromagnetic radiation from common GSM mobile

Studies that show **Cell Phone** Health Effects

phones, (Global System for Mobile Telecommunications) with a carrier frequency at 900 MHz, “modulated” by human voice, (speaking emission) decreases the reproductive capacity of the insect *Drosophila melanogaster* by 50%–60%, whereas the corresponding “nonmodulated” field (nonspeaking emission) decreases the reproductive capacity by 15%–20%. The insects were exposed to the near field of the mobile phone antenna for 6 min per day during the first 2–5 days of their adult lives. The GSM field is found to affect both females and males. Our results suggest that this field-radiation decreases the rate of cellular processes during gonad development in insects.

Nittby H, Moghadam MK, Sun W, Malmgren L, Eberhardt J, Persson BR, Salford LG. Analgetic effects of non-thermal GSM-1900 radiofrequency electromagnetic fields in the land snail *Helix pomatia*. *Int J Radiat Biol.* ;88(3):245-252, 2012

Abstract. Abstract Purpose: To investigate whether mobile phone radiation might affect snail nociception, employing radiofrequency (RF) electromagnetic fields (EMF) which, to our knowledge, have hitherto not been studied in a snail model. Exposure to extremely low frequency (ELF) magnetic fields has however been shown to significantly affect nociceptive responses. Materials and methods: In the present study, we exposed 29 land snails of the strain *Helix pomatia* to global system for mobile communications (GSM) EMF at 1900 MHz at the non-thermal level 48 mW/kg for 1 hour each and 29 snails were sham controls. The experiments took place during the onset of summer, with all snails being well out of hibernation. Before and after GSM or sham exposure, the snails were subjected to thermal pain by being placed on a hot plate. The reaction time for retraction from the hot plate was measured by two blinded observers. Results: Comparing the reaction pattern of each snail before and after exposure, the GSM exposed snails were less sensitive to thermal pain as compared to the sham controls, indicating that RF exposure induces a significant analgesia (Mann-Whitney $p < 0.001$). Conclusion: This study might support earlier findings, describing beneficial effects of EMF exposure upon nociception.

Markkanen A, Penttinen P, Naarala J, Pelkonen J, Sihvonen A-P, Juutilainen J. Apoptosis induced by ultraviolet radiation is enhanced by amplitude modulated radiofrequency radiation in mutant yeast cells *Bioelectromagnetics* 25:127-133, 2004.

The aim of this study was to investigate whether radiofrequency (RF) electromagnetic field (EMF) exposure affects cell death processes of yeast cells. *Saccharomyces cerevisiae* yeast cells of the strains KFY417 (wild-type) and KFY437 (*cdc48*-mutant) were exposed to 900 or 872 MHz RF fields, with or without exposure to ultraviolet (UV) radiation, and incubated simultaneously with elevated temperature (+37°C) to induce apoptosis in the *cdc48*-mutated strain. The RF exposure was carried out in a special waveguide exposure chamber where the temperature of the cell cultures can be precisely controlled. Apoptosis was analyzed using the annexin V-FITC method utilizing flow cytometry. Amplitude modulated (217 pulses per second) RF exposure significantly enhanced UV induced apoptosis in *cdc48*-mutated cells, but no effect was observed in cells exposed to unmodulated fields at identical time-average specific absorption rates

Studies that show **Cell Phone** Health Effects

(SAR, 0.4 or 3.0 W/kg). The findings suggest that amplitude modulated RF fields, together with known damaging agents, can affect the cell death process in mutated yeast cells.

Kumar NR, Sangwan S, Badotra P. Exposure to cell phone radiations produces biochemical changes in worker honey bees. Toxicol Int. 18(1):70-72, 2011.

The present study was carried out to find the effect of cell phone radiations on various biomolecules in the adult workers of *Apis mellifera* L. The results of the treated adults were analyzed and compared with the control. Radiation from the cell phone influences honey bees' behavior and physiology. There was reduced motor activity of the worker bees on the comb initially, followed by en masse migration and movement toward "talk mode" cell phone. The initial quiet period was characterized by rise in concentration of biomolecules including proteins, carbohydrates and lipids, perhaps due to stimulation of body mechanism to fight the stressful condition created by the radiations. At later stages of exposure, there was a slight decline in the concentration of biomolecules probably because the body had adapted to the stimulus

Geronikolou S, Zimeras S, Davos CH, Michalopoulos I, Tsitomenas S. Diverse Radiofrequency Sensitivity and Radiofrequency Effects of Mobile or Cordless Phone near Fields Exposure in *Drosophila melanogaster*. PLoS One. 2014 Nov 17;9(11):e112139. doi: 10.1371/journal.pone.0112139. eCollection 2014.

INTRODUCTION: The impact of electromagnetic fields on health is of increasing scientific interest. The aim of this study was to examine how the *Drosophila melanogaster* animal model is affected when exposed to portable or mobile phone fields. **METHODS/RESULTS:** Two experiments have been designed and performed in the same laboratory conditions. Insect cultures were exposed to the near field of a 2G mobile phone (the GSM 2G networks support and complement in parallel the 3G wide band or in other words the transmission of information via voice signals is served by the 2G technology in both mobile phones generations) and a 1880 MHz cordless phone both digitally modulated by human voice. Comparison with advanced statistics of the egg laying of the second generation exposed and non-exposed cultures showed limited statistical significance for the cordless phone exposed culture and statistical significance for the 900 MHz exposed insects. We calculated by physics, simulated and illustrated in three dimensional figures the calculated near fields of radiation inside the experimenting vials and their difference. Comparison of the power of the two fields showed that the difference between them becomes null when the experimental cylinder radius and the height of the antenna increase. **CONCLUSIONS/SIGNIFICANCE:** Our results suggest a possible radiofrequency sensitivity difference in insects which may be due to the distance from the antenna or to unexplored intimate factors. Comparing the near fields of the two frequencies bands, we see similar not identical geometry in length and height from the antenna and that lower frequencies tend to drive to increased radiofrequency effects.

Grigor'ev IuG. [Biological effects of mobile phone electromagnetic field on chick

Studies that show **Cell Phone** Health Effects

embryo (risk assessment using the mortality rate)] Radiats Biol Radioecol. 43(5):541-543, 2003. [Article in Russian]

Chicken embryos were exposed to EMF from GSM mobile phone during the embryonic development (21 days). As a result the embryo mortality rate in the incubation period increased to 75% (versus 16% in control group).

Favre D. Mobile phone-induced honeybee worker piping Apidologie 42:270–279, 2011.

The worldwide maintenance of the honeybee has major ecological, economic, and political implications. In the present study, electromagnetic waves originating from mobile phones were tested for potential effects on honeybee behavior. Mobile phone handsets were placed in the close vicinity of honeybees. The sound made by the bees was recorded and analyzed. The audiograms and spectrograms revealed that active mobile phone handsets have a dramatic impact on the behavior of the bees, namely by inducing the worker piping signal. In natural conditions, worker piping either announces the swarming process of the bee colony or is a signal of a disturbed bee colony.

Chen G, Lu D, Chiang H, Leszczynski D, Xu Z. Using model organism *Saccharomyces cerevisiae* to evaluate the effects of ELF-MF and RF-EMF exposure on global gene expression. Bioelectromagnetics. 33(7):550-560, 2012.

The potential health hazard of exposure to electromagnetic fields (EMF) continues to cause public concern. However, the possibility of biological and health effects of exposure to EMF remains controversial and their biophysical mechanisms are unknown. In the present study, we used *Saccharomyces cerevisiae* to identify genes responding to extremely low frequency magnetic fields (ELF-MF) and to radiofrequency EMF (RF-EMF) exposures. The yeast cells were exposed for 6 h to either 0.4 mT 50 Hz ELF-MF or 1800 MHz RF-EMF at a specific absorption rate of 4.7 W/kg. Gene expression was analyzed by microarray screening and confirmed using real-time reverse transcription-polymerase chain reaction (RT-PCR). We were unable to confirm microarray-detected changes in three of the ELF-MF responsive candidate genes using RT-PCR ($P > 0.05$). On the other hand, out of the 40 potential RF-EMF responsive genes, only the expressions of structural maintenance of chromosomes 3 (SMC3) and aquaporin 2 (AQY2 (m)) were confirmed, while three other genes, that is, halotolerance protein 9 (HAL9), yet another kinase 1 (YAK1) and one function-unknown gene (open reading frame: YJL171C), showed opposite changes in expression compared to the microarray data ($P < 0.05$). In conclusion, the results of this study suggest that the yeast cells did not alter gene expression in response to 50 Hz ELF-MF and that the response to RF-EMF is limited to only a very small number of genes. The possible biological consequences of the gene expression changes induced by RF-EMF await further investigation.

Aksoy U, Sahin S, Ozkoc S, Ergor G. The effect of electromagnetic waves on the growth of *Entamoeba histolytica* and *Entamoeba dispar*. Saudi Med J. 26(9):1388-1390, 2005.

OBJECTIVE: The aim of this study was to investigate the influence of electromagnetic radiation of a digital Global System for Mobile Communication mobile telephone (900

Studies that show **Cell Phone** Health Effects

MHz) on *Entamoeba histolytica* (*E. histolytica*) and *Entamoeba dispar* (*E. dispar*) (cysts or trophozoites, or both) in a 24-hour period. **METHODS:** This study was carried out from April 2004 to May 2004 at the Department of Parasitology, Medical Faculty of Dokuz Eylul University in Izmir, Turkey. The cultivated isolate tubes, which were exposed to electromagnetic field at 37OC, were evaluated as study group, whereas the tubes without exposure were assessed as control group. Finally, only living parasites in all tubes were counted using a hemacytometer. The effect of the temperature was evaluated for both control and study groups. **RESULTS:** The influence of electromagnetic field and temperature was assessed separately for the study group. The parasite number of *E. histolytica* decreased after exposure at 37OC and room temperature ($p=0.009$) compared to the decrease in the control group ($p=0.009$). The parasite number of *E. dispar* also decreased after exposure at 37OC and room temperature ($p=0.009$). In comparison to control tubes, this was a significant decrease ($p=0.008$). In the case of exposure of *E. histolytica* the results did not reveal any significant difference between temperature degrees to magnetic field ($p=0.459$) and *E. dispar* ($p=0.172$). **CONCLUSION:** Our findings show that exposure to electromagnetic field for a certain period of time may cause damage that can lead to death in single-cell organisms.

Cammaerts MC, Debeir O, Cammaerts R. Changes in *Paramecium caudatum* (protozoa) near a switched-on GSM telephone. *Electromagn Biol Med.* 30(1):57-66, 2011.

The protozoan *Paramecium caudatum* was examined under normal conditions versus aside a switched-on GSM telephone (900 MHz; 2 Watts). Exposed individuals moved more slowly and more sinuously than usual. Their physiology was affected: they became broader, their cytopharynx appeared broader, their pulse vesicles had difficult in expelling their content outside the cell, their cilia less efficiently moved, and trichocysts became more visible. All these effects might result from some bad functioning or damage of the cellular membrane. The first target of communication electromagnetic waves might thus be the cellular membrane.

Cammaerts MC, Rachidi Z, Bellens F, De Doncker P. Food collection and response to pheromones in an ant species exposed to electromagnetic radiation. *Electromagn Biol Med.* 2013 Jan 15. [Epub ahead of print]

We used the ant species *Myrmica sabuleti* as a model to study the impact of electromagnetic waves on social insects' response to their pheromones and their food collection. We quantified *M. sabuleti* workers' response to their trail, area marking and alarm pheromone under normal conditions. Then, we quantified the same responses while under the influence of electromagnetic waves. Under such an influence, ants followed trails for only short distances, no longer arrived at marked areas and no longer orientated themselves to a source of alarm pheromone. Also when exposed to electromagnetic waves, ants became unable to return to their nest and recruit congeners; therefore, the number of ants collecting food increases only slightly and slowly. After 180 h of exposure, their colonies deteriorated. Electromagnetic radiation obviously affects social insects' behavior and physiology.

Studies that show **Cell Phone** Health Effects

Cammaerts M-C, Vandenbosch GAE, Volski V. Effect of short-term GSM radiation at representative levels in society on a biological model: the ant *Myrmica sabuleti*. J Insect Beh. 27(4):514-526. 2014.

Well-controlled electromagnetic exposure conditions were set up at a representative societal GSM radiation intensity level, 1.5 V/m, which is the legally allowed level in Brussels. Two nests of the ant species *Myrmica sabuleti* were repeatedly irradiated during 10 min. before their behavior was observed, based on the analysis of the ant trajectories. Under these exposure conditions, behavioral effects were detected. The ants' locomotion slightly changed. The ants' orientation towards their attractive alarm pheromone statistically became of lower quality. The ants still presented their trail following behavior but less efficiently. In this controversial issue, ants could be considered as possible bioindicators.

Miscellaneous Effects

Yariktas M, Doner F, Ozguner F, Gokalp O, Dogru H, Delibas N. Nitric oxide level in the nasal and sinus mucosa after exposure to electromagnetic field. Otolaryngol Head Neck Surg. 132(5):713-716, 2005.

OBJECTIVE: The purpose of this study was to examine the changes in nitric oxide (NO) level in the nasal and paranasal sinus mucosa after exposure radiofrequency electromagnetic fields (EMF). STUDY DESIGN AND SETTING: Thirty male Sprague-Dawley rats were randomly grouped as follows: EMF group (group I; n, 10), EMF group in which melatonin received (group II; n, 10) and the control (sham operated) group (group III; n, 10). Groups I and II were exposed to a 900 MHz. Oral melatonin was given in group II. Control rats (group III) were also placed in the tube as the exposure groups, but without exposure to EMF. At the end of 2 weeks, the rats were sacrificed, and the nasal and paranasal sinus mucosa dissected. NO was measured in nasal and paranasal mucosa. RESULTS: The nasal and paranasal sinus mucosa NO levels of group I were significantly higher than those of the control group (group III) ($P < 0.05$). However, there was no statistically significant difference between group II and the control group (group III) regarding NO output ($P > 0.05$). CONCLUSION: Exposure to EMF released by mobile phones (900 MHz) increase NO levels in the sinus and nasal mucosa. SIGNIFICANCE: Increased NO levels may act as a defense mechanism and presumably related to tissue damage. In addition, melatonin may have beneficial effect to prevent these changes in the mucosa.

Taberski K, Klose M, Grote K, El Ouardi A, Streckert J, Hansen VW, Lerchl A. Noninvasive Assessment of Metabolic Effects of Exposure to 900 MHz Electromagnetic Fields on Djungarian Hamsters (*Phodopus sungorus*). Radiat Res. 181(6):617-622, 2014.

Sixteen male Djungarian hamsters, serving as their own controls, were individually exposed to RF-EMF (900 MHz, GSM modulation) at 0 (sham), 0.08, 0.4 or 4 W/kg specific absorption rate (SAR) in specially constructed rectangular waveguides. Exposure

Studies that show **Cell Phone** Health Effects

duration was one week per condition, followed by one week without exposure. Once per day, the temperatures of the hamsters' back fur (a surrogate for skin temperature) and the cornea of the eye (a surrogate for body temperature), were measured by infrared thermography. Oxygen, carbon dioxide and humidity were measured continuously in the ambient and exhaled air. Food and water consumption, as well as body weight were recorded once per week. Only at the highest SAR level were the following effects observed: fur temperatures were elevated by approximately 0.5°C (P < 0.001), while the temperatures of the eyes' surface were not affected; food consumption was lowered (P < 0.05), while water consumption and body weight were not affected; the production of carbon dioxide was lowered during the day (P < 0.01) and unaffected during the night, while oxygen consumption levels remained unaffected and finally the respiratory quotient (carbon dioxide production divided by oxygen consumption) was lower during the day (P < 0.05) and also somewhat lower during the night (not significant). The results demonstrate the usefulness of our methods for experiments dealing with metabolic effects of RF-EMF exposure in rodents. They also confirm the assumption that even though the metabolism is reduced at high SAR levels, the body core temperature is being kept constant by the energy uptake from the RF-EMF exposure which is able to physiologically compensate for the reduced metabolism.

Urbiniello D, Rösli M. Impact of one's own mobile phone in stand-by mode on personal radiofrequency electromagnetic field exposure. J Expo Sci Environ Epidemiol. 23:545-548, 2013.

When moving around, mobile phones in stand-by mode periodically send data about their positions. The aim of this paper is to evaluate how personal radiofrequency electromagnetic field (RF-EMF) measurements are affected by such location updates. Exposure from a mobile phone handset (uplink) was measured during commuting by using a randomized cross-over study with three different scenarios: disabled mobile phone (reference), an activated dual-band phone and a quad-band phone. In the reference scenario, uplink exposure was highest during train rides (1.19 mW/m(2)) and lowest during car rides in rural areas (0.001 mW/m(2)). In public transports, the impact of one's own mobile phone on personal RF-EMF measurements was not observable because of high background uplink radiation from other people's mobile phone. In a car, uplink exposure with an activated phone was orders of magnitude higher compared with the reference scenario. This study demonstrates that personal RF-EMF exposure is affected by one's own mobile phone in stand-by mode because of its regular location update. Further dosimetric studies should quantify the contribution of location updates to the total RF-EMF exposure in order to clarify whether the duration of mobile phone use, the most common exposure surrogate in the epidemiological RF-EMF research, is actually an adequate exposure proxy.

Weisbrot D, Lin H, Ye L, Blank M, Goodman R. Effects of mobile phone radiation on reproduction and development in *Drosophila melanogaster*. J Cell Biochem 89(1):48-55, 2003.

In this report we examined the effects of a discontinuous radio frequency (RF) signal

Studies that show **Cell Phone** Health Effects

produced by a GSM multiband mobile phone (900/1,900 MHz; SAR approximately 1.4 W/kg) on *Drosophila melanogaster*, during the 10-day developmental period from egg laying through pupation. As found earlier with low frequency exposures, the non-thermal radiation from the GSM mobile phone increased numbers of offspring, elevated hsp70 levels, increased serum response element (SRE) DNA-binding and induced the phosphorylation of the nuclear transcription factor, ELK-1. The rapid induction of hsp70 within minutes, by a non-thermal stress, together with identified components of signal transduction pathways, provide sensitive and reliable biomarkers that could serve as the basis for realistic mobile phone safety guidelines.

Paredi P, Kharitonov SA, Hanazawa T, Barnes PJ, Local vasodilator response to mobile phones. *Laryngoscope* 111(1):159-162, 2001.

OBJECTIVES: The use of mobile phones with the resulting generation of potentially harmful electromagnetic fields (EMF) is the focus of public interest. Heat generation and the activation of the inducible form of nitric oxide (NO) synthase may be possible causes of the biological effects of EMF exposure. We investigated if a mobile telephone conversation can modify skin temperature, NO, and nasal resistance. **METHODS:** We studied the effect of an EMF (900 MHz) generated by a commercially available cellular phone during a 30-minute telephone conversation on skin temperature, nasal NO measured by chemiluminescence, and nasal minimal cross-sectional area (MCA) measured by rhinometry. Eleven normal subjects (mean age +/- standard error of mean [SEM], 32 +/- 5 y; 10 male) were studied. **RESULTS:** There was a similar and significant increase in skin temperature of the nostril and occipital area on the same side as the telephone (maximal increase 2.3 +/- 0.2 degrees C at 6 min) as well as a tendency for higher nasal NO levels (maximal increase 12.9 +/- 4.9% at 10 min), whereas the MCA was significantly reduced (maximal decrease -27 +/- 6% at 15 min). Such changes were not recorded when an earpiece was used to avoid the direct exposure to the electromagnetic field. There were no changes in the skin temperature and nasal NO measured on the opposite side to the mobile phone, whereas the MCA was significantly increased (38 +/- 10%). **CONCLUSIONS:** Exposure to EMF produced by a mobile phone produces biological effects that can be easily measured. Microwaves may increase skin temperature and therefore cause vasodilation and reduce MCA. Further studies are needed to study the long-term effects of mobile phone use and the relation among NO production, vasodilation, and temperature.

Panagopoulos, D. J., Johansson O. & Carlo G.L. Polarization: A Key Difference between Man-made and Natural Electromagnetic Fields, in regard to Biological Activity. *Sci. Rep.* 5, 14914; doi: 10.1038/srep14914 (2015). Published online Oct 12, 2015.

In the present study we analyze the role of polarization in the biological activity of Electromagnetic Fields (EMFs)/Electromagnetic Radiation (EMR). All types of man-made EMFs/EMR - in contrast to natural EMFs/EMR - are polarized. Polarized EMFs/EMR can have increased biological activity, due to: 1) Ability to produce constructive interference effects and amplify their intensities at many locations. 2) Ability to force all charged/polar molecules and especially free ions within and around all living cells to

Studies that show **Cell Phone** Health Effects

oscillate on parallel planes and in phase with the applied polarized field. Such ionic forced-oscillations exert additive electrostatic forces on the sensors of cell membrane electro-sensitive ion channels, resulting in their irregular gating and consequent disruption of the cell's electrochemical balance. These features render man-made EMFs/EMR more bioactive than natural non-ionizing EMFs/EMR. This explains the increasing number of biological effects discovered during the past few decades to be induced by man-made EMFs, in contrast to natural EMFs in the terrestrial environment which have always been present throughout evolution, although human exposure to the latter ones is normally of significantly higher intensities/energy and longer durations. Thus, polarization seems to be a trigger that significantly increases the probability for the initiation of biological/health effects.

Lepp A, Barkley JE, Sanders GJ, Rebold M, Gates P. The relationship between cell phone use, physical and sedentary activity, and cardiorespiratory fitness in a sample of U.S. college students. *Int J Behav Nutr Phys Act.* 2013 Jun 21;10:79. doi: 10.1186/1479-5868-10-79.

BACKGROUND: Today's cell phones increase opportunities for activities traditionally defined as sedentary behaviors (e.g., surfing the internet, playing video games). People who participate in large amounts of sedentary behaviors, relative to those who do not, tend to be less physically active, less physically fit, and at greater risk for health problems. However, cell phone use does not have to be a sedentary behavior as these devices are portable. It can occur while standing or during mild-to-moderate intensity physical activity. Thus, the relationship between cell phone use, physical and sedentary activity, and physical fitness is unclear. The purpose of this study was to investigate these relationships among a sample of healthy college students. **METHODS:** Participants were first interviewed about their physical activity behavior and cell phone use. Then body composition was assessed and the validated self-efficacy survey for exercise behaviors completed. This was followed by a progressive exercise test on a treadmill to exhaustion. Peak oxygen consumption (VO₂ peak) during exercise was used to measure cardiorespiratory fitness. Hierarchical regression was used to assess the relationship between cell phone use and cardiorespiratory fitness after controlling for sex, self-efficacy, and percent body fat. Interview data was transcribed, coded, and Chi-square analysis was used to compare the responses of low and high frequency cell phone users. **RESULTS:** Cell phone use was significantly ($p = 0.047$) and negatively ($\beta = -0.25$) related to cardio respiratory fitness independent of sex, self-efficacy, and percent fat which were also significant predictors ($p < 0.05$). Interview data offered several possible explanations for this relationship. First, high frequency users were more likely than low frequency users to report forgoing opportunities for physical activity in order to use their cell phones for sedentary behaviors. Second, low frequency users were more likely to report being connected to active peer groups through their cell phones and to cite this as a motivation for physical activity. Third, high levels of cell phone use indicated a broader pattern of sedentary behaviors apart from cell phone use, such as watching television. **CONCLUSION:** Cell phone use, like traditional sedentary behaviors, may disrupt physical activity and reduce cardiorespiratory fitness.

Studies that show **Cell Phone** Health Effects

Lee KS, Choi JS, Hong SY, Son TH, Yu K. Mobile phone electromagnetic radiation activates MAPK signaling and regulates viability in Drosophila. Bioelectromagnetics.29(5):371-379, 2008.

Mobile phones are widely used in the modern world. However, biological effects of electromagnetic radiation produced by mobile phones are largely unknown. In this report, we show biological effects of the mobile phone 835 MHz electromagnetic field (EMF) in the Drosophila model system. When flies were exposed to the specific absorption rate (SAR) 1.6 W/kg, which is the proposed exposure limit by the American National Standards Institute (ANSI), more than 90% of the flies were viable even after the 30 h exposure. However, in the SAR 4.0 W/kg strong EMF exposure, viability dropped from the 12 h exposure. These EMF exposures triggered stress response and increased the production of reactive oxygen species. The EMF exposures also activated extracellular signal regulated kinase (ERK) and c-Jun N-terminal kinase (JNK) signaling, but not p38 kinase signaling. Interestingly, SAR 1.6 W/kg activated mainly ERK signaling and expression of an anti-apoptotic gene, whereas SAR 4.0 W/kg strongly activated JNK signaling and expression of apoptotic genes. In addition, SAR 4.0 W/kg amplified the number of apoptotic cells in the fly brain. These findings demonstrate that the exposure limit on electromagnetic radiation proposed by ANSI triggered ERK-survival signaling but the strong electromagnetic radiation activated JNK-apoptotic signaling in Drosophila.

Aly AA, Cheema MI, Tambawala M, Laterza R, Zhou E, Rathnabharathi K, Barnes FS. Effects of 900-MHz Radio Frequencies on the Chemotaxis of Human Neutrophils in Vitro. IEEE Transactions on Biomedical Engineering, 55(2): 795-797, 2008.

Summary: The effects of radio frequency (RF) fields on the ability of human neutrophils to follow concentration gradients of Cyclic Adenosine 3', 5'-Monophosphate (C-AMP) are reported. Blood from healthy adult donors was exposed in vitro to different temperatures and 900-MHz RF field at approximately 0.4 V/m. It was observed that the neutrophils' speed increased with increasing temperatures from 35 °C to 40 °C where it peaked and then decreased above 40 °C without RF exposure. When 900-MHz RF field was applied, the speed increased above the value observed at the same temperature, and the maximum speed exceeded that measured value at any temperature by approximately 50%. The calculated temperature change resulting from the RF exposure was less than one microdegree. The direction of motion changed from along the concentration gradient and the electrical field lines to motion at right angles to the concentration gradient and the electric field. The average time for the neutrophils to respond to the effect of RF radiation was about 2.5 min.

Aweda MA, Ajekigbe AT, Ibitoye AZ, Evwahierhurhoma BO, Eletu OB. Potential health risks due to telecommunications radiofrequency radiation exposures in Lagos State Nigeria. Nig Q J Hosp Med. 19(1):6-14, 2009.

BACKGROUND: The global system mobile telecommunications system (GSM) which was recently introduced in Nigeria is now being used by over 40 million people in Nigeria. The use of GSM is accompanied with exposure of the users to radiofrequency radiation

Studies that show Cell Phone Health Effects

(RFR), which if significant, may produce health hazards. This is the reason why many relevant national and international organizations recommended exposure limits to RFR and why it is made compulsory for GSM handsets to indicate the maximum power output as a guide to potential consumers. **OBJECTIVE:** This study was conducted to measure the RFR output power densities (S) from the most commonly used GSM handsets used in Lagos State and compare with the limit recommended for safety assessment. **METHODS:** Over 1100 most commonly used handsets of different makes and models as well as wireless phones were sampled and studied in all over the local government areas of the State. An RFR meter, Electrosmog from LESSEMF USA was used for the measurements. The handsets were assessed for health risks using the reference value of $9 \text{ Wm}(-2)$ as recommended by the International Commission on Non-Ionizing Radiation Protection (ICNIRP). **RESULTS:** The range of the S-values obtained varied from a minimum of $1.294 \text{ Wm}(-2)$ with Siemens model R228 to a maximum of $16.813 \pm 0.094 \text{ Wm}(-2)$ with Samsung model C140*. The results from wireless telephones showed very low S-values ranging from a minimum of $0.024 \pm 0.001 \text{ Wm}(-2)$ with HUAWEI and ST CDMA 1 to a maximum of $0.093 \pm 0.002 \text{ Wm}(-2)$ with HISENSE. **CONCLUSION:** The results showed that the population in Lagos State may be at risk due to significant RFR exposures resulting principally from the use of GSM. Quite a number of handsets emit power above the ICNIRP recommended value. Measured RFR power close to Radio and Television masts and transmitters are within tolerable limits in most cases, only that the public should not reside or work close to RFR installations. Phone calls with GSM should be restricted to essential ones while youths and children that are more susceptible to RFR hazards should be supervised in their use of GSM. Wireless phones are quite safe.

Ayata A, Mollaoglu H, Yilmaz HR, Akturk O, Ozguner F, Altuntas I. Oxidative stress-mediated skin damage in an experimental mobile phone model can be prevented by melatonin. J Dermatol. 31(11):878-883, 2004.

Most mobile phones emit 900 MHz of radiation that is mainly absorbed by the external organs. The effects of 900 MHz of radiation on fibrosis, lipid peroxidation, and antioxidant enzymes and the ameliorating effects of melatonin (Mel) were evaluated in rat skin. Thirty Wistar-Albino rats were used in the study. The experimental groups were the control group, the irradiated group (IR), and the irradiated+Mel treated group (IR+Mel). A dose of 900 MHz, 2 W radiation was applied to the IR group every day for 10 days (30 min/day). The IR+Mel group received 10 mg/kg/day melatonin in tap water for 10 days before the irradiation. At the end of the 10th day, a skin specimen was excised from the thoracoabdominal area. The levels of malondialdehyde (MDA) and hydroxyproline and the activities of superoxide dismutase (SOD), glutathione peroxidase (GSH-Px), and catalase (CAT) were studied in the skin samples. MDA and hydroxyproline levels and activities of CAT and GSH-Px were increased significantly in the IR group compared to the control group ($p < 0.05$) and decreased significantly in the IR+Mel group ($p < 0.05$). SOD activity was decreased significantly in the IR group and this decrease was not prevented by the Mel treatment. These results suggest that rats irradiated with 900 MHz suffer from increased fibrosis and lipid peroxidation (LPO). Mel

Studies that show **Cell Phone** Health Effects

treatment can reduce the fibrosis and LPO caused by radiation.

Barteri M, De Carolis R, Marinelli F, Tomassetti G, Montemiglio LC. Effects of microwaves (900 MHz) on peroxidase systems: a comparison between lactoperoxidase and horseradish peroxidase. Electromagn Biol Med. 2015 Jan 12:1-7. [Epub ahead of print]

This work shows the effects of exposure to an electromagnetic field at 900 MHz on the catalytic activity of the enzymes lactoperoxidase (LPO) and horseradish peroxidase (HRP). Experimental evidence that irradiation causes conformational changes of the active sites and influences the formation and stability of the intermediate free radicals is documented by measurements of enzyme kinetics, circular dichroism spectroscopy (CD) and cyclic voltammetry.

Driving Impairment Using a Cell Phone

Violanti JM, Cellular phones and traffic accidents. Public Health 111(6):423-428, 1997.

Cellular phone use in motor vehicles is becoming an increasing world-wide phenomenon. Using data obtained from traffic accidents reported between 1992 and 1995 in the state of Oklahoma, USA, this study examined statistical rate-ratios of accident characteristics between drivers with or without cellular phones. Rates were calculated between cellular phone involvement and reported accident causes, types of collision, driver actions immediately prior to the accident, location of the accident, the extent of fatalities, and age and gender of drivers. Results indicated a significant increased rate among drivers with cellular phones for inattention, unsafe speed, driving on wrong side of road, striking a fixed object, overturning their vehicle, swerving prior to the accident, and running off the roadway. People with phones stood an increased risk of being killed in an accident over persons without phones. Males with phones had a significantly higher rate than females for many of accident characteristics mentioned above. Rate-ratios of some accident characteristics and fatalities increased as age increased, with the exception of drivers under age 20 yrs, who had the highest fatality rate. *Limitations* of the study and possible prevention alternatives are discussed.

Violanti JM, Marshall JR, Cellular phones and traffic accidents: an epidemiological approach. Accid Anal Prev 28(2):265-270, 1996.

Using epidemiological case-control design and logistic regression techniques, this study examined the association of cellular phone use in motor vehicles and traffic accident risk. The amount of time per month spent talking on a cellular phone and 18 other driver inattention factors were examined. Data were obtained from: (1) a case group of 100 randomly selected drivers involved in accidents within the past 2 years, and (2) a control group of 100 randomly selected

Studies that show **Cell Phone** Health Effects

licensed drivers not involved in accidents within the past 10 years. Groups were matched on geographic residence. Approximately 13% (N = 7) of the accident and 9% (N = 7) of the non-accident group reported use of cellular phones while driving. Data was obtained from Department of Motor Vehicles accident reports and survey information from study subjects. We hypothesized that increased use of cellular phones while driving was associated with increased odds of a traffic accident. Results indicated that talking more than 50 minutes per month on cellular phones in a vehicle was associated with a 5.59-fold increased risk in a traffic accident. The combined use of cellular phones and motor and cognitive activities while driving were also associated with increased traffic accident risk. Readers should be cautioned that this study: (1) consists of a small sample, (2) reveals statistical associations and not causal relationships, and (3) does not conclude that talking on cellular phones while driving is inherently dangerous.

Violanti JM, Cellular phones and fatal traffic collisions. *Accid Anal Prev* 30(4):519-524, 1998.

A case-control study was conducted to determine statistical associations between traffic fatalities and the use or presence of a cellular phone, given involvement in a collision. The hypothesis of this study does not imply that cellular phones directly affect fatalities, but that phones increase the risk of certain accident characteristics in fatal collisions more than those same characteristics in non-fatal collisions. Analysis employed data from 223,137 traffic accidents occurring between 1992 and 1995. Information on collision characteristics and cellular phone involvement for each fatality was compared with the same information for each non-fatality (controls). Statistically adjusting for other collision variables (age, gender, alcohol use, speed, inattention and driving left of center), an approximate nine-fold increased risk was found for a fatality given the use of a cellular phone. An approximate two-fold increased risk for a fatality was found given the presence of a cellular phone in the vehicle. Combined effects of reported phone use, driving to the left of center and inattention increased the risk of a fatal collision more than phone use did by itself. This analysis implies a statistical, but not necessarily a causal, relationship. A multitude of factors are involved in any traffic collision, and the exact cause of an accident and its severity level is difficult to disentangle.

Strayer DL, Johnston WA. Driven to distraction: dual-Task studies of simulated driving and conversing on a cellular telephone. *Psychol Sci* 12(6):462-466, 2001.

Dual-task studies assessed the effects of cellular-phone conversations on performance of a simulated driving task. Performance was not disrupted by listening to radio broadcasts or listening to a book on tape. Nor was it disrupted by a continuous shadowing task using a handheld phone, ruling out, in this case, dual-task interpretations associated with holding the phone, listening, or speaking. However

Studies that show **Cell Phone** Health Effects

significant interference was observed in a word-generation variant of the shadowing task, and this deficit increased with the difficulty of driving. Moreover unconstrained conversations using either a handheld or a hands-free cell phone resulted in a twofold increase in the failure to detect simulated traffic signals and slower reactions to those signals that were detected. We suggest that cellular-phone use disrupts performance by diverting attention to an engaging cognitive context other than the one immediately associated with driving.

Strayer DL, Drews FA, Johnston WA. Cell phone-induced failures of visual attention during simulated driving. J Exp Psychol Appl 9(1):23-32, 2003.

This research examined the effects of hands-free cell phone conversations on simulated driving. The authors found that these conversations impaired driver's reactions to vehicles braking in front of them. The authors assessed whether this impairment could be attributed to a withdrawal of attention from the visual scene, yielding a form of inattention blindness. Cell phone conversations impaired explicit recognition memory for roadside billboards. Eye-tracking data indicated that this was due to reduced attention to foveal information. This interpretation was bolstered by data showing that cell phone conversations impaired implicit perceptual memory for items presented at fixation. The data suggest that the impairment of driving performance produced by cell phone conversations is mediated, at least in part, by reduced attention to visual inputs.

Jenness JW, Lattanzio RJ, O'Toole M, Taylor N, Pax C. Effects of manual versus voice-activated dialing during simulated driving. Percept Mot Skills 94(2):363-379, 2002.

We measured driving performance (lane-keeping errors, driving times, and glances away from the road scene) in a video driving simulator for 24 volunteers who each drove alone on a 10.6-km multicurved course while simultaneously placing calls on a mobile phone subscribed to a voice-activated dialing system. Driving performance also was measured for the same distance while participants manually dialed phone numbers and while they drove without dialing. There were 22% fewer lane-keeping errors ($p < .01$) and 56% fewer glances away from the road scene ($p < .01$) when they used voice-activated dialing as compared to manual dialing. Significantly longer driving times in both of the dialing conditions as compared to the No Dialing condition are discussed in terms of the hypothesis that drivers decrease driving speed to compensate for the demands of the secondary phone tasks.

Zajdel R, Zajdel J, Zwolińska A, Smigielski J, Beling P, Cegliński T, Nowak D. The sound of a mobile phone ringing affects the complex reaction time of its owner. Arch Med Sci. 8(5):892-898, 2012.

INTRODUCTION: Mobile phone conversation decreases the ability to concentrate and impairs the attention necessary to perform complex activities, such as driving a car. Does the ringing sound of a mobile phone affect the driver's ability to perform complex sensory-motor activities? We compared a subject's reaction time while performing a test either with a mobile phone ringing or without. MATERIAL AND METHODS: The examination was performed on a PC-based reaction time self-constructed system

Studies that show **Cell Phone** Health Effects

Reactor. The study group consisted of 42 healthy students. The protocol included instruction, control without phone and a proper session with subject's mobile phone ringing. The terms of the study were standardised. RESULTS: There were significant differences ($p < 0.001$) in reaction time in control (597 ms), mobile (633 ms) and instruction session (673 ms). The differences in female subpopulation were also significant ($p < 0.01$). Women revealed the longest reaction time in instruction session (707 ms), were significantly quicker in mobile (657 ms, $p < 0.01$) and in control session (612 ms, $p < 0.001$). In men, the significant difference was recorded only between instruction (622 ms) and control session (573 ms, $p < 0.01$). The other differences were not significant ($p > 0.08$). Men proved to complete significantly quicker than women in instruction ($p < 0.01$) and in mobile session ($p < 0.05$). Differences amongst the genders in control session was not significant ($p > 0.05$). CONCLUSIONS: The results obtained proofed the ringing of a phone exerts a significant influence on complex reaction time and quality of performed task.

Strayer DL, Drews FA. Profiles in driver distraction: effects of cell phone conversations on younger and older drivers. Hum Factors. 46(4):640-649, 2004.

Our research examined the effects of hands-free cell phone conversations on simulated driving. We found that driving performance of both younger and older adults was influenced by cell phone conversations. Compared with single-task (i.e., driving-only) conditions, when drivers used cell phones their reactions were 18% slower, their following distance was 12% greater, and they took 17% longer to recover the speed that was lost following braking. There was also a twofold increase in the number of rear-end collisions when drivers were conversing on a cell phone. These cell-phone-induced effects were equivalent for younger and older adults, suggesting that older adults do not suffer a significantly greater penalty for talking on a cell phone while driving than compared with their younger counterparts. Interestingly, the net effect of having younger drivers converse on a cell phone was to make their average reactions equivalent to those of older drivers who were not using a cell phone. Actual or potential applications of this research include providing guidance for recommendations and regulations concerning the use of mobile technology while driving.

Redelmeier DA, Tibshirani RJ, Association between cellular-telephone calls and motor vehicle collisions. N Engl J Med 13;336(7):453-458, 1997.

BACKGROUND: Because of a belief that the use of cellular telephones while driving may cause collisions, several countries have restricted their use in motor vehicles, and others are considering such regulations. We used an epidemiologic method, the case-crossover design, to study whether using a cellular telephone while driving increases the risk of a motor vehicle collision. METHODS: We studied 699 drivers who had cellular telephones and who were involved in motor vehicle collisions resulting in substantial property damage but no personal injury. Each person's cellular-telephone calls on the day of the collision and during the previous week were analyzed through the use of detailed billing records. RESULTS: A total of 26,798 cellular-telephone calls were made during the 14-month study period. The risk of a collision when using a cellular telephone was four

Studies that show **Cell Phone** Health Effects

times higher than the risk when a cellular telephone was not being used (relative risk, 4.3; 95 percent confidence interval, 3.0 to 6.5). The relative risk was similar for drivers who differed in personal characteristics such as age and driving experience; calls close to the time of the collision were particularly hazardous (relative risk, 4.8 for calls placed within 5 minutes of the accident, as compared with 1.3 for calls placed more than 15 minutes before the accident; $P < 0.001$); and units that allowed the hands to be free (relative risk, 5.9) offered no safety advantage over hand-held units (relative risk, 3.9; P not significant). Thirty-nine percent of the drivers called emergency services after the collision, suggesting that having a cellular telephone may have had advantages in the aftermath of an event. CONCLUSIONS: The use of cellular telephones in motor vehicles is associated with a quadrupling of the risk of a collision during the brief time interval involving a call. Decisions about regulation of such telephones, however, need to take into account the benefits of the technology and the role of individual responsibility.

Tornros JE, Bolling AK. Mobile phone use-Effects of handheld and handsfree phones on driving performance. *Accid Anal Prev.* 37(5):902-909, 2005.

The study was concerned with effects of handsfree and handheld mobile phone dialling and conversation in simulated driving. In the main experiment dealing with conversation, 48 participants drove a distance of about 70km on a route which led through urban and rural environments. In the dialling experiment, the participants drove a distance of 15km on a rural two-lane road. The experimental design was mixed with phone mode as a between-subjects factor and phone use (yes/no) as a within-subjects factor. Performance on a peripheral detection task (PDT) while driving was impaired by dialling and conversation for both phone modes, interpreted as an increase in mental workload. Driving performance was impaired by dialling-lateral position deviation increased in a similar way for both phone modes. Conversation had, however, opposite effects-lateral position deviation decreased in a similar way for both phone modes. Driving speed decreased as an effect of dialling with the greatest effect for handsfree phone mode. Conversation also caused reduced speed, but only for handheld phone mode. The effects on speed can be interpreted as a compensatory effort for the increased mental workload. In spite of the compensatory behaviour, mental workload was still markedly increased by phone use. Subjective effects of dialling and conversation were also analysed. Most participants reported a speed decrease but no effect on lateral position deviation as an effect of dialling or conversation. In the conversation experiment, driving performance was rated better for handsfree than for handheld mode. In the dialling experiment, no difference between the two phone modes appeared.

Oommen BS, Stahl JS. Inhibited head movements: A risk of combining phoning with other activities? *Neurology* 65(5):754-756, 2005.

Abstract-- Studies of cellular phone use while driving have attributed impaired performance to the distractions of conversation. We determined that holding an inactive phone to the ear reduces the probability of eccentric head positions, potentially indicating reduced ability to monitor the visual surround. This effect may constitute a

Studies that show **Cell Phone** Health Effects

risk of cellular phone use independent of conversation and peculiar to handheld models.

Miller G, Zhu G, Wright MJ, Hansell NK, Martin, NG. The Heritability and Genetic Correlates of Mobile Phone Use: A Twin Study of Consumer Behavior. Twin Research and Human Genetics / Volume 15 / Issue 01 / February 2012, pp 97-106.

There has been almost no overlap between behavior genetics and consumer behavior research, despite each field's importance in understanding society. In particular, both have neglected to study genetic influences on consumer adoption and usage of new technologies — even technologies as important as the mobile phone, now used by 5.8 out of 7.0 billion people on earth. To start filling this gap, we analyzed self-reported mobile phone use, intelligence, and personality traits in two samples of Australian teenaged twins (mean ages 14.2 and 15.6 years), totaling 1,036 individuals.

ACE modeling using Mx software showed substantial heritabilities for how often teens make voice calls (.60 and .34 in samples 1 and 2, respectively) and for how often they send text messages (.53 and .50). Shared family environment — including neighborhood, social class, parental education, and parental income (i.e., the generosity of calling plans that parents can afford for their teens) — had much weaker effects. Multivariate modeling based on cross-twin, cross-trait correlations showed negative genetic correlations between talking/texting frequency and intelligence (around $-.17$), and positive genetic correlations between talking/texting frequency and extraversion (about $.20$ to $.40$).

Our results have implications for assessing the risks of mobile phone use such as radiofrequency field (RF) exposure and driving accidents, for studying adoption and use of other emerging technologies, for understanding the genetic architecture of the cognitive and personality traits that predict consumer behavior, and for challenging the common assumption that consumer behavior is shaped entirely by culture, media, and family environment.

McEvoy SP, Stevenson MR, McCartt AT, Woodward M, Haworth C, Palamara P, Cercarelli R. Role of mobile phones in motor vehicle crashes resulting in hospital attendance: a case-crossover study. BMJ 331(7514):428, 2005.

OBJECTIVES: To explore the effect of drivers' use of mobile (cell) phones on road safety. DESIGN: A case-crossover study. SETTING: Perth, Western Australia. PARTICIPANTS: 456 drivers aged ≥ 17 years who owned or used mobile phones and had been involved in road crashes necessitating hospital attendance between April 2002 and July 2004. MAIN OUTCOME MEASURE: Driver's use of mobile phone at estimated time of crash and on trips at the same time of day in the week before the crash. Interviews with drivers in hospital and phone company's records of phone use. RESULTS: Driver's use of a mobile phone up to 10 minutes before a crash was associated with a fourfold increased likelihood of crashing (odds ratio 4.1, 95% confidence interval 2.2 to 7.7, $P < 0.001$). Risk was raised irrespective of whether or not a hands-free device was used (hands-free: 3.8, 1.8 to 8.0, $P < 0.001$; hand held: 4.9, 1.6 to 15.5, $P = 0.003$). Increased risk was similar in

Studies that show **Cell Phone** Health Effects

men and women and in drivers aged ≥ 30 and < 30 years. A third ($n=21$) of calls before crashes and on trips during the previous week were reportedly on hand held phones.

CONCLUSIONS: When drivers use a mobile phone there is an increased likelihood of a crash resulting in injury. Using a hands-free phone is not any safer.

Matthews R, Legg S, Charlton S. The effect of cell phone type on drivers subjective workload during concurrent driving and conversing. *Accid Anal Prev* 35(4):451-457, 2003.

The effect of three types of cell phones (hand held, hands free with an external speaker and personal hands free) on total subjective workload (including its constituent components; mental demand, physical demand, temporal demand, performance, effort and frustration) and intelligibility was measured using the NASA-task load index (TLX) and the modified rhyme test (MRT), respectively in 13 experienced drivers (nine male, four female, age range 28-65 years), whilst driving on a rural highway. The drivers rated all components of workload for each type of cell phone to be significantly higher than for a control condition in which no cell phone was used. The mean (standard deviation) total workload was lowest for the personal hands free cell phone (7.1(3.65)) and highest for the hands free speaker phone (10.8 (3.63)) ($P<0.001$). The mean (standard deviation) intelligibility score was highest for the personal hands free cell phone (74.1 (7.9)) and lowest for the hands free speaker phone (55.0 (10.4)) ($P<0.001$). Frustration was significantly correlated with total workload (0.60, $P<0.001$) and intelligibility was significantly correlated with frustration (-0.35, $P<0.05$). Physical demand was not a high contributor to total workload. It is concluded that a personal hands free cell phone would interfere least with the cognitive demands of driving.

Lesch MF, Hancock PA. Driving performance during concurrent cell-phone use: are drivers aware of their performance decrements? *Accid Anal Prev.* 36(3):471-480, 2004.

Prior research has documented the manner in which a variety of driving performance measures are impacted by concurrent cell-phone use as well as the influence of age and gender of the driver. This current study examined the extent to which different driver groups are aware of their associated performance decrements. Subjects' confidence in dealing with distractors while driving and their ratings of task performance and demand were compared with their actual driving performance in the presence of a cell-phone task. While high confidence ratings appeared to be predictive of better driving performance for male drivers (as confidence increased, the size of the distraction effects decreased), this relationship did not hold for females; in fact, for older females, as confidence increased, performance decreased. Additionally, when drivers were matched in terms of confidence level, brake responses of older females were slowed to a much greater extent (0.38s) than were brake responses of any other group (0.10s for younger males and females and 0.07s for older males). Finally, females also rated the driving task as less demanding than males, even though their performance was more greatly affected by distraction. These results suggest that many drivers may not be aware of their decreased performance while using cell-phones and that it may be particularly important to target educational campaigns on driver distraction towards female drivers

Studies that show **Cell Phone** Health Effects

for whom there tended to be a greater discrepancy between driver perceptions and actual performance.

Lam LT. Distractions and the risk of car crash injury: the effect of drivers' age. J Safety Res 33(3):411-419, 2002.

PROBLEM: Motor-vehicle accidents are one of the major causes of injury in most motorized countries. Driver distractions have been suggested as a contributor to traffic accidents. Moreover, age of the driver seems to have a role in the relationship between distractions and car crashes. But very few studies have investigated the effect of driver's age on this relationship. This exploratory study investigated the association between distractions, both inside and outside the vehicle, and the increased risk of car crash injury among drivers across different ages. METHOD: This study used a case series design to analyze data routinely collected by the NSW police in Australia. A special focus of this study was on how drivers' age affects the risk of car crash injury, which was determined by using a well-documented risk estimation methodology. RESULTS: The results obtained indicated that drivers of all ages, on the whole, are more susceptible to distractions inside the vehicle than distractions coming from outside. Age was shown to affect the relationship between in-vehicle distraction and the risk of car crash injury. A separate analysis was also conducted on hand-held phone usage while driving with results supplementing previous findings reported in the literature. IMPACT TO INDUSTRY: Safety strategies to countermeasure in-vehicle distractions have been suggested and discussed.

Lamble D, Kauranen T, Laakso M, Summala H, Cognitive load and detection thresholds in car following situations: safety implications for using mobile (cellular) telephones while driving. Accid Anal Pre 31(6):617-623, 1999.

This study was aimed at investigating drivers' ability to detect a car ahead decelerating, while doing mobile phone related tasks. Nineteen participants aged between 20 and 29 years, (2000-125000 km driving experience) drove at 80 km/h, 50 m behind a lead car, on a 30 km section of motorway in normal traffic. During each trial the lead car started to decelerate at an average of 0.47 m/s² while the participant either looked at the car in front (control), continuously dialed series of three random integers on a numeric keypad (divided visual attention), or performed a memory and addition task (non-visual attention). The results indicated that drivers' detection ability was impaired by about 0.5 s in terms of brake reaction time and almost 1 s in terms of time-to-collision, when they were doing the non-visual task whilst driving. This impairment was similar to when the drivers were dividing their visual attention between the road ahead and dialing numbers on the keypad. It was concluded that neither a hands-free option nor a voice controlled interface removes the safety problems associated with the use of mobile phones in a car.

Laberge-Nadeau C, Maag U, Bellavance F, Lapierre SD, Desjardins D, Messier S, Sai;di A. Wireless telephones and the risk of road crashes. Accid Anal Prev. 35(5):649-660, 2003.

Studies that show **Cell Phone** Health Effects

In light of the rapidly increasing development of the cell phone market, the use of such equipment while driving raises the question of whether it is associated with an increased accident risk; and if so, what is its magnitude. This research is an epidemiological study on two large cohorts, namely users and non-users of cell phones, with the objective of verifying whether an association exists between cell phone use and road crashes, separating those with injuries. The Societe de l'Assurance Automobile du Quebec (SAAQ) mailed a questionnaire and letter of consent to 175000 licence holders for passenger vehicles. The questionnaire asked about exposure to risk, driving habits, opinions about activities likely to be detrimental to driving and accidents within the last 24 months. For cell phone users, questions pertaining to the use of the telephone were added. We received 36078 completed questionnaires, with a signed letter of consent. Four wireless phone companies provided the files on cell phone activity, and the SAAQ the files for 4 years of drivers' records and police reports. The three data sources were merged using an anonymized identification number. The statistical methods include logistic-normal regression models to estimate the strength of the links between the explanatory variables and crashes. The relative risk of all accidents and of accidents with injuries is higher for users of cell phones than for non-users. The relative risks (RR) for injury collisions and also for all collisions is 38% higher for men and women cell phone users. These risks diminish to 1.1 for men and 1.2 for women if other variables, such as the kilometres driven and driving habits are incorporated into the models. Similar results hold for several sub-groups. The most significant finding is a dose-response relationship between the frequency of cell phone use, and crash risks. The adjusted relative risks for heavy users are at least two compared to those making minimal use of cell phones; the latter show similar collision rates as do the non-users.

Korpinen L, Pääkkönen R. Accidents and close call situations connected to the use of mobile phones. *Accid Anal Prev.* 45(2):75-82, 2012.

Abstract. The aim of our work was to study the accidents and close call situations connected to the use of mobile phones. We have analyzed how the accidents/close call situations are connected to background information, in particular age, gender and self-reported symptoms. The study was carried out as a cross-sectional study by posting the questionnaire to 15,000 working-age Finns. The responses (6121) were analyzed using the logistic regression models. Altogether 13.7% of respondents had close call situations and 2.4% had accidents at leisure, in which the mobile phone had a partial effect, and at work the amounts were 4.5% and 0.4% respectively, during the last 12 months. Essentially, we found that: (1) men tend to have more close calls and accidents while on a mobile phone, (2) younger people tend to have more accidents and close calls while on a mobile phone, but it does not appear to be large enough to warrant intervention, (3) employed people tend to have more problems with mobile phone usage and accidents/close calls, and (4) there was a slight increase in mobile-phone-related accidents/close calls if the respondent also reported sleep disturbances and minor aches and pains. In the future, it is important to take into account and study how symptoms can increase the risk of accidents or close call situations in which a mobile phone has a partial effect.

Studies that show **Cell Phone** Health Effects

Hunton J, Rose JM. Cellular telephones and driving performance: the effects of attentional demands on motor vehicle crash risk. Risk Anal. 25(4):855-866, 2005.

This study examines the effects of conversation mode and split-attention communication training on driving performance. The study is based on an experiment where drivers with and without communication training (pilots vs. nonpilots) completed a simulated driving course while involved in one of three conversation modes: no conversation, conversation with passenger, or conversation on a hands-free cellular telephone. Results indicate that cellular telephone conversations consume more attention and interfere more with driving than passenger conversations. Cell phone conversations lack the nonverbal cues available during close-contact conversations and conversation participants expend significant cognitive resources to compensate for the lack of such cues. The results also demonstrate that communication training may reduce the hazardous effects of cell phone conversations on driving performance.

Hladky, A, Musil, J, Roth, Z, Urban, P, Blazkova, V, Acute effects of using a mobile phone on CNS functions. Cent Eur J Public Health 7(4):165-167. 1999.

Twenty volunteers participated in two experiments exploring the acute effects of using the mobile phone Motorola GSM 8700 on the functions of the CNS. When speaking (5 minutes reading a text from daily newspapers) the electromagnetic fields from the mobile apparatus did not affect the visual evoked potentials. Also a 6-min exposure did not reveal any effect of electromagnetic fields on the results in two tests (memory and attention) performed while speaking into the mobile. On the other hand the phone call itself strongly influenced the performance in a secondary task applying a test of switching attention which is a good model for driving a car. The response and decision speed were significantly worse. This is a proof that even a slight psychological stress involved in calling while driving can be a great risk.

Horberry T, Anderson J, Regan MA, Triggs TJ, Brown J. Driver distraction: The effects of concurrent in-vehicle tasks, road environment complexity and age on driving performance. Accid Anal Prev. 38(1):185-191. 2006.

This paper presents the findings of a simulator study that examined the effects of distraction upon driving performance for drivers in three age groups. There were two in-vehicle distracter tasks: operating the vehicle entertainment system and conducting a simulated hands-free mobile phone conversation. The effect of visual clutter was examined by requiring participants to drive in simple and complex road environments. Overall measures of driving performance were collected, together with responses to roadway hazards and subjective measures of driver perceived workload. The two in-vehicle distraction tasks degraded overall driving performance, degraded responses to hazards and increased subjective workload. The performance decrements that occurred as a result of in-vehicle distraction were observed in both the simple and complex highway environments and for drivers in different age groups. One key difference was that older drivers traveled at lower mean speeds in the complex highway environment compared with younger drivers. The conclusions of the research are that both in-vehicle tasks impaired several aspects of driving performance, with the entertainment system

Studies that show **Cell Phone** Health Effects

distracter having the greatest negative impact on performance, and that these findings were relatively stable across different driver age groups and different environmental complexities.

Jenness JW, Lattanzio RJ, O'Toole M, Taylor N, Pax C. Effects of manual versus voice-activated dialing during simulated driving. Percept Mot Skills 94(2):363-379, 2002.

We measured driving performance (lane-keeping errors, driving times, and glances away from the road scene) in a video driving simulator for 24 volunteers who each drove alone on a 10.6-km multicurved course while simultaneously placing calls on a mobile phone subscribed to a voice-activated dialing system. Driving performance also was measured for the same distance while participants manually dialed phone numbers and while they drove without dialing. There were 22% fewer lane-keeping errors ($p < .01$) and 56% fewer glances away from the road scene ($p < .01$) when they used voice-activated dialing as compared to manual dialing. Significantly longer driving times in both of the dialing conditions as compared to the No Dialing condition are discussed in terms of the hypothesis that drivers decrease driving speed to compensate for the demands of the secondary phone tasks.

Dreyer NA, Loughlin JE, Rothman KJ, Cause-specific mortality in cellular telephone users. JAMA 282(19):1814-1816, 1999.

A survey of standardized mortality rates (from cancer, circulatory diseases, and motor vehicle collisions) of 285,561 analog telephone users with known age, sex, and telephone type, showed that the only category of cause of death for which there was an indication of increasing risk with increasing minutes of phone use was motor vehicle collisions. Similar results were found for number of telephone calls per day. collisions were particularly hazardous (relative risk, 4.8 for calls placed within 5 minutes of the accident, as compared with 1.3 for calls placed more than 15 minutes before the accident; $P < 0.001$); and units that allowed the hands to be free (relative risk, 5.9) offered no safety advantage over hand-held units (relative risk, 3.9; P not significant). Thirty-nine percent of the drivers called emergency services after the collision, suggesting that having a cellular telephone may have had advantages in the aftermath of an event. CONCLUSIONS: The use of cellular telephones in motor vehicles is associated with a quadrupling of the risk of a collision during the brief time interval involving a call. Decisions about regulation of such telephones, however, need to take into account the benefits of the technology and the role of individual responsibility.

Cooper PJ, Zheng Y. Turning gap acceptance decision-making: the impact of driver distraction. J Safety Res 33(3):321-335, 2002.

PROBLEM: A number of studies have found that use of in-car phones by drivers can interfere with the cognitive processing necessary for making appropriate and timely vehicle control decisions. However, the specific linkage between communication-based distraction and unsafe decision-making has not been sufficiently explored. METHOD: In a closed-course driving experiment, 39 subjects were exposed to approximately 100 gaps each in a circulating traffic stream of eight vehicles on an instrumented test track

Studies that show **Cell Phone** Health Effects

that was wet about half the time. The subjects were at the controls of an instrumented car, which was oriented in a typical left-turn configuration (traffic-crossing situation in North America) and with parking brake on and the transmission in neutral. The subjects were instructed to press on the accelerator pedal when they felt that a gap was safe to accept. Their performances were monitored and incentives were provided for balancing safe decision-making with expeditious completion of the task. For half of the gap exposures (randomly assigned), each subject was required to listen and respond to a complex verbal message. RESULTS: When not distracted, the subjects' gap acceptance judgment was found to be significantly influenced by their age, the gap size, the speed of the trailing vehicle, the level of "indecision," and the condition of the track surface. However, when distracted, the subjects did not factor pavement surface condition into the decision process. On wet pavement, the subjects were judged to have initiated twice the level of potential collisions when distracted by the messages that they did when not distracted. DISCUSSION: Listening/responding to verbal messages may reduce the capacity of drivers to process adequately all the important information necessary for safe decision-making. The effects of the messages in our study seemed to cause the subjects to misjudge gap size and speed information when operating under the additional disadvantage of adverse pavement condition. SUMMARY: Attention to complex messages while making decisions about turning through gaps in an on-coming vehicle stream was associated with significantly increased unsafe decision making by subjects in our experiment when the additional complexity of wet surface condition was introduced. IMPACT ON INDUSTRY: While the results reflected a somewhat artificial situation where the measure was signaled intention to act rather than the act itself, nevertheless, they do strongly suggest a scenario in which mental distraction could contribute to crash risk. With the rapid proliferation of telematics in the vehicle market, even with the laudable objectives represented by the Intelligent Transportation Systems initiative, there is a danger of the primary task of the driver being subordinated to a perceived need to enhance information flow to/from the external "world." Industry and governments need to work together to ensure that apparently desirable in-vehicle communication improvements do not compromise safety.

Consiglio W, Driscoll P, Witte M, Berg WP. Effect of cellular telephone conversations and other potential interference on reaction time in a braking response. *Accid Anal Prev* 35(4):495-500, 2003.

This experiment studied the effect of phone conversations and other potential interference on reaction time (RT) in a braking response. Using a laboratory station which simulated the foot activity in driving, 22 research participants were requested to release the accelerator pedal and depress the brake pedal as quickly as possible following the activation of a red brake lamp. Mean reaction time was determined for five conditions: (a) control, (b) listening to a radio, (c) conversing with a passenger, (d) conversing using a hand-held phone, and (e) conversing using a hands-free phone. Results indicated that conversation, whether conducted in-person or via a cellular phone caused RT to slow, whereas listening to music on the radio did not.

Studies that show **Cell Phone** Health Effects

Charlton SG. Perceptual and attentional effects on drivers' speed selection at curves. *Accid Anal Prev.* 36(5):877-884, 2004.

This paper describes an experiment comparing the relative effectiveness of various types of warnings on drivers' speed selection at curves. The experiment compared three types of curve warnings across three different curve types in a driving simulator. All of the warnings worked reasonably well for severe curves (45 km/h), regardless of demands from a secondary (cell phone) task. For less demanding curves, only those warnings with a strong perceptual component (i.e., implicit cues) were effective in reducing drivers' curve speeds in the presence of the cell phone task. The design implications of these data appear straightforward; curve warnings that contain perceptual components or emphasise the physical features of the curve work best, particularly in cognitively demanding situations. The cell phone task added to driver workload and drivers became less responsive to primary task demands (i.e., speeds were elevated and reaction times were longer).

Astrain I, Bernaus J, Claverol J, Escobar A, Godoy P. [Prevalence of mobile phone use while driving vehicles] *Gac Sanit* 17(1):66-69, 2003. [Article in Spanish]

Objective: To estimate the prevalence of mobile telephone use while driving vehicles in the city of Lleida (Spain). Methods: A random sample of 1536 cars passing through six intersections regulated by traffic lights in Lleida were selected (three with urban traffic and three with interurban traffic). Cyclists, motorcyclists and driving school cars were excluded. The variables studied were mobile telephone use, age, (18-40; 41-60; >61), sex, the presence of passengers, type of intersection (urban traffic/interurban traffic), day of the week (working day/weekend or holiday) and hour of the day (rush hour/non-rush hour). The prevalence of mobile telephone use was calculated in percentages with a 95% CI. The relationship among the dependent variable (mobile telephone use) and the other independent variables was studied using odds ratios (OR) and 95% CI. Results: A total of 1536 direct observations were made and mobile telephone use was detected in 50 drivers. The prevalence was 3.3 (95% CI, 2.4-4.3). The prevalence was higher in men (OR = 2.2; 95% CI, 1.0-5.7), in drivers aged more than 60 years old (OR = 2.2; 95% CI, 0.5-8.4) and in those aged 18-40 years old (OR = 1.5; 95% CI, 0.8-3.0), in unaccompanied drivers (OR = 3.0; 95% CI, 1.5-6.3), in urban intersections (OR = 2.7; 95% CI, 1.2-5.9), on workdays (OR = 2.0; 95% CI, 0.9-4.4) and at the rush hour (OR = 1.4; 95% CI, 0.8-2.4). Conclusions: The prevalence of mobile telephone use while driving vehicles can be considered high, because of the increase in car accidents. The profile of drivers using mobile telephones corresponds to men aged 18-40 years or more than 61 years, in urban intersections, without passengers, during workdays and at the rush hour. We recommend the implementation of measures to decrease the use of mobile telephones while driving

Asbridge M, Brubacher JR, Chan H. Cell phone use and traffic crash risk: a culpability analysis. *Int J Epidemiol.* 2012 Nov 18. [Epub ahead of print]

BACKGROUND: The use of a cell phone or communication device while driving is illegal in many jurisdictions, yet evidence evaluating the crash risk associated with cell phone

Studies that show **Cell Phone** Health Effects

use in naturalistic settings is limited. This article aims to determine whether cell phone use while driving increases motor vehicle crash culpability. **METHOD:** Drivers involved in crashes where police reported cell phone use (n = 312) and propensity matched drivers (age, sex, suspect alcohol/drug impairment, crash type, date, time of day, geographical location) without cell phone use (n = 936) were drawn from Insurance Corporation of British Columbia Traffic Accident System data. A standardized scoring tool, modified to account for Canadian driving conditions, was used to determine crash culpability from police reports on all drivers from the crashes. The association between crash culpability and cell phone use was determined, with additional subgroup analyses based on crash severity, driver characteristics and type of licence. **RESULTS:** A comparison of crashes with vs without cell phones revealed an odds ratio of 1.70 (95% confidence interval 1.22-2.36; P = 0.002). This association was consistent after adjustment for matching variables and other covariates. Subgroup analyses demonstrated an association for male drivers, unimpaired drivers, injured and non-injured drivers, and for drivers aged between 26 and 65 years. **CONCLUSIONS:** Crash culpability was found to be significantly associated with cell phone use by drivers, increasing the odds of a culpable crash by 70% compared with drivers who did not use a cell phone. This increased risk was particularly high for middle-aged drivers.

Atchley P, Dressel J. Conversation limits the functional field of view. Hum Factors. 46(4):664-673, 2004.

The purpose of these two experiments is to investigate one possible mechanism that might account for an increase in crash risk with in-car phone use: a reduction in the functional field of view. In two between-subjects experiments, college undergraduates performed a task designed to measure the functional field of view in isolation and while performing a hands-free conversational task. In both experiments, the addition of the conversational task led to large reductions in the functional field of view. Because similar reductions have been shown to increase crash risk, reductions in the functional field of view by conversation may be an important mechanism involved in increased risk for crashes with in-car phone use. Actual or potential applications of this research include improving driver performance.

Barkana Y, Zadok D, Morad Y, Avni I. Visual field attention is reduced by concomitant hands-free conversation on a cellular telephone. Am J Ophthalmol. 138(3):347-353, 2004.

PURPOSE: To quantify the central attention-diverting effect of hands-free cellular phone conversation on visual field awareness. **DESIGN:** Experimental study. **METHODS:** Twenty male and 21 female healthy participants performed a pretest and baseline Esterman visual field examinations with the Humphrey Systems Visual Field Analyzer II. During the consequent third examination, each participant engaged in a hands-free conversation using a cellular phone. The conversation was the same for all participants. Visual field performance parameters were compared between the second (baseline) examination, and the third (test) examination for each eye. **RESULTS:** During phone conversation, missed points increased from mean 1.0 +/- 1.5 to 2.6 +/- 3.4 (P <= .001) in the right eye

Studies that show **Cell Phone** Health Effects

and from 1.1 +/- 1.53 to 3.0 +/- 3.4 ($P < .001$) in the left eye. Fixation loss increased from mean 7.8% to 27.4% ($P < .0001$) and from 7.2% to 34.8% ($P < .0001$) for the right and left eyes, respectively. Test duration increased by a mean of 0.28 seconds (15%) per stimulus ($P < .0001$). Approximately half of missed points were inside the central 30 degrees. There was no significant difference in the performance of male and female participants. **CONCLUSION:** We describe a new model for the quantification of the attention-diverting effect of cellular-phone conversation on the visual field. In the current study, cellular hands-free conversation caused some subjects to miss significantly more points, react slower to each stimulus, and perform with reduced precision. Legislative restrictions on concomitant cellular-phone conversation and driving may need to be based on individual performance rather than a general ban on cellular phone usage.

Beede KE, Kass SJ. Engrossed in conversation: The impact of cell phones on simulated driving performance. *Accid Anal Prev*38(2):415-21, 2006.

The current study examined the effects of cognitively distracting tasks on various measures of driving performance. Thirty-six college students with a median of 6 years of driving experience completed a driving history questionnaire and four simulated driving scenarios. The distraction tasks consisted of responding to a signal detection task and engaging in a simulated cell phone conversation. Driving performance was measured in terms of four categories of behavior: traffic violations (e.g., speeding, running stop signs), driving maintenance (e.g., standard deviation of lane position), attention lapses (e.g., stops at green lights, failure to visually scan for intersection traffic), and response time (e.g., time to step on brake in response to a pop-up event). Performance was significantly impacted in all four categories when drivers were concurrently talking on a hands-free phone. Performance on the signal detection task was poor and not significantly impacted by the phone task, suggesting that considerably less attention was paid to detecting these peripheral signals. However, the signal detection task did interact with the phone task on measures of average speed, speed variability, attention lapses, and reaction time. The findings lend further empirical support of the dangers of drivers being distracted by cell phone conversations.

Overlooked Health Effects in Science

Marino AA, Carrubba S. The effects of mobile-phone electromagnetic fields on brain electrical activity: a critical analysis of the literature. *Electromagn Biol Med.* 28(3):250-274, 2009.

We analyzed the reports in which human brain electrical activity was compared between the presence and absence of radio-frequency and low-frequency electromagnetic fields (EMFs) from mobile phones, or between pre- and post-exposure to the EMFs. Of 55 reports, 37 claimed and 18 denied an EMF-induced effect on either the baseline electroencephalogram (EEG), or on cognitive processing of visual or auditory stimuli as reflected in changes in event-related potentials. The positive reports

Studies that show **Cell Phone** Health Effects

did not adequately consider the family-wise error rate, the presence of spike artifacts in the EEG, or the confounding role of the two different EMFs. The negative reports contained neither positive controls nor power analyses. Almost all reports were based on the incorrect assumption that the brain was in equilibrium with its surroundings. Overall, the doubt regarding the existence of reproducible mobile-phone EMFs on brain activity created by the reports appeared to legitimate the knowledge claims of the mobile-phone industry. However, it funded, partly or wholly, at least 87% of the reports. From an analysis of their cognitive framework, the common use of disclaimers, the absence of information concerning conflicts of interest, and the industry's donations to the principal EMF journal, we inferred that the doubt was manufactured by the industry. The crucial scientific question of the pathophysiology of mobile-phone EMFs as reflected in measurements of brain electrical activity remains unanswered, and essentially unaddressed.

Wilen J, Sandstrom M, Hansson Mild K. Subjective symptoms among mobile phone users-A consequence of absorption of radiofrequency fields? Bioelectromagnetics 24(3):152-159, 2003.

In a previous epidemiological study, where we studied the prevalence of subjective symptoms among mobile phone (MP) users, we found as an interesting side finding that the prevalence of many of the subjective symptoms increased with increasing calling time and number of calls per day. In this extrapolative study, we have selected 2402 people from the epidemiological study who used any of the four most common GSM MP. We used the information about the prevalence of symptoms, calling time per day, and number of calls per day and combined it with measurements of the Specific Absorption Rate (SAR). We defined three volumes in the head and measured the maximum SAR averaged over a cube of 1 g tissue (SAR(1g)) in each volume. Two new exposure parameters Specific Absorption per Day (SAD) and Specific Absorption per Call (SAC) have been devised and are obtained as combinations of SAR, calling time per day, and number of calls per day, respectively. The results indicates that SAR values >0.5 W/kg may be an important factor for the prevalence of some of the symptoms, especially in combination with long calling times per day.

Marjanović AM, Pavičić I, Trošić I Biological indicators in response to radiofrequency/microwave exposure. Arh Hig Rada Toksikol. 2012 Sep 25;63(3):407-416, 2012.

Over the years, due to rapid technological progress, radiation from man-made sources exceeded that of natural origin. There is a general concern regarding a growing number of appliances that use radiofrequency/ microwave (RF/MW) radiation with particular emphasis on mobile communication systems. Since nonthermal biological effects and mechanisms of RF/MW radiation are still uncertain, laboratory studies on animal models, tissues, cells, and cell free system are of extraordinary importance in bioelectromagnetic research. We believe that such investigations play a supporting role in public risk assessment. Cellular systems with the potential for a clear response to RF/MW exposures should be used in those studies. It is known that organism is a

Studies that show **Cell Phone** Health Effects

complex electrochemical system where processes of oxidation and reduction regularly occur. One of the plausible mechanisms is connected with generation of reactive oxygen species (ROS). Depending on concentration, ROS can have both beneficial and deleterious effects. Positive effects are connected with cell signalling, defence against infectious agents, and proliferative cell ability. On the other hand, excessive production, which overloads antioxidant defence mechanism, leads to cellular damage with serious potential for disease development. ROS concentration increase within the cell caused by RF/MW radiation seems to be a biologically relevant hypothesis to give clear insight into the RF/MW action at non-thermal level of radiation. In order to better understand the exact mechanism of action and its consequences, further research is needed in the field. We would like to present current knowledge on possible biological mechanisms of RF/MW actions.

Lauer O, Frei P, Gosselin MC, Joseph W, Rösli M, Fröhlich J. Combining near- and far-field exposure for an organ-specific and whole-body RF-EMF proxy for epidemiological research: A reference case. *Bioelectromagnetics*. 2013 Feb 15. doi: 10.1002/bem.21782. [Epub ahead of print]

A framework for the combination of near-field (NF) and far-field (FF) radio frequency electromagnetic exposure sources to the average organ and whole-body specific absorption rates (SARs) is presented. As a reference case, values based on numerically derived SARs for whole-body and individual organs and tissues are combined with realistic exposure data, which have been collected using personal exposure meters during the Swiss Qualifex study. The framework presented can be applied to any study region where exposure data is collected by appropriate measurement equipment. Based on results derived from the data for the region of Basel, Switzerland, the relative importance of NF and FF sources to the personal exposure is examined for three different study groups. The results show that a 24-h whole-body averaged exposure of a typical mobile phone user is dominated by the use of his or her own mobile phone when a Global System for Mobile Communications (GSM) 900 or GSM 1800 phone is used. If only Universal Mobile Telecommunications System (UMTS) phones are used, the user would experience a lower exposure level on average caused by the lower average output power of UMTS phones. Data presented clearly indicate the necessity of collecting band-selective exposure data in epidemiological studies related to electromagnetic fields.

Huss A, Egger M, Hug K, Huwiler-Müntener K, Rösli M, Gomes D, Da Ros MA Source of funding and results of studies of health effects of mobile phone use: systematic review of experimental studies. *Cien Saude Colet*. 13(3):1005-1012, 2008.

There is concern regarding the possible health effects of cellular telephone use. We conducted a systematic review of studies of controlled exposure to radiofrequency radiation with health-related outcomes (electroencephalogram, cognitive or cardiovascular function, hormone levels, symptoms, and subjective well-being). We searched Embase, Medline, and a specialist database in February 2005 and scrutinized reference lists from relevant publications. Data on the source of funding, study design,

Studies that show **Cell Phone** Health Effects

methodologic quality, and other study characteristics were extracted. The primary outcome was the reporting of at least one statistically significant association between the exposure and a health-related outcome. Data were analyzed using logistic regression models. Of 59 studies, 12 (20%) were funded exclusively by the telecommunications industry, 11 (19%) were funded by public agencies or charities, 14 (24%) had mixed funding (including industry), and in 22 (37%) the source of funding was not reported. Studies funded exclusively by industry reported the largest number of outcomes, but were least likely to report a statistically significant result. The interpretation of results from studies of health effects of radiofrequency radiation should take sponsorship into account.

Hutter HP, Moshammer H, Wallner P, Kundi M. Public perception of risk concerning cell towers and mobile phones. *Soz Präventivmed.* 49(1):62-66, 2004.

OBJECTIVE: The controversy about health risks of electromagnetic fields (EMF) has contributed in raising fears concerning emissions from celltowers. The study was to examine whether or not neighbours of celltowers are particularly concerned about adverse health effects of mobile phones and their base stations. METHODS: Prior to information delivered by medical doctors of the Institute of Environmental Health at public hearings a questionnaire was handed out to participants asking for their personal rating of several environmental health risks including those of mobile telecommunication (n = 123, response rate approx. 48%). Medical students (n = 366) served as a contrast group. RESULTS: Participants rated health risk for both, mobile phones and celltowers higher as students. A trend for higher ratings was also seen with older subjects and female sex. The risk ratings of both exposures correlated well with each other. The magnitude of the perceived risks, however, resembled that of other ubiquitous exposures like traffic noise and air pollution. CONCLUSION: Contrary to the claims of the telecommunication industry, opponents of celltowers generally do not express unusual fears concerning electromagnetic field exposure. The outcome of our study indicates that the risk rating is comparable with other perceived common hazards of the civilised world. It is hypothesised that offering information and participation to the concerned population will be efficient in reducing exaggerated fears.

Redmayne M, Inyang I, Dimitriadis C, Benke G, Abramson MJ. Cordless telephone use: implications for mobile phone research. *J Environ Monit.* 12(4):809-812, 2010.

Cordless and mobile (cellular) telephone use has increased substantially in recent years causing concerns about possible health effects. This has led to much epidemiological research, but the usual focus is on mobile telephone radiofrequency (RF) exposure only despite cordless RF being very similar. Access to and use of cordless phones were included in the Mobile Radiofrequency Phone Exposed Users Study (MoRPhEUS) of 317 Year 7 students recruited from Melbourne, Australia. Participants completed an exposure questionnaire-87% had a cordless phone at home and 77% owned a mobile phone. There was a statistically significant positive relationship ($r = 0.38$, $p < 0.01$) between cordless and mobile phone use. Taken together, this increases total RF exposure and its ratio in high-to-low mobile users. Therefore, the design and analysis of

Studies that show **Cell Phone** Health Effects

future epidemiological telecommunication studies need to assess cordless phone exposure to accurately evaluate total RF telephone exposure effects.

Studies that show **Cell Tower** Health Effects

Table of Contents

78 Studies Showing Health Effects from Cell Tower Radio Frequency Radiation

Health Effects	Pages
Cancer	Pages 2-4
Neurological Effects	Pages 5-7
Effects on Reproductive Organs and Fertility Effects	Pages 7-9
Effects on Wellbeing	Pages 9-14
Effects on Sleep	Pages 14-15
Effects on Cells	Pages 16
Effects on Eyes	Pages 17-18
Oxidative Stress	Pages 18-20
Effects on Blood	Pages 20-21
Death	Page 22-23
Effects on Hormones	Pages 23-24
Genetic Effects	Pages 24-29
Effects on Glands	Pages 29
Effects on Animals and Environment	Page 29-32
Effects on Skin	Page 32
Effects on Protein	Pages 33
Effects on Immune Function	Page 33-34
Electromagnetic Hyper-Sensitivity	Pages 34
RF Levels in Cities	Pages 34-36
Effects on Water	Page 36
RF Levels From Cell Towers	Pages 36-37
Danger Perception and Symptoms	Pages 37-38
Reason Why Not A Lot of Studies on Cell Towers	Page 38-39
Miscellaneous Effects	Pages 39

Studies that show **Cell Tower** Health Effects

Cancer

Wolf R, Wolf D. Increased incidence of cancer near a cell-phone transmitter station. Inter J Cancer Prev 1(2):123-128, 2004.

Significant concern has been raised about possible health effects from exposure to radiofrequency (RF) electromagnetic fields, especially after the rapid introduction of mobile telecommunication systems. Parents are especially concerned with the possibility that children might develop cancer after exposure to the RF emissions from mobile telephone base stations erected in or near schools. The few epidemiologic studies that did report on cancer incidence in relation to RF radiation have generally presented negative or inconsistent results, and thus emphasized the need for more studies that should investigate cohorts with high RF exposure for changes in cancer incidence. The aim of this study is to investigate whether there is an increased cancer incidence in populations, living in a small area, and exposed to RF radiation from a cell-phone transmitter station.

This is an epidemiologic assessment, to determine whether the incidence of cancer cases among individuals exposed to a cell-phone transmitter station is different from that expected in Israel, in Netanya, or as compared to people who lived in a nearby area. Participants are people (n=622) living in the area near a cell-phone transmitter station for 3-7 years who were patients of one health clinic (of DW). The exposure began 1 year before the start of the study when the station first came into service. A second cohort of individuals (n=1222) who get their medical services in a clinic located nearby with very closely matched, environment, workplace and occupational characteristics was used for comparison.

In the area of exposure (area) eight cases of different kinds of cancer were diagnosed in a period of only one year. This rate of cancers was compared both with the rate of 31 cases per 10,000 per year in the general population and the 2/1222 rate recorded in the nearby clinic (area B). Relative cancer rates for female were 10.5 for area A. 0.6 for area B and 1 for the whole town of Netanya. Cancer incidence of women in area A was thus significantly higher ($p < 0.0001$) compared with that of area B and the whole city. A comparison of the relative risk revealed that there were 4.15 times more cases in area than in the entire population. The study indicates an association between increased incidence of cancer and living in proximity to a cell-phone transmitter station.

Yakymenko I, Sidorik E, Kyrylenko S, Chekhun V. Long-term exposure to microwave radiation provokes cancer growth: evidences from radars and mobile communication systems. Exp Oncol. 33(2):62-70, 2011.

In this review we discuss alarming epidemiological and experimental data on possible carcinogenic effects of long term exposure to low intensity microwave (MW) radiation. Recently, a number of reports revealed that under certain conditions the irradiation by low intensity MW can substantially induce cancer progression in humans and in animal models. The carcinogenic effect of MW irradiation is typically manifested after long term (up to 10 years and more) exposure. Nevertheless, even a year of operation of a

Studies that show **Cell Tower** Health Effects

powerful base transmitting station for mobile communication reportedly resulted in a dramatic increase of cancer incidence among population living nearby. In addition, model studies in rodents unveiled a significant increase in carcinogenesis after 17-24 months of MW exposure both in tumor-prone and intact animals. To that, such metabolic changes, as overproduction of reactive oxygen species, 8-hydroxi-2-deoxyguanosine formation, or ornithine decarboxylase activation under exposure to low intensity MW confirm a stress impact of this factor on living cells. We also address the issue of standards for assessment of biological effects of irradiation. It is now becoming increasingly evident that assessment of biological effects of non-ionizing radiation based on physical (thermal) approach used in recommendations of current regulatory bodies, including the International Commission on Non-Ionizing Radiation Protection (ICNIRP) Guidelines, requires urgent reevaluation. We conclude that recent data strongly point to the need for re-elaboration of the current safety limits for non-ionizing radiation using recently obtained knowledge. We also emphasize that the everyday exposure of both occupational and general public to MW radiation should be regulated based on a precautionary principles which imply maximum restriction of excessive exposure.

Michelozzi P, Ancona C, Fusco D, Forastiere F, Perucci CA, Risk of leukemia and residence near a radio transmitter in Italy. Epidemiology 9 (Suppl) 354p, 1998.

We conducted a small area study to investigate a cluster of leukemia near a high power radio-transmitter in a peripheral area of Rome. The leukemia mortality within 3.5 km (5,863 inhabitants) was higher than expected (SMR=2.5, 95% confident interval 1.07-4.83); the excess was due to a significant higher mortality among men (7 cases observed, SMR=3.5). The results of the Stone's test, after adjusting for socio-economic confounding, showed a significant decline in risk with distance from the transmitter only among men (p=0.005), whereas the p-value for both sexes was p=0.07.

Lourencini da Silva R , Albano F, Lopes dos Santos LR , Tavares AD, Felzenszwalb I, The effect of electromagnetic field exposure on the formation of DNA lesions. Redox Rep 5(5):299-301, 2000.

In an attempt to determine whether electromagnetic field (EMF) exposure might lead to DNA damage, we exposed SnCl₂-treated pBR322 plasmids to EMF and analysed the resulting conformational changes using agarose gel electrophoresis. An EMF-dependent potentiation of DNA scission (i.e. the appearance of relaxed plasmids) was observed. In confirmation of this, plasmids pre-exposed to EMF also were less capable of transforming Escherichia coli. The results indicate that EMF, in the presence of a transition metal, is capable of causing DNA damage. These observations support the idea that EMF, probably through secondary generation of reactive oxygen species, can be clastogenic and provide a possible explanation for the observed correlation between EMF exposure and the frequency of certain types of cancers in humans.

Li CY, Liu CC, Chang YH, Chou LP, Ko MC. A population-based case-control study of radiofrequency exposure in relation to childhood neoplasm. Sci Total Environ. 435-436:472-478, 2012.

Studies that show **Cell Tower** Health Effects

This population-based case-control study in Taiwan considered incident cases aged 15 years or less and admitted in 2003 to 2007 for all neoplasm (ICD-9-CM: 140-239) (n=2606), including 939 leukemia and 394 brain neoplasm cases. Controls were randomly selected, with a case/control ratio of 1:30 and matched on year of birth, from all non-neoplasm children insured in the same year when the index case was admitted. Annual summarized power (ASP, watt-year) was calculated for each of the 71,185 mobile phone base stations (MPBS) in service between 1998 and 2007. Then, the **annual power density (APD, watt-year/km(2))** of each township (n=367) was computed as a ratio of the total ASP of all MPBS in a township to the area of that particular township. Exposure of each study subject to radio frequency (RF) was indicated by the averaged APD within 5 years prior to the neoplasm diagnosis (cases) or July 1st of the year when the index case was admitted (controls) in the township where the subject lived. Unconditional logistic regression model with generalized estimation equation was employed to calculate the covariate-**adjusted odds ratio [AOR]** of childhood neoplasm in relation to RF exposure. A higher than median averaged APD (approximately 168 WYs/km(2)) was significantly associated with an increased AOR for all neoplasms (1.13; 1.01 to 1.28), but not for leukemia (1.23; 0.99 to 1.52) or brain neoplasm (1.14, 0.83 to 1.55). This study noted a significantly increased risk of all neoplasms in children with higher-than-median RF exposure to MPBS. The slightly elevated risk was seen for leukemia and brain neoplasm, but was not statistically significant. These results may occur due to several methodological limitations.

Johnson EH, Chima SC, Muirhead DE, A cerebral primitive neuroectodermal tumor in a squirrel monkey (*Saimiri sciureus*). J Med Primatol 28(2):91-96, 1999.

An adult squirrel monkey with a history of long-term exposure to microwave radiation was found at necropsy to have a malignant tumor of the right cerebral cortex. Gross examination revealed a mass with expanding borders in the right frontoparietal cortex with compression of the adjacent lateral ventricle. Microscopy revealed a tumor composed of sheets of moderate-sized cells, resembling an oligodendroglioma, with clear cytoplasm and central nuclei interrupted by delicate vasculature. Malignant features were present in the form of marked nuclear pleomorphism, frequent mitotic figures, and focal necrosis. A neuronal cell origin for this tumor was supported by immunohistochemical analysis, which revealed immunopositivity for neurofilament proteins and neuron-specific enolase. Staining for vimentin and glial fibrillary acid protein was negative, except in reactive astrocytes at the tumor margins and adjacent to intra-tumoral blood vessels. Antibody activity against Ki-67 antigen, a marker of rapidly proliferating tumor cells, and p53 oncoprotein was strongly positive, indicative of the aggressive and malignant nature of this tumor. The tumor was diagnosed as a cerebral primitive neuroectodermal tumor.

Neurological Effects

Khurana VG, Hardell L, Everaert J, Bortkiewicz A, Carlberg M, Ahonen M.

Studies that show **Cell Tower** Health Effects

Epidemiological evidence for a health risk from mobile phone base stations. *Int J Occup Environ Health.* 16(3):263-267, 2010.

Human populations are increasingly exposed to microwave/radiofrequency (RF) emissions from wireless communication technology, including mobile phones and their base stations. By searching PubMed, we identified a total of 10 epidemiological studies that assessed for putative health effects of mobile phone base stations. Seven of these studies explored the association between base station proximity and neurobehavioral effects and three investigated cancer. We found that eight of the 10 studies reported increased prevalence of adverse neurobehavioral symptoms or cancer in populations living at distances < 500 meters from base stations. None of the studies reported exposure above accepted international guidelines, suggesting that current guidelines may be inadequate in protecting the health of human populations. We believe that comprehensive epidemiological studies of long-term mobile phone base station exposure are urgently required to more definitively understand its health impact.

Hocking B, Westerman R. Neurological abnormalities associated with CDMA exposure. *Occup Med (Lond)* 51(6):410-413, 2001.

Dysaesthesiae of the scalp and neurological abnormality after mobile phone use have been reported previously, but the roles of the phone per se or the radiations in causing these findings have been questioned. We report finding a neurological abnormality in a patient after accidental exposure of the left side of the face to mobile phone radiation [code division multiple access (CDMA)] from a down-powered mobile phone base station antenna. He had headaches, unilateral left blurred vision and pupil constriction, unilateral altered sensation on the forehead, and abnormalities of current perception thresholds on testing the left trigeminal ophthalmic nerve. His nerve function recovered during 6 months follow-up. His exposure was 0.015-0.06 mW/cm(2) over 1-2 h. The implications regarding health effects of radiofrequency radiation are discussed.

Abdel-Rassoul G, El-Fateh OA, Salem MA, Michael A, Farahat F, El-Batanouny M, Salem E. Neurobehavioral effects among inhabitants around mobile phone base stations. *Neurotoxicology.* 28(2):434-40, 2007.

BACKGROUND: There is a general concern on the possible hazardous health effects of exposure to radiofrequency electromagnetic radiations (RFR) emitted from mobile phone base station antennas on the human nervous system. AIM: To identify the possible neurobehavioral deficits among inhabitants living nearby mobile phone base stations. METHODS: A cross-sectional study was conducted on (85) inhabitants living nearby the first mobile phone station antenna in Menoufiya governorate, Egypt, 37 are living in a building under the station antenna while 48 opposite the station. A control group (80) participants were matched with the exposed for age, sex, occupation and educational level. All participants completed a structured questionnaire containing: personal, educational and medical histories; general and neurological examinations; neurobehavioral test battery (NBTB) [involving tests for visuomotor speed, problem solving, attention and memory]; in addition to Eysenck personality questionnaire (EPQ). RESULTS: The prevalence of neuropsychiatric complaints as headache (23.5%), memory

Studies that show **Cell Tower** Health Effects

changes (28.2%), dizziness (18.8%), tremors (9.4%), depressive symptoms (21.7%), and sleep disturbance (23.5%) were significantly higher among exposed inhabitants than controls: (10%), (5%), (5%), (0%), (8.8%) and (10%), respectively (P<0.05). The NBTB indicated that the exposed inhabitants exhibited a significantly lower performance than controls in one of the tests of attention and short-term auditory memory [Paced Auditory Serial Addition Test (PASAT)]. Also, the inhabitants opposite the station exhibited a lower performance in the problem solving test (block design) than those under the station. All inhabitants exhibited a better performance in the two tests of visuomotor speed (Digit symbol and Trailmaking B) and one test of attention (Trailmaking A) than controls. The last available measures of RFR emitted from the first mobile phone base station antennas in Menoufiya governorate were less than the allowable standard level. CONCLUSIONS AND RECOMMENDATIONS: Inhabitants living nearby mobile phone base stations are at risk for developing neuropsychiatric problems and some changes in the performance of neurobehavioral functions either by facilitation or inhibition. So, revision of standard guidelines for public exposure to RER from mobile phone base station antennas and using of NBTB for regular assessment and early detection of biological effects among inhabitants around the stations are recommended.

Akbari A, Jelodar G, Nazifi S. Vitamin C protects rat cerebellum and encephalon from oxidative stress following exposure to radiofrequency wave generated by a BTS antenna model. Toxicol Mech Methods. 24(5):347-352, 2014.

Radio frequency wave (RFW) generated by base transceiver station has been reported to produce deleterious effects on the central nervous system function, possibly through oxidative stress. This study was conducted to evaluate the effect of RFW-induced oxidative stress in the cerebellum and encephalon and the prophylactic effect of vitamin C on these tissues by measuring the antioxidant enzymes activity, including: glutathione peroxidase, superoxide dismutase, catalase, and malondialdehyde (MDA). Thirty-two adult male Sprague-Dawley rats were randomly divided into four equal groups. The control group; the control-vitamin C group received L-ascorbic acid (200 mg/kg of body weight/day by gavage) for 45 days. The RFW group was exposed to RFW and the RFW+ vitamin C group was exposed to RFW and received vitamin C. At the end of the experiment, all groups were killed and encephalon and cerebellum of all rats were removed and stored at -70 °C for measurement of antioxidant enzymes activity and MDA. The results indicate that exposure to RFW in the test group decreased antioxidant enzymes activity and increased MDA compared with the control groups (p < 0.05). The protective role of vitamin C in the treated group improved antioxidant enzymes activity and reduced MDA compared with the test group (p < 0.05). It can be concluded that RFW causes oxidative stress in the brain and vitamin C improves the antioxidant enzymes activity and decreases MDA.

Bak M, Dudarewicz A, Zmyślony M, Sliwinska-Kowalska M. Effects of GSM signals during exposure to event related potentials (ERPs). Int J Occup Med Environ Health. 23(2):191-199, 2010.

Studies that show **Cell Tower** Health Effects

Objectives: The primary aim of this work was to assess the effect of electromagnetic field (EMF) from the GSM mobile phone system on human brain function. The assessment was based on the assay of event related potentials (ERPs). **Material and Methods:** The study group consisted of 15 volunteers, including 7 men and 8 women. The test protocol comprised determination of P300 wave in each volunteer during exposure to the EMF. To eliminate possible effects of the applied test procedure on the final result, the test was repeated without EMF exposure. P300 latency, amplitude, and latency of the N1, N2, P2 waves were analysed. **Results:** The statistical analysis revealed an effect of EMF on P300 amplitude. In the experiment with EMF exposure, lower P300 amplitudes were observed only at the time in which the volunteers were exposed to EMF; when the exposure was discontinued, the values of the amplitude were the same as those observed before EMF application. No such change was observed when the experiment was repeated with sham exposure, which may be considered as an indirect proof that lower P300 amplitude values were due to EMF exposure. No statistically significant changes were noted in the latencies of the N1, N2, P2 waves that precede the P300 wave, nor in the latency of the P300 itself. **Conclusions:** The results suggest that exposure to GSM EMF exerts some effects on CNS, including effects on long latency ERPs.

Hinrichs H, Heinze HJ. Effects of GSM electromagnetic field on the MEG during an encoding-retrieval task. *Neuroreport*. 15(7):1191-1194, 2004.

Potential effects of GSM 1800 electromagnetic fields (EMF) on verbal memory encoding were investigated by recording event-related magnetic fields (ERMF) from the brain during subsequent memory retrieval. Twelve normal subjects participated in the study. After encoding words from a study list presented in the first phase they had to discriminate old from new words mixed together in a test list presented during the second phase. All subjects performed two experimental sessions, one with exposure to EMF during the study phase, and one without. Exposure to EMF changed an early (350-400 ms) task-specific component of the ERMF indicating an interference of EMF and item encoding. Behavioural measures were not significantly affected. Adverse health effects cannot be derived from these data.

Effects on Reproductive Organs and Fertility Effects

Magras, IN, Xenos, TD, RF radiation-induced changes in the prenatal development of mice. *Bioelectromagnetics* 18(6):455-461, 1997.

The possible effects of radiofrequency (RF) radiation on prenatal development has been investigated in mice. This study consisted of RF level measurements and in vivo experiments at several places around an "antenna park." At these locations RF power densities between 168 nW/cm² and 1053 nW/cm² were measured. Twelve pairs of mice, divided in two groups, were placed in locations of different power densities and were repeatedly mated five times. One hundred eighteen newborns were collected. They were measured, weighed, and examined macro- and microscopically. A progressive decrease in the number of newborns

Studies that show **Cell Tower** Health Effects

per dam was observed, which ended in irreversible infertility. The prenatal development of the newborns, however, evaluated by the crown-rump length, the body weight, and the number of the lumbar, sacral, and coccygeal vertebrae, was improved.

Jelodar G, Nazifi S, Akbari A. The prophylactic effect of vitamin C on induced oxidative stress in rat testis following exposure to 900 MHz radio frequency wave generated by a BTS antenna model. Electromagn Biol Med. 2013 Jan 16. [Epub ahead of print]
Radio frequency wave (RFW) generated by base transceiver station (BTS) has been reported to make deleterious effects on reproduction, possibly through oxidative stress. This study was conducted to evaluate the effect of RFW generated by BTS on oxidative stress in testis and the prophylactic effect of vitamin C by measuring the antioxidant enzymes activity, including glutathione peroxidase, superoxide dismutase (SOD) and catalase, and malondialdehyde (MDA). Thirty-two adult male Sprague-Dawley rats were randomly divided into four experimental groups and treated daily for 45 days as follows: sham, sham+vitamin C (l-ascorbic acid 200 mg/kg of body weight/day by gavage), RFW (exposed to 900 MHz RFW) 'sham' and 'RFW' animals were given the vehicle, i.e., distilled water and the RFW+vitamin C group (received vitamin C in addition to exposure to RFW). At the end of the experiment, all the rats were sacrificed and their testes were removed and used for measurement of antioxidant enzymes and MDA activity. The results indicate that exposure to RFW in the test group decreased antioxidant enzymes activity and increased MDA compared with the control groups ($p < 0.05$). In the treated group, vitamin C improved antioxidant enzymes activity and reduced MDA compared with the test group ($p < 0.05$). It can be concluded that RFW causes oxidative stress in testis and vitamin C improves the antioxidant enzymes activity and decreases MDA.

Lukac N, Massanyi P, Roychoudhury S, Capcarova M, Tvrda E, Knazicka Z, Kolesarova A, Danko J. In vitro effects of radiofrequency electromagnetic waves on bovine spermatozoa motility. J Environ Sci Health A Tox Hazard Subst Environ Eng. 46(12):1417-1423, 2011.

In this study the effects of 1800 MHz GSM-like radiofrequency electromagnetic waves (RF-EMW) exposure on bovine semen was monitored. The experimental samples were analyzed in vitro in four time periods (0, 30, 120 and 420 min) and compared with unexposed samples (control). Spermatozoa motility was determined by computer assisted semen analyzer (CASA). Evaluation of the percentage of motile spermatozoa showed significant ($P < 0.001$) decrease in experimental groups after 120 and 420 min of culture when exposed to microwaves, in comparison to control. Similar spermatozoa motility inhibition was detected for the percentage of progressively motile spermatozoa, too. Average path distance decreased significantly ($p < 0.001$) in experimental groups after 30 and 420 min of culture. Path velocity increased in the experimental groups exposed to RF-EMW after 30 minutes of culture, but subsequently decreased after 420 min of culture, in comparison to control. This indicates a possible initial stimulation and subsequent velocity inhibition of bovine spermatozoa under RF-EMW exposure. Changes in spermatozoa motility were also detected for some fine parameters, too. A significant decrease ($P < 0.001$) was noted for amplitude of lateral

Studies that show **Cell Tower** Health Effects

head displacement in the experimental group after 420 minutes of culture. Detailed in vitro motility analysis of bovine spermatozoa exposed to microwave radiation suggested that the parameters of path and velocity at the beginning of the culture significantly increase, but after longer culture (420 minutes) a significant decrease occur in the experimental group as compared to control. In general, results of this experiment indicate a negative time-dependent effect of 1800 MHz RF-EMW radiation on bovine spermatozoa motility.

Effects on Wellbeing

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. Study of the health of people living in the vicinity of mobile phone base stations: I. Influence of distance and sex. Pathol Biol (Paris) 50(6):369-373, 2002. [Article in French]

A survey study using questionnaire was conducted in 530 people (270 men, 260 women) living or not in vicinity of cellular phone base stations, on 18 Non Specific Health Symptoms. Comparisons of complaints frequencies (CHI-SQUARE test with Yates correction) in relation with distance from base station and sex, show significant ($p < 0.05$) increase as compared to people living > 300 m or not exposed to base station, till 300 m for tiredness, 200 m for headache, sleep disturbance, discomfort, etc. 100 m for irritability, depression, loss of memory, dizziness, libido decrease, etc. Women significantly more often than men ($p < 0.05$) complained of headache, nausea, loss of appetite, sleep disturbance, depression, discomfort and visual perturbations. This first study on symptoms experienced by people living in vicinity of base stations shows that, in view of radioprotection, minimal distance of people from cellular phone base stations should not be < 300 m.

Santini R, Santini P, Le Ruz P, Danze JM, Seigne M, Survey study of people living in the vicinity of cellular phone base stations. Electromag Biol Med 22:41-49, 2003.

A survey study was conducted, using a questionnaire, on 530 people (270 men, 260 women) living or not in proximity to cellular phone base stations. Eighteen different symptoms (Non Specific Health Symptoms-NSHS), described as radiofrequency sickness, were studied by means of the chi-square test with Yates correction. The results that were obtained underline that certain complaints are experienced only in the immediate vicinity of base stations (up to 10 m for nausea, loss of appetite, visual disturbances), and other at greater distances from base stations (up to 100 m for irritability, depressive tendencies, lowering of libido, and up to 200 m for headaches, sleep disturbance, feeling of discomfort). In the 200 m to 300 m zone, only the complaint of fatigue is experienced significantly more often when compared with subjects residing at more than 300 m or not exposed (reference group). For seven of the studied symptoms and for the distance up to 300 m, the frequency of reported complaints is significantly higher ($P < 0.05$) for women in comparison to men. Significant differences are also observed in relation to the ages of subjects, and for the location of subjects in relation to the antennas and to other electromagnetic factors.

Studies that show **Cell Tower** Health Effects

Navarro EA, Sequera J, Portoles M, Gomez-Perretta de Mateo C. The Microwave Syndrome: A Preliminary Study in Spain. *Electromag Biol Med* 22:161-169, 2003.

A health survey was carried out in Murcia, Spain, in the vicinity of a Cellular Phone Base Station working in DCS-1800 MHz. This survey contained health items related to "microwave sickness" or "RF syndrome." The microwave power density was measured at the respondents' homes. Statistical analysis showed significant correlation between the declared severity of the symptoms and the measured power density. The separation of respondents into two different exposure groups also showed an increase of the declared severity in the group with the higher exposure.

Lerchl A, Krüger H, Niehaus M, Streckert JR, Bitz AK, Hansen V. Effects of mobile phone electromagnetic fields at nonthermal SAR values on melatonin and body weight of Djungarian hamsters (*Phodopus sungorus*). *J Pineal Res.* 44(3):267-272, 2008.

In three experiments, adult male Djungarian hamsters (*Phodopus sungorus*) were exposed 24 hr/day for 60 days to radio frequency electromagnetic fields (RF-EMF) at 383, 900, and 1800 MHz, modulated according to the TETRA (383 MHz) and GSM standards (900 and 1800 MHz), respectively. A radial waveguide system ensured a well defined and uniform exposure at whole-body averaged specific absorption rates of 80 mW/kg, which is equal to the upper limit of whole-body exposure of the general population in Germany and other countries. For each experiment, using two identical waveguides, hamsters were exposed (n = 120) and sham-exposed (n = 120) in a blind fashion. In all experiments, pineal and serum melatonin levels as well as the weights of testes, brain, kidneys, and liver were not affected. At 383 MHz, exposure resulted in a significant transient increase in body weight up to 4%, while at 900 MHz this body weight increase was more pronounced (up to 6%) and not transient. At 1800 MHz, no effect on body weight was seen. The results corroborate earlier findings which have shown no effects of RF-EMF on melatonin levels in vivo and in vitro. The data are in accordance with the hypothesis that absorbed RF energy may result in metabolic changes which eventually cause body weight increases in exposed animals. The data support the notion that metabolic effects of RF-EMFs need to be investigated in more detail in future studies.

Kato Y, Johansson O. Reported functional impairments of electrosensitive Japanese: A questionnaire survey. *Pathophysiology*.19(2) 95-100, 2012.

An increasing number of people worldwide complain that they have become electromagnetic hypersensitive (EHS). We conducted a questionnaire survey of EHS persons in Japan. The aim was to identify electromagnetic fields (EMF) and plausible EMF sources that caused their symptoms. Postal questionnaires were distributed via a self-help group, and 75 participants (95% women) responded. Reported major complaints were "fatigue/tiredness" (85%), "headache", "concentration, memory, and thinking" difficulty (81%, respectively). Seventy-two per cent used some form of complementary/alternative therapy. The most plausible trigger of EHS onset was a mobile phone base station or personal handy-phone system (37%). Sixty-five percent experienced health problems to be due to the radiation from other passengers' mobile

Studies that show **Cell Tower** Health Effects

phones in trains or buses, and 12% reported that they could not use public transportation at all. Fifty-three percent had a job before the onset, but most had lost their work and/or experienced a decrease in income. Moreover, 85.3% had to take measures to protect themselves from EMF, such as moving to low EMF areas, or buying low EMF electric appliances. EHS persons were suffering not only from their symptoms, but also from economical and social problems.

Hutter HP, Moshammer H, Wallner P, Kundi M. Subjective symptoms, sleeping problems, and cognitive performance in subjects living near mobile phone base stations. *Occup Environ Med.* 63(5):307-313, 2006.

BACKGROUND: The erection of mobile telephone base stations in inhabited areas has raised concerns about possible health effects caused by emitted microwaves.

METHODS: In a cross-sectional study of randomly selected inhabitants living in urban and rural areas for more than one year near to 10 selected base stations, 365 subjects were investigated. Several cognitive tests were performed, and wellbeing and sleep quality were assessed. Field strength of high-frequency electromagnetic fields (HF-EMF) was measured in the bedrooms of 336 households. RESULTS: Total HF-EMF and exposure related to mobile telecommunication were far below recommended levels (max. 4.1 mW/m²). Distance from antennae was 24-600 m in the rural area and 20-250 m in the urban area. Average power density was slightly higher in the rural area (0.05 mW/m²) than in the urban area (0.02 mW/m²). Despite the influence of confounding variables, including fear of adverse effects from exposure to HF-EMF from the base station, there was a significant relation of some symptoms to measured power density; this was highest for headaches. Perceptual speed increased, while accuracy decreased insignificantly with increasing exposure levels. There was no significant effect on sleep quality. CONCLUSION: Despite very low exposure to HF-EMF, effects on wellbeing and performance cannot be ruled out, as shown by recently obtained experimental results; however, mechanisms of action at these low levels are unknown.

Bortkiewicz A, Zmyslony M, Szyjowska A, Gadzicka E. [Subjective symptoms reported by people living in the vicinity of cellular phone base stations: a review of the studies] *Med Pr.* 55(4):345-351, 2004. [Article in Polish]

The problem of health effects of electromagnetic fields (EMF) emitted by cellular phone base stations evokes much interest in view of the fact that people living in their vicinity are fated to continuous exposure to EMF. None of the studies carried out throughout the world have revealed excessive values of standards adopted by the International Commission on Non-Ionizing Radiation Protection (ICNIRP). A questionnaire was used as a study tool. The results of the questionnaire survey reveal that people living in the vicinity of base stations report various complaints mostly of the circulatory system, but also of sleep disturbances, irritability, depression, blurred vision, concentration difficulties, nausea, lack of appetite, headache and vertigo. The performed studies showed the relationship between the incidence of individual symptoms, the level of exposure, and the distance between a residential area and a base station. This association was observed in both groups of persons, those who linked their complaints

Studies that show **Cell Tower** Health Effects

with the presence of the base station and those who did not notice such a relation. Further studies, clinical and those based on questionnaires, are needed to explain the background of reported complaints.

Bortkiewicz A, Gadzicka E, Szykowska A, Politański P, Mamrot P, Szymczak W, Zmysłony M. Subjective complaints of people living near mobile phone base stations in Poland. *Int J Occup Med Environ Health*. 25(1):31-40, 2012.

OBJECTIVES: The aim of our study was to assess the health conditions and subjective symptoms of the inhabitants living in the base stations vicinity and to analyse the relationship between the complaints and level of exposure to electromagnetic fields (EMF). MATERIALS AND METHODS: Our study was performed in housing estates located in five regions of Łódź. The electric field measurements were performed in the buildings located closest to the azimuth of the antennas. Respondents were selected by trained interviewers using an uniform procedure. The number of the households to be examined was set at a minimum of 420. The questionnaire contained: demographic data, occupational and environmental exposure to EMF, health condition, subjective complaints. Results were adjusted for confounders (age, gender, EMF at the workplace and EMF emitted by household equipment) using multiple regression model. RESULTS: 181 men and 319 women from 500 households were examined. Electric field above 0.8 V/m was recorded in 12% of flats. There was no significant correlation between electric field strength and the distance of examined flats from the base stations. To make possible comparison with relevant literature, we analysed also the frequency of the reported symptoms vs. the distance. Headache was declared by 57% people, most frequently (36.4%) living 100-150 m away from the base station compared to people living at longer distances ($p = 0.013$). 24.4% subjects, mostly living at a distance above 150 m, declared impaired memory. Difference was statistically significant in comparison with people living at other distances ($p = 0.004$). CONCLUSIONS: The explanation why we did not find any correlation between the electric field strength and frequency of subjective symptoms but found a correlation between subjective symptoms and distance from base station needs further studies. Maybe new metrics of exposure assessment should be adopted for this purpose.

Augner C, Florian M, Pauser G, Oberfeld G, Hacker GW. GSM base stations: Short-term effects on well-being. *Bioelectromagnetics*. 30:73-80, 2009.

The purpose of this study was to examine the effects of short-term GSM (Global System for Mobile Communications) cellular phone base station RF-EMF (radiofrequency electromagnetic fields) exposure on psychological symptoms (good mood, alertness, calmness) as measured by a standardized well-being questionnaire. Fifty-seven participants were selected and randomly assigned to one of three different exposure scenarios. Each of those scenarios subjected participants to five 50-min exposure sessions, with only the first four relevant for the study of psychological symptoms. Three exposure levels were created by shielding devices in a field laboratory, which could be installed or removed during the breaks between sessions such that double-blinded conditions prevailed. The overall median power flux densities were 5.2 microW/m² during "low," 153.6 microW/m² during "medium," and 2126.8 microW/m² during

Studies that show **Cell Tower** Health Effects

"high" exposure sessions. For scenario HM and MH, the first and third sessions were "low" exposure. The second session was "high" and the fourth was "medium" in scenario HM; and vice versa for scenario MH. Scenario LL had four successive "low" exposure sessions constituting the reference condition. Participants in scenarios HM and MH (high and medium exposure) were significantly calmer during those sessions than participants in scenario LL (low exposure throughout) ($P = 0.042$). However, no significant differences between exposure scenarios in the "good mood" or "alertness" factors were obtained. We conclude that short-term exposure to GSM base station signals may have an impact on well-being by reducing psychological arousal.

Augner C, Hacker GW. Are people living next to mobile phone base stations more strained? Relationship of health concerns, self-estimated distance to base station, and psychological parameters. *Indian J Occup Environ Med.* 13(3):141-145, 2009.

BACKGROUND AND AIMS: Coeval with the expansion of mobile phone technology and the associated obvious presence of mobile phone base stations, some people living close to these masts reported symptoms they attributed to electromagnetic fields (EMF). Public and scientific discussions arose with regard to whether these symptoms were due to EMF or were nocebo effects. The aim of this study was to find out if people who believe that they live close to base stations show psychological or psychobiological differences that would indicate more strain or stress. Furthermore, we wanted to detect the relevant connections linking self-estimated distance between home and the next mobile phone base station (DBS), daily use of mobile phone (MPU), EMF-health concerns, electromagnetic hypersensitivity, and psychological strain parameters. DESIGN, MATERIALS AND METHODS: Fifty-seven participants completed standardized and non-standardized questionnaires that focused on the relevant parameters. In addition, saliva samples were used as an indication to determine the psychobiological strain by concentration of alpha-amylase, cortisol, immunoglobulin A (IgA), and substance P. RESULTS: Self-declared base station neighbors (DBS \leq 100 meters) had significantly higher concentrations of alpha-amylase in their saliva, higher rates in symptom checklist subscales (SCL) somatization, obsessive-compulsive, anxiety, phobic anxiety, and global strain index PST (Positive Symptom Total). There were no differences in EMF-related health concern scales. CONCLUSIONS: We conclude that self-declared base station neighbors are more strained than others. EMF-related health concerns cannot explain these findings. Further research should identify if actual EMF exposure or other factors are responsible for these results.

Blettner M, Schlehofer B, Breckenkamp J, Kowall B, Schmiedel S, Reis U, Potthoff P, Schuez J, Berg-Beckhoff G. Mobile phone base stations and adverse health effects: Phase 1: A population-based cross-sectional study in Germany. *Occup Environ Med.* 66(2):118-123. 2009.

Abstract OBJECTIVE: The aim of this first phase of a cross-sectional study from Germany was to investigate whether proximity of residence to mobile phone base stations as well as risk perception is associated with health complaints. METHODS: We conducted a population-based multi-phase cross-sectional study within the context of a large panel

Studies that show **Cell Tower** Health Effects

survey regularly carried out by a private research institute in Germany. In the initial phase, which we will report on in this paper, 30,047 persons from a total of 51,444 who took part in the nationwide survey also answered questions on how mobile phone base stations affect their health. A list of 38 health complaints was used. A multiple linear regression model was used to identify predictors of health complaints including proximity of residence to mobile phone base stations and risk perception. RESULTS: Of the 30,047 participants (response rate 58.6%), 18.7% of participants were concerned about adverse health effects of mobile phone base stations, while an additional 10.3% attributed their personal adverse health effects to the exposure from them. Participants who are concerned about or attribute adverse health effects to mobile phone base stations and those living in the vicinity of a mobile phone base station (500 m) reported slightly more health complaints than others. CONCLUSIONS: A substantial proportion of the German population is concerned about adverse health effects caused by exposure from mobile phone base stations. The observed slightly higher prevalence of health complaints near base stations can however not be fully explained by attributions or concerns.

Effects on Sleep

Mohammed HS, Fahmy HM, Radwah NM, Elsayed AA. Non-thermal continuous and modulated electromagnetic radiation fields effects on sleep EEG of rats. J Adv Res 4(2) 181-187, 2013.

In the present study, the alteration in the sleep EEG in rats due to chronic exposure to low-level non-thermal electromagnetic radiation was investigated. Two types of radiation fields were used; 900 MHz unmodulated wave and 900 MHz modulated at 8 and 16 Hz waves. Animals has exposed to radiation fields for 1 month (1 h/day). EEG power spectral analyses of exposed and control animals during slow wave sleep (SWS) and rapid eye movement sleep (REM sleep) revealed that the REM sleep is more susceptible to modulated radiofrequency radiation fields (RFR) than the SWS. The latency of REM sleep increased due to radiation exposure indicating a change in the ultradian rhythm of normal sleep cycles. The cumulative and irreversible effect of radiation exposure was proposed and the interaction of the extremely low frequency radiation with the similar EEG frequencies was suggested.

Liu H, Chen G, Pan Y, Chen Z, Jin W, Sun C, Chen C, Dong X, Chen K, Xu Z, Zhang S, Yu Y. (2014) Occupational Electromagnetic Field Exposures Associated with Sleep Quality: A Cross-Sectional Study. PLoS ONE 9(10): e110825. doi:10.1371/journal.pone.0110825.

BACKGROUND: Exposure to electromagnetic field (EMF) emitted by mobile phone and other machineries concerns half the world's population and raises the problem of their impact on human health. The present study aims to explore the effects of electromagnetic field exposures on sleep quality and sleep duration among workers from electric power plant. METHODS: A cross-sectional study was conducted in an

Studies that show **Cell Tower** Health Effects

electric power plant of Zhejiang Province, China. A total of 854 participants were included in the final analysis. The detailed information of participants was obtained by trained investigators using a structured questionnaire, which including socio-demographic characteristics, lifestyle variables, sleep variables and electromagnetic exposures. Physical examination and venous blood collection were also carried out for every study subject. **RESULTS:** After grouping daily occupational electromagnetic exposure into three categories, subjects with long daily exposure time had a significantly higher risk of poor sleep quality in comparison to those with short daily exposure time. The adjusted odds ratios were 1.68 (95%CI: 1.18, 2.39) and 1.57 (95%CI: 1.10, 2.24) across tertiles. Additionally, among the subjects with long-term occupational exposure, the longer daily occupational time apparently increased the risk of poor sleep quality (OR (95%CI): 2.12 (1.23~3.66) in the second tertile; 1.83 (1.07~3.15) in the third tertile). There was no significant association of long-term occupational exposure duration, monthly electric fee or years of mobile-phone use with sleep quality or sleep duration. **CONCLUSIONS:** The findings showed that daily occupational EMF exposure was positively associated with poor sleep quality. It implies EMF exposure may damage human sleep quality rather than sleep duration.

Hung CS, Anderson C, Horne JA, McEvoy P. Mobile phone 'talk-mode' signal delays EEG-determined sleep onset. *Neurosci Lett.* 421: 82-86, 2007.

Mobile phones signals are pulse-modulated microwaves, and EEG studies suggest that the extremely low-frequency (ELF) pulse modulation has sleep effects. However, 'talk', 'listen' and 'standby' modes differ in the ELF (2, 8, and 217Hz) spectral components and specific absorption rates, but no sleep study has differentiated these modes. We used a GSM900 mobile phone controlled by a base-station simulator and a test SIM card to simulate these three specific modes, transmitted at 12.5% (23dBm) of maximum power. At weekly intervals, 10 healthy young adults, sleep restricted to 6h, were randomly and single-blind exposed to one of: talk, listen, standby and sham (nil signal) modes, for 30min, at 13:30h, whilst lying in a sound-proof, lit bedroom, with a thermally insulated silent phone beside the right ear. Bipolar EEGs were recorded continuously, and subjective ratings of sleepiness obtained every 3min (before, during and after exposure). After exposure the phone and base-station were switched off, the bedroom darkened, and a 90min sleep opportunity followed. We report on sleep onset using: (i) visually scored latency to onset of stage 2 sleep, (ii) EEG power spectral analysis. There was no condition effect for subjective sleepiness. Post-exposure, sleep latency after talk mode was markedly and significantly delayed beyond listen and sham modes. This condition effect over time was also quite evident in 1-4Hz EEG frontal power, which is a frequency range particularly sensitive to sleep onset. It is possible that 2, 8, 217Hz modulation may differentially affect sleep onset.

Studies that show **Cell Tower** Health Effects

Effects on Cells

Neshev NN, Kirilova EI, Environmental-health aspects of pulse-modulated microwaves. Rev Environ Health 11(1-2):85-88, 1996.

Our theoretical model describes the potential influence of irradiation with pulse-modulated microwaves on the conformational oscillations of enzymes in living organisms. Certain values of pulse-repetition time, determined by the period of conformational oscillations of the corresponding type of enzyme, can produce the effect at extremely low power levels. Synchronized oscillations in identical enzyme molecules produce in turn large-scale oscillations within living cells. Thus, short periods of exposure to pulse-modulated microwaves could be beneficial to cellular function, whereas maintaining the amplitude of such oscillations at a maximum for long periods may have a stressful effect on biochemical processes. The model discloses the possible environmental-health risks of long-term exposure in ambient fields that are created by radar, navigation, and communication systems.

Kwee S, Raskmark P, Changes in cell proliferation due to environmental non-ionizing radiation 2. Microwave radiation. Bioelectrochem Bioenerg 44(2) 251-255, 1998.

Due to the use of mobile telephones, there is an increased exposure of the environment to weak radiofrequency (RF) electromagnetic fields, emitted by these devices. This study was undertaken to investigate if the microwave radiation from these fields will have a similar effect on cell proliferation as weak electromagnetic (ELF) fields. The field was generated by signal simulation of the Global System for Mobile communications (GSM) of 960 MHz. Cell cultures, growing in microtiter plates, were exposed in a specially constructed chamber, a Transverse Electromagnetic (TEM) cell. The Specific Absorption Rate (SAR) values for each cell well were calculated for this exposure system. Experiments were performed on cell cultures of transformed human epithelial amnion cells (AMA), which were exposed to 960 MHz microwave fields at three different power levels and three different exposure times, respectively. It was found that cell growth in the exposed cells was decreased in comparison to that in the control and sham exposed cells. Cell proliferation during the period following exposure varied not only with the various SAR levels, but also with the length of exposure time. On the other hand, repeated periods of exposure did not seem to change the effects. There was a general linear correlation between power level and growth change. However, the exposure time required to obtain the maximum effect was not the same for the various power levels. It turned out that at low power level, a maximum effect was first reached after a longer exposure time than at higher power level. A similar phenomenon was registered in the studies on ELF electromagnetic fields. Here, it was found that there was a linear correlation between the length of exposure time to obtain maximum effect and field strength.

Studies that show **Cell Tower** Health Effects

Effects on Eyes

Lu L, Xu H, Wang X, Guo G. Increased nitric oxide synthase activity is essential for electromagnetic-pulse-induced blood-retinal barrier breakdown in vivo. *Brain Res.* 1264:104-10, 2009.

PURPOSE: To examine whether electromagnetic pulses (EMPs) affected the permeability of the **blood-retinal barrier (BRB)**, gene expression of occludin and activity of nitric oxide synthase (NOS). **METHODS:** Sprague-Dawley (SD) rats were used and randomized into EMP and control groups. Retinas were removed immediately, and 2 h or 24 h after EMP radiation. BRB permeability was analyzed by transmission electron microscopy and Evans Blue staining. Retinal NOS activity and concentrations of nitrite and nitrate were measured. Occludin mRNA and protein levels were detected by RT-PCR and Western blotting. **RESULTS:** Exposure of SD rats to EMP resulted in increased BRB permeability, with the greatest decrease in occludin at 24 h. Moreover, this permeability defect was also correlated with significant increases in the formation of NO and induction of NOS activity in SD rats. Furthermore, we found that treatment with NOS inhibitor N-nitro-L-arginine methyl ester (L-NAME) blocked BRB breakdown and prevented the increase in NO formation and induction of NOS activity, as well as the decrease in occluding expression. **CONCLUSION:** Taken together, these results support the view that NOS-dependent NO production is an important factor that contributes to EMP-induced BRB dysfunction, and suggests that NOS induction may play an important role in BRB breakdown.

Hässig M, Jud F, Naegeli H, Kupper J, Spiess B. Prevalence of nuclear cataract in Swiss veal calves and its possible association with mobile telephone antenna base stations. *Schweiz Arch Tierheilkd.* 151(10):471-478, 2009.

The purpose of this study was to evaluate the prevalence of nuclear cataract in veal calves and to elucidate a possible impact by mobile phone base stations (MPBS). For this experiment a cohort study was conducted. A follow-up of the geographical location of each dam and its calf from conception through the fetal period up to slaughter was performed. The first trimester of gestation (organogenesis) was particularly emphasized. The activities of selected protective antioxidants (superoxide dismutase, catalase, glutathione peroxidase [GPx]) were assessed in aqueous humor of the eye to evaluate the redox status. Of 253 calves, 79 (32 %) had various degrees of nuclear cataract, but only 9 (3.6 %) calves had severe nuclear cataract. Results demonstrate a relation between the location of veal calves with nuclear cataracts in the first trimester of gestation and the strength of antennas. The number of antennas within 100 to 199 meters was associated with oxidative stress and there was an association between oxidative stress and the distance to the nearest MPBS. Oxidative stress was increased in eyes with cataract (OR per kilometer: 0.80, confidence interval 95 % 0.62, 0.93). It has not been shown that the antennas actually affected stress. Hosmer-Lemeshow statistics showed an accuracy of 100 % in negative cases with low radiation, and only 11.11 % accuracy in positive cases with high radiation. This reflects, that there are a lot of other

Studies that show **Cell Tower** Health Effects

possibilities for nuclear cataract beside MPBS. Further studies on the influence of electromagnetic fields during embryonic development animal or person at risk are indicated.

Hässig M, Jud F, Spiess B. [Increased occurrence of nuclear cataract in the calf after erection of a mobile phone base station]. *Schweiz Arch Tierheilkd.* 154(2):82-86, 2012.[Article in German]

We examined and monitored a dairy farm in which a large number of calves were born with nuclear cataracts after a mobile phone base station had been erected in the vicinity of the barn. Calves showed a 3.5 times higher risk for heavy cataract if born there compared to Swiss average. All usual causes such as infection or poisoning, common in Switzerland, could be excluded. The real cause of the increased incidence of cataracts remains unknown.

Jelodar G, Akbari A, Nazifi S. The prophylactic Effect of Vitamin C on Oxidative Stress Indexes in Rat Eyes Following Exposure to Radiofrequency Wave Generated by a BTS Antenna Model. *Int J Radiat Biol.* 89(2):128-131, 2013.

Purpose: This study was conducted to evaluate the effect of radiofrequency wave (RFW)-induced oxidative stress in the eye and the prophylactic effect of vitamin C on this organ by measuring the antioxidant enzymes activity including: glutathione peroxidase (GPx), superoxide dismutase (SOD) and catalase (CAT), and malondialdehyde (MDA). Materials and methods: Thirty-two adult male Sprague-Dawley rats were randomly divided into four experimental groups and treated daily for 45 days as follows: control, vitamin C (L-ascorbic acid 200 mg/kg of body weight/day by gavage), test (exposed to 900 MHz RFW) and the treated group (received vitamin C in addition to exposure to RFW). At the end of the experiment all animals were killed, their eyes were removed and were used for measurement of antioxidant enzymes and MDA activity. Results: The results indicate that exposure to RFW in the test group decreased antioxidant enzymes activity and increased MDA compared with the control groups (P<0.05). In the treated group vitamin C improved antioxidant enzymes activity and reduced MDA compared to the test group (P<0.05). Conclusions: It can be concluded that RFW causes oxidative stress in the eyes and vitamin C improves the antioxidant enzymes activity and decreases MDA.

Oxidative Stress

Achudume A, Onibere B, Aina F, Tchokossa P. Induction of oxidative stress in male rats subchronically exposed to electromagnetic fields at non-thermal intensities. *J Electromagnetic Analysis and Applications* 2(8), 482-487, 2010.

To investigate the oxidative stress-inducing potential of non-thermal electromagnetic fields in rats. Male Wister rats were exposed to electrical field intensity of $2.3 \pm 0.82 \mu\text{V/m}$. Exposure was in three forms: continuous waves, or modulated at 900 MHz or modulated GSM-nonDTX. The radio frequency radiation (RFR) was 1800 MHz, specific

Studies that show **Cell Tower** Health Effects

absorption radiation (SAR) (0.95-3.9 W/kg) for 40 and/or 60 days continuously. Control animals were located > 300 m from base station, while sham control animals were located in a similar environmental conditions, but in the vicinity of a non-functional base station. The rats were assessed for thiobarbituric and reactive species (TBARS), reduced glutathione (GSH) content, catalase activity, glutathione reductase (GR) and glucose residue after 40 and 60 days of exposure. At 40 days, electromagnetic radiation failed to induce any significant alterations. However, at 60 days of exposure various attributes evaluated decreased. The respective decreases in both nicotinamide adenine dinucleotide phosphate (NADPH) and Ascorbate- linked lipid peroxidation (LPO) with concomitant diminution in enzymatic antioxidative defense systems resulted in decreased glucose residue. The present studies showed some biochemical changes that may be associated with a prolong exposure to electromagnetic fields and its relationship to the activity of antioxidant system in rat Regular assessment and early detection of antioxidative defense system among people working around the base stations are recommended.

Augner C, Hacker GW, Oberfeld G, Florian M, Hitzl W, Hutter J, Pauser G. Effects of Exposure to GSM mobile phone base station signals on salivary cortisol, alpha-amylase, and Immunoglobulin A. *Biomed Environ Sci.* 23(3):199-207, 2010.

OBJECTIVE: The present study aimed to test whether exposure to radiofrequency electromagnetic fields (RF-EMF) emitted by mobile phone base stations may have effects on salivary alpha-amylase, immunoglobulin A (IgA), and cortisol levels. METHODS: Fifty seven participants were randomly allocated to one of three different experimental scenarios (22 participants to scenario 1, 26 to scenario 2, and 9 to scenario 3). Each participant went through five 50-minute exposure sessions. The main RF-EMF source was a GSM-900-MHz antenna located at the outer wall of the building. In scenarios 1 and 2, the first, third, and fifth sessions were "low" (median power flux density 5.2 $\mu\text{W}/\text{m}^2$) exposure. The second session was "high" (2126.8 $\mu\text{W}/\text{m}^2$), and the fourth session was "medium" (153.6 $\mu\text{W}/\text{m}^2$) in scenario 1, and vice versa in scenario 2. Scenario 3 had four "low" exposure conditions, followed by a "high" exposure condition. Biomedical parameters were collected by saliva samples three times a session. Exposure levels were created by shielding curtains. RESULTS: In scenario 3 from session 4 to session 5 (from "low" to "high" exposure), an increase of cortisol was detected, while in scenarios 1 and 2, a higher concentration of alpha-amylase related to the baseline was identified as compared to that in scenario 3. IgA concentration was not significantly related to the exposure. CONCLUSIONS: RF-EMF in considerably lower field densities than ICNIRP-guidelines may influence certain psychobiological stress markers.

Marzook EA, Abd El Moneim AE, Elhadary AA. Prootective role of seame oil against mobile phone base station-induced oxidative stress. *J Rad Res Appl Sci* 7(1):1-6, 2014.

The present study was undertaken to shed the light on the environmental threats associated with the wireless revolution and the health hazards associated with exposure to mobile base station (MBS). Besides, studying the possible protective role of sesame oil (SO) as an antioxidant against oxidative stress. Therefore, the present work was

Studies that show **Cell Tower** Health Effects

designed to study the effect of chronic exposure to electromagnetic radiations (EMR), produced by a cellular tower for mobile phone and the possible protective role of sesame oil on glutathione reductase (GSH-Rx), superoxide dismutase (SOD), catalase (CAT), total testosterone and lipid profile (total cholesterol (Tch), triglycerides (TG), low density lipoprotein cholesterol (LDL-c) and high density lipoprotein cholesterol (HDL-c) in male albino rats. Rats were arranged into four groups: the control unexposed, the exposed untreated and the exposed treated groups (1.5 and 3 ml oil). Exposed groups were subjected to electromagnetic field at frequency of 900 MHz, for 24 h/day for 8 weeks, at the same time both treated groups were supplied with oral injection of sesame oil three times per week. At the end of the experiment, blood samples were obtained for determination of the above mentioned variables in serum. The results obtained revealed that TG and testosterone were raised significantly over control in all groups and the significant increase in oil groups occurred in dose dependent manner. SOD and CAT activities were reduced significantly in exposed rats than control and increased significantly in sesame oil groups as the dose of oil increased. Total cholesterol only showed remarkable reduction in the group treated with 3 ml sesame oil. Also, in this latter group, significant elevation of GSH-Rx was recorded. Changes in serum HDL-c and LDL-c followed an opposite trend in exposed and sesame oil groups reflecting their affection by EMR or sesame oil. In conclusion, all results of the current study proved that sesame oil can be used as an edible oil to attenuate the oxidative stress which could be yielded as a result of chronic exposure to EMR.

Effects on Blood

Kismali G, Ozgur E, Guler G, Akcay A, Sel T, Seyhan N. The influence of 1800 MHz GSM-like signals on blood chemistry and oxidative stress in non-pregnant and pregnant rabbits. Int J Radiat Biol. 88(5):414-419, 2012.

PURPOSE: Environmental electromagnetic fields originate from man-made sources, such as mobile phones and base stations, and have led to increasing public concern about their possible adverse health effects. We aimed to investigate the possible effects of radiofrequency radiation (RFR) generated from these devices on oversensitive animals, such as pregnant rabbits. **MATERIALS AND METHODS:** In the present study, the effects of whole body 1800 MHz Global System for Mobile Communications (GSM)-like RFR exposure for 15 min/day for seven days on blood chemistry and lipid peroxidation levels in both non-pregnant and pregnant New Zealand White rabbits were investigated. Thirteen-month-old rabbits were studied in the following four groups: Non-pregnant control, non-pregnant RFR-exposed, pregnant control and pregnant RFR-exposed. **RESULTS:** Lipid peroxidation, namely malondialdehyde (MDA) levels, did not change after RFR exposure. However, blood chemistry parameters, such as cholesterol (CHO), total protein (TP), albumin (ALB), uric acid, creatinin and **creatine kinase (CK)** and **creatine kinase-myocardial band isoenzyme (CK-MB)** changed due to both pregnancy and RFR exposure. **CONCLUSION:** Our investigations have been shown that no indication for oxidative stress was detected in the blood of pregnant rabbits upon RF exposure at specific conditions employed in the present study. Minor

Studies that show **Cell Tower** Health Effects

changes in some blood chemistry parameters were detected but CK-MB and CK increases were found remarkable. Studies on RFR exposure during pregnancy will help establish international standards for the protection of pregnant women from environmental RFR.

Yurekli AI, Ozkan M, Kalkan T, Saybasili H, Tuncel H, Atukeren P, Gumustas K, Seker S. GSM Base Station Electromagnetic Radiation and Oxidative Stress in Rats. *Electromagn Biol Med.* 2006;25(3):177-188, 2006.

The ever increasing use of cellular phones and the increasing number of associated base stations are becoming a widespread source of nonionizing electromagnetic radiation. Some biological effects are likely to occur even at low-level EM fields. In this study, a gigahertz transverse electromagnetic (GTEM) cell was used as an exposure environment for plane wave conditions of far-field free space EM field propagation at the GSM base transceiver station (BTS) frequency of 945 MHz, and effects on oxidative stress in rats were investigated. When EM fields at a power density of 3.67 W/m² (specific absorption rate = 11.3 mW/kg), which is well below current exposure limits, were applied, MDA (malondialdehyde) level was found to increase and GSH (reduced glutathione) concentration was found to decrease significantly ($p < 0.0001$). Additionally, there was a less significant ($p = 0.0190$) increase in SOD (superoxide dismutase) activity under EM exposure.

Jin YB, Lee HJ, Seon Lee J, Pack JK, Kim N, Lee YS. One-year, simultaneous combined exposure of CDMA and WCDMA radiofrequency electromagnetic fields to rats. *Int J Radiat Biol.* 87(4):416-423, 2011.

PURPOSE: We investigated whether one-year, long-term, simultaneous exposure to code division multiple access (CDMA; 849 MHz) and wideband code division multiple access (WCDMA; 1.95 GHz) radiofrequencies (RF) would induce chronic illness in Sprague-Dawley (SD) rats. **MATERIALS AND METHODS:** Two groups of 40 SD rats (50% males and females in sham and exposed groups) were exposed to CDMA and WCDMA RF simultaneously at 2.0 W/kg for 45 min/day (total 4.0 W/kg), 5 days per week for a total of one year. Body and organ weight measurements, urinalysis, haematological and blood biochemical analysis, and histopathological evaluations were performed.

RESULTS: The mortality patterns in male and female rats exposed to RF were compared with those found in gender-matched sham control animals. No significant alteration in body weight was observed with the simultaneous combined RF exposure. Most RF-exposed rats showed no significant alteration, based on urinalysis, haematology, blood biochemistry, or histopathology. However, some altered parameters of the complete blood count and serum chemistry were seen in RF-exposed rats. The total tumour incidence was not different between sham-exposed and RF-exposed animals.

CONCLUSIONS: Our results suggest that one-year chronic exposure to CDMA (849 MHz) and WCDMA (1.95 GHz) RF simultaneously at 2.0 W/kg for 45-min RF exposure periods (total, 4 W/kg) did not increase chronic illness in rats, although there were some altered parameters in the complete blood count and serum chemistry.

Studies that show **Cell Tower** Health Effects

Death

Dode AC, Leão MM, Tejo Fde A, Gomes AC, Dode DC, Dode MC, Moreira CW, Condessa VA, Albinatti C, Caiaffa WT. Mortality by neoplasia and cellular telephone base stations in the Belo Horizonte municipality, Minas Gerais state, Brazil. Sci Total Environ. 409(19):3649-3665, 2011.

Pollution caused by the electromagnetic fields (EMFs) of radio frequencies (RF) generated by the telecommunication system is one of the greatest environmental problems of the twentieth century. The purpose of this research was to verify the existence of a spatial correlation between **base station (BS)** clusters and cases of deaths by neoplasia in the Belo Horizonte municipality, Minas Gerais state, Brazil, from 1996 to 2006 and to measure the human exposure levels to EMF where there is a major concentration of cellular telephone transmitter antennas. A descriptive spatial analysis of the BSs and the cases of death by neoplasia identified in the municipality was performed through an ecological-epidemiological approach, using georeferencing. The database employed in the survey was composed of three data banks: 1. death by neoplasia documented by the Health Municipal Department; 2. BSs documented in ANATEL ("Agência Nacional de Telecomunicações": 'Telecommunications National Agency'); and 3. census and demographic city population data obtained from official archives provided by IBGE ("Instituto Brasileiro de Geografia e Estatística": 'Brazilian Institute of Geography and Statistics'). The results show that approximately 856 BSs were installed through December 2006. Most (39.60%) of the BSs were located in the "Centro-Sul" ('Central-Southern') region of the municipality. Between 1996 and 2006, 7191 deaths by neoplasia occurred and within an area of 500 m from the BS, the mortality rate was 34.76 per 10,000 inhabitants. Outside of this area, a decrease in the number of deaths by neoplasia occurred. The greatest accumulated incidence was 5.83 per 1000 in the Central-Southern region and the lowest incidence was 2.05 per 1000 in the Barreiro region. During the environmental monitoring, the largest accumulated electric field measured was 12.4 V/m and the smallest was 0.4 V/m. The largest density power was 40.78 $\mu\text{W}/\text{cm}^2$, and the smallest was 0.04 $\mu\text{W}/\text{cm}^2$.

Adang D, Remacle C, Vorst AV Results of a long-term low-level microwave exposure of rats. IEEE Trans Microwave Theor Tech 57: 2488-2497, 2009.

This paper summarizes the results of experimental research on biological effects induced by electromagnetic exposure to low-level microwaves. We exposed four-month-old Wistar albino rats during 21 months to two different microwave frequencies and exposure modes, 2 h a day, seven days a week. In order to assess possible biological effects of microwaves, we selected among others the following parameters: leucocytes, erythrocytes, monocytes, neutrophils, lymphocytes, hemoglobin, mean corpuscular hemoglobin concentration, and mortality rate. After three and eight months of exposure, we found a statistically significant difference of about 20% between the 970-MHz continuous wave group and sham-exposed group regarding the monocytes in both considered periods. After 14 and 18 months of exposure, we observed a significant

Studies that show **Cell Tower** Health Effects

increase in white blood cells and neutrophils of about 15% and 25%, respectively. Lymphocytes fell down after 18 months of exposure with about 15% compared to the sham-exposed group. No other statistically significant differences were found, except for minor changes with little biological significance. The most obvious effect we detected is the increase in mortality rate of the exposed groups with respect to the sham-exposed group after 21 months of exposure at the age of 25 months. This increase even increases when observing rats until the age of 28 months: mortality in exposed groups then reaches almost twice the value observed in the sham-exposed group.

Effects on Hormones

Eskander EF, Estefan SF, Abd-Rabou AA. How does long term exposure to base stations and mobile phones affect human hormone profiles? Clin Biochem. 45(1-2):157-161, 2012

OBJECTIVES: This study is concerned with assessing the role of exposure to radio frequency radiation (RFR) emitted either from mobiles or base stations and its relations with human's hormone profiles. **DESIGN AND METHODS:** All volunteers' samples were collected for hormonal analysis. **RESULTS:** This study showed significant decrease in volunteers' ACTH, cortisol, thyroid hormones, prolactin for young females, and testosterone levels. **CONCLUSION:** The present study revealed that high RFR effects on pituitary-adrenal axis.

Eşmekaya MA, Seyhan N, Omeroğlu S. Pulse modulated 900 MHz radiation induces hypothyroidism and apoptosis in thyroid cells: A light, electron microscopy and immunohistochemical study. Int J Radiat Biol. 86(12):1106-1116, 2010.

Purpose: In the present study we investigated the possible histopathological effects of pulse modulated Radiofrequency (RF) fields on the thyroid gland using light microscopy, electron microscopy and immunohistochemical methods. Materials and methods: Two months old male Wistar rats were exposed to a 900 MHz pulse-modulated RF radiation at a specific absorption rate (SAR) of 1.35 Watt/kg for 20 min/day for three weeks. The RF signals were pulse modulated by rectangular pulses with a repetition frequency of 217 Hz and a duty cycle of 1:8 (pulse width 0.576 ms). To assess thyroid endocrine disruption and estimate the degree of the pathology of the gland, we analysed structural alterations in follicular and colloidal diameters and areas, colloid content of the follicles, and height of the follicular epithelium. Apoptosis was confirmed by Transmission Electron Microscopy and assessing the activities of an initiator (caspase-9) and an effector (caspase-3) caspases that are important markers of cells undergoing apoptosis. Results: Morphological analyses revealed hypothyrophy of the gland in the 900 MHz RF exposure group. The results indicated that thyroid hormone secretion was inhibited by the RF radiation. In addition, we also observed formation of apoptotic bodies and increased caspase-3 and caspase-9 activities in thyroid cells of the rats that were exposed to modulated RF fields. Conclusion: The overall findings indicated that whole body exposure to pulse-modulated RF radiation that is similar to that emitted by

Studies that show **Cell Tower** Health Effects

global system for mobile communications (GSM) mobile phones can cause pathological changes in the thyroid gland by altering the gland structure and enhancing caspase-dependent pathways of apoptosis.

Genetic Effects

Gandhi G, Kaur G, Nisar U. A cross-sectional case control study on genetic damage in individuals residing in the vicinity of a mobile phone base station. Electromagn Biol Med. 2014 Jul 9:1-11. [Epub ahead of print]

Mobile phone base stations facilitate good communication, but the continuously emitting radiations from these stations have raised health concerns. Hence in this study, genetic damage using the single cell gel electrophoresis (comet) assay was assessed in peripheral blood leukocytes of individuals residing in the vicinity of a mobile phone base station and comparing it to that in healthy controls. The power density in the area within 300 m from the base station exceeded the permissive limits and was significantly ($p = 0.000$) higher compared to the area from where control samples were collected. The study participants comprised 63 persons with residences near a mobile phone tower, and 28 healthy controls matched for gender, age, alcohol drinking and occupational sub-groups. Genetic damage parameters of DNA migration length, damage frequency (DF) and damage index were significantly ($p = 0.000$) elevated in the sample group compared to respective values in healthy controls. The female residents ($n = 25$) of the sample group had significantly ($p = 0.004$) elevated DF than the male residents ($n = 38$). The linear regression analysis further revealed daily mobile phone usage, location of residence and power density as significant predictors of genetic damage. The genetic damage evident in the participants of this study needs to be addressed against future disease-risk, which in addition to neurodegenerative disorders, may lead to cancer.

Fucic A, Garaj-Vrhovac V, Skara M, Dimitrovic B, X-rays, microwaves and vinyl chloride monomer: their clastogenic and aneugenic activity, using the micronucleus assay on human lymphocytes. Mutat Res 282(4):265-271, 1992.

Chromosome aberration assays, sister-chromatid exchange techniques and micronucleus assays are commonly used methods for biomonitoring genetic material damaged by chemical or physical agents. On the other hand, their aneugenic activity, which can lead to hypoploidy and may also be associated with carcinogenesis, has not been thoroughly investigated. In our study we chose the micronucleus assay with a new mathematical approach to separate clastogenic from aneugenic activity of three well-known mutagens (vinyl chloride monomer, X-rays and microwaves) on the genome of human somatic cells. The comparison of frequencies of size distribution of micronuclei in the lymphocytes of humans exposed to each of these three mutagens showed that X-rays and microwaves were preferentially clastogens while vinyl chloride monomer showed aneugenic activity as well. Microwaves possess some mutagenic characteristics typical of chemical mutagens.

Studies that show **Cell Tower** Health Effects

Esmekaya MA, Aytekin E, Ozgur E, Güler G, Ergun MA, Omeroğlu S, Seyhan N. Mutagenic and morphologic impacts of 1.8GHz radiofrequency radiation on human peripheral blood lymphocytes (hPBLs) and possible protective role of pre-treatment with Ginkgo biloba (EGb 761). Sci Total Environ. 410-411:59-64, 2011.

The mutagenic and morphologic effects of 1.8GHz Global System for Mobile Communications (GSM) modulated RF (radiofrequency) radiation alone and in combination with Ginkgo biloba (EGb 761) pre-treatment in human peripheral blood lymphocytes (hPBLs) were investigated in this study using Sister Chromatid Exchange (SCE) and electron microscopy. Cell viability was assessed with 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide (MTT) reduction assay. The lymphocyte cultures were exposed to GSM modulated RF radiation at 1.8GHz for 6, 8, 24 and 48h with and without EGb 761. We observed morphological changes in pulse-modulated RF radiated lymphocytes. Longer exposure periods led to destruction of organelle and nucleus structures. Chromatin change and the loss of mitochondrial crista occurred in cells exposed to RF for 8h and 24h and were more pronounced in cells exposed for 48h. Cytoplasmic lysis and destruction of membrane integrity of cells and nuclei were also seen in 48h RF exposed cells. There was a significant increase ($p < 0.05$) in SCE frequency in RF exposed lymphocytes compared to sham controls. EGb 761 pre-treatment significantly decreased SCE from RF radiation. RF radiation also inhibited cell viability in a time dependent manner. The inhibitory effects of RF radiation on the growth of lymphocytes were marked in longer exposure periods. EGb 761 pre-treatment significantly increased cell viability in RF+EGb 761 treated groups at 8 and 24h when compared to RF exposed groups alone. The results of our study showed that RF radiation affects cell morphology, increases SCE and inhibits cell proliferation. However, EGb 761 has a protective role against RF induced mutagenity. We concluded that RF radiation induces chromosomal damage in hPBLs but this damage may be reduced by EGb 761 pre-treatment.

Kim JY, Hong SY, Lee YM, Yu SA, Koh WS, Hong JR, Son T, Chang SK, Lee M. In vitro assessment of clastogenicity of mobile-phone radiation (835 MHz) using the alkaline comet assay and chromosomal aberration test. Environ Toxicol. 23(3):319-327, 2008.

Recently we demonstrated that 835-MHz radiofrequency radiation electromagnetic fields (RF-EMF) neither affected the reverse mutation frequency nor accelerated DNA degradation in vitro. Here, two kinds of cytogenetic endpoints were further investigated on mammalian cells exposed to 835-MHz RF-EMF (the most widely used communication frequency band in Korean CDMA mobile phone networks) alone and in combination with model clastogens: in vitro alkaline comet assay and in vitro chromosome aberration (CA) test. No direct cytogenetic effect of 835-MHz RF-EMF was found in the in vitro CA test. The combined exposure of the cells to RF-EMF in the presence of ethylmethanesulfonate (EMS) revealed a weak and insignificant cytogenetic effect when compared to cells exposed to EMS alone in CA test. Also, the comet assay results to evaluate the ability of RF-EMF alone to damage DNA were nearly negative, although showing a small increase in tail moment. However, the applied RF-EMF had potentiation

Studies that show **Cell Tower** Health Effects

effect in comet assay when administered in combination with model clastogens (cyclophosphamide or 4-nitroquinoline 1-oxide). Thus, our results imply that we cannot confidently exclude any possibility of an increased risk of genetic damage, with important implications for the possible health effects of exposure to 835-MHz electromagnetic fields.

Nikolova T, Czyz J, Rolletschek A, Blyszczuk P, Fuchs J, Jovtchev G, Schuderer J, Kuster N, Wobus AM. Electromagnetic fields affect transcript levels of apoptosis-related genes in embryonic stem cell-derived neural progenitor cells. ASEB J. 19(12):1686-1688, 2005.

Mouse embryonic stem (ES) cells were used as an experimental model to study the effects of electromagnetic fields (EMF). ES-derived nestin-positive neural progenitor cells were exposed to extremely low frequency EMF simulating power line magnetic fields at 50 Hz (ELF-EMF) and to radiofrequency EMF simulating the Global System for Mobile Communication (GSM) signals at 1.71 GHz (RF-EMF). Following EMF exposure, cells were analyzed for transcript levels of cell cycle regulatory, apoptosis-related, and neural-specific genes and proteins; changes in proliferation; apoptosis; and cytogenetic effects. Quantitative RT-PCR analysis revealed that ELF-EMF exposure to ES-derived neural cells significantly affected transcript levels of the apoptosis-related bcl-2, bax, and cell cycle regulatory "growth arrest DNA damage inducible" GADD45 genes, whereas mRNA levels of neural-specific genes were not affected. RF-EMF exposure of neural progenitor cells resulted in down-regulation of neural-specific Nurr1 and in up-regulation of bax and GADD45 mRNA levels. Short-term RF-EMF exposure for 6 h, but not for 48 h, resulted in a low and transient increase of DNA double-strand breaks. No effects of ELF- and RF-EMF on mitochondrial function, nuclear apoptosis, cell proliferation, and chromosomal alterations were observed. We may conclude that EMF exposure of ES-derived neural progenitor cells transiently affects the transcript level of genes related to apoptosis and cell cycle control. However, these responses are not associated with detectable changes of cell physiology, suggesting compensatory mechanisms at the translational and posttranslational level.

Maes A, Collier M, Slaets D, Verschaeve L, 954 MHz microwaves enhance the mutagenic properties of mitomycin C. Environ Mol Mutagen 28(1):26-30, 1996.

This paper focuses on the combined effects of microwaves from mobile communication frequencies and a chemical DNA damaging agent mitomycin C (MMC). The investigation was performed in vitro by exposing whole blood samples to a 954 MHz emitting antenna from a GSM (Global System for Mobile Communication) base station, followed by lymphocyte cultivation in the presence of MMC. A highly reproducible synergistic effect was observed as based on the frequencies of sister chromatid exchanges in metaphase figures.

Fritze K, Wiessner C, Kuster N, Sommer C, Gass P, Hermann DM, Kiessling M, Hossmann KA, Effect of global system for mobile communication microwave exposure on the genomic response of the rat brain. Neuroscience 81(3):627-639, 1997.

Studies that show **Cell Tower** Health Effects

The acute effect of global system for mobile communication (GSM) microwave exposure on the genomic response of the central nervous system was studied in rats by measuring changes in the messenger RNAs of hsp70, the transcription factor genes c-fos and c-jun and the glial structural gene GFAP using in situ hybridization histochemistry. Protein products of transcription factors, stress proteins and marker proteins of astroglial and microglial activation were assessed by immunocytochemistry. Cell proliferation was evaluated by bromodeoxyuridine incorporation. A special GSM radiofrequency test set, connected to a commercial cellular phone operating in the discontinuous transmission mode, was used to simulate GSM exposure. The study was conducted at time averaged and brain averaged specific absorption rates of 0.3 W/kg (GSM exposure), 1.5 W/kg (GSM exposure) and 7.5 W/kg (continuous wave exposure), respectively. Immediately after exposure, in situ hybridization revealed slight induction of hsp70 messenger RNA in the cerebellum and hippocampus after 7.5 W/kg exposure, but not at lower intensities. A slightly increased expression of c-fos messenger RNA was observed in the cerebellum, neocortex and piriform cortex of all groups subjected to immobilization, but no differences were found amongst different exposure conditions. C-jun and GFAP messenger RNAs did not increase in any of the experimental groups. 24 h after exposure, immunocytochemical analysis of FOS and JUN proteins (c-FOS, FOS B, c-JUN JUN B, JUN D), of HSP70 or of KROX-20 and -24 did not reveal any alterations. Seven days after exposure, neither increased cell proliferation nor altered expression of astroglial and microglial marker proteins were observed. In conclusion, acute high intensity microwave exposure of immobilized rats may induce some minor stress response but does not result in lasting adaptive or reactive changes of the brain.

Baohong Wang, Jiliang H, Lifen J, Deqiang L, Wei Z, Jianlin L, Hongping D. Studying the synergistic damage effects induced by 1.8GHz radiofrequency field radiation (RFR) with four chemical mutagens on human lymphocyte DNA using comet assay in vitro. *Mutat Res.* 578(1-2):149-157, 2005.

The aim of this investigation was to study the synergistic DNA damage effects in human lymphocytes induced by 1.8GHz radiofrequency field radiation (RFR, SAR of 3W/kg) with four chemical mutagens, i.e. mitomycin C (MMC, DNA crosslinker), bleomycin (BLM, radiomimetic agent), methyl methanesulfonate (MMS, alkylating agent), and 4-nitroquinoline-1-oxide (4NQO, UV-mimetic agent). The DNA damage of lymphocytes exposed to RFR and/or with chemical mutagens was detected at two incubation time (0 or 21h) after treatment with comet assay in vitro. Three combinative exposure ways were used. Cells were exposed to RFR and chemical mutagens for 2 and 3h, respectively. Tail length (TL) and tail moment (TM) were utilized as DNA damage indexes. The results showed no difference of DNA damage indexes between RFR group and control group at 0 and 21h incubation after exposure ($P>0.05$). There were significant difference of DNA damage indexes between MMC group and RFR+MMC co-exposure group at 0 and 21h incubation after treatment ($P<0.01$). Also the significant difference of DNA damage indexes between 4NQO group and RFR+4NQO co-exposure group at 0 and 21h incubation after treatment was observed ($P<0.05$ or $P<0.01$). The DNA damage in RFR+BLM co-exposure groups and RFR+MMS co-exposure groups was not significantly

Studies that show **Cell Tower** Health Effects

increased, as compared with corresponding BLM and MMS groups ($P>0.05$). The experimental results indicated 1.8GHz RFR (SAR, 3W/kg) for 2h did not induce the human lymphocyte DNA damage effects in vitro, but could enhance the human lymphocyte DNA damage effects induced by MMC and 4NQO. The synergistic DNA damage effects of 1.8GHz RFR with BLM or MMS were not obvious.

Baohong W, Lifen J, Lanjuan L, Jianlin L, Deqiang L, Wei Z, Jiliang H. Evaluating the combinative effects on human lymphocyte DNA damage induced by ultraviolet ray C plus 1.8GHz microwaves using comet assay in vitro. *Toxicology*. 232(3):311-316, 2007.

The objective of this study was to observe whether 1.8GHz microwaves (MW) (SAR, 3 W/kg) exposure can influence human lymphocyte DNA damage induced by ultraviolet ray C (UVC). The lymphocytes, which were from three young healthy donors, were exposed to 254 nm UVC at the doses of 0.25, 0.5, 0.75, 1.0, 1.5 and 2.0 J m⁻², respectively. The lymphocytes were irradiated by 1.8GHz MW (SAR, 3 W/kg) for 0, 1.5 and 4 h. The combinative exposure of UVC plus MW was conducted. The treated cells were incubated for 0, 1.5 and 4 h. Finally, comet assay was used to measure DNA damage of above treated lymphocytes. The results indicated that the difference of DNA damage induced between MW group and control group was not significant ($P>0.05$). The MTLs induced by UVC were 1.71 \pm 0.09, 2.02 \pm 0.08, 2.27 \pm 0.17, 2.27 \pm 0.06, 2.25 \pm 0.12, 2.24 \pm 0.11 microm, respectively, which were significantly higher than that (0.96 \pm 0.05 microm) of control ($P<0.01$). MTLs of some sub-groups in combinative exposure groups at 1.5-h incubation were significantly lower than those of corresponding UVC sub-groups ($P<0.01$ or $P<0.05$). However, MTLs of some sub-groups in combinative exposure groups at 4-h incubation were significantly higher than those of corresponding UVC sub-groups ($P<0.01$ or $P<0.05$). In this experiment it was found that 1.8GHz (SAR, 3 W/kg) MW exposure for 1.5 and 4 h did not enhance significantly human lymphocyte DNA damage, but could reduce and increase DNA damage of human lymphocytes induced by UVC at 1.5-h and 4-h incubation, respectively.

Canseven AG, Esmekaya MA, Kayhan H, Tuysuz MZ, Seyhan N. Effects of microwave exposure and Gemcitabine treatment on apoptotic activity in Burkitt's lymphoma (Raji) cells. *Electromagn Biol Med*. 2014 Jun 5:1-5. [Epub ahead of print]

We investigated the effects of 1.8 MHz Global System for Mobile Communications (GSM)-modulated microwave (MW) radiation on apoptotic level and cell viability of Burkitt's lymphoma (Raji) cells with or without Gemcitabine, which exhibits cell phase specificity, primarily killing cells undergoing DNA synthesis (S-phase). Raji cells were exposed to 1.8 GHz GSM-modulated MW radiation at a specific absorption rate (SAR) of 0.350 W/kg in a CO₂ incubator. The duration of the exposure was 24 h. The amount of apoptotic cells was analyzed using Annexin V-FITC and propidium iodide (PI) staining with flow cytometer. The apoptotic activity of MW exposed Raji cells was increased significantly. In addition, cell viability of exposed samples was significantly decreased. Combined exposure of MW and Gemcitabine increased the amount of apoptotic cells than MW radiation alone. Moreover, viability of MW + Gemcitabine exposed cells was lower than that of cells exposed only to MW. These results demonstrated that MW

Studies that show **Cell Tower** Health Effects

radiation exposure and Gemcitabine treatment have a synergistic effect on apoptotic activity of Raji cells.

Effects on Glands

Aydogan F, Unlu I, Aydin E, Yumusak N, Devrim E, Samim EE, Ozgur E, Unsal V, Tomruk A, Ozturk GG, Seyhan N. The effect of 2100 MHz radiofrequency radiation of a 3G mobile phone on the parotid gland of rats. Am J Otolaryngol. 2014 Oct 5. pii: S0196-0709(14)00207-5. doi: 10.1016/j.amjoto.2014.10.001. [Epub ahead of print]

PURPOSE: We aimed to evaluate the effect of 2100 MHz radiofrequency radiation on the parotid gland of rats in short and relatively long terms. **MATERIAL AND METHODS:** Thirty Wistar albino rats were divided into four groups. Groups A and B served as the control groups (for 10 days and 40 days, respectively), and each group included six rats. Groups C and D were composed of nine rats each, and they were the exposure groups. The rats were exposed to 2100 MHz radiofrequency radiation emitted by a generator, simulating a third generation mobile phone for 6 hours/day, 5 days/week, for 10 or 40 days. Following exposure, the rats were sacrificed and parotid glands were removed. Histopathological and biochemical examinations were performed. **RESULTS:** Although there were no histopathological changes in the control groups except for two animals in group A and three animals in group B, the exposure groups C (10 days) and D (40 days) showed numerous histopathological changes regarding salivary gland damage including acinar epithelial cells, interstitial space, ductal system, vascular system, nucleus, amount of cytoplasm and variations in cell size. The histopathological changes were more prominent in group D compared to group C. There was statistically significant different parameter regarding variation in cell size between the groups B and D ($p=0.036$). **CONCLUSION:** The parotid gland of rats showed numerous histopathological changes after exposure to 2100 MHz radiofrequency radiation, both in the short and relatively long terms. Increased exposure duration led to an increase in the histopathological changes.

Effects on Animals and Environment

Panagopoulos DJ, Chavdoula ED, Margaritis LH. Bioeffects of mobile telephony radiation in relation to its intensity or distance from the antenna. Int J Radiat Biol. 86(5):345-357, 2010.

PURPOSE: To examine the bioactivity of GSM 900 and 1800 (Global System for Mobile Telecommunications) radiations, in relation to the distance from the antenna or to the radiation-field intensities. **MATERIALS AND METHODS:** *Drosophila melanogaster* adult insects were exposed to the radiation of a GSM 900/1800 mobile phone antenna at different distances ranging from 0 to 100 cm, and the effect on their reproductive capacity and cell death induction in the gonads by the use of TUNEL (Terminal

Studies that show **Cell Tower** Health Effects

deoxynucleotide transferase dUTP Nick End Labeling) assay, was studied. RESULTS: These radiations/fields decreased the reproductive capacity by cell death induction, at all the different distances tested. The effect diminished with the distance/decreasing intensities. An increased bioactivity 'window' was revealed at distances of 20-30 cm from the mobile phone antenna, (radiation intensity around 10 microW/cm(2)) where the effect became highest, in relation to smaller or longer distances. The effect diminished considerably for distances longer than 40-50 cm and became not evident for distances longer than 1 m or radiation intensities smaller than 1 microW/cm(2). CONCLUSIONS: GSM bioactivity is highest for intensities down to less than 10 microW/cm(2) and still evident until 1 microW/cm(2) exhibiting 'window' effects

Loscher W, Kas G, Extraordinary behavior disorders in cows in proximity to transmission stations. Der Praktische Tierarz 79:437-444, 1998. (Article in German) In addition to reduction of milk yield and increased health problems, behavioral abnormalities were observed over a period of two years in a herd of dairy cows maintained in close proximity to a TV and cell phone transmitting antenna. Evaluation of possible factors which could explain the abnormalities in the live stock did not disclose any factors other than the high-frequency electromagnetic fields. An experiment in which a cow with abnormal behavior was brought to a stable 20 km away from the antenna resulted in a complete normalization of the cow within five days, whereas symptoms returned when the cow was brought back to the stable nearby the antenna. In view of the previous described effects of electromagnetic fields, it might be possible that the observed abnormalities in cows are related to electromagnetic field exposure. (power densities measured 0.02-7 mW/m2).

Koldayev VM, Shchepin YV, Effects of electromagnetic radiation on embryos of sea-urchins. Bioelectrochem Bioenerg 43:161-164, 1997. Electromagnetic radiation (EMR) causes a decrease in the number of fertilized eggs and an increase in the number of zygotes with abnormal fertilization envelopes in sea-urchins. The microstructural impairments of the cellular surface, the increase of lipid peroxidation and the changes of amino acid metabolism show that the impairments of the development of embryos exposed to EMR are caused by the damages of the membrane structures.

Cammaerts MC, De Doncker P, Patris X, Bellens F, Rachidi Z, Cammaerts D. GSM 900 MHz radiation inhibits ants' association between food sites and encountered cues. Electromagn Biol Med. 31(2):151-165, 2012.

The kinetics of the acquisition and loss of the use of olfactory and visual cues were previously obtained in six experimental colonies of the ant *Myrmica sabuleti* meinert 1861, under normal conditions. In the present work, the same experiments were conducted on six other naive identical colonies of *M. sabuleti*, under electromagnetic radiation similar to those surrounding GSM and communication masts. In this situation, no association between food and either olfactory or visual cues occurred. After a recovery period, the ants were able to make such an association but never reached the

Studies that show **Cell Tower** Health Effects

expected score. Such ants having acquired a weaker olfactory or visual score and still undergoing olfactory or visual training were again submitted to electromagnetic waves. Not only did they lose all that they had memorized, but also they lost it in a few hours instead of in a few days (as under normal conditions when no longer trained). They kept no visual memory at all (instead of keeping 10% of it as they normally do). The impact of GSM 900 MHz radiation was greater on the visual memory than on the olfactory one. These communication waves may have such a disastrous impact on a wide range of insects using olfactory and/or visual memory, i.e., on bees.

Senavirathna MD, Asaeda T, Thilakarathne BL, Kadono H. Nanometer-scale elongation rate fluctuations in the *Myriophyllum aquaticum* (Parrot feather) stem were altered by radio-frequency electromagnetic radiation. *Plant Signal Behav.* 2014 Mar 26;9(3). pii: e28590. [Epub ahead of print]

The emission of radio-frequency electromagnetic radiation (EMR) by various wireless communication base stations has increased in recent years. While there is wide concern about the effects of EMR on humans and animals, the influence of EMR on plants is not well understood. In this study, we investigated the effect of EMR on the growth dynamics of *Myriophyllum aquaticum* (Parrot feather) by measuring the nanometric elongation rate fluctuation (NERF) using a statistical interferometry technique. Plants were exposed to 2 GHz EMR at a maximum of 1.42 Wm⁻² for 1 h. After continuous exposure to EMR, *M. aquaticum* plants exhibited a statistically significant 51 ± 16% reduction in NERF standard deviation. Temperature observations revealed that EMR exposure did not cause dielectric heating of the plants. Therefore, the reduced NERF was due to a non-thermal effect caused by EMR exposure. The alteration in NERF continued for at least 2.5 h after EMR exposure and no significant recovery was found in post-EMR NERF during the experimental period.

Balmori A. Mobile Phone Mast Effects on Common Frog (*Rana temporaria*) Tadpoles: The City Turned into a Laboratory. *Electromagn Biol Med.* 29(1-2):31-35, 2010.

An experiment has been made exposing eggs and tadpoles of the common frog (*Rana temporaria*) to electromagnetic radiation from several mobile (cell) phone antennae located at a distance of 140 meters. The experiment lasted two months, from the egg phase until an advanced phase of tadpole prior to metamorphosis. Measurements of electric field intensity (radiofrequencies and microwaves) in V/m obtained with three different devices were 1.8 to 3.5 V/m. In the exposed group (n = 70), low coordination of movements, an asynchronous growth, resulting in both big and small tadpoles, and a high mortality (90%) was observed. Regarding the control group (n = 70) under the same conditions but inside a Faraday cage, the coordination of movements was normal, the development was synchronous, and a mortality of 4.2% was obtained. These results indicate that radiation emitted by phone masts in a real situation may affect the development and may cause an increase in mortality of exposed tadpoles. This research may have huge implications for the natural world, which is now exposed to high microwave radiation levels from a multitude of phone masts.

Studies that show **Cell Tower** Health Effects

Balode, Z, Assessment of radio-frequency electromagnetic radiation by the micronucleus test in bovine peripheral erythrocytes. Sci Total Environ 180(1):81-85, 1996.

Previous bioindicative studies in the Skrunda Radio Location Station area have focused on the somatic influence of electromagnetic radiation on plants, but it is also important to study genetic effects. We have chosen cows as test animals for cytogenetical evaluation because they live in the same general exposure area as humans, are confined to specific locations and are chronically exposed to radiation. Blood samples were obtained from female Latvian Brown cows from a farm close to and in front of the Skrunda Radar and from cows in a control area. A simplified alternative to the Schiff method of DNA staining for identification of micronuclei in peripheral erythrocytes was applied. Microscopically, micronuclei in peripheral blood erythrocytes were round in shape and exhibited a strong red colour. They are easily detectable as the only coloured bodies in the uncoloured erythrocytes. From each individual animal 2000 erythrocytes were examined at a magnification of x 1000 for the presence of micronuclei. The counting of micronuclei in peripheral erythrocytes gave low average incidences, 0.6 per 1000 in the exposed group and 0.1 per 1000 in the control, but statistically significant ($P < 0.01$) differences were found in the frequency distribution between the control and exposed groups.

Effects on Skin

Cam ST, Seyhan N, Kavaklı C, Celikbıçak O. Effects of 900 MHz Radiofrequency Radiation on Skin Hydroxyproline Contents. Cell Biochem Biophys. 2014 Apr 24. [Epub ahead of print]

The present study aimed to investigate the possible effect of pulse-modulated radiofrequency radiation (RFR) on rat skin hydroxyproline content, since skin is the first target of external electromagnetic fields. Skin hydroxyproline content was measured using liquid chromatography mass spectrometer method. Two months old male wistar rats were exposed to a 900 MHz pulse-modulated RFR at an average whole body specific absorption rate (SAR) of 1.35 W/kg for 20 min/day for 3 weeks. The radiofrequency (RF) signals were pulse modulated by rectangular pulses with a repetition frequency of 217 Hz and a duty cycle of 1:8 (pulse width 0.576 ms). A skin biopsy was taken at the upper part of the abdominal costa after the exposure. The data indicated that whole body exposure to a pulse-modulated RF radiation that is similar to that emitted by the global system for mobile communications (GSM) mobile phones caused a statistically significant increase in the skin hydroxyproline level ($p = 0.049$, Mann-Whitney U test). Under our experimental conditions, at a SAR less than the International Commission on Non-Ionizing Radiation Protection safety limit recommendation, there was evidence that GSM signals could alter hydroxyproline concentration in the rat skin.

Studies that show **Cell Tower** Health Effects

Effects on Protein

Hässig M, Wullschleger M, Naegeli HP, Kupper J, Spiess B, Kuster N, Capstick M, Murbach M. Influence of non ionizing radiation of base stations on the activity of redox proteins in bovines. *BMC Vet Res.* 2014 Jun 19;10(1):136. [Epub ahead of print]

BACKGROUND: The influence of electromagnetic fields on the health of humans and animals is still an intensively discussed and scientifically investigated issue (Prakt Tierarzt 11:15-20, 2003; Umwelt Medizin Gesellschaft 17:326-332, 2004; J Toxicol Environment Health, Part B 12:572-597, 2009). We are surrounded by numerous electromagnetic fields of variable strength, coming from electronic equipment and its power cords, from high-voltage power lines and from antennas for radio, television and mobile communication. Particularly the latter cause's controversy, as everyone likes to have good mobile reception at anytime and anywhere, whereas nobody wants to have such a base station antenna in their proximity. **RESULTS:** In this experiment, the **non-ionizing radiation (NIR)** has resulted in changes in the enzyme activities. Certain enzymes were disabled, others enabled by NIR. Furthermore, individual behavior patterns were observed. While certain cows reacted to NIR, others did not react at all, or even inversely. **CONCLUSION:** The present results coincide with the information from the literature, according to which NIR leads to changes in redox proteins, and that there are individuals who are sensitive to radiation and others that are not. However, the latter could not be distinctly attributed - there are cows that react clearly with one enzyme while they do not react with another enzyme at all, or even the inverse. The study approach of testing ten cows each ten times during three phases has proven to be appropriate. Future studies should however set the post-exposure phase later on.

Effects on Immune Function

Li CY, Liao MH, Lin CW, Tsai WS, Huang CC, Tang TK. Inhibitory Effects of Microwave Radiation on LPS-Induced NFκB Expression in THP-1 Monocytes. *Chin J Physiol.* 55(6):421-427, 2012.

Microwave radiations can be encountered regularly in daily lives. When WHO announced that microwave radiations were a kind of environmental energy which interfere with the physiological functions of the human body, great concerns have been raised over the damages microwave frequencies can do to human physiology. The immunological performance and the activities of the cellular inflammatory factor NFκB have been closely related in monocyte. Due to the effect of phorbol 12-myristate 13-acetate (PMA) on THP-1 monocytes, THP-1 monocytes would differentiate into macrophages and would then react with lipopolysaccharides (LPS), and the amount of NFκB increased in the THP-1 monocytes. Expression of cytokine is affected when cells are exposed to a frequency of 2450 MHz and at 900 W. Thus, in our experiments, an observation was made when THP-1 monocytes were stimulated with PMA and LPS to differentiate into macrophage, the amount of NFκB in cells increased exponentially, and

Studies that show **Cell Tower** Health Effects

the levels of NFκB expression were decreased by the exposure of microwave radiation. In conclusion, microwave radiations were found to inhibit the activity functions of THP-1 monocytes stimulated with PMA and LPS.

Electro Hypersensitivity

Nordin S, Neely G, Olsson D, Sandström M. Odor and Noise Intolerance in Persons with Self-Reported Electromagnetic Hypersensitivity. Int J Environ Res Public Health. 11(9):8794-8805, 2014.

Lack of confirmation of symptoms attributed to electromagnetic fields (EMF) and triggered by EMF exposure has highlighted the role of individual factors. Prior observations indicate intolerance to other types of environmental exposures among persons with electromagnetic hypersensitivity (EHS). This study assessed differences in odor and noise intolerance between persons with EHS and healthy controls by use of subscales and global measures of the Chemical Sensitivity Scale (CSS) and the Noise Sensitivity Scale (NSS). The EHS group scored significantly higher than the controls on all CSS and NSS scales. Correlation coefficients between CSS and NSS scores ranged from 0.60 to 0.65 across measures. The findings suggest an association between EHS and odor and noise intolerance, encouraging further investigation of individual factors for understanding EMF-related symptoms.

RF Levels in Cities

Urbiniello D, Huss A, Beekhuizen J, Vermeulen R, Rösli M. Use of portable exposure meters for comparing mobile phone base station radiation in different types of areas in the cities of Basel and Amsterdam. Sci Total Environ. 468-469:1028-1033, 2014.

BACKGROUND: Radiofrequency electromagnetic fields (RF-EMF) are highly variable and differ considerably within as well as between areas. Exposure assessment studies characterizing spatial and temporal variation are limited so far. Our objective was to evaluate sources of data variability and the repeatability of daily measurements using portable exposure meters (PEMs). METHODS: Data were collected at 12 days between November 2010 and January 2011 with PEMs in four different types of urban areas in the cities of Basel (BSL) and Amsterdam (AMS). RESULTS: Exposure from mobile phone base stations ranged from 0.30 to 0.53 V/m in downtown and business areas and in residential areas from 0.09 to 0.41 V/m. Analysis of variance (ANOVA) demonstrated that measurements from various days were highly reproducible (measurement duration of approximately 30 min) with only 0.6% of the variance of all measurements from mobile phone base station radiation being explained by the measurement day and only 0.2% by the measurement time (morning, noon, afternoon), whereas type of area (30%) and city (50%) explained most of the data variability. CONCLUSIONS: We conclude that mobile monitoring of exposure from mobile phone base station radiation with PEMs is

Studies that show **Cell Tower** Health Effects

useful due to the high repeatability of mobile phone base station exposure levels, despite the high spatial variation.

Urbiniello D, Joseph W, Verloock L, Martens L, Rösli M. Temporal trends of radio-frequency electromagnetic field (RF-EMF) exposure in everyday environments across European cities. Environ Res. 2014 Aug 12;134C:134-142. doi: 10.1016/j.envres.2014.07.003. [Epub ahead of print]

BACKGROUND: The rapid development and increased use of wireless telecommunication technologies led to a substantial change of radio-frequency electromagnetic field (RF-EMF) exposure in the general population but little is known about temporal trends of RF-EMF in our everyday environment. **OBJECTIVES:** The objective of our study is to evaluate temporal trends of RF-EMF exposure levels in different microenvironments of three European cities using a common measurement protocol. **METHODS:** We performed measurements in the cities of Basel (Switzerland), Ghent and Brussels (Belgium) during one year, between April 2011 and March 2012. RF-EMF exposure in 11 different frequency bands ranging from FM (Frequency Modulation, 88MHz) to WLAN (Wireless Local Area Network, 2.5GHz) was quantified with portable measurement devices (exposimeters) in various microenvironments: outdoor areas (residential areas, downtown and suburb), public transports (train, bus and tram or metro rides) and indoor places (airport, railway station and shopping centers). Measurements were collected every 4s during 10-50min per environment and measurement day. Linear temporal trends were analyzed by mixed linear regression models. **RESULTS:** Highest total RF-EMF exposure levels occurred in public transports (all public transports combined) with arithmetic mean values of 0.84V/m in Brussels, 0.72V/m in Ghent, and 0.59V/m in Basel. In all outdoor areas combined, mean exposure levels were 0.41V/m in Brussels, 0.31V/m in Ghent and 0.26V/m in Basel. Within one year, total RF-EMF exposure levels in all outdoor areas in combination increased by 57.1% (p<0.001) in Basel by 20.1% in Ghent (p=0.053) and by 38.2% (p=0.012) in Brussels. Exposure increase was most consistently observed in outdoor areas due to emissions from mobile phone base stations. In public transports RF-EMF levels tended also to increase but mostly without statistical significance. **DISCUSSION:** An increase of RF-EMF exposure levels has been observed between April 2011 and March 2012 in various microenvironments of three European cities. Nevertheless, exposure levels were still far below regulatory limits of each country. A continuous monitoring is needed to identify high exposure areas and to anticipate critical development of RF-EMF exposure at public places.

Estenberg J, Augustsson T. Extensive frequency selective measurements of radiofrequency fields in outdoor environments performed with a novel mobile monitoring system. Bioelectromagnetics. 2013 Dec 27. doi: 10.1002/bem.21830. [Epub ahead of print]

A novel, car based, measuring system for estimation of general public outdoor exposure to radiofrequency fields (RF) has been developed. The system enables fast, large area, isotropic spectral measurements with a bandwidth covering the frequency range of

Studies that show **Cell Tower** Health Effects

30 MHz to 3 GHz. Measurements have shown that complete mapping of a town with 15000 inhabitants and a path length of 115 km is possible to perform within 1 day. The measured areas were chosen to represent typical rural, urban and city areas of Sweden. The data sets consist of more than 70000 measurements. All measurements were performed during the daytime. The median power density was $16 \mu\text{W}/\text{m}^2$ in rural areas, $270 \mu\text{W}/\text{m}^2$ in urban areas, and $2400 \mu\text{W}/\text{m}^2$ in city areas. In urban and city areas, base stations for mobile phones were clearly the dominating sources of exposure.

Effects on Water

Hinrikus H, Lass J, Karai D, Pilt K, Bachmann M. Microwave effect on diffusion: a possible mechanism for non-thermal effect. Electromagn Biol Med. 23:1-7, 2014.

In this study, we assume that microwave radiation affects hydrogen bonding between dipolar water molecules and through that diffusion in water at constant temperature. The experimental study was performed on the setup of two identical reservoirs filled with pure water and 0.9% NaCl solution and connected by a thin tube. Alterations of NaCl concentration in the reservoir initially filled with pure water were measured using the resistance of the solution as an indicator. The applied 450 MHz continuous-wave microwave field had the maximal specific absorption rate of 0.4 W/kg on the connecting tube. The standard deviation of water temperature in the setup was 0.02 °C during an experiment. Our experimental data demonstrated that microwave exposure makes faster the process of diffusion in water. The time required for reduction of initial resistance of the solution by 10% was 1.7 times shorter with microwave. This result is consistent with the proposed mechanism of low-level microwave effect: microwave radiation, rotating dipolar water molecules, causes high-frequency alterations of hydrogen bonds between water molecules, thereby affects its viscosity and makes faster diffusion.

RF Levels From Cell Towers

Martinez-Burdalo M, Martin A, Anguiano M, Villar R. On the safety assessment of human exposure in the proximity of cellular communications base-station antennas at 900, 1800 and 2170 MHz. Phys Med Biol. 50(17):4125-4137, 2005.

In this work, the procedures for safety assessment in the close proximity of cellular communications base-station antennas at three different frequencies (900, 1800 and 2170 MHz) are analysed. For each operating frequency, we have obtained and compared the distances to the antenna from the exposure places where electromagnetic fields are below reference levels and the distances where the specific absorption rate (SAR) values in an exposed person are below the basic restrictions, according to the European safety guidelines. A high-resolution human body model has been located, in front of each base-station antenna as a worst case, at different distances, to compute whole body averaged SAR and maximum 10 g averaged SAR

Studies that show **Cell Tower** Health Effects

inside the exposed body. The finite-difference time-domain method has been used for both electromagnetic fields and SAR calculations. This paper shows that, for antenna-body distances in the near zone of the antenna, the fact that averaged field values be below the reference levels could, at certain frequencies, not guarantee guidelines compliance based on basic restrictions.

Hu J, Lu Y, Zhang H, Xie H, Yang X. [Level of microwave radiation from mobile phone base stations built in residential districts] Wei Sheng Yan Jiu. 38(6):712-716, 2009.
[Article in Chinese]

OBJECTIVE: To investigate the condition of microwave radiation pollution from mobile phone base station built in populated area. **METHODS:** Random selected 18 residential districts where had base station and 10 residential districts where had no base stations. A TES-92 electromagnetic radiation monitor were used to measure the intensity of microwave radiation in external and internal living environment. **RESULTS:** The intensities of microwave radiation in the exposure residential districts were more higher than those of the control residential districts ($p < 0.05$). There was a intensity peak at about 10 m from the station, it would gradually weaken with the increase of the distance. The level of microwave radiation in antenna main lobe region is not certainly more higher than the side lobe direction, and the side lobe direction also is not more lower. At the same district, where there were two base stations, the electromagnetic field nestification would take place in someplace. The intensities of microwave radiation outside the exposure windows in the resident room not only changed with distance but also with the height of the floor. The intensities of microwave radiation inside the aluminum alloys security net were more lower than those of outside the aluminum alloys security net ($p < 0.05$), but the inside or outside of glass-window appears almost no change ($p > 0.05$). **CONCLUSIONS:** Although all the measure dates on the ground around the base station could be below the primary standard in "environment electromagnetic wave hygienic standard" (GB9175-88), there were still a minorities of windows which exposed to the base station were higher, and the outside or inside of a few window was even higher beyond the primary safe level defined standard. The aluminum alloys security net can partly shield the microwave radiation from the mobile phone base station.

Danger Perception and Symptoms

Hutter HP, Moshammer H, Wallner P, Kundi M. Public perception of risk concerning cell towers and mobile phones. Soz Praventivmed. 49(1):62-66, 2004.

OBJECTIVE: The controversy about health risks of electromagnetic fields (EMF) has contributed in raising fears concerning emissions from celltowers. The study was to examine whether or not neighbours of celltowers are particularly concerned about adverse health effects of mobile phones and their base stations. **METHODS:** Prior to information delivered by medical doctors of the Institute of Environmental Health at public hearings a questionnaire was handed out to participants asking for their personal

Studies that show **Cell Tower** Health Effects

rating of several environmental health risks including those of mobile telecommunication (n = 123, response rate approx. 48%). Medical students (n = 366) served as a contrast group. RESULTS: Participants rated health risk for both, mobile phones and celltowers higher as students. A trend for higher ratings was also seen with older subjects and female sex. The risk ratings of both exposures correlated well with each other. The magnitude of the perceived risks, however, resembled that of other ubiquitous exposures like traffic noise and air pollution. CONCLUSION: Contrary to the claims of the telecommunication industry, opponents of celltowers generally do not express unusual fears concerning electromagnetic field exposure. The outcome of our study indicates that the risk rating is comparable with other perceived common hazards of the civilised world. It is hypothesised that offering information and participation to the concerned population will be efficient in reducing exaggerated fears.

Reason Why Not A Lot Of Studies on Cell Towers

Kundi M, Hutter HP. Mobile phone base stations-Effects on wellbeing and health. Pathophysiology. 16(2-3):123-135, 2009.

Studying effects of mobile phone base station signals on health have been discouraged by authoritative bodies like WHO International EMF Project and COST 281. WHO recommended studies around base stations in 2003 but again stated in 2006 that studies on cancer in relation to base station exposure are of low priority. As a result only few investigations of effects of base station exposure on health and wellbeing exist. Cross-sectional investigations of subjective health as a function of distance or measured field strength, despite differences in methods and robustness of study design, found indications for an effect of exposure that is likely independent of concerns and attributions. Experimental studies applying short-term exposure to base station signals gave various results, but there is weak evidence that UMTS and to a lesser degree GSM signals reduce wellbeing in persons that report to be sensitive to such exposures. Two ecological studies of cancer in the vicinity of base stations report both a strong increase of incidence within a radius of 350 and 400m respectively. Due to the limitations inherent in this design no firm conclusions can be drawn, but the results underline the urgent need for a comprehensive investigation of this issue. Animal and in vitro studies are inconclusive to date. An increased incidence of DMBA induced mammary tumors in rats at a SAR of 1.4W/kg in one experiment could not be replicated in a second trial. Indications of oxidative stress after low-level in vivo exposure of rats could not be supported by in vitro studies of human fibroblasts and glioblastoma cells. From available evidence it is impossible to delineate a threshold below which no effect occurs, however, given the fact that studies reporting low exposure were invariably negative it is suggested that power densities around 0.5-1mW/m² must be exceeded in order to observe an effect. The meager data base must be extended in the coming years. The difficulties of investigating long-term effects of base station exposure have been exaggerated, considering that base station and handset exposure have almost nothing

Studies that show **Cell Tower** Health Effects

in common both needs to be studied independently. It cannot be accepted that studying base stations is postponed until there is firm evidence for mobile phones.

Miscellaneous Effects

Panagopoulos, D. J., Johansson O. & Carlo G.L. Polarization: A Key Difference between Man-made and Natural Electromagnetic Fields, in regard to Biological Activity. *Sci. Rep.* 5, 14914; doi: 10.1038/srep14914 (2015). Published online Oct 12, 2015.

In the present study we analyze the role of polarization in the biological activity of Electromagnetic Fields (EMFs)/Electromagnetic Radiation (EMR). All types of man-made EMFs/EMR - in contrast to natural EMFs/EMR - are polarized. Polarized EMFs/EMR can have increased biological activity, due to: 1) Ability to produce constructive interference effects and amplify their intensities at many locations. 2) Ability to force all charged/polar molecules and especially free ions within and around all living cells to oscillate on parallel planes and in phase with the applied polarized field. Such ionic forced-oscillations exert additive electrostatic forces on the sensors of cell membrane electro-sensitive ion channels, resulting in their irregular gating and consequent disruption of the cell's electrochemical balance. These features render man-made EMFs/EMR more bioactive than natural non-ionizing EMFs/EMR. This explains the increasing number of biological effects discovered during the past few decades to be induced by man-made EMFs, in contrast to natural EMFs in the terrestrial environment which have always been present throughout evolution, although human exposure to the latter ones is normally of significantly higher intensities/energy and longer durations. Thus, polarization seems to be a trigger that significantly increases the probability for the initiation of biological/health effects.

Dhami AK. Study of electromagnetic radiation pollution in an Indian city. *Environ Monit Assess.*184(11):6507-6512, 2012.

Abstract. Electromagnetic radiation emitted by cell phone towers is a form of environmental pollution and is a new health hazard, especially to children and patients. The present studies were taken to estimate the microwave/RF pollution by measuring radiation power densities near schools and hospitals of Chandigarh city in India. The cell phone radiations were measured using a handheld portable power density meter TES 593 and specific absorption rates were estimated from the measured values. These values of electromagnetic radiation in the environment were compared with the levels at which biological system of humans and animals starts getting affected. The values were also compared with the international exposure limits set by the International Commission on Non-Ionizing Radiation Protection (ICNIRP). The highest measured power density was 11.48 mW/m(2) which is 1,148% of the biological limit. The results indicated that the exposure levels in the city were below the ICNIRP limit, but much above the biological limit.

December 5, 2016,

Dear Maryland Children's Environmental Health and Protection Advisory Council Members,

As a parent of two elementary school girls in Montgomery County, I am deeply concerned about their daily exposure to Wi-Fi radiation. At home, I can take measures to reduce such radiation, by using hard-wired computers and limiting the amount of time they use wireless devices such as mobile phones and tablets.

However, like many elementary school students throughout the county, my children are seated in a classroom with a wireless access point right above their heads, which emits Wi-Fi radiation the entire school day. Moreover, when they are using their Chromebooks, the students are exposed to even higher levels of radiation emitted from Chromebook antennae. Many of these students are permitted to use Chromebooks on their laps or in other ways that are not in compliance with user manuals (which typically instruct that Chromebooks be used on table tops a good distance away from the user's body). In fact, my daughter's Fifth Grade English classroom is an "alternative design" without desks, and the students regularly (i.e., *daily*) sit on the floor or in beanbags with the Chromebooks on their laps for lengthy periods of time.

Exposure to Wi-Fi radiation is cumulative. Multiply the number of hours in a school day (6.5 hours) by the number of school days in a year (180 days), by the number of years they will have been using Wi-Fi based technology by the time they graduate (for my own children, it will be 11 years), and you arrive at 12,870 cumulative hours. This is a staggering figure to contemplate. Given the recent push to one-to-one computing in schools, my children will be among the first generation to be exposed to Wi-Fi radiation for this long of a period.

I am writing to urge the Council to support strong recommendations to Maryland schools to reduce children's exposure to Wi-Fi radiation. Students and staff should know that wireless devices emit Wi-Fi radiation and that solutions exist to minimize this exposure. Please issue specific recommendations to schools to decrease exposures to as low as possible.

Technology should be hardwired (not wireless) to significantly reduce radiation exposures in the classroom. At a minimum, Wi-Fi should be shut off in classrooms when not in use. The location of wireless access points should not be located above children's heads and developing brains. Children should be using computers on desks, not their laps, and the equipment should be as far away from their bodies as possible.

We are at a critical juncture. More communities throughout the world are recognizing that unfettered access to Wi-Fi is a health hazard, especially when it comes to children's brain development and growing bodies. Please encourage a safer environment in Maryland schools by issuing recommendations that protect the health of our children.

Sincerely yours,

Leesa Klepper

December 5, 2016,

Dear Maryland Children's Environmental Health and Protection Advisory Council Members,

As a parent of two elementary school girls in Montgomery County, I am deeply concerned about their daily exposure to Wi-Fi radiation. At home, I can take measures to reduce such radiation, by using hard-wired computers and limiting the amount of time they use wireless devices such as mobile phones and tablets.

However, like many elementary school students throughout the county, my children are seated in a classroom with a wireless access point right above their heads, which emits Wi-Fi radiation the entire school day. Moreover, when they are using their Chromebooks, the students are exposed to even higher levels of radiation emitted from Chromebook antennae. Many of these students are permitted to use Chromebooks on their laps or in other ways that are not in compliance with user manuals (which typically instruct that Chromebooks be used on table tops a good distance away from the user's body). In fact, my daughter's Fifth Grade English classroom is an "alternative design" without desks, and the students regularly (i.e., *daily*) sit on the floor or in beanbags with the Chromebooks on their laps for lengthy periods of time.

Exposure to Wi-Fi radiation is cumulative. Multiply the number of hours in a school day (6.5 hours) by the number of school days in a year (180 days), by the number of years they will have been using Wi-Fi based technology by the time they graduate (for my own children, it will be 11 years), and you arrive at 12,870 cumulative hours. This is a staggering figure to contemplate. Given the recent push to one-to-one computing in schools, my children will be among the first generation to be exposed to Wi-Fi radiation for this long of a period.

I am writing to urge the Council to support strong recommendations to Maryland schools to reduce children's exposure to Wi-Fi radiation. Students and staff should know that wireless devices emit Wi-Fi radiation and that solutions exist to minimize this exposure. Please issue specific recommendations to schools to decrease exposures to as low as possible.

Technology should be hardwired (not wireless) to significantly reduce radiation exposures in the classroom. At a minimum, Wi-Fi should be shut off in classrooms when not in use. The location of wireless access points should not be located above children's heads and developing brains. Children should be using computers on desks, not their laps, and the equipment should be as far away from their bodies as possible.


We are at a critical juncture. More communities throughout the world are recognizing that unfettered access to Wi-Fi is a health hazard, especially when it comes to children's brain development and growing bodies. Please encourage a safer environment in Maryland schools by issuing recommendations that protect the health of our children.

Sincerely yours,

Leesa Klepper

HARVARD MEDICAL SCHOOL

Martha R. Herbert, Ph.D., M.D.
Assistant Professor, Neurology
Director, TRANSCEND Research Program
www.transcendresearch.org
transcend@partners.org



MASSACHUSETTS GENERAL HOSPITAL

Martinos Center for Biomedical Imaging
149 13th Street, Room 10.043
Charlestown (Boston), Massachusetts 02129
martha.herbert@mgh.harvard.edu
<https://connects.catalyst.harvard.edu/profiles/display/Person/47629>

December 12, 2015

Montgomery County Schools
Carver Educational Services Center
850 Hungerford Drive
Rockville, MD 20850

cc Montgomery County City Council

Dear Montgomery County School District,

I am a pediatric neurologist and neuroscientist on the faculty of Harvard Medical School and on staff at the Massachusetts General Hospital. I am Board Certified in Neurology with Special Competency in Child Neurology, and Subspecialty Certification in Neurodevelopmental Disorders.

I have an extensive history of research and clinical practice in neurodevelopmental disorders, particularly autism spectrum disorders. I have published papers in brain imaging research, in physiological abnormalities in autism spectrum disorders, and in environmental influences on neurodevelopmental disorders such as autism and on brain development and function.

A few years ago I accepted an invitation to review literature pertinent to a potential link between Autism Spectrum Disorders and Electromagnetic Frequencies (EMF) and Radiofrequency Radiation(RFR). I set out to write a paper of modest length, but found much more literature than I had anticipated to review. I ended up producing a 60 page single spaced paper with over 550 citations. It is available at http://www.biointitiative.org/report/wp-content/uploads/pdfs/sec20_2012_Findings_in_Autism.pdf and it was published in a revised and somewhat shortened form in two parts in the peer reviewed indexed journal *Pathophysiology* (2013) with the title: "Autism and EMF? Plausibility of a pathophysiological link." Please also see the appendix to this letter which contains a summary of this material and includes substantial scientific citations.

More recently I published an article entitled ["Connections in Our Environment: Sizing up Electromagnetic Fields."](#) in *Autism Notebook Spring 2015* edition in which I summarized and personalized the information in the . In this article I describe how here is a whole series of problems at the cellular, sub-cellular and metabolic levels and immune levels that have been identified in autism. And interestingly, for every single one of those problems, there's literature about how EMFs can create those kinds of problems.

The argument I made in these articles is not that EMF is proven to cause autism, but rather, that EMF can certainly contribute to degrading the physiological integrity of the system at the cellular and molecular level" – and this in turn appears to contribute to the pathogenesis/causation not only of autism but of many highly common chronic illnesses, including cancer, obesity, diabetes and heart disease.. Please see this article on page 24-25 at the link <http://virtualpublications.soloprinting.com/publication/?i=252361>

In fact, there are thousands of papers that have accumulated over decades –and are now accumulating at an accelerating pace, as our ability to measure impacts become more sensitive – that document adverse health and neurological impacts of EMF/RFR. Children are more vulnerable than adults, and children with chronic illnesses and/or neurodevelopmental disabilities are even more vulnerable. Elderly or chronically ill adults are more vulnerable than healthy adults.

Current technologies were designed and promulgated without taking account of biological impacts other than thermal impacts. We now know that there are a large array of impacts that have nothing to do with the heating of tissue. The claim from wifi proponents that the only concern is thermal impacts is now definitively outdated scientifically.

Radiofrequency electromagnetic radiation from wifi and cell towers can exert a disorganizing effect on the ability to learn and remember, and can also be destabilizing to immune and metabolic function. This will make it harder for some children to learn, particularly those who are already having learning or medical problems in the first place. And since half of the children in this country have some kind of chronic illness, this means that a lot of people are more vulnerable than you might expect to these issues.

Powerful industrial entities have a vested interest in leading the public to believe that EMF/RFR, which we cannot see, taste or touch, is harmless, but this is not true. Please do the right and precautionary thing for our children.

I urge you to opt for wired technologies in Montgomery County classrooms, particularly for those subpopulations that are most sensitive. It will be easier for you to make a healthier decision now than to undo misguided decisions later.

Thank you.



Martha Herbert, PhD, MD

Selected pertinent publications

[Connections in our Environment: Sizing up Electromagnetic Fields](#) by M.R. Herbert (published in Autism Notebook Spring 2015, pp. 24-25) reviews in two pages key points of the more technical Herbert & Sage Autism-EMF paper

Herbert, M.R. and Sage, C. “Autism and EMF? Plausibility of a Pathophysiological Link”. [Part 1: Pathophysiology, 2013, Jun;20\(3\):191-209](#), epub Oct 4, PMID 24095003. [Pubmed abstract for Part 1](#). [Part II: Pathophysiology, 2013 Jun;20\(3\):211-34](#). Epub 2013 Oct 8, PMID 24113318. [Pubmed abstract for Part II](#).

APPENDIX: MORE DETAILED SUMMARY OF THE PATHOPHYSIOLOGY

I became interested in the health and brain effects of electromagnetic frequency (EMF) and radiofrequency radiation (RFR) exposures in relation to my brain research because I was interested in how such exposures might alter brain function. In order to familiarize myself in more detail existing literature on the pathophysiological impacts of EMF/RFR, I coauthored a 40,000 word chapter in the 2012 update of the Bioinitiative, ¹ and published an updated 30,000 word version of that paper (“Autism and EMF? Plausibility of a Pathophysiological Link”) in 2013 in two parts in the peer reviewed journal *Pathophysiology*. ^{2,3} My intention was to assess the plausibility of an association between increasing incidence of autism spectrum disorder and increasing EMF/RFR exposures. Rather than directly address the epidemiological issues, I looked at the parallels between the pathophysiological features documented in autism and the pathophysiological impacts of EMF/RFR documented in the peer-reviewed published scientific literature.

I will include here a brief summary of the paper (prepared for a lay audience) of the features of EMF/RFR that I reviewed (with citations at the end of this letter):

- EMF/RFR stresses cells. It lead to cellular stress, such as production of heat shock proteins, even when The EMF/RFR isn't intense enough to cause measurable heat increase. ⁴⁻⁶
- EMF/RFR damages cell membranes, and make them leaky, which makes it hard for them to maintain important chemical and electrical differences between what is inside and outside the membrane. This degrades metabolism in many ways – makes it inefficient. ⁷⁻¹⁵
- EMF/RFR damages mitochondria. Mitochondria are the energy factories of our cells. Mitochondria conduct their chemical reactions on their membranes. When those membranes get damaged, the mitochondria struggle to do their work and don't do it so well. Mitochondria can also be damaged through direct hits to steps in their chemical assembly line. When mitochondria get inefficient, so do we. This can hit our brains especially hard, since electrical communication and synapses in the brain demands huge amounts of energy.
- EMF/RFR creates “oxidative stress.” Oxidative stress is something that occurs when the system can't keep up with the stress caused by utilizing oxygen, because the price we pay for using oxygen is that it generates free radicals. These are generated in the normal course of events, and they are “quenched” by antioxidants like we get

in fresh fruits and vegetables; but when the antioxidants can't keep up or the damage is too great, the free radicals start damaging things.

- EMF/RFR is genotoxic and damages proteins, with a major mechanism being EMF/RFR-created free radicals which damage cell membranes, DNA, proteins, anything they touch. When free radicals damage DNA they can cause mutations. This is one of the main ways that EMF/RFR is genotoxic – toxic to the genes. When they damage proteins they can cause them to fold up in peculiar ways. We are learning that diseases like Alzheimer's are related to the accumulation of mis-folded proteins, and the failure of the brain to clear out this biological trash from its tissues and fluids.
- EMF/RFR depletes glutathione, which is the body's premier antioxidant and detoxification substance. So on the one hand EMF/RFR creates damage that increases the need for antioxidants, and on the other hand they deplete those very antioxidants.^{1, 16}
- EMF/RFR damages vital barriers in the body, particularly the blood-brain barrier, which protects the brain from things in the blood that might hurt the brain. When the blood-brain barrier gets leaky, cells inside the brain suffer, be damaged, and get killed.^{1, 16, 17}
- EMF/RFR can alter the function of calcium channels, which are openings in the cell membranes that play a huge number of vital roles in brain and body.¹⁸⁻²⁷
- EMF/RFR degrades the rich, complex integration of brainwaves, and increase the "entropy" or disorganization of signals in the brain – this means that they can become less synchronized or coordinated; such reduced brain coordination has been measured in autism.²⁸⁻⁴⁰
- EMF/RFR can interfere with sleep and the brain's production of melatonin.⁴¹⁻⁴³
- EMF/RFR can contribute to immune problems.⁴⁴⁻⁵⁰
- EMF/RFR contribute to increasing stress at the chemical, immune and electrical levels, which we experience psychologically.^{51-57 17, 58-62 63-68}

Please note that:

1. There are a lot of other things that can create similar damaging effects, such as thousands of "xenobiotic" substances that we call toxicants. Significantly, toxic chemicals (including those that contain naturally occurring toxic elements such as lead and mercury) cause damage through many of the same mechanisms outlined above.
2. In many of the experimental studies with EMF/RFR, damage could be diminished by improving nutrient status, particularly by adding antioxidants and melatonin.⁶⁹⁻⁷²

I understand that the concept of electromagnetic hypersensitivity is not always well understood in the medical and scientific communities. Indeed, the inter-individual variability is perplexing to those who would expect a more consistent set of features.

But given the range of challenges I have listed that EMF/RFR poses to core processes in biological systems, and given the inter-individually variable vulnerability across these symptoms, it is really not surprising that there would be subgroups with different combinations of symptom clusters.

It also appears to be the case that the onset and duration of symptoms or even brain response to EMR/RFR can be variable. This again is to be expected given the mediation of these symptoms through a variety of the above-listed pathophysiological processes, many of which differ in scale (ranging from molecular to cellular to tissue and organ) and time course of impact. The different parts of the body also absorb this energy differently, both

because of their biophysical properties and as a function of their state of health or compromise thereof.

Here is a list of subgroups of symptom clusters identified by a group of German physicians, t exemplifies these variability issues:

- Group 1** no symptoms
- Group 2** sleep disturbance, tiredness, depressive mood
- Group 3** headaches, restlessness, dazed state, irritability, disturbance of concentration, forgetfulness, learning difficulties, difficulty finding words
- Group 4** frequent infections, sinusitis, lymph node swellings, joint and limb pains, nerve and soft tissue pains, numbness or tingling, allergies
- Group 5** tinnitus, hearing loss, sudden hearing loss, giddiness, impaired balance, visual disturbances, eye inflammation, dry eyes
- Group 6** tachycardia, episodic hypertension, collapse
- Group 7** other symptoms: hormonal disturbances, thyroid disease, night sweats, frequent urge to urinate, weight increase, nausea, loss of appetite, nose bleeds, skin complaints, tumors, diabetes

CITATIONS

1. Herbert MR, Sage C. Findings in autism spectrum disorders consistent with electromagnetic frequencies (emf) and radiofrequency radiation (rfr). *BioInitiative Update*. 2012
2. Herbert MR, Sage C. Autism and emf? Plausibility of a pathophysiological link - part i. *Pathophysiology*. 2013;20:191-209
3. Herbert MR, Sage C. Autism and emf? Plausibility of a pathophysiological link - part ii. *Pathophysiology*. 2013;20:211-234
4. Blank M. Electromagnetic fields. *Pathophysiology*. 2009;16 (2-3)
5. Blank M. Evidence for stress response (stress proteins) (section 7). *The BioInitiative Report 2012: A Rationale for a Biologically-based Public Exposure Standard for Electromagnetic Fields (ELF and RF)*. 2012:<http://www.bioinitiative.org/table-of-contents/>
6. Evers M, Cunningham-Rundles C, Hollander E. Heat shock protein 90 antibodies in autism. *Mol Psychiatry*. 2002;7 Suppl 2:S26-28
7. Desai NR, Kesari KK, Agarwal A. Pathophysiology of cell phone radiation: Oxidative stress and carcinogenesis with focus on male reproductive system. *Reprod Biol Endocrinol*. 2009;7:114
8. Phelan AM, Lange DG, Kues HA, Luty GA. Modification of membrane fluidity in melanin-containing cells by low-level microwave radiation. *Bioelectromagnetics*. 1992;13:131-146
9. Beneduci A, Filippelli L, Cosentino K, Calabrese ML, Massa R, Chidichimo G. Microwave induced shift of the main phase transition in phosphatidylcholine membranes. *Bioelectrochemistry*. 2012;84:18-24
10. El-Ansary A, Al-Ayadhi L. Lipid mediators in plasma of autism spectrum disorders. *Lipids Health Dis*. 2012;11:160
11. El-Ansary AK, Bacha AG, Al-Ayahdi LY. Plasma fatty acids as diagnostic markers in autistic patients from saudi arabia. *Lipids Health Dis*. 2011;10:62
12. Chauhan A, Chauhan V, Brown WT, Cohen I. Oxidative stress in autism: Increased lipid peroxidation and reduced serum levels of ceruloplasmin and transferrin--the antioxidant proteins. *Life Sci*. 2004;75:2539-2549
13. Pecorelli A, Leoncini S, De Felice C, Signorini C, Cerrone C, Valacchi G, et al. Non-protein-bound iron and 4-hydroxynonenal protein adducts in classic autism. *Brain Dev*. 2012:epub.
14. Ming X, Stein TP, Brimacombe M, Johnson WG, Lambert GH, Wagner GC. Increased excretion of a lipid peroxidation biomarker in autism. *Prostaglandins Leukot Essent Fatty Acids*. 2005;73:379-384
15. Yao Y, Walsh WJ, McGinnis WR, Pratico D. Altered vascular phenotype in autism: Correlation with oxidative stress. *Arch Neurol*. 2006;63:1161-1164
16. Herbert MR, Sage C. Autism and emf? Plausibility of a pathophysiological link, parts i and ii. *Pathophysiology*. In press

17. Salford LG, Nittby H, Persson BR. Effects of emf from wireless communication upon the blood-brain barrier. *BioInitiative 2012: A Rationale for a Biologically-based Public Exposure Standard for Electromagnetic Fields (ELF and RF)*. 2012;www.bioinitiative.org, Section 10
18. Pall ML. Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. *J Cell Mol Med*. 2013
19. Nesin V, Bowman AM, Xiao S, Pakhomov AG. Cell permeabilization and inhibition of voltage-gated Ca^{2+} and Na^{+} channel currents by nanosecond pulsed electric field. *Bioelectromagnetics*. 2012;33:394-404
20. Maskey D, Kim HJ, Kim HG, Kim MJ. Calcium-binding proteins and gfap immunoreactivity alterations in murine hippocampus after 1 month of exposure to 835 mhz radiofrequency at sar values of 1.6 and 4.0 w/kg. *Neurosci Lett*. 2012;506:292-296
21. Maskey D, Kim M, Aryal B, Pradhan J, Choi IY, Park KS, et al. Effect of 835 mhz radiofrequency radiation exposure on calcium binding proteins in the hippocampus of the mouse brain. *Brain Res*. 2010;1313:232-241
22. Kittel A, Siklos L, Thuroczy G, Somosy Z. Qualitative enzyme histochemistry and microanalysis reveals changes in ultrastructural distribution of calcium and calcium-activated atpases after microwave irradiation of the medial habenula. *Acta Neuropathol*. 1996;92:362-368
23. Dutta SK, Das K, Ghosh B, Blackman CF. Dose dependence of acetylcholinesterase activity in neuroblastoma cells exposed to modulated radio-frequency electromagnetic radiation. *Bioelectromagnetics*. 1992;13:317-322
24. Palmieri L, Persico AM. Mitochondrial dysfunction in autism spectrum disorders: Cause or effect? *Biochim Biophys Acta*. 2010;1797:1130-1137
25. Peng TI, Jou MJ. Oxidative stress caused by mitochondrial calcium overload. *Ann N Y Acad Sci*. 2010;1201:183-188
26. Pessah IN, Lein PJ. Evidence for environmental susceptibility in autism: What we need to know about gene x environment interactions. *Humana*; 2008.
27. Stamou M, Streifel KM, Goines PE, Lein PJ. Neuronal connectivity as a convergent target of gene-environment interactions that confer risk for autism spectrum disorders. *Neurotoxicol Teratol*. 2012
28. Bachmann M, Lass J, Kalda J, Sakki M, Tomson R, Tuulik V, et al. Integration of differences in eeg analysis reveals changes in human eeg caused by microwave. *Conf Proc IEEE Eng Med Biol Soc*. 2006;1:1597-1600
29. Marino AA, Nilsen E, Frilot C. Nonlinear changes in brain electrical activity due to cell phone radiation. *Bioelectromagnetics*. 2003;24:339-346
30. Marino AA, Carrubba S. The effects of mobile-phone electromagnetic fields on brain electrical activity: A critical analysis of the literature. *Electromagn Biol Med*. 2009;28:250-274
31. Vecchio F, Babiloni C, Ferreri F, Curcio G, Fini R, Del Percio C, et al. Mobile phone emission modulates interhemispheric functional coupling of eeg alpha rhythms. *Eur J Neurosci*. 2007;25:1908-1913
32. Hountala CD, Maganioti AE, Papageorgiou CC, Nanou ED, Kyprianou MA, Tsiafakis VG, et al. The spectral power coherence of the eeg under different emf conditions. *Neurosci Lett*. 2008;441:188-192
33. Just MA, Cherkassky VL, Keller TA, Minshew NJ. Cortical activation and synchronization during sentence comprehension in high-functioning autism: Evidence of underconnectivity. *Brain*. 2004;127:1811-1821
34. Muller RA, Shih P, Keehn B, Deyoe JR, Leyden KM, Shukla DK. Underconnected, but how? A survey of functional connectivity mri studies in autism spectrum disorders. *Cereb Cortex*. 2011;21:2233-2243
35. Wass S. Distortions and disconnections: Disrupted brain connectivity in autism. *Brain Cogn*. 2011;75:18-28
36. Duffy FH, Als H. A stable pattern of eeg spectral coherence distinguishes children with autism from neuro-typical controls - a large case control study. *BMC Med*. 2012;10:64
37. Isler JR, Martien KM, Grieve PG, Stark RI, Herbert MR. Reduced functional connectivity in visual evoked potentials in children with autism spectrum disorder. *Clin Neurophysiol*. 2010
38. Murias M, Swanson JM, Srinivasan R. Functional connectivity of frontal cortex in healthy and adhd children reflected in eeg coherence. *Cereb Cortex*. 2007;17:1788-1799

39. Murias M, Webb SJ, Greenson J, Dawson G. Resting state cortical connectivity reflected in eeg coherence in individuals with autism. *Biol Psychiatry*. 2007;62:270-273
40. Coben R, Clarke AR, Hudspeth W, Barry RJ. Eeg power and coherence in autistic spectrum disorder. *Clin Neurophysiol*. 2008;119:1002-1009
41. Rossignol DA, Frye RE. Melatonin in autism spectrum disorders: A systematic review and meta-analysis. *Dev Med Child Neurol*. 2011;53:783-792
42. Buckley AW, Rodriguez AJ, Jennison K, Buckley J, Thurm A, Sato S, et al. Rapid eye movement sleep percentage in children with autism compared with children with developmental delay and typical development. *Arch Pediatr Adolesc Med*. 2010;164:1032-1037
43. Giannotti F, Cortesi F, Cerquiglini A, Vagnoni C, Valente D. Sleep in children with autism with and without autistic regression. *J Sleep Res*. 2011;20:338-347
44. Johansson O. Disturbance of the immune system by electromagnetic fields-a potentially underlying cause for cellular damage and tissue repair reduction which could lead to disease and impairment. *Pathophysiology*. 2009;16:157-177
45. Johansson O. Evidence for effects on immune function. *BioInitiative Report: A Rationale for a Biologically-based Public Exposure Standard for Electromagnetic Fields (ELF and RF)*. 2007;<http://bioinitiative.org/freeaccess/report/index.htm>
46. Bilbo SD, Jones JP, Parker W. Is autism a member of a family of diseases resulting from genetic/cultural mismatches? Implications for treatment and prevention. *Autism Res Treat*. 2012;2012:910946
47. Persico AM, Van de Water J, Pardo CA. Autism: Where genetics meets the immune system. *Autism Res Treat*. 2012;2012:486359
48. Kong SW, Collins CD, Shimizu-Motohashi Y, Holm IA, Campbell MG, Lee IH, et al. Characteristics and predictive value of blood transcriptome signature in males with autism spectrum disorders. *PLoS One*. 2012;7:e49475
49. Waly MI, Hornig M, Trivedi M, Hodgson N, Kini R, Ohta A, et al. Prenatal and postnatal epigenetic programming: Implications for gi, immune, and neuronal function in autism. *Autism Res Treat*. 2012;2012:190930
50. Lintas C, Sacco R, Persico AM. Genome-wide expression studies in autism spectrum disorder, rett syndrome, and down syndrome. *Neurobiol Dis*. 2012;45:57-68
51. Andrzejak R, Poreba R, Poreba M, Derkacz A, Skalik R, Gac P, et al. The influence of the call with a mobile phone on heart rate variability parameters in healthy volunteers. *Ind Health*. 2008;46:409-417
52. Szmigielski S, Bortkiewicz A, Gadzicka E, Zmyslony M, Kubacki R. Alteration of diurnal rhythms of blood pressure and heart rate to workers exposed to radiofrequency electromagnetic fields. *Blood Press Monit*. 1998;3:323-330
53. Bortkiewicz A, Gadzicka E, Zmyslony M, Szymczak W. Neurovegetative disturbances in workers exposed to 50 hz electromagnetic fields. *Int J Occup Med Environ Health*. 2006;19:53-60
54. Graham C, Cook MR, Sastre A, Gerkovich MM, Kavet R. Cardiac autonomic control mechanisms in power-frequency magnetic fields: A multistudy analysis. *Environ Health Perspect*. 2000;108:737-742
55. Saunders RD, Jefferys JG. A neurobiological basis for elf guidelines. *Health Phys*. 2007;92:596-603
56. Buchner K, Eger H. Changes of clinically important neurotransmitters under the influence of modulated rf fields—a long-term study under real-life conditions (translated; original study in german). *Umwelt-Medizin-Gesellschaft* 2011;24:44-57
57. Bellieni CV, Acampa M, Maffei M, Maffei S, Perrone S, Pinto I, et al. Electromagnetic fields produced by incubators influence heart rate variability in newborns. *Arch Dis Child Fetal Neonatal Ed*. 2008;93:F298-301
58. Narayanan A, White CA, Saklayen S, Scaduto MJ, Carpenter AL, Abduljalil A, et al. Effect of propranolol on functional connectivity in autism spectrum disorder—a pilot study. *Brain Imaging Behav*. 2010;4:189-197
59. Anderson CJ, Colombo J. Larger tonic pupil size in young children with autism spectrum disorder. *Dev Psychobiol*. 2009;51:207-211
60. Anderson CJ, Colombo J, Unruh KE. Pupil and salivary indicators of autonomic dysfunction in autism spectrum disorder. *Dev Psychobiol*. 2012

61. Daluwatte C, Miles JH, Christ SE, Beversdorf DQ, Takahashi TN, Yao G. Atypical pupillary light reflex and heart rate variability in children with autism spectrum disorder. *J Autism Dev Disord*. 2012
62. Ming X, Bain JM, Smith D, Brimacombe M, Gold von-Simson G, Axelrod FB. Assessing autonomic dysfunction symptoms in children: A pilot study. *J Child Neurol*. 2011;26:420-427
63. Hirstein W, Iversen P, Ramachandran VS. Autonomic responses of autistic children to people and objects. *Proc Biol Sci*. 2001;268:1883-1888
64. Toichi M, Kamio Y. Paradoxical autonomic response to mental tasks in autism. *J Autism Dev Disord*. 2003;33:417-426
65. Ming X, Julu PO, Brimacombe M, Connor S, Daniels ML. Reduced cardiac parasympathetic activity in children with autism. *Brain Dev*. 2005;27:509-516
66. Mathewson KJ, Drmic IE, Jetha MK, Bryson SE, Goldberg JO, Hall GB, et al. Behavioral and cardiac responses to emotional stroop in adults with autism spectrum disorders: Influence of medication. *Autism Res*. 2011;4:98-108
67. Cheshire WP. Highlights in clinical autonomic neuroscience: New insights into autonomic dysfunction in autism. *Auton Neurosci*. 2012;171:4-7
68. Chang MC, Parham LD, Blanche EI, Schell A, Chou CP, Dawson M, et al. Autonomic and behavioral responses of children with autism to auditory stimuli. *Am J Occup Ther*. 2012;66:567-576
69. Kesari KK, Kumar S, Behari J. 900-mhz microwave radiation promotes oxidation in rat brain. *Electromagn Biol Med*. 2011;30:219-234
70. Oktem F, Ozguner F, Mollaoglu H, Koyu A, Uz E. Oxidative damage in the kidney induced by 900-mhz-emitted mobile phone: Protection by melatonin. *Arch Med Res*. 2005;36:350-355
71. Lai H, Singh NP. Melatonin and a spin-trap compound block radiofrequency electromagnetic radiation-induced DNA strand breaks in rat brain cells. *Bioelectromagnetics*. 1997;18:446-454
72. Xu S, Zhou Z, Zhang L, Yu Z, Zhang W, Wang Y, et al. Exposure to 1800 mhz radiofrequency radiation induces oxidative damage to mitochondrial DNA in primary cultured neurons. *Brain Res*. 2010;1311:189-196



3 August 2016

Petaluma City Schools
District Office
200 Douglas Street
Petaluma, California 94952

Dear Sirs/Madams:

I am a public health physician who served as the Co-Editor of the Bioinitiative Report, published in 2007 as a comprehensive review of the adverse health effects of radiofrequency electromagnetic fields.

There is strong and consistent evidence that excessive exposure to radiofrequency electromagnetic fields has adverse human health effects. Of particular concern is the clear evidence that children are more vulnerable than adults. The best-documented adverse effects are an increase in risk of cancer, but cancers do not appear immediately upon exposure but rather come years later. The National Toxicology Program has within the past couple of months reported that even rats exposed to radiofrequency radiation develop brain cancer! Within a school setting there is increasing evidence that excessive exposures reduce learning ability, which is the last thing one wants in a school. Some children will also develop a syndrome of electrohypersensitivity, where they get headaches and reduced ability to pay attention and learn. While these effects are not nearly as well documented as those relating to cancer, they are particularly important within a school. This is especially the case in a wireless computer classroom, where exposure can be very high. However there will be essentially no exposure in a wired computer classroom.

The exposure levels of the Federal Communications Commission are totally outdated and do not protect the health of the public, especially of children. I urge you to abandon any plans for wireless communication within schools. It is of course critical that all children have access to the Internet, but when this is done through wired connections they will not be exposed to excessive electromagnetic fields.

Yours sincerely,

David O. Carpenter, M.D.
Director, Institute for Health and the Environment
University at Albany

District Office
200 Douglas Street
Petaluma, California
94952
USA

4 August, 2016

Dear Petaluma City Schools;
Superintendent Gary Callahan and Board of Trustees

Regarding: Wireless technology should not be used in schools or pre-schools due to health risks for children and employees

We have been asked to declare our opinion about wireless technology in schools by parents that are concerned about their children.

Based on current published scientific studies, we urge your administration to educate themselves on the potential risks from wireless technologies in schools, and to choose wired teaching technologies. The well-being and educational potential of children depends on it.

High-speed connectivity to schools is important but it can be a wired connection instead of Wi-Fi. Wireless classroom infrastructure and wireless devices for schoolchildren should be avoided for these reasons:

- Wireless radiofrequency (RF) radiation emissions were classified as a Possible Human Carcinogen (group 2B) by the World Health Organization International Agency for Research on Cancer (IARC) in May 2011. One of the signers, Dr Hardell, was part of the evaluation group.
- The IARC classification holds for *all forms of radio frequency radiation* including RF-EMF emissions from wireless transmitters (access points), tablets and laptops.
- Epidemiological studies show links between RF radiation exposure and cancer, neurological disorders, hormonal changes, symptoms of electrical hypersensitivity (EHS) and more. Laboratory studies show that RF radiation exposure increases risk of cancer, abnormal sperm, learning and memory deficits, and heart irregularities. Foetal exposures in both animal and human studies may result in altered brain development in the young offspring, with disruption in learning, memory and behaviour.
- Recently a report was released from The National Toxicology Program (NTP) under the National Institutes of Health (NIH) in USA on the largest ever animal study on cell phone RF radiation and cancer (<http://biorxiv.org/content/biorxiv/early/2016/05/26/055699.full.pdf>). An increased incidence of glioma and malignant schwannoma in the heart was found. Interestingly our research group and others have in epidemiological studies shown that persons using wireless phones (both mobile phones and cordless phones; DECT) have an increased risk for glioma and acoustic neuroma. Acoustic neuroma or vestibular schwannoma is the same type of tumour as the one found in the heart, although benign.
- The research showing increased brain cancer risk in humans *has strengthened* since the IARC 2011 classification as new research has been published which repeatedly shows a significant association after RF radiation exposure. In addition, tumour

promotion studies have now been replicated showing cancer promotion after exposures at low levels.

- It is our opinion and that of many colleagues that the current IARC cancer risk classification should move to an *even higher* risk group. The carcinogenic effect has been shown in human and animal studies. Several laboratory studies have shown mechanistic effects in carcinogenesis such as oxidative stress, down regulation of mRNA, DNA damage with single strand breaks.
- In summary RF radiation should be classified as Carcinogenic to Humans, Group 1 according to the IARC classification. This classification should have a major impact on prevention.

The evidence for these statements is based on hundreds of published, peer-reviewed scientific studies that report adverse health effects at levels much lower than current ICNIRP and FCC public safety limits. Compliance with government regulations does not mean that the school wireless environment is safe for children and staff (especially pregnant staff).

As researchers in cancer epidemiology and RF radiation exposures, we have published extensively in this area and it is our opinion that schools should choose wired Internet connections. Multiple epidemiological research studies show that exposures equivalent to 30 minutes a day of cell phone use over ten years results in a significantly increased brain cancer risk.

What will be the health effect for a child exposed all day long in school for 12 years? Wireless networks in schools result in full body low level RF radiation exposures that can have a cumulative effect on the developing body of a child. No safe level of this radiation has been determined by any health agency and therefore we have no safety assurances. Cancers can have long latency periods (time from first exposure until diagnosis) and it will take decades before we know the full extent of health impacts from this radiation. The statistics and effects will be borne by the children you serve.

Wi-Fi in schools, in contrast to wired Internet connections, will increase risk of neurologic impairment and long-term risk of cancer in students. Promoting wireless technology in schools disregards the current health warnings from international science and public health experts in this field.

We recommend that your school district install wired Internet connections and develop curriculum that teaches students at all ages safer ways to use their technology devices. If cell phones and other wireless devices are used in the school curriculum (as many schools are now doing with Bring your Own Device Policy) then there should be educational curriculum in place and well posted instructions in classrooms so that the students and staff use these devices in ways that reduce exposure to the radiation as much as possible.

Supporting wired educational technologies is the safe solution in contrast to potentially hazardous exposures from wireless radiation.

Respectfully submitted

Lennart Hardell, MD, PhD
Department of Oncology,
Örebro University Hospital,

Michael Carlberg, MSc
Department of Oncology,
Örebro University Hospital,

SE-701 85 Örebro, Sweden
E-mail: lennart.hardell@regionorebrolan.se

SE-701 85 Örebro, Sweden
michael.carlberg@regionorebrolan.se

Lena Hedendahl, MD
Östra Skolgatan 12,
SE-972 53 Luleå, Sweden
E-mail: lenahedendahl@telia.com

References and additional reading:

Coureau G, Bouvier G, Lebaillly P, Fabbro-Peray P, Gruber A, Leffondre K, Guillamo JS, Loiseau H, Mathoulin-Pélissier S, Salamon R, Baldi I. **Mobile phone use and brain tumours in the CERENAT case-control study.** Occup Environ Med. 2014;71(7):514-22.

Carlberg M, Hardell L. **Decreased survival of glioma patients with astrocytoma grade IV (glioblastoma multiforme) associated with long-term use of mobile and cordless phones.** Int J Environ Res Public Health. 2014;11(10):10790-805.

Carlberg M, Hedendahl L, Ahonen M, Koppel T, Hardell L. **Increasing incidence of thyroid cancer in the Nordic countries with main focus on Swedish data.** BMC Cancer. 2016 Jul 7;16:426. doi: 10.1186/s12885-016-2429-4.

Hardell L, Carlberg M. **Mobile phone and cordless phone use and the risk for glioma - Analysis of pooled case-control studies in Sweden, 1997-2003 and 2007-2009.** Pathophysiology. 2015;22(1):1-13.

Hardell L, Carlberg M, Söderqvist F, Hansson Mild K. **Case-control study of the association between malignant brain tumours diagnosed between 2007 and 2009 and mobile and cordless phone use.** Int J Oncol. 2013;43(6):1833-45.

Hardell L, Carlberg M, Hansson Mild K. **Pooled analysis of two case-control studies on the use of cellular and cordless telephones and the risk of benign brain tumours diagnosed during 1997-2003.** Int J Oncol. 2006;28(2):509-18.

Hardell L, Carlberg M, Söderqvist F, Hansson Mild K. **Pooled analysis of case-control studies on acoustic neuroma diagnosed 1997-2003 and 2007-2009 and use of mobile and cordless phones.** Int J Oncol. 2013;43(4): 1036-44.

Hardell L, Carlberg M, Hansson Mild K. **Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects.** Int J Oncol. 2011;38(5):1465-74.

Hardell L, Carlberg M. **Using the Hill viewpoints from 1965 for evaluating strengths of evidence of the risk for brain tumors associated with use of mobile and cordless phones.** Rev Environ Health. 2013;28(2-3):97-106.

Hardell L, Carlberg M, Hansson Mild K. **Use of mobile phones and cordless phones is associated with increased risk for glioma and acoustic neuroma.** Pathophysiology. 2013;20(2):85-110.

Hedendahl L, Carlberg M, Hardell L. **Electromagnetic hypersensitivity - an increasing challenge to the medical profession.** Rev Environ Health 2015;30:209-215.

International Agency for Research on Cancer (IARC). **Non-ionizing radiation, Part II: Radiofrequency electromagnetic fields.** IARC Monogr Eval Carcinog Risks Hum. 2011;102(2):1-460.

Lerchl A, Klose M, Grote K, Wilhelm AF, Spathmann O, Fiedler T, Streckert J, Hansen V, Clemens M. **Tumor promotion by exposure to radiofrequency electromagnetic fields below exposure limits for humans.** Biochem Biophys Res Commun. 2015;459(4):585-90.

Wyde M, Cesta M, Blystone C, Elmore S, Foster P, Hooth M, Kissling G, Malarkey D, Sills R, Stout M, *et al*: **Report of Partial Findings from the National Toxicology Program Carcinogenesis Studies of Cell Phone Radiofrequency Radiation in Hsd: Sprague Dawley® SD rats (Whole Body Exposures).** Draft 5-19-2016. US National Toxicology Program (NTP), 2016. doi: <http://dx.doi.org/10.1101/055699>. Available online: <http://biorxiv.org/content/biorxiv/early/2016/05/26/055699.full.pdf>

Yakymenko I, Tsybulin O, Sidorik E, Henshel D, Kyrylenko O, Kyrylenko S. **Oxidative mechanisms of biological activity of low-intensity radiofrequency radiation.** Electromagn Biol Med. 2015;19:1-16.

Dr. Anthony B. Miller

3800 Yonge Street, Suite 406,
Toronto, ON, M4N 3P7
Telephone 416 487 5825
Email: ab.miller@sympatico.ca

August 4, 2016

Petaluma City Schools
District Office
200 Douglas Street
Petaluma, California
94952

Re: Adverse Effects of Radiofrequency fields

I am writing to express my concern over the increasing exposure of children in schools to Radiofrequency Fields (e.g. from wi-fi, as required for cell phones and iPads, and emitted by cell towers) and the lack of concern expressed by many councils, governments and School Boards on this issue. In particular, justification for the “safety” of radiofrequency fields is placed upon the use of outdated safety standards, based upon tissue heating, whereas it has now been well demonstrated that adverse biological effects occur at far lower levels of radiofrequency fields that do not induce tissue heating, including a recent animal study performed by the National Toxicology Program in the United States which found an increased incidence of brain cancers and other cancers in rats exposed to prolonged Radiofrequency fields.

I am a physician and epidemiologist specializing in cancer etiology, prevention, and screening, expert in epidemiology, and particularly causes of human cancer. I have performed research on ionizing radiation and cancer, electromagnetic fields and cancer, and have served on many committees assessing the carcinogenicity of various exposures, including working groups of the International Agency for Research on Cancer (IARC), widely regarded as providing unbiased assessment on the carcinogenicity of chemicals and other exposure to humans.

In 2011, an IARC working group designated radiofrequency fields as a class 2B carcinogen, a possible human carcinogen. Since that review a number of additional studies have been reported. One of the most important was a large case-control study in France, which found a doubling of risk of glioma, the most malignant form of brain cancer, after two years of exposure to cell phones. After five years exposure the risk was five-fold. They also found that in those who lived in urban environments the risk was even higher. In my view, and that of many colleagues who have written papers on this issue, these studies provide evidence that radiofrequency fields are not just a possible human carcinogen but a probable human carcinogen, i.e. IARC category 2A. It would be impossible to ignore such an assessment in regulatory approaches.

It is important to recognize that there are no safe levels of exposure to human carcinogens. Risk increases with increasing intensity of exposure, and for many carcinogens, even more with increasing duration of exposure. The only way to avoid the carcinogenic risk is to avoid exposure altogether. This is why we ban known carcinogens from the environment and why much effort is taken to get people, particularly young people, not to smoke. We now recognize that exposure to carcinogens in childhood can increase the risk of cancer in adulthood many years later. Further, people vary in their genetic makeup, and certain genes can make some people more susceptible than others to the effect of carcinogens. It is the young and those who are susceptible we should protect.

As an epidemiologist who has done a great deal of work on breast cancer, I have been concerned by a series of case reports from California and elsewhere of women who developed unusual breast cancers in the exact position where they kept cell phones in their bras. These are unusual cancers. They are multifocal, mirroring where the cell phone was kept. Thus in these relatively young women the radiofrequency radiation from very close contact with a cell phone has caused breast cancer.

Not only brain and breast cancers but parotid gland tumors, tumors of the salivary gland, have been associated with prolonged exposure to cell phones.

Given the long natural history of cancer and the fact that human populations have not been exposed for a sufficient length of time to reveal the full adverse effects of radiofrequency fields, it is extremely important to adopt a precautionary approach to the exposure of humans to such fields. An individual, if appropriately informed, can reduce her or his exposure to radiofrequency fields from devices that use wi-fi, but in the case of cell towers, smart meters and wi-fi in schools, the exposure they receive is outside their control. Then, with the people who manufacture these devices and those who promote wi-fi failing to issue adequate health warnings, we are reaching a situation where schools, work places and homes are being saturated with radiofrequency fields.

Thus to avoid a potential epidemic of cancer caused by radiofrequency fields from wi-fi and other devices, we should introduce means to reduce exposure as much as reasonably achievable, use hard wire connections to the internet and strengthen the codes that are meant to protect the public.

Yours sincerely

A handwritten signature in black ink, appearing to read 'A. B. Miller', with a stylized, cursive script.

Anthony B. Miller, MD, FRCP(C), FRCP, FACE
Professor Emeritus
Dalla Lana School of Public Health, University of Toronto, Ontario, Canada



Karolinska Institutet

Department of Neuroscience
Experimental Dermatology Unit

Stockholm, December 8, 2015

To:

MCPS CEO Dr. Andrew Zuckerman [Andrew_Zuckerman@mcpsmd.org]
MCPS Superintendent Mr. Larry Bowers [Larry_Bowers@mcpsmd.org]
MCPS Chief Technology Officer Mr. Sherwin Collette [Sherwin_Collette@mcpsmd.org]
MCPS Board of Education [boe@mcpsmd.org]
840 Hungerford Drive
Rockville, MD 20850, USA

cc:

Montgomery County Council [county.council@montgomerycountymd.gov]

Dear Madame or Sir,

My name is Olle Johansson, and I am an associate professor, heading the Experimental Dermatology Unit at Sweden's Karolinska Institute in the Department of Neuroscience. I understand you have recently made public pronouncements regarding the safety of Wi-Fi. As a neuroscientist who has been studying the biophysical and epidemiological effects of electromagnetic fields (EMFs) for over 30 years, I believe this designation is short-sighted.

Wireless communication is now being implemented in our daily life in a very fast way. At the same time, it is becoming more and more obvious that the exposure to electromagnetic fields not only may induce acute thermal effects to living organisms, but also non-thermal effects, the latter often after longer exposures. This has been demonstrated in a very large number of **non-ionizing radiation** studies and includes cellular DNA-damage, disruptions and alterations of cellular functions like increases in intracellular stimulatory pathways and calcium handling, disruption of tissue structures like the blood-brain barrier, impact on vessel and immune functions, and loss of fertility. Whereas scientists can observe and reproduce these effects in controlled laboratory experiments, epidemiological and ecological data derived from long-term exposures in well-designed case-control studies reflect this link all the way from molecular and cellular effects to the living organism up to the induction and proliferation of diseases observed in humans. It should be noted that we are not the only species at jeopardy; practically all animals, plants and bacteria may be at stake. Although epidemiological and ecological investigations as such never demonstrate causative effects, due to the vast number of confounders, they confirm the relevance of the controlled observations in the laboratories.

Many times since the early 1980s I have pointed out that the public's usage of cell phones has become the largest full-scale biological and medical experiment ever with mankind, and I was also the first person to firmly point out that this involuntary exposure violates the Nuremberg Code's principles for human experimentation, which clearly states that voluntary

Mailing address
Experimental Dermatology Unit
Department of Neuroscience
Karolinska Institutet
171 77 Stockholm
Sweden

Visiting address
Retziuslaboratoriet
Retzius väg 8
Solna

Direct	468-52 48 70 58
Switchboard	468-52 48 64 00
Fax	468-30 39 04
Fax (KI)	468-31 11 01



consent of human subjects is absolutely essential. Among many effects seen, the very serious one is the deterioration of the genome. Such an effect - if seen in a food item under development or in a potential pharmaceutical drug - immediately would completely ban it from further marketing and sale; genotoxic effects are not to be allowed or spread. For these reasons above, we, scientists, can not accept that children undergo an enormous health risk for their present and future, by being exposed to WI-FI in kindergardens or schools (even if the WI-FI masts/routers are not in the children's classroom). The precautionary principle has to be respected. Furthermore, when men place cell phones in their front pocket, or laptops on their laps, it should be noted that experimental studies have demonstrated that after similar exposures there is a decrease in sperm count as well as in the quality of sperm, which is a phenomenon that could affect society's overall ability to procreate in the future. Experiments in mice point to that it may be true already in 5 generations time.

Many other states including France, Russia, Israel and Germany, have employed various precautionary steps and their responses (including labelling cell phones and other transmitting devices with SAR ratings, discouraging the use of cell phones and other wireless gadgets by children, warning parents of the risks, and removing or restricting WiFi in schools and replacing it with hard-wired ethernet) as a result of the *WHO/IARC classification of radiofrequency electromagnetic radiation in 2011 as a Class 2B carcinogen as well as the earlier classification of power-frequent magnetic fields in 2001 also as a Class 2B carcinogen*, the information summarized in the Bioinitiative Reports of 2007 and 2012, and the other considerable international and independent research and reviews, that show adverse biological effects from electromagnetic fields, including heart palpitations, headaches, skin rashes, damage to DNA, mental health effects, impaired concentration, decreased problem-solving capacity, electrohypersensitivity, etc., are about to set a new standard for educational quality with due respect to children's and staff's health.

In the case of "protection from exposure to electromagnetic fields", it is thus of paramount importance to act from a prudence avoidance/precautionary principle point of view. Anything else would be highly hazardous. Total transparency of information is the key sentence here, as I believe the public does not appreciate having the complete truth revealed years after a certain catastrophe already has taken place. For instance, it shall be noted, that today's recommended values for wireless systems, such as the SAR-values, are just recommendations, and not safety levels. Since scientists observe biological effects at as low as 20 microWatts/kg, can it truly be stated that it is safe to allow irradiation of humans at SAR 2 W/kg, or at 100,000 times stronger levels of radiation?

IMBALANCED REPORTING

Another misunderstanding is the use of scientific publications (as the tobacco industry did for many years) as 'weights' to balance each other. But one can NEVER balance a report showing a negative health effect with one showing no effect. This is a misunderstanding which, unfortunately, is very often used both by the industrial representatives as well as official authorities to the detriment of the general public. True balance would be reports showing negative health effects against *exact replications* showing no or positive effects. However, this is not what the public has been led to believe.



NEED FOR INDEPENDENT RESEARCH

In many commentaries, debate articles and public lectures - for the last 20-30 years – I have urged that completely independent research projects must be inaugurated immediately to ensure our public health. These projects must be entirely independent of all types of commercial interests; public health can not have a price-tag! It is also of paramount importance that scientists involved in such projects must be free of any carrier considerations and that the funding needed is covered to 100%, not 99% or less. This is the clear responsibility of the democratically elected body of every country.

WHO/INTERNATIONAL AGENCY FOR RESEARCH ON CANCER (IARC), 2011

Very recently (in Lyon, France, May 31, 2011) the WHO/International Agency for Research on Cancer (IARC) has classified radiofrequency electromagnetic fields as possibly carcinogenic to humans (Group 2B), based on an increased risk for glioma, a malignant type of brain cancer. This should be added to the previous (2001) 2B classification of power-frequent (ELF) electromagnetic fields – emitted at high levels from handheld gadgets, such as eReaders and mobile phones – as a risk factor for childhood leukemia. Given the 2001 very close votes (9 to 11) for moving it to 2A and all the new knowledge that has accumulated since 2001, today the association between childhood leukemia and power-frequent (ELF) electromagnetic fields would definitely be signed into the much more serious 2A (“probably carcinogenic”) category. So, the ‘red flag’ is – unfortunately – flying very high.

INVOLUNTARY EXPOSURE

According to Article 24 of the UNICEF’s Child Convention “children have the right to ... a clean and safe environment, and information to help them stay healthy”. We must all ensure that this article never is violated. This is about our social responsibility, and is very much a public health issue.

In summary, electromagnetic fields may be among the most serious and overlooked health issues today, and having these fields checked and reduced/removed from schools and kindergardens may be essential for health protection and restoration, and is a must for persons with the functional impairment electrohypersensitivity as for children who are more fragile (cf. Belyaev I, Dean A, Eger H, Hubmann G, Jandrisovits R, Johansson O, Kern M, Kundi M, Lercher P, Mosgöller W, Moshhammer H, Müller K, Oberfeld G, Ohnsorge P, Pelzmann P, Scheingraber C, Thill R, "EUROPAEM EMF Guideline 2015 for the prevention, diagnosis and treatment of EMF-related health problems and illnesses", Rev Environ Health 2015; 30: 337–371). In addition, as recently discussed in a think-tank group here in Stockholm, it is very important to constantly educate oneself and participate in the general debate and public discussions to keep the information build-up active. Thus, it is of paramount importance to keep the "kettle boiling", never blindly trusting or accepting given 'facts', but only read and think for yourself and for your loved ones. Only so you can arrive at a genuinely working precautionary principle.



Karolinska Institutet
Department of Neuroscience
Experimental Dermatology Unit

CONCLUSION

In conclusion, wireless systems, such as Wi-Fi routers or cell towers, and their electromagnetic fields, can not be regarded as safe in schools, but must be deemed highly hazardous and unsafe for the children as well as for the staff.

I encourage governments and local health and educational bodies to adopt a framework of guidelines for public and occupational EMF exposure that reflect the Precautionary Principle. As noted, the Precautionary Principle states when there are indications of possible adverse effects, though they remain uncertain, the risks from doing nothing may be far greater than the risks of taking action to control these exposures. The Precautionary Principle shifts the burden of proof from those suspecting a risk to those who discount it — as some nations have already done. Precautionary strategies should be based on design and performance standards and may not necessarily define numerical thresholds because such thresholds may erroneously be interpreted as levels below which no adverse effect can occur.

Some 100 years back, we learned the hard lessons of ionizing radiation and the need for strict health protections – now we must openly face the possibility that we must take a seat in life's school and learn again. This time it is about non-ionizing radiation.

Based on all of the above, I strongly urge you to reconsider your public stance on the safety of Wi-Fi, cell towers, and similar systems in schools as their non-ionizing radiation emissions very likely are hazardous and unsafe for students, staff and teachers.

With my very best regards
Yours sincerely
Olle Johansson

(Olle Johansson, associate professor
The Experimental Dermatology Unit
Department of Neuroscience
Karolinska Institute
171 77 Stockholm
Sweden)

Mailing address
Experimental Dermatology Unit
Department of Neuroscience
Karolinska Institutet
171 77 Stockholm
Sweden

Visiting address
Retziuslaboratoriet
Retzius väg 8
Solna

Direct	468-52 48 70 58
Switchboard	468-52 48 64 00
Fax	468-30 39 04
Fax (KI)	468-31 11 01

Martin L. Pall, Professor Emeritus
Biochemistry and Basic Medical Sciences,
Washington State University,
638 NE 41st Ave., Portland, OR
97232-3312
503-232-3883
martin_pall@wsu.edu

MCPS COO Dr. Andrew Zuckerman
MCPS Interim Superintendent Larry Bowers
MCPS Board of Education
MCPS Office of Technology
Montgomery County Schools
Carver Educational Services Center
850 Hungerford Drive
Rockville, MD 20850

January 3, 2016

Dear Montgomery County COO Dr. Andrew Zuckerman, Interim Superintendent Larry Bowers,
Board of Education and Office of Technology;

I have been asked to comment on the [MCPS Statement Concerning Deployment of Wireless Computing Technologies](#). I am happy to do so.

The first paragraph in that statement is not relevant to the issue at hand because it is perfectly possible to use wired communication for such education. This document is being produced on a computer on which I only use wired communication, connecting to the internet, connecting to my printer and for other purposes, as well.

The 2nd and 3rd paragraphs of your statement may well be technically correct. However these give us no assurance whatsoever of safety of Wi-Fi fields. The FCC guidelines as are many other such guidelines, are based on the assumption that only heating effects of microwave/lower frequency EMFs can have biological effects. However that assumption has been falsified by thousands of studies published from the 1950s to the present, each showing that non-thermal levels of exposure often produce biological effects. For example, in 1971, the U.S. Office of Naval Medical Research produced a document reporting over 100 different non-thermal effects [1], listing 40 apparent neuropsychiatric changes produced by non-thermal microwave frequency exposures, including 5 central/peripheral nervous system (NS) changes, 9 central NS effects, 4 autonomic system effects, 17 psychological disorders, 4 behavioral changes and 2 misc. effects [1]. It also listed cardiac effects including ECG changes and cardiac necrosis as well as both hypotension and hypertension, and also 8 different endocrine effects.

Changes affecting fertility included tubular degeneration in the testis, decreased spermatogenesis, altered sex ratio, altered menstrual activity, altered fetal development, programmed cell death (what is now known as apoptosis) and decreased lactation. Many other non-thermal changes were also listed for a total of over 100 non-thermal effects. They also provided [1] approximately 2000 citations documenting these various health effects. That was almost 45 years ago and is only the beginning of the evidence for the existence of non-thermal effects. My own recent paper [2] shows that widespread neuropsychiatric effects are caused by non-thermal exposures to many different microwave frequency electromagnetic fields (EMFs).

Tolgskaya and Gordon [3] in 1973 published a long and detailed review of effects of microwave and lower frequency EMFs on experimental animals, mostly rodents. They report that non-thermal exposures impact many tissues, with the nervous system being the most sensitive organ in the body, based on histological studies, followed by the heart and the testis. They also report effects of non-thermal exposures on liver, kidney, endocrine and many other organs. The nervous system effects are very extensive and include changes many changes in cell structure, disfunction of synaptic connections between neurons and programmed cell death and are discussed in Refs. [2,3] and more modern studies reporting extensive effects of such non-thermal EMF exposures on the brain are also cited in [2]. There are also many modern studies showing effects of non-thermal exposures on fertility in animals.

The Raines 1981 National Aeronautics and Space Administration (NASA) report [4] reviewed an extensive literature based on occupational exposures to non-thermal microwave EMFs. Based on multiple studies, Raines [4] reports that 19 neuropsychiatric effects are associated with occupational microwave/ radiofrequency EMFs, as well as cardiac effects, endocrine including neuroendocrine effects and several other effects.

I reviewed many other scientific reviews on this topic, each of which clearly supports the view that there are various non-thermal health impacts of these EMFs [5]. In 2015, 206 international scientists signed [a statement](#) sent to the United Nations Secretary General and to member states, stating that international safety guidelines and standards are inadequate to protect human health [6]. Each of these 206 scientists from 40 countries had scientific publications on biological effects of such EMFs and therefore each is well qualified to judge this. ***It can be seen from this statement to the UN, that there is a strong scientific consensus that current safety guidelines and standards are inadequate because they do not take into consideration all of the non-thermal health effects produced by various EMF exposures.***

That scientific consensus also rejects, therefore, the FCC EMF guidelines, guidelines that cannot be defended despite your own attempt to do so in MCPS Statement Concerning Deployment of Wireless Computing Technologies.

It can be seen from the previous paragraphs, that the following non-thermal effects of EMF exposures are well documented:

- Ø Widespread neuropsychiatric effects
- Ø Several types of endocrine (that is hormonal) effects
- Ø Cardiac effects impacting the electrocardiogram (Note: these are often associated with occurrence of sudden cardiac death)
- Ø Male infertility

However, there are many additional types of biological changes produced by non-thermal EMF exposures (reviewed in 5,7] including:

- Ø Oxidative stress
- Ø Changes in calcium fluxes and calcium signaling
- Ø Several types of DNA damage to the cells of the body, including single strand and double strand DNA breaks and 8-OH-guanine in DNA
- Ø Cancer (which is undoubtedly caused, in part, by such DNA damage)
- Ø Female infertility
- Ø Lowered melatonin; sleep disruption
- Ø Therapeutic effects of EMFs when they are highly controlled and focused on a specific part of the body

It can be seen from the above, that each of the things that we most value as individuals and as a species are being attacked by non-thermal microwave frequency EMFs [5.7]:

§ **Our Health**

§ **Our brain function**

§ **The integrity of our genomes**

§ **Our ability to produce healthy offspring**

I want to emphasize that the specific health effects listed above are **not** the only things that are likely to be impacted by non-thermal EMF exposures, they are however the best documented such effects.

While it has been clear for many years that there are many non-thermal health effects of microwave frequency EMFs, it has not been clear until about 2 ½ years ago, how these effects are produced by such exposures. I stumbled onto the mechanism in 2012 and published on it in mid-2013. This 2013 paper [8] was honored by being placed on the Global Medical Discovery web site as one of the most important medical papers of 2013. At this writing, it has been cited 61 times according to the Google Scholar database, with over 2/3rds of those citations during 2015. So clearly it is having a substantial and rapidly increasing impact on the scientific literature. I have given 26 professional talks, in part or in whole on EMF effects in 10 different countries over the last 2 1/4 years. So it is clear that there has been a tremendous amount of interest in this research.

What the 2013 study showed [8], was that in 24 different studies (and there are now 2 more that can now be added [2]), effects of low-intensity EMFs, both microwave frequency and lower frequency EMFs could be blocked by calcium channel blockers, drugs that block what are called voltage-gated calcium channels (VGCCs). There were a total of 5 different types of calcium

channel blocker drugs used in these studies, with each type acting on a different site on the VGCCs and each thought to be highly specific for blocking VGCCs. What these studies tell us is that these EMFs act to produce non-thermal effects by activating the VGCCs. Where several effects were studied, when one of them was blocked or greatly lowered, each other effect studied was also blocked or greatly lowered. This tells us that the role of VGCC activation is quite wide – many effects go through that mechanism, possibly even all non-thermal effects in mammals. There are a number of other types of evidence confirming this mechanism of action of microwave frequency EMFs [2,]. Each of the 11 health impacts caused by non-thermal EMF exposures can be explained as being produced by indirect effects of VGCC activation [5,7].

It is now apparent [7] that these EMFs act directly on the voltage sensor of the VGCCs, the part of the VGCC protein that detects electrical changes and can open the channel in response to electrical changes. The voltage sensor (and this is shown on pp. 102-104 in [7]) is predicted, because of its structure and its location in the plasma membrane of the cell, to be extraordinarily sensitive to activation by these EMFs, about 7.2 million times more sensitive than are single charged groups elsewhere in the cell. What this means is that arguments that EMFs produced by particular devices are too weak to produce biological effects, are immediately highly suspect because the actual target, the voltage sensor of the VGCCs is extremely sensitive to these EMFs. **Because heating is mostly produced by forces on these singly charged groups elsewhere in the cell, limiting safety guidelines to heating effects means that these guideline allow exposures that are something like 7.2 million times too high.**

Why then does the FCC stick with these totally unscientific safety guidelines? That is the 64 billion dollar question. The FCC has been shown, in a long detailed document published by Harvard University Center for Ethics, to be a “captured agency”, that is captured by the telecommunications industry that the FCC is supposed to be regulating [9; can be obtained full text from web site listed in 9]. So perhaps the failure of the FCC to follow the extensive science in this important area, can be understood. Of course, what that means is that the FCC is completely failing in its role of protecting the public and it is a major blunder, therefore for either you or any other organization to depend on the FCC guideline as a reliable predictor of impacts of EMFs in humans.

So what is known about health impacts of Wi-Fi EMFs?

Table 1. The following Table summarizes various health impacts of Wi-Fi EMF exposures:

Citation(s)	Health Effects
[10,11,12,13,14,15,16]	Sperm/testicular damage, male infertility
[10,15,17,18,19,20]	Oxidative stress
[20]	Calcium overload

[11,12,20]	Apoptosis (programmed cell death)
[17]	Melatonin lowering; sleep disruption
[10,13]	Cellular DNA damage
[21]	MicroRNA expression (brain)
[18]	Disrupts development of teeth
[22]	Cardiac changes, blood pressure disruption; erythrocyte damage; catecholamine elevation
[23,24]	Neuropsych changes including EEG
[25]	Growth stimulation of adipose stem cells (role in obesity?)

Each of the effects reported above in 2 to 7 studies have an extensive literature for their occurring in response to various other microwave frequency EMFs so it should be clear that these observations on Wi-Fi exposures are highly probable to be correct. These include (see Table 1) findings that Wi-Fi exposures produce impacts on the testes leading to lowered male fertility; oxidative stress; intracellular calcium overload; apoptosis (a process that has an important causal role in neurodegenerative diseases); cellular DNA damage; neuropsychiatric changes including EEG changes. Each of these are very serious and oxidative stress has causal roles in many different human diseases; intracellular calcium overload has many different consequences – for example, it has a central role in causing neurodegenerative diseases; cellular DNA damage can cause cancer and produce mutations that impact future generations (if there are any). Other Wi-Fi effects each only documented by a single study are also effects where a variety of other non-thermal microwave EMFs also cause these, as shown by extensive literature on each of them. These include: melatonin lowering and sleep disruption; and the effects reported by Sali et al [22] cardiac changes, blood pressure disruption; erythrocyte damage; catecholamine elevation. So these may well be correct observations as well despite having only a single Wi-Fi specific study for each.

Summary:

1. The EMF safety guidelines supported by the FCC and others assume that only heating effects need be of concern. These assumptions have been known to be false for at least 45 years and there is a scientific consensus on this, that has lead to the petition by 206 highly qualified international scientists to the UN stating that current safety guidelines are inadequate.
2. We now know that low intensity non-thermal exposures work via VGCC activation and that indirect effects of such VGCC activation can produce each of the health effects that have been widely reported to occur in response to such EMF exposures for something like 60 years.

These attack:

a. Our health

- b. Our brain function**
- c. The integrity of our genomes**
- d. Our ability to produce healthy offspring**

3. The voltage sensor of the VGCCs is stunningly sensitive to such low intensity EMFs, about 7.2 million times more sensitive than are singly charge groups elsewhere in our cells. The consequence of this is that safety guidelines allow exposures that are very roughly 7.2 million times too high.

4. The FCC has been shown, in a detailed Harvard University study, to be a Captured Agency, captured by the industry that it is supposed to be regulating. This provides an additional reason to be very highly skeptical about all FCC safety guidelines.

5. 15 studies have each shown health effects of Wi-Fi, most of which have also been shown to occur in response to low intensity exposures to other types of microwave frequency EMFs. These are likely to have massive health effects by producing male infertility (female infertility has not been studied in response to Wi-Fi), oxidative stress (involved in dozens of human diseases), cellular DNA damage (possibly leading to both cancer and mutations in future generations), life threatening cardiac effects, cellular apoptosis and also intracellular calcium overload (with both of these possibly leading to neurodegenerative diseases), various neuropsychiatric changes and many others.

It is my view that it is sheer insanity to fail to see the threat to our and to all human civilization by continuing to ignore the threats from such EMFs, starting with Wi-Fi.

Martin L. Pall, Professor Emeritus
Biochemistry and Basic Medical Sciences,
Washington State University,
martin_pall@wsu.edu

Literature cited:

[1] Naval Medical Research Institute Research Report, June 1971. Bibliography of Reported Biological Phenomena ("Effects") and Clinical Manifestations Attributed to Microwave and Radio-Frequency Radiation. Report No. 2 Revised.

[2] Pall ML. 2015. Microwave frequency electromagnetic fields (EMFs) produce widespread neuropsychiatric effects including depression. J. Chem. Neuroanat. 2015 Aug 20. pii: S0891-0618(15)00059-9.doi: 10.1016/j.jchemneu.2015.08.001. [Epub ahead of print] Review.

[3] Tolgskaya MS, Gordon ZV. 1973. Pathological Effects of Radio Waves, Translated from Russian by B Haigh. Consultants Bureau, New York/London, 146 pages.

[4] Raines JK. 1981. Electromagnetic Field Interactions with the Human Body: Observed Effects and Theories. Greenbelt, Maryland: National Aeronautics and Space Administration 1981; 116 p.

- [5] Pall ML. 2015. How to approach the challenge of minimizing non-thermal health effects of microwave radiation from electrical devices. *Int J Innovative Research Engineering Management (IJIREM)* ISSN: 2350-0557, Volume-2, Issue -5, September 2015; 71-76.
- [6] <https://emfscientist.org/index.php/emf-scientist-appeal>
- [7] Pall ML. 2015 Scientific evidence contradicts findings and assumptions of Canadian Safety Panel 6: microwaves act through voltage-gated calcium channel activation to induce biological impacts at non-thermal levels, supporting a paradigm shift for microwave/lower frequency electromagnetic field action. *Rev Environ Health* 30:99-116.
- [8] Pall ML. 2013 Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects. *J Cell Mol Med* 17:958-65.
- [9] Alster N. 2015 Captured Agency: How the Federal Communications Commission Is Dominated by the Industries It Presumably Regulates. Edmond J. Safra Center for Ethics Harvard University 124 Mount Auburn Street, Suite 520N Cambridge, MA 02138 USA http://www.ethics.harvard.edu/http://www.ethics.harvard.edu/files/center-for-ethics/files/capturedagency_alster.pdf
- [10] Atasoy HI, Gunal MY, Atasoy P, Elgund S, Bugdayci G. 2013 Immunopathologic demonstration of deleterious effects on growing rat testes of radiofrequency waves emitted from conventional Wi-Fi devices. *J Pediatr Urol* 9:223-229.
- [11] Shokri S, Soltani A, Kazemi M, Sardari D, Mofrad FB. 2015 Effects of Wi-Fi (2.45 GHz) exposure on apoptosis, sperm parameters and testicular histomorphology in rats: a time course study. *Cell J* 17:322-31.
- [12] Dasdag S, Tas M, Akdag MZ, Yegin K. 2015 Effect of long-term exposure of 2.4 GHz radiofrequency radiation emitted from Wi-Fi equipment on testes functions. *Electromagn Biol Med* 34:37-42.
- [13] Avendaño C, Mata A, Sanchez Sarmiento CA, Doncel GF. 2012 Use of laptop computers connected to the internet through Wi-Fi decreases human sperm motility and increases sperm DNA fragmentation. *Fertil Steril* 97:39-45.
- [14] Yildiring ME, Kaynar M, Badem H, Cavis M, Karatus OF, Cimentepe E. 2015 What is harmful for male fertility: Cell phone or wireless internet? *Kaosiung J Med Sci* 31:480-4.
- [15] Özorak A1, Nazıroğlu M, Çelik Ö, Yüksel M, Özçelik D, Özkaya MO, Çetin H, Kahya MC, Kose SA. 2013 Wi-Fi (2.45 GHz)- and mobile phone (900 and 1800 MHz)-induced risks on oxidative stress and elements in kidney and testis of rats during pregnancy and the development of offspring. *Biol Trace Elem Res* 156:221-9.
- [16] Oni OM, Amuda DB, Gilbert CE. 2011 Effects of radiofrequency radiation from WiFi devices on human ejaculated sperm. *Int J Res Reve Appl Sci* 9: Article 13; 2011.
- [17] Aynali G, Nazıroğlu M, Çelik Ö, Doğan M, Yarıktaş M, Yasan H. 2013 Modulation of wireless (2.45 GHz)-induced oxidative toxicity in laryngotracheal mucosa of rat by melatonin. *Eur Arch Otorhinolaryngol* 2013;270:1695-700.
- [18] Çiftçi ZZ, Kırzioğlu Z, Nazıroğlu M, Özmen Ö. 2015 Effects of prenatal and postnatal exposure of Wi-Fi on development of teeth and changes in teeth element concentration in rats. [corrected]. *Biol Trace Elem Res* 163:193-201
- [19] Tök L, Nazıroğlu M1, Doğan S, Kahya MC, Tök O. 2014 Effects of melatonin on Wi-Fi-induced oxidative stress in lens of rats. *Indian J Ophthalmol* 62:12-5.

- [20] Çiğ B, Nazıroğlu M. 2015 Investigation of the effects of distance from sources on apoptosis, oxidative stress and cytosolic calcium accumulation via TRPV1 channels induced by mobile phones and Wi-Fi in breast cancer cells. *Biochim Biophys Acta* 1848(10 Pt B):2756-65.
- [21] Dasdag S, Akdag MZ, Erdal ME, Ay O, Ay ME, Yilmaz SG, Tasdelen B, Yegin K. 2015 Effects of 2.3 GHz radiofrequency radiation emitted from Wi-Fi equipment on microRNA expression in brain tissue. *Int J Radiat Biol* 91:555-61.
- [22] Saili L, Hanini A, Smirani C, Azzouz I, Azzouz A, Sakly M, Abdelmelek H, Bouslama Z. 2015 Effects of acute WiFi signals (2.45 GHz) on heart variability and blood pressure in albino rabbits. *Environ Toxicol Pharmacol* 40:600-5.
- [23] Papageorgiou CC, Hountala CD, Maganioti AE, Kiprianou MA, Rabavilas ASD, Papademitriou GN, Capalis CN. 2011 Effects of Wi-Fi signals on the P300 component or event-related potentials during an auditory hayling task. *J Integr Neurosci* 10:189-202.
- [24] Maganioti AE, Papageorgiou CC, Hountala CD, Kiprianou MA, Rabavilas AD, Papademitriou GN, Capalis CN 2010 Wi-Fi electromagnetic fields exert gender related alterations on EEG. 6th International Workshop on Biological Effects of Electromagnetic Fields. https://www.researchgate.net/profile/Miltiades_Kyprianou3/publication/267816859_WI-FI_ELECTROMAGNETIC_FIELDS_EXERT_GENDER_RELATED_ALTERATIONS_ON_EEG/links/550ab8670cf265693ced8e9c.pdf
- [25] Lee SS, Kim HR, Kim MS, Park SH, Kim DW. 2014 Influence of smart phone Wi-Fi signals on adipose-derived stem cells. *Ja J Cranofac Surg* 25:1902-7.

Columbia University, College of Physicians and Surgeons
Department of Physiology and Cellular Biophysics

Board Member
Los Angeles Unified School District,
Board of Education

Re: Health effects of cell tower radiation


As an active researcher on biological effects of electromagnetic fields (EMF) for over twenty five years at Columbia University, as well as one of the organizers of the 2007 online Bioinitiative Report on the subject, I am writing in support of a limit on the construction of cell towers in the vicinity of schools.

There is now sufficient scientific data about the biological effects of EMF, and in particular about radiofrequency (RF) radiation, to argue for adoption of precautionary measures. We can state unequivocally that EMF can cause single and double strand DNA breakage at exposure levels that are considered safe under the FCC guidelines in the USA. As I shall illustrate below, there are also epidemiology studies that show an increased risk of cancers associated with exposure to RF. Since we know that an accumulation of changes or mutations in DNA is associated with cancer, there is good reason to believe that the elevated rates of cancers among persons living near RF towers are probably linked to DNA damage caused by EMF. Because of the nature of EMF exposure and the length of time it takes for most cancers to develop, one cannot expect ‘conclusive proof’ such as the link between helicobacter pylori and gastric ulcer. (That link was recently demonstrated by the Australian doctor who proved a link conclusively by swallowing the bacteria and getting the disease.) However, there is enough evidence of a plausible mechanism to link EMF exposure to increased risk of cancer, and therefore of a need to limit exposure, especially of children.

EMF have been shown to cause other potentially harmful biological effects, such as leakage of the blood brain barrier that can lead to damage of neurons in the brain, increased micronuclei (DNA fragments) in human blood lymphocytes, all at EMF exposures well below the limits in the current FCC guidelines. Probably the most convincing evidence of potential harm comes from living cells themselves when they start to manufacture stress proteins upon exposure to EMF. The stress response occurs with a number of potentially harmful environmental factors, such as elevated temperature, changes in pH, toxic metals, etc. This means that ***when stress protein synthesis is stimulated by radiofrequency or power frequency EMF, the body is telling us in its own language that RF exposure is potentially harmful.***

There have been several attempts to measure the health risks associated with exposure to RF, and I can best summarize the findings with a graph from the study by Dr. Neil Cherry of all childhood cancers around the Sutro Tower in San Francisco between the years 1937 and 1988. Similar studies with similar results were done around broadcasting antennas in Sydney, Australia and Rome, Italy, and there are now studies of effects of cellphones on brain cancer. The Sutro tower contains antennas for broadcasting FM (54.7 kW), TV (616 kW) and UHF (18.3 MW) signals over a fairly wide area, and while the fields are not uniform, and also vary during the day, the fields were measured and average values estimated, so that one could associate the cancer risk with the degree of EMF exposure.

The data in the figure are the risk ratios (RR) for a total of 123 cases of childhood cancer from a population of 50,686 children, and include a 51 cases of leukaemia, 35 cases of brain cancer and 37 cases of lymphatic cancer. It is clear from the results that the risk ratio for all childhood cancers is elevated in the area studied, and while the risk falls off with radial distance from the antennas, as expected, it is still above a risk ratio of 5 even at a distance of 3km where the field was $1\mu\text{W}/\text{cm}^2$. This figure is what we can expect from prolonged RF exposure. In the Bioinitiative Report, we recommended $0.1\mu\text{W}/\text{cm}^2$ as a desirable precautionary level based on this and related studies, including recent studies of brain cancer and cellphone exposure.



As I mentioned above, many potentially harmful effects, such as the stress response and DNA strand breaks, occur at nonthermal levels (field strengths that do not cause a temperature increase) and are therefore considered safe. It is obvious that the safety standards must be revised downward to take into account the nonthermal as well as thermal biological responses that occur at much lower intensities. Since we cannot rely on the current standards, it is best to act according to the precautionary principle, the approach advocated by the European Union and the scientists involved in the Bioinitiative report. In light of the current evidence, the precautionary approach appears to be the most reasonable for those who must protect the health and welfare of the public and especially its most vulnerable members, children of school-age.

Sincerely yours,

Martin Blank, Ph.D.

Associate Professor of Physiology and Cellular Biophysics



6840 Seaview Road, Sechelt, BC V0N 3A4

www.zonein.ca

MCPS COO Dr. Andrew Zuckerman
MCPS Interim Superintendent Larry Bowers
MCPS Board of Education
MCPS Office of Technology
Montgomery County Schools
Carver Educational Services Center
850 Hungerford Drive
Rockville, MD 20850

December 13, 2015

Dear Montgomery County COO Dr. Andrew Zuckerman, Interim Superintendent Larry Bowers, Board of Education and Office of Technology;

In my capacity as a pediatric occupational therapist, biologist, international speaker, and author on the subject of the impact of technology on child development and learning, I'm writing to you on behalf of students, teachers, and parents requesting you reconsider the use of devices which operate using wireless radiation.

Please find below guiding principles regarding managed balance between technology and healthy activity, as well as information on wireless radiation. More judicious use of educational based technologies in a safe manner, will serve to ensure sustainable futures for all children. Reversion to Ethernet or fiber optic cable devices, until such time as the World Health Organization deems wireless to not be harmful to young children, is recommended.

Guiding principles for the use of educational based technology in school environments.

Minimize Risk and Maximize Safety.

- Wireless radiation has not been proven safe (WHO 2011).
- Recent research indicates wireless radiation causes harmful effects to adult humans (Avendano 2012, Hardell 2013).
- Long term effects of wireless radiation on children are unknown at this time (AAP 2013).
- Children have thinner skulls, more aqueous bodies, and have rapidly developing cells, indicating they are exceedingly more vulnerable to harmful effects from wireless radiation than adults (AAP 2013, C4ST 2015).
- The American Academy of Pediatrics and the Canadian Pediatric Society recommends no more than 1-2 hours total technology use per day, including

educational technology. Many schools exceed these expert guidelines (AAP 2014).

Weigh Risk vs. Benefit.

- Education technology is not evidence based and is laden with conflict of interest e.g. manufacturers claims are financially motivated, and are not substantiated by university level research.
- Traditional and standardized teaching methods have substantive research support and evidence, yet are being rapidly replaced with education technology.

Ensure adequate foundational skills prior to use of technology.

Children need to balance the following 4 critical factors with technology, to optimize development and learning. Time spent with technology adversely affects these factors.

- *Movement*: stimulates vestibular, proprioceptive and cardiovascular systems.
- *Touch*: stimulates parasympathetic system for lowered cortisol and adrenalin.
- *Human Connection*: activates parasympathetic system; a life sustaining force.
- *Nature*: attention restorative, improves learning, erases effects of technology.
- See video: [Message to Schools on EdTech](#)

Risks associated with the use of technology by children are as follows:

- *Sedentary nature* of technology use is causally related to the recent rise in obesity/diabetes, developmental delay and learning difficulties (Tremblay 2011, HELP EDI Mapping 2009/13, Ratey 2008, PISA 2012).
- *Isolating factor* of technology use is associated with escalation in social impairments, mental illnesses (including adhd and autism), and self-regulation difficulties (Houtrow 2014).
- *Overstimulation* from technology use is a causal factor in rise in attention deficit, aggression, sleep disturbance, and chronic stress from hyper-arousal of the sympathetic nervous system (Christakis 2004, Gentile 2009, Markman 2010, Bristol University 2010).
- *Neglect* of students by teachers and support staff who are engaged in their own personal technology, is unfortunately common.
- Consequently, the risks associated with using education technology far outweigh the dubious benefits.

When In Doubt, Act With Caution.

- Existing research on harmful effects of wireless radiation on *adults*, indicates taking a cautionary approach when considering same radiation exposure to *children* (AAP 2014).

- Rapid cell turnover in children creates particular concern regarding potential DNA damage from wireless radiation, and consequent susceptibility to cancer. While rise in cancer incidence is becoming more apparent, rise in rates of cancer in children will not be observable until adulthood.
- Removal of wireless radiation and reversion to Ethernet cabled devices, will ensure immediate and long term safety to all students, teachers, and support staff.
- Defaulting to a remote authority regarding removing wireless radiation from schools, is not acting in the best interests of students and staff, and may not be defensible in a court of law.

Montgomery County's statement that the radiofrequency levels in schools "is compliant" with federal regulations *does not* assure safety to the students in your care. The current proposed technology plan to further increase the use of screens in classrooms on a daily basis, clearly does not support children's healthy development.

The implications of failure of schools to act with caution now regarding wireless radiation and technology, could potentially be horrific in both scope and magnitude, and may constitute neglect of children. Please act now to safeguard your children's future.

Respectfully,

CRowan

Cris Rowan, BScBi, BScOT, SIPT, AOTA Approved Provider
CEO Zone'in Programs Inc. and Sunshine Coast Occupational Therapy Inc.
crowan@zonein.ca email
Websites: www.zonein.ca, www.suncoastot.com, www.virtualchild.ca;
Blog: www.movingtolearn.ca

Research References

American Academy of Pediatrics, Letter to Federal Communications Commission on August 29, 2013; retrieved on February 26, 2014 at
<http://apps.fcc.gov/ecfs/document/view?id=7520941318>.

American Academy of Pediatrics, Policy Statement – *Children, Adolescents, and the Media*, Council on Communications and Media; retrieved on February 26, 2014 from
<http://pediatrics.aappublications.org/content/132/5/958.full>

Avendano, C., Mata, A., Sanchez Sarmiento C. A., Doncel, C. F. Use of laptop computers connected to internet through Wi-Fi decreases human sperm motility and increases sperm DNA fragmentation. *Fertility and Sterility*. January 2012; 97(1): 39-45. Retrieved on March 18, 2014 from [http://www.fertstert.org/article/S0015-0282\(11\)02678-1/abstract](http://www.fertstert.org/article/S0015-0282(11)02678-1/abstract)

Bristol University: School for Policy Studies News (2010). Available at: <http://www.bristol.ac.uk/sps/news/2010/107.html>

C4ST: Dr. Oz Alert – Why Cell Phones Could Cause Cancer in Kids (2015). Available at: <http://www.c4st.org/news/educate-yourself/dr-oz-alert-why-cell-phones-could-cause-cancer-in-kids.html>

Christakis DA, Zimmerman FJ, DiGiuseppe DL, McCarty CA. Early television exposure and subsequent attentional problems in children. *Pediatrics*. 2004; 113 (4): 708-713.

Gentile, D. Pathological Video-Game Use Among Youth Ages 8 to 18: A National Study. *Psychological Science*. 2009 May;20(5):594-602. doi: 10.1111/j.1467-9280.2009.02340.x.

Hardell, L., Carlberg, M., Soderqvist, F., Hansson, K. Pooled analysis of case-control studies on acoustic neuroma diagnosed 1997-2003 and 2007-2009 and use of mobile and cordless phones *International Journal of Oncology*. 2013; 43(4); 1036-1044. DOI: 10.3892/ijo.2013.2025.

Healthy Early Learning Partnership – Early Development Inventory Maps for British Columbia, University of British Columbia; retrieved on February 26, 2014 from <http://earlylearning.ubc.ca/maps/edi/bc/>.

Houtrow, A. J., Larson K., Olson, L. M., Newacheck, P. W., Halfon, N. Changing Trends of Childhood Disability, 2001-2011. *Pediatrics*. Available at <http://pediatrics.aappublications.org/content/early/2014/08/12/peds.2014-0594.abstract>

Markman, A. (2010) Ulterior Motives- How goals seen and unseen drive behaviours. *Psychology Today*. Retrieved from <http://www.psychologytoday.com/blog/ulterior-motives/201003/the-broad-view-research-video-games-and-aggression> on March 11, 2014.



6840 Seaview Road, Sechelt, BC V0N 3A4

www.zonein.ca

Ratey JJ, Hagerman E (2008). Spark: The Revolutionary New Science of Exercise and the Brain. Little, Brown and Company, New York.

Tremblay, M.S., LeBlanc, A.G., Kho, M.E., Saunders, T.J., Larouche, R., Colley, R.C., Goldfield,

G., Gorber, S.C. (2011) Systematic review of sedentary behaviour and health indicators in school-aged children and youth. *International Journal of Behavioral Nutrition and Physical Activity* 2011, 8:98 doi:10.1186/1479-5868-8-98

World Health Organization – International Agency for Research on Cancer, Press Release No. 208, May 31, 2011. Retrieved on June 1, 2015 at http://www.iarc.fr/en/media-centre/pr/2011/pdfs/pr208_E.pdf

© Cris Rowan, Zone'in Programs Inc. 2015



P.O. Box 58
Teton Village, WY 83025
www.ehtrust.org

Montgomery County Board of Education
Montgomery County Schools
Carver Educational Services Center
850 Hungerford Drive
Rockville, MD 20850

January 20, 2016

Dear Montgomery County Board of Education,

Concerned parents in your school district have asked me to write to you regarding the health risks of wireless radiofrequency radiation exposure in the classroom. Based on what I have been told, I want to urge you to halt programs that currently have students use their own phones in ways that expose their eyes and brains to levels of radiation that have never been tested for safety.

I was Founding Director of the Board on Environmental Studies and Toxicology of the U.S. National Research Council, and Founding Director of the Center for Environmental Oncology at the University of Pittsburgh Cancer Institute. President Clinton appointed me to the Chemical Safety and Hazard Investigation Board, and I am former Senior Advisor to the Assistant Secretary for Health in the Department of Health and Human Services. I founded the non-profit Environmental Health Trust in 2007 to provide basic research and education about environmental health hazards. Our scientific team is currently focusing on the health risks of radiofrequency radiation as an important public health issue.

Many people are unaware that cell phones and wireless laptops and tablets function as two-way microwave radios. A typical classroom might have the following scenario: every student has a laptop--which is typically tested for use 8 inches from an adult male body--a cell phone in the pocket--which is also tested at a minimum distance from an adult male body-- and a network transmitter on the ceiling and possibly a cell tower outside next to the sports field. All these devices emit microwave radiation which can be readily absorbed into children's bodies and brains.

Manufacturers specifically recommend that cell phones be used "as tested"—at this little-known minimum distance from the body. Recently, [*Consumer Reports*](#) in November advised that people should not keep phones in the pocket—advice that few children or adults appreciate. *These devices have never been tested for safety with children.* Accumulating research indicates that long-term exposure to low levels over long lifetimes could pose a serious risk to our health.

Regarding tested distances for using laptops, the Federal Communications Commission (FCC) states that laptops and computers are “mobile devices are transmitters designed to be used in such a way that a separation distance of at least 20 centimeters is normally maintained between the transmitter's radiating structure(s) and the body of the user or nearby persons.” The body in this instance refers to a large male weighing more than 200 pounds and standing six feet tall.

As the county is preparing to increase student use of Chromebooks, please be aware that the Samsung [Chromebook manual](#) states:

“United States of America USA and Canada Safety Requirements and Notices

- Do not touch or move antenna while the unit is transmitting or receiving.
- Do not hold any component containing the radio such that the antenna is very close or touching any exposed parts of the body, especially the face or eyes, while transmitting.
- Regardless of the power levels, care should be taken to minimize human contact during normal operation.
- This device should be used more than 20 cm (8 inches) from the body when wireless devices are on and transmitting.
- FCC Statement for Wireless LAN use: *“While installing and operating this transmitter and antenna combination the radio frequency exposure limit of 1mW/cm² may be exceeded at distances close to the antenna installed. Therefore, the user must maintain a minimum distance of 20cm from the antenna at all times.”*

As one of the leaders in educational policy of this nation, your school district has an opportunity to set an example for school districts nationwide by installing safer technology in classrooms and educating students, teachers and staff about tested distances that devices should be used to reduce radiation. A number of public and private schools have already implemented such policies. Just as we provide children with seat belts and bike helmets, a precautionary approach to wireless is recommended by many scientists and governments worldwide.

For more information about all of these issues, please read cell phone instructions for various models at <http://showthefineprint.org>. Our [newly posted Ebook](#) also details fine print safety instructions in wireless device user manuals.

When children use these devices close to their bodies, they are exceeding these safety instructions, and exposing themselves to radiofrequency (RF) radiation levels which can exceed our government FCC RF radiation exposure limits. The FCC RF exposure limit was designed to protect the public from the thermal (heating) effects of acute exposure to RF energy. The FCC states, “Tissue damage in humans could occur during exposure to high RF levels because of the body's inability to cope with or dissipate the excessive heat that could be generated. Two areas of the body, the eyes and the testes, are particularly vulnerable to RF heating because of the relative lack of available blood flow to dissipate the excess heat load.”

CHILDREN ABSORB MORE RADIATION THAN ADULTS

Our recently published research in the [IEEE Spectrum](#) with investigators at the Federal Universities of Brazil provides new state-of-the-art radiation exposure brain modeling which confirms that substantially higher radiofrequency radiation doses occur in younger children as compared to adults even where products comply with tested guidelines developed for adults.

FCC REGULATIONS ARE OUTDATED

FCC exposure limits were set more than 19 years ago and were based on decades-old research. The Government Accountability Office published a [2012 Report](#) that calls on the FCC to formally reassess their current RF energy (microwave) exposure limits, stating that the “FCC RF energy exposure limit *may not* reflect the latest research.” I encourage you to read scientific submissions to FCC Proceeding Number 13-84 at <http://bit.ly/1aGxQiq>. It is unknown when the FCC will make a ruling, however, *until that time* the current outdated FCC limits are *not reflective* of the current state of science.

FCC REGULATIONS DO NOT PROTECT THE PUBLIC FROM BIOLOGICAL EFFECTS

As the California Medical Association states in their [2014 Resolution](#) calling for updated FCC Regulations, “peer reviewed research has demonstrated adverse biological effects of wireless EMF [electromagnetic fields] including single and double stranded DNA breaks, creation of reactive oxygen species, immune dysfunction, cognitive processing effects, stress protein synthesis in the brain, altered brain development, sleep and memory disturbances, ADHD, abnormal behavior, sperm dysfunction, and brain tumors.”

In May 2015, over 200 scientists who have authored more than 2,000 articles on this topic appealed to the United Nations to address “the emerging public health crisis” related to cellphones and other wireless devices, urging that the United Nations Environmental Programme (UNEP) initiate an assessment of alternatives to current exposure standards and practices that could substantially lower human exposures to non-ionizing radiation. These scientists state that “the ICNIRP guidelines do not cover long-term exposure and low-intensity effects, “ and are “ insufficient to protect public health.” They also state that “the various agencies setting safety standards have failed to impose sufficient guidelines to protect the general public, particularly children who are more vulnerable to the effects of EMF.” Please see their website at <https://emfscientist.org>.

INCREASED CANCER RISK

Wireless radiofrequency radiation was classified as a Class 2B “Possible Human Carcinogen” by the World Health Organization’s International Agency for Research on Cancer in 2011. According to many scientists, evidence *has increased* since 2011, indicating that cell phone and wireless radiation should be classified as a “probable carcinogen.” Those exposed at younger ages show four to eight times increased cancer risk. [Replicated research](#) just published in Biochemical and Biophysical Research Communications indicates that radiofrequency acts as a *tumor promoter* at low to moderate levels.


CONCERN FOR PREGNANT STUDENTS AND STAFF

Pregnant students and staff are especially at risk from wireless because the fetus is the most vulnerable to toxic exposures. Several experimental studies are showing irreversible changes after prenatal exposure to cell phone and wireless radiation such as altered brain functioning, decreased brain cells and altered reproductive organ development. More than 100 physicians, scientists and public health professionals joined together to express their concern about the risk that wireless radiation poses to pregnancy and now *urge pregnant women to limit their exposures*. Please read these scientists' [BabySafe Joint Statement](#)

VIRTUAL TECHNOLOGY RESULTS IN HIGHER EXPOSURES TO THE EYE AND BRAIN

Most recently, I was contacted by a parent in your district about the virtual reality devices now used in MCPS classrooms to go on a virtual “field trip.” As indicated by online instructions, this experience involves using smartphones placed directly in front of the child’s eyes so that they can directly watch a fascinating video of faraway lands. The smartphone is streaming radiation throughout the classroom from the teacher's iPad for the entire “field trip.”

Please be aware that FCC regulations set decades ago did not utilize science that looks at the effects from cell phones on different body tissues such as the eyes. Upon hearing about this issue, I contacted EHT-associated scientists at federal universities of Brazil who do state-of-the-art computer modeling. I asked them to position the phone as it would be in the virtual reality cardboard for use in front of the child’s eyes and assess the microwave radiation. The yellow and orange color show the highest exposures.



My colleagues and I are sharing this work with you today because we believe you should have more information about microwave radiation exposures that will take place through this system.

This research image above utilizes [a sophisticated computer system](#) that the U.S. Food and Drug Administration (FDA) currently applies to evaluate medical devices. It simulates the radiation absorption into *anatomically correct models*--something that currently used systems for testing phones and devices cannot do. [In a study from Memorial Sloan-Kettering Cancer Center](#), radiation physicist David Gultekin, working with Bell Labs electrical engineer Lothar Moeller, reported that normal working cell phones can create tiny hotspots within brain tissue. Unlike other organs, [eyes](#) do not have circulation to effectively carry away heat.

In addition to the impact from the microwave radiation, there could also be impacts to a child’s retina from the blue light emitted by the screen. Youths under the age of 20, and especially very young children,

have little or no yellowing of the lens (which helps protect the adult eye). Therefore, blue light (or UV) which enters the eye is unfiltered in children and strikes the retina at full-strength exposing not only the retina, but the lens to possible damage over the long time. Such injury may not be evident until later in time.

In 2010, [Andreas Christ and team](#) reported that children's hippocampus and hypothalamus absorbs 1.6–3.1 times higher and the cerebellum absorbs 2.5 times higher microwave radiation compared to adults; children's bone marrow of the skull absorbs 10 times higher microwave radiation than in adults, *and children's eyes absorb much higher microwave radiation than adults*. A recent [Deans' Lecture](#) I delivered to University of Melbourne provides an overview on this research.

SIMPLE STEPS WILL PROTECT CHILDREN

Compelling research raises the possibility of very serious harm to children from radiofrequency radiation exposures well below “FCC compliant” levels. Legal does not mean safe. Based on the preliminary work that I share with you here, I urge you to forgo the use of such devices such as virtual reality cardboard as there is no research that has considered their impact on children's eyes. At this time, the smart choice for school decision makers is to act now and reduce radiofrequency wireless exposures. In fact, many countries (over 20) and health authorities worldwide recommend reducing radiofrequency radiation to children.

More recently, the Cyprus Government's National Committee on Environment and Children's Health released a [video about reducing wireless](#) and I invite you to watch this excellent example of responsible action at this link <https://www.youtube.com/watch?v=H43IKNjTvRM>.

I understand that your county has a Bring Your Own Device policy whereby cell phones are not only allowed *in* the classroom but are actively used in the curriculum. As I have been told, students in film class might use their cell phones to take footage to create a movie, and in some math classes they use their cell phones as a calculator. Advice should be routinely provided to any student using a wireless device at school about *how to reduce exposures*. For example, if phones are used on airplane mode, and wireless is turned off on computers then these devices will neither send nor receive microwave radiation.

When powered on, phones undergo short bursts of microwave radiation up to 900 times per minute, *whether or not the phone is being used for talking*. Once teachers and students are educated on how they can simply turn their phone onto airplane mode, then they can use the phone in the classroom *without* being exposed to unnecessary radiofrequency radiation.

Likewise, laptops such as Chromebooks are also emitting constant radiation and at much higher levels when a student is streaming video or using cloud based applications. Laptops can easily be hardwired to ethernet so that students can safely use the internet without radiation emissions. Please review the [Best Practices for Low EMF in Schools developed by the Northeast Collaborative For High Performing Schools](#) which details how schools can reduce exposure to radiofrequency fields and still have full internet connectivity.

Along with [the recommendation](#) of over 200 scientists (see <https://emfscientist.org>) and health authorities worldwide, I recommend that the best course of action is to take simple precautions—as many nations already currently advise. *Children’s exposures to wireless radiation should be reduced as much as possible.* We have a responsibility to act now to reduce children’s exposure to radiofrequency radiation. Children’s nervous, immune and reproductive systems are rapidly developing and, along with pregnant women, children deserve an abundance of caution.

As several colleagues and I wrote in [a letter](#) to the U.S. Secretary of Education just a few months ago, we recommend your school district do the following:

1. **Raise school community awareness through new educational curriculum:** Students, teachers and their families should be given information on wireless health risks and simple precautionary steps they can take to protect their health. It is important to teach children how to use technology both safely and more responsibly in order to protect their health and wellbeing.
2. **Install a safe communication and information technology infrastructure in schools to meet educational needs:** Solutions exist to reduce exposures to wireless emissions and mitigate the health risk. Low-EMF Best Practices have been developed, allowing educational needs to be met with safer, hard-wired Internet connections, which are also faster and more secure.

Low-EMF Best Practices are the solution that allows for full communication, information access and learning tools use in the classroom while minimizing unnecessary health risks. Your district can thoughtfully integrate safe technology into every classroom while responsibly safeguarding the health of every generation.

I fully understand that this information has not been widely understood. I would be happy to provide or develop an online technical briefing to your senior staff to assist you as you make decisions today that will affect the health of students for the rest of their lives.

Yours respectfully,



Devra Davis, PhD MPH
President and Founder
Environmental Health Trust
Visiting Professor of Medicine
The Hebrew University, Hadassah Medical Center
Associate Editor, Frontiers in Radiation and Health
ehtrust.org



July 28, 2014

Board of Trustees
Fay School
48 Main Street
Southborough, MA 01772

Re: Advisability of WiFi in schools

Dear Sirs/Madams:

This is concerning potential adverse health effects associated with exposure to radiofrequency/microwave (RF/MW) radiation, specifically that from wireless routers and wireless computers. I am writing to express concern that students at your school are experiencing electrosensitivity symptoms from these technologies.

I am a public health physician who has been involved in issues related to electromagnetic fields (EMFs) for several decades. I served as the Executive Secretary for the New York Powerline Project in the 1980s, a program of research that showed that children living in homes with elevated magnetic fields coming from powerlines suffered from an elevated risk of developing leukemia. I served as Director of the Wadsworth Laboratory of the New York State Department of Health, as well as Dean of the School of Public Health at the University at Albany/SUNY. I have edited two books on effects of EMFs, ranging from low frequency fields to radiofrequency/ microwave radiation, or the kind emitted by WiFi routers, cell phones, neighborhood antennas and wireless computer equipment. I served as the co-editor of the BioInitiative Report 2012 (Bioinitiative.org), a comprehensive review of the literature showing biological effects at non-thermal levels of exposure, much of which has since been published in the peer-reviewed journal, *Pathophysiology* (attached). Also, I served on the President's Cancer Panel that examined radiation exposures as they relate to cancer risk, in 2009, and a report from that testimony is also attached. Thus, this is a subject which I know well, and one on which I take a public health approach rooted in the fundamental principle of the need to protect against risk of disease, even when one may not have all the information that would be desirable.

There is clear and strong evidence that intensive use of cell phones increases the risk of brain cancer, tumors of the auditory nerve and cancer of the parotid gland, the salivary gland in the cheek by the ear. The evidence for this conclusion is detailed in the attached publications. The WHO's International Agency for Research on Cancer has also classified the radiation from both cell phones and WiFi as a Class 2B "Possible Carcinogen" (2011). WiFi uses similar radio-frequency radiation as cell phones (in the 1.8 to 5.0 GHz range). The difference between a cell phone and a WiFi environment, however, is that while the cell phone is used only intermittently, and at higher power, a WiFi environment is continuous, and transmitting even when not being used. In addition, WiFi transmitters are indoors, where people (and in this case, children) may be very close by, or certainly close to devices using the WiFi, such as wireless computers, iPads and smart boards, the radiation from which can be intolerable to sensitive people.

Furthermore, commercial routers, like those in schools, operate at much higher wattage than consumer routers. They are designed to penetrate through materials like cement, wood and brick, to handle dozens to hundreds of users, and to reach into outdoor areas, so industrial grade routers are of much greater concern.

An additional consideration to appreciate is that it is not only the power of wireless radiation that causes biological dysregulation, but the frequencies, pulsing, amplitude, and the quantity and kind of information being transmitted that can have effects as well. These 'non-thermal effects' have been shown in thousands of studies to be biologically active, and may be more important than the effects from the power. Thus, while a router may be in the ceiling, or not right next to a student, teacher or administrator, the known biological and health effects, particularly the non-thermal ones, are still very much occurring.

Finally, while acute electrosensitivity symptoms, like the ones I understand your students are experiencing, are of course of great concern (such as cognitive effects impairing attention, memory, energy levels, and concentration; cardiac irregularities, including in children; or, headaches or other symptoms in students wearing braces), the full effects for society from chronic and cumulative exposures are not known at this time. Given what we do know, however, including the DNA effects, I must, as a public health physician, advise minimizing these exposures as much as possible. Indications are that cell phones and wireless technologies may turn out to be a serious public health issue, comparable to tobacco, asbestos, DDT, PCBs, pesticides and lead paint, or possibly worse given the ubiquitous nature of the exposures. While unfortunately we must wait for federal regulation to catch up with the science, the prudent thing to do in the interim would be to exercise precaution at every opportunity.

Computers and the world-wide web have tremendous value in education, but the value also depends on how these are used in numerous respects. As wired internet connections do not pose radiation risk, are readily available, are faster and more secure than WiFi, and are now even available for certain tablets, I highly recommend you factor the risks I have described into your technology planning. At the same time, I would urge you to take the complaints of your students very seriously, and potentially involve the school nurse and teachers in helping to assess the extent of the electrosensitivity problem among students at the school.

An excellent reference on the EMF and electrosensitivity science is "Electrosensitivity and Electrohypersensitivity—A Summary" (2013) authored by M.J. Bevington and available through Electrosensitivity-U.K. (www.es-uk.info/)

If I can be of further help, please do not hesitate to call.

Yours sincerely,

A handwritten signature in blue ink, appearing to read "David O. Carpenter".

David O. Carpenter, M.D.
Director, Institute for Health and the Environment
University at Albany

Enclosures

Martin Blank, PhD
Department of Physiology and Cellular Biophysics
Columbia University
New York, NY 10032

July 25, 2014

Mr. Thomas McKean, President, Board of Trustees
Mr. James Shay, President-Elect, Board of Trustees
Fay School
48 Main Street
Southborough, MA01772

To the Board of Trustees,

It has been brought to my attention that school children have become symptomatic at your school after installation of WiFi. I am writing to express my concern and to encourage you to review the independent science on this matter.

I can say with conviction, in light of the science, and in particular in light of the cellular and DNA science, which has been my focus at Columbia University for several decades, putting radiating antennas in schools (and in close proximity to developing children) is an uninformed choice. Assurances that the antennas are within 'FCC guidelines' is meaningless today, given that it is now widely understood that the methodology used to assess exposure levels only accounts for one type of risk from antennas, the thermal effect from the power, not the other known risks, such as non-thermal frequencies, pulsing, signal characteristics, etc. They fail also to consider multiple simultaneous exposures from a variety of sources in the environment, and cumulative exposures over a lifetime. Compliance with FCC guidelines, thus, unfortunately, is not in any way an assurance of safety today, as the guidelines are fundamentally flawed. Until the guidelines and advisories in the U.S. are updated, the intelligent thing for your Board of Trustees to do is to exercise the Precautionary Principle and hard wire all internet connections.


I know this might be disappointing to hear, as I understand you have invested in the WiFi. But there is no amount of money that could justify the added physiological stress from wireless antenna radiation and its many consequences, most in particular for children. Our research has shown that the cellular stress response, a protective reaction that is indicative of cellular damage, occurs at levels that are deemed 'safe'. Many other harmful reactions have been reported, such as the impairment of DNA processes that can account for the observed increased risk of cancer, as well as the potential cognitive decline, and sleep effects that may be due to impairment of the blood brain barrier. The DNA effects are of particular concern for future generations, an area of research that is just beginning to raise alarms. As with other environmental toxic exposures, children are far more vulnerable than adults, and they will have longer lifetimes of exposure.

The science showing reasons for concern about the microwave radiation emitted by antennas is abundant and there will be a day of reckoning. As I explain in my recent book,

Overpowered, The Precautionary Principle instructs us that in the face of serious threats, a lack of scientific 'certainty' never justifies inaction. The changes occurring at the molecular level, and known associations with many diseases, are sufficient at this time to give us pause and to recommend minimizing exposures to these fields, in our homes, schools, neighborhoods and workplaces. There is significant potential for risk, and to very large numbers of people, and the effects are occurring nonetheless whether or not we are noticing them.

I recommend you hardwire the internet connections at your school, and also encourage students to use hard wired connections at home for internet access, as well as for all computer equipment connections and voice communications.


Sincerely yours,



Martin Blank, PhD
mb32@columbia.edu,



Martin Blank, PhD, Special Lecturer and (ret.) Associate Professor, Columbia University, Department of Physiology and Cellular Biophysics. Dr. Blank is a leading expert in the effects of electromagnetic fields on DNA and biology, and Past President of the Bioelectromagnetics Society. He holds two PhDs, in physical chemistry and in colloid science, an interdisciplinary field involving chemistry, physics and nanoscience. Dr. Blank was author of the BioInitiative Report's section on the impact of electromagnetic fields on Stress Proteins; Editor of the journal *Pathophysiology's* special issue on Electromagnetic Fields (2009); and co-author of "Electromagnetic fields and health: DNA based dosimetry" (2012), which recommends a new way of assessing the biological impact of electromagnetic fields across the spectrum, using DNA. Dr. Blank's book, *"Overpowered—What Science Tells Us About the Dangers of Cell Phones and Other WiFi-Age Devices"*, was published in 2014.



Wi-Fi in Schools

Are We Playing It Safe With Our Kids?

“Current FCC standards do not account for the unique vulnerability and use patterns specific to pregnant women and children. It is essential that any new standard for cell phones or other wireless devices be based on protecting the youngest and most vulnerable populations to ensure they are safeguarded throughout their lifetimes.” American Academy of Pediatrics Letter to FCC August 29, 2013 (20)

By Cindy Russell, MD

VP of Community Health, SCCMA

Industry has been quite successful in creating magically useful wireless technologies such as cell phones, Ipads, Wi-Fi, and now wearable tech devices such as Google glasses, we all love. Many of these handy gadgets have now reached the typical classroom across the globe. It has become apparent, however, that there are substantial downsides to being too connected to technology and as safety concerns mount, governments such as France and Israel are backing away from the blind adoption of wireless technology in schools, especially for young children.

These devices are cool and convenient, however there remains nagging questions of overuse and safety as the application of these devices has increased to the point we are literally exposed 24 hours a day to this radiation. Wireless microwaves come from many sources both at work and at home.

An increasing number of physicians, scientists, and parents are concerned about long term health effects from Wi-Fi in schools. (42)(43)(44)(49) As any parent knows, computers now are as ubiquitous in schools as they are at work. From kindergarteners on up kids are required to learn computer skills in order to take core testing online. There is a push to enable students to be connected to the internet 24/7 to take photos, email documents, and research a topic. In schools, wired connections for computers have been rapidly being eliminated to install wireless systems that connect students both indoors and outdoors on campus.

Europe and some schools in the U.S. are taking a different more precautionary approach and going back to the future with wired plug in computers. Studies have also cast doubt on some of the benefits of classroom computers and warned of the new age of “Digital Dementia” which has now crept into Korean youth due to the heavy use of electronic gadgets. (17)(48)

Professors in college are banning computers during lectures and finding students learn more. (38) (39)

CHILDREN ARE MORE VULNERABLE THUS NEED MORE PROTECTION

Children have several organ systems that are immature at birth and are thus much more sensitive to toxic exposures. The human brain, one of the top vital organs, is far from being a finished product in youth. Long-term structural maturation of the nervous system is required for successful development of cognitive, motor, and sensory functions. Neuronal axons – long thin projections from the nerve cell – act as electronic transmission lines. Axons in major pathways of the brain continue to develop throughout childhood and adolescence. Myelin is the insulation surrounding individual nerves protecting it from outside electrical charges. The process of myelination is much faster the first two years but continues into adulthood. (16) Children have thinner skulls (29), their immune systems are undeveloped, their cells are dividing more rapidly, thus, they are more vulnerable to EMF radiation and other carcinogens. They also have a longer cumulative exposure to all toxins including EMF radiation.

CURRENT WIRELESS SAFETY STANDARDS AND MICROWAVING POTATOES

Wireless devices work on high frequency microwaves similar to the microwave you use to cook food with. It is with less power but substantial research (1)(2)(3)(4) demonstrates that even at low power within the current safety standards these microwaves can cause biologic harm to plants, animals, and cellular structures. Current Federal Communications Commission (FCC) standards are based only on heat generated by the device, not on adverse biological effects seen in hundreds of studies and at much lower levels.

Our own CMA supports reassessment of EMF standards. The California Medical Association, in 2014, passed a resolution as follows:

“Resolved 1: That CMA supports efforts to re-evaluate microwave safety exposure levels associated with wireless communication devices, including consideration

Continued on page 18

of adverse nonthermal biologic and health effects from non-ionizing electromagnetic radiation used in wireless communications and be it further

Resolved 2: That CMA support efforts to implement new safety limits for wireless devices to levels that do not cause human or environmental harm based on scientific research.

ADVERSE EFFECTS DEMONSTRATED IN PEER REVIEWED PUBLISHED RESEARCH (2)

- DNA with single and double stranded breaks
- Leakage of the blood brain barrier (two hours of cell phone exposure causes 7+ days of albumin leakage)
- Stress protein production in the body indicating injury
- Infertility/reproductive harm
- Neurologic harm with direct damage to brain cells
- Lowering of melatonin levels
- Immune dysfunction
- Inflammation/oxidation.

PLAUSIBLE MECHANISM FOUND FOR EMF MICROWAVE EFFECTS

Dr. Martin Pall, Professor Emeritus of Biochemistry, Washington State University has studied how electromagnetic fields impact the cells of our bodies. His 2013 paper on this subject highlights a major biological mechanism of action of EMF microwave radiation on cell structure. His work, along with two dozen prior studies, demonstrated that EMF microwave radiation effects cellular calcium channels and this can be inhibited with calcium channel blockers. "A whole series of biological changes reportedly produced by microwave exposures can now be explained in terms of this new paradigm of EMF actions via Voltage Gated Calcium Channels (VGCC) activation." (14)(15)

EMF AFFECTS ON WILDLIFE: BIRDS, BEES, AND TOMATO PLANTS

Bird researchers in Germany found that their migratory European Robins lost their sense of navigation when in the city. (5) This was found to be due to the EMF radiation interfering with the bird's special internal magnetic compass. They replicated the experiment over seven years before publishing the results in the prestigious journal *Nature*.

John Phillips and others have found that newts, sea turtles, and migratory birds use a magnetic compass to navigate long distances and this can be interrupted by low levels of EMF. (6)(7) A review of effects on cell towers and wireless devices showed that beehives can have rapid colony collapse with exposure to cell phone radiation. (8)

Plants have been shown to have stress response to EMF from wireless devices. (9)(10) (22) In tomatoes exposed for short duration, the stress response seen by exposure to EMF was prevented by administration of calcium counteracting drugs. (11) Even simple high school science experiments document abnormal seed growth near Wi-Fi routers. (19) There appear to be adverse biological effects of this seemingly harmless radiation.

HUMAN ELECTROSENSITIVITY: IS IT REAL?

There is varied opinion about those who state they are sensitive to EMF. Scientific research has not given a definitive answer, nevertheless, many seem to suffer from vague and often disabling symptoms they feel in the presence of EMF. Exposure to EMF radiation in some people reportedly causes headaches, memory problems, fatigue, sleep disorders, depression. This is so significant for some people that they have to live in a very low EMF environment to feel normal. (25)

Sweden recognizes electro-sensitivity as a functional impairment and estimates that about 3% of the population suffers from this. (23)(24) Dr. Magda Havas found in replicated studies that some EMF sensitive individuals heart rates increased with wireless devices turned on in double blind study. (12)(26) Researchers at Louisiana State University, in 2011, studied a self reported EMF sensitive physician and found "In a double-blinded EMF provocation procedure specifically designed to minimize unintentional sensory cues, the subject developed temporal pain, headache, muscle twitching, and skipped heartbeats within 100 s after initiation of EMF exposure ($p < .05$)." They concluded that "EMF hypersensitivity can occur as a bona fide environmentally inducible neurological syndrome." (27)

Genius and Lipp reviewed the current literature on EHS, in 2011, and point to several explanations for this multisystem phenomenon, including toxicant induced loss of tolerance as many with EHS symptoms had high levels of PCB's possibly causing immune dysfunction. Scientific research also identifies an inflammatory response with cytokine production. Another aspect of research points to catecholamine and adrenal gland dysfunction. In addition, heavy metal toxicity has also been proposed as contributing to EHS. (28)

The Austrian Medical Association feels Electrohypersensitivity is a real phenomenon and in 2012 published Guidelines for EMF and Electro-hypersensitivity. They state the primary method of treatment should consist in the prevention or reduction of EMF exposure, taking care to reduce or eliminate all sources of EMF if possible. (32)

GOVERNMENT ACTIONS ON WI-FI IN SCHOOLS

While much of the U.S. is marching forward with Wi-Fi in schools, Europe is changing direction, as indicated by the policies listed below. (45) Internationally there is wide disagreement in standards. The U.S. and Canadian limits are 1000 microwatts/cm². China and Russia are 10 microwatts/cm². Belgium is 2.4 microwatts/cm², and Austria is 0.001 microwatts/cm². The Bioinitiative Report 2012 recommendation for "No Observable Effect" is 0.0003 microwatts/cm². Cosmic background EMF we evolved with is <0.0000000001 microwatts/cm². (2)

COUNCIL OF EUROPE PARLIAMENT ASSEMBLY 2011 EMF MICROWAVE POLICY : "THE POTENTIAL DANGERS OF ELECTROMAGNETIC FIELDS AND THEIR EFFECT ON THE ENVIRONMENT"

The report notes "other non-ionizing frequencies, whether from ex-

In May 2011, the International Agency for Research on Cancer (IARC) classified radiofrequency electromagnetic fields as possibly carcinogenic to humans (Group 2B).(30)

tremely low frequencies, power lines or certain high frequency waves used in the fields of radar, telecommunications, and mobile telephony, appear to have more or less potentially harmful, non-thermal, biological effects on plants, insects, and animals, as well as the human body, even when exposed to levels that are below the official threshold values.”

The Council calls for a number of measures to protect humans and the environment, especially from high-frequency electromagnetic fields. One of the recommendations is to “take all reasonable measures to reduce exposure to electromagnetic fields, especially to radio frequencies from mobile phones, and particularly the exposure to children and young people who seem to be most at risk from head tumors”. (37)

IN FRANCE: A NEW NATIONAL LAW BANS WI-FI IN NURSERY SCHOOLS

In January 2015, France passed a landmark law that calls for precaution with wireless devices for children and the general public. (34)(35) It calls for:

1. Wi-Fi banned in nursery schools.
2. Wi-Fi routers should be turned off in school when not in use.
3. Schools are informed when new tech equipment is installed.
4. Citizens will have access to environmental cell tower radiation measurements near homes.
5. There will be continued research conducted into health effects of wireless communications.
6. Information on reducing exposure to EMF radiation is mandatory in the contents of the cell phone package.
7. Wi-Fi hotspots are labeled.

ISRAELI MINISTRY OF EDUCATION ISSUE GUIDELINES TO LIMIT WI-FI IN SCHOOLS

On August 27, 2013, the Israeli Ministry of Education issued new guidelines regarding Wi-Fi use in schools.

(33) The guidelines will:

1. Stop the installation of wireless networks in classrooms in kindergarten.
2. Limit the use of Wi-Fi between first and third grades. In the first grade, students will be limited to use Wi-Fi to study for one hour per day and no more than three days per week. Between the first and third grades, students will be limited to use Wi-Fi up to two hours per day for no more than four days per week.
3. To limit unnecessary exposure teachers will be required to turn off mobile phones and Wi-Fi routers when they are not in use for educational purposes.
4. All Wi-Fi equipment be tested for compliance with safety limits before and after installation in an Israeli school.
5. Desktop computers and power supplies be kept at least 20 cm from students.

2012 THE RUSSIAN COMMITTEE ON NON-IONIZING RADIATION PROTECTION



OFFICIALLY RECOMMENDED THAT WI-FI NOT BE USED IN SCHOOLS.

2011 THE RUSSIAN COMMITTEE ON NON-IONIZING RADIATION PROTECTION (RNCNIRP) RELEASED THEIR RESOLUTION ENTITLED “ELECTROMAGNETIC FIELDS FROM MOBILE PHONES: HEALTH EFFECTS ON CHILDREN AND TEENAGERS.”

According to the opinion of the Russian National Committee on Non-Ionizing Radiation Protection, the following health hazards are likely to be faced by the children mobile phone users in the nearest future: disruption of memory, decline of attention, diminishing learning and cognitive abilities, increased irritability, sleep problems, increase in sensitivity to the stress, increased epileptic readiness. (36)

Expected (possible) remote health risks: brain tumors, tumors of acoustical and vestibular nerves (in the age of 25-30 years), Alzheimer’s

Continued on page 20

disease, “got dementia”, depressive syndrome, and the other types of degeneration of the nervous structures of the brain (in the age of 50 to 60).

PLAYING IT SAFE FOR OUR KIDS

A healthy and safe learning environment is a cornerstone of education. Current FCC standards are obsolete and inappropriate as they are based only on heat effects, not biological effects. They give us a false sense of security. There may be higher EMF levels at school than at home as routers are more powerful. Cumulative Effects on DNA or cell structures are not taken into consideration in any safety standard. Because of the long-term exposure to EMF microwave radiation this generation is experiencing, they will be at higher risk for potential health problems. We will not know what happens to our progeny’s DNA until our grandchildren are born.

Considering there has been a more precautionary approach internationally to microwave radiation exposure and the trend is toward less exposure in schools, especially to vulnerable populations such as children, it makes sense to re-evaluate our wireless schools. We buckle our seat belts and wear a helmet when we ride bikes even though we don’t know if we will get in an accident. Although not all the issues of wireless microwaves are understood, there is enough science to understand it acts as a toxicant at even low levels that fall within current safety standards. We also know

3. **Limit Wi-Fi use**, especially in younger grades.
4. **Cell phones stay off and in the backpacks during class** and on the campus during school hours.
5. **Have EMF and electrical measurements done by one or more qualified, experienced consultants before and after any installation.** Understand you may need to increase your knowledge of low and high frequency electromagnetic fields and limits to accurately interpret the reports. The Bioinitiative Report is a very useful compendium that has recommendations for safer levels.
6. **Support efforts by governments to provide independent standardized transparent research to define safe limits in all the different wireless frequencies used commercially.** This could lead to less EMF emissions and safer wireless devices.

REFERENCES

1. “Overpowered” by Dr. Martin Blank, 2014
2. **Bioinitiative Report.** 2012 <http://www.bioinitiative.org/>
3. **EMF Portal.** <http://www.emf-portal.de/>
4. **Why Fi? : Is Wireless Communication Hazardous to Your Health?** <http://www.sccma-mcsm.org/Portals/19/assets/docs/Why%20Fi.pdf>
5. **Electronics’ noise disorients migratory birds.** Man-made

“Certain high frequency waves used in the fields of radar, telecommunications, and mobile telephony, appear to have more or less potentially harmful, non-thermal, biological effects on plants, insects, and animals, as well as the human body, even when exposed to levels that are below the official threshold values.”

that decades of research precedes meaningful regulation in the area of toxins, thus the only reasonable approach is precautionary.

In addition, we need to be thoughtful about how much our kids should use computers and what this is doing not only to them, but to our society as a whole. We get starry eyed with every new wireless gadget, however, in “Alone Together” Sherry Turkle expertly addresses the rise in isolation, loneliness, lack of privacy, and increasing pressure on students in this age of invasive technology. Her thorough and non-judgmental scientific investigation of the psychological effects of computers makes us aware that we need to take care that we do not replace real human connection with a “virtual reality” that will redirect us in an unhealthy direction.

As physicians and parents, we understand that decisions we make today may have far reaching consequences in the future for our kids. Let’s play it safe for them right now.

RECOMMENDATIONS FOR SCHOOLS

1. **Wired internet connections** like we used to have are the safest and possibly cheapest option – all the benefits of the internet without the risk.
2. **Wireless devices**, but with an on/off switch in each room so teachers can use only when needed for educational purposes.

electromagnetic radiation disrupts robins’ internal magnetic compasses. May 7, 2014 <http://www.nature.com/news/electronics-noise-disorients-migratory-birds-1.15176>

6. **True Navigation: Sensory Bases of Gradient Maps.** Phillips, J. 2006 <http://www.pigeon.psy.tufts.edu/asc/Phillips/>
7. **A behavioral perspective on the biophysics of the light-dependent magnetic compass: a link between directional and spatial perception?** Phillips, J. Journal of Experimental Biology. June 17, 2010. <http://jeb.biologists.org/content/213/19/3247.full.pdf>
8. **Impacts of radio-frequency electromagnetic field (RF-EMF) from cell phone towers and wireless devices on biosystem and ecosystem – a review.** Sivani, S. *Biology and Medicine*. Dec 3, 2012 http://www.emrpolicy.org/regulation/united_states/Exhibit_19_EMRI_Sivani_Bio_&_Med_2012.pdf
9. **Effects of Microwaves on the Trees and Other Plants.** Balmori, A. 2003. http://www.hese-project.org/de/emf/WissenschaftForschung/Balmori_Dr._Alfonso/showDoc.php?lang=de&header=Dr.%20Balmori&file=THE%20EFFECTS%20OF%20MICROWAVES%20ON%20THE%20TREES%20AND%20OTHER%20PLANTS.html&back=../showAuthor.php?target=Balmori_Dr._Alfonso
10. **Effects of Electromagnetic Waves Emitted by Mobile Phones on**

- Germination, Root Growth, and Root Tip Cell Mitotic Division.** Pol J of Envir Studies. Nov 2010. <http://www.pjoes.com/pdf/21.1/Pol.J.Envir.Stud.Vol.21.No.1.23-29.pdf>
11. **Intercellular Communication in Plants: Evidence for an EMF-Generated Signal that Evokes Local and Systemic Transcriptional Responses in Tomato.** 2013. http://link.springer.com/chapter/10.1007/978-3-642-36470-9_16
 12. **Provocation Study using heart rate variability shows microwaves radiation from 2.4 Ghz cordless phones.** *European Journal of Oncology*. January 16, 2011. <http://www.bemri.org/publications/dect/341-provocation-study-using-heart-rate-variability-shows-microwave-radiation-from-2-4ghz-cordless-phone.html?path=>
 13. **Is newborn melatonin production influenced by magnetic fields produced by incubators?** *Early Hum Dev* 2012; 88 (8): 707 – 710 <http://www.emf-portal.de/viewer.php?aid=20376&l=e>
 14. **Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects.** Pall, ML. *J Cell Mol Med*. 2013 Aug;17(8):958-65. <http://www.ncbi.nlm.nih.gov/pubmed/23802593>
 15. **Microwave Electromagnetic Fields Act by Activating Voltage-Gated Calcium Channels: Why the Current International Safety Standards Do Not Predict Biological Hazard.** Professor Dr. Martin L. Pall <http://apps.fcc.gov/ecfs/document/view?id=7521102473>
 16. **Communication Studies UCLA.** <http://cogweb.ucla.edu/CogSci/Myelinate.html>
 17. **“Digital Dementia” on the Rise.** June 24, 2013 <http://koreajoongangdaily.joins.com/news/article/article.aspx?aid=2973527>
 18. **International Precautionary Actions EMF.** <http://ehtrust.org/international-policy-actions-on-wireless/>
 19. **Student Science Experiment Finds Plants won’t Grow near Wi-Fi Router** <http://www.globalresearch.ca/student-science-experiment-finds-plants-wont-grow-near-wi-fi-router/5336877>
 20. **American Academy of Pediatrics Letter to FCC and FDA regarding EMF Exposure Policies and Standards.** <http://www.wirelesswatchblog.org/wp-content/uploads/2011/06/FCC-13-84-09-03-2013-American-Academy-of-Pediatrics-7520941318-1.pdf>
 21. **Transient DNA damage induced by high-frequency electromagnetic fields (GSM 1.8 GHz) in the human trophoblast HTR-8/SVneo cell line evaluated with the alkaline comet assay.** *Mutat Res*. 2010 Jan 5. <http://www.ncbi.nlm.nih.gov/pubmed/19822160>
 22. **High frequency (900 MHz) low amplitude (5 V m-1) electromagnetic field: a genuine environmental stimulus that affects transcription, translation, calcium and energy charge in tomato.** Roux, D. *Planta*. 2008 Mar <http://www.ncbi.nlm.nih.gov/pubmed/18026987>
 23. **Electrosensitivity in Sweden.** <http://www.emfacts.com/2009/02/1014-electrosensitivity-in-sweden-by-olle-johansson/>
 24. **Towards Better Health: Switzerland TV Program on Electrohypersensitivity.** 2009. <http://mieuxprevenir.blogspot.com/2013/03/swiss-tv-program-from-2009-on.html>
 25. **‘Wi-fi refugees’ shelter in West Virginia mountains.** 2011. <http://www.bbc.com/news/world-us-canada-14887428>
 26. **Replication of heart rate variability provocation study with 2.4-GHz cordless phone confirms original findings.** Havas, M. *Electromagn Biol Med*. 2013 Jun <http://www.ncbi.nlm.nih.gov/pubmed/23675629>
 27. **Electromagnetic hypersensitivity: evidence for a novel neurological syndrome.** *Int J Neurosci*. 2011 Dec;121(12):670-6. <http://www.ncbi.nlm.nih.gov/pubmed/21793784>
 28. **Review: Electromagnetic hypersensitivity: Fact or Fiction?** Genius, S, Lipp, C. *Science of Total Environment*. Sept 2011. http://www.academia.edu/4125616/Genuis_EHS_paper
 29. **Electromagnetic absorption in the human head and neck for mobile telephones at 835 and 1900 MHz.** Gandi, O. *Microwave Theory and Techniques*. Vol 44, Oct, 1996. <http://ieeexplore.ieee.org/xpl/articleDetails.jsp?reload=true&arnumber=539947>
 30. **International Agency for Research on Cancer. IARC Classifies Radiofrequency Electromagnetic Fields as Possibly Carcinogenic to Humans.** http://www.iarc.fr/en/media-centre/pr/2011/pdfs/pr208_E.pdf
 31. **Alone Together.** Sherry Turkle. 2012. <http://www.npr.org/2012/10/18/163098594/in-constant-digital-contact-we-feel-alone-together>
 32. **Austrian Medical Association Guidelines- EMF and Electrohypersensitivity.** <http://freiburger-appell-2012.info/media/EMF%20Guideline%20OAK-AG%20%202012%2003%2003.pdf>
 33. **Israeli Ministry of Education issue guidelines to limit Wi-Fi in schools.** 2013 <http://www.gsma.com/publicpolicy/israeli-ministry-of-education-issue-guidelines-to-limit-wi-fi-in-schools>
 34. **France: New National Law Bans Wi-Fi in Nursery School!** <http://ehtrust.org/france-new-national-law-bans-wifi-nursery-school/>
 35. **French government bans advertising of mobiles to children.** 2009 <http://www.independent.co.uk/life-style/gadgets-and-tech/news/french-government-bans-advertising-of-mobiles-to-children-1299673.html>
 36. **Electromagnetic Fields From Mobile Phones: Health Effects on Children and Teenagers.** April 2011. http://www.magdahavas.com/wordpress/wp-content/uploads/2011/06/Russia_20110514-rncnirp_resolution.pdf
 37. **Parliament Assembly European Council EMF Microwave Policy.** 2011. <http://assembly.coe.int/Mainf.asp?link=/Documents/AdoptedText/tal1/ERES1815.htm>
 38. **Why A Leading Professor of New Media Just Banned Technology Use in Class.** Sept 25, 2014. *Washington Post*. <http://www.washingtonpost.com/blogs/answer-sheet/wp/2014/09/25/why-a-leading-professor-of-new-media-just-banned-technology-use-in-class/>
 39. **Princeton/UCLA Study: Its Time to Ban Laptops in Law School Classrooms.** Feb 5, 2015. http://taxprof.typepad.com/taxprof_blog/2015/02/princetonucla-study-.html
 40. **Brain Cancer – Mobile phone and cordless phone use and the risk for glioma – Analysis of pooled case-control studies in Sweden, 1997-2003 and 2007-2009.** Hardell. *Pathophysiology*. 2014 Oct 29. <http://www.ncbi.nlm.nih.gov/pubmed/25466607>
 41. **Brain Cancer – Mobile phone use and brain tumors in the CERENAT case-control study.** Coureau G. *Occup. Environ Med*. 2014 Jul; <http://www.ncbi.nlm.nih.gov/pubmed/24816517>



STEPHEN T. SINATRA M.D., F.A.C.C.

F.A.C.N., C.N.S., C.B.T.,
Integrative Metabolic Cardiology

July 16, 2014

Chairman and Trustees
Fay School
48 Main Street
Southborough, MA 01772

RE: Wi-Fi in Schools

Dear Chairman and Trustees:

I am writing this letter on behalf of concerned parents of children who are attending schools with Wi-Fi technology. I'm a cardiologist and co-founder of Doctors for Safer Schools, an organization dedicated to informing teachers, parents and superintendents about the uncertainty and possible environmental health hazards of Wi-Fi technologies.

The heart is a delicate and complex electromagnetic organ that can be adversely affected by exogenous signals from wireless technology and microwave radiation. For this reason it is unwise to expose students and teachers to Wi-Fi radiation for internet access, especially when safer alternative wired options are available. Children are particularly vulnerable to this radiation and the incidents of cardiovascular events including sudden cardiac arrest, seems to be increasing, especially among young athletes (up to the age of 19). In some cases this is due to undetected heart defects, blunt trauma to the heart in contact sports, and heat stress during strenuous exercise, but in instances these irregularities may be exacerbated by or due to microwave signals interfering with the autonomic nervous system that regulates the heart.

I know this because I am a board certified cardiologist and have been a Fellow of the American College of Cardiology since 1977. At the Manchester Memorial Hospital in Connecticut, I served in several roles, including Chief of Cardiology, Director of Cardiac Rehabilitation, and Director of Medical Education.

In both Canada and the United States a large number of students are complaining that they feel unwell in classrooms that have Wi-Fi technology. These complaints have been investigated and what emerges is the following:

1. Symptoms common among these students include headaches, dizziness, nausea, feeling faint, pulsing sensations or pressure in the head, chest pain or pressure, difficulty

concentrating, weakness, fatigue, and a racing or irregular heart accompanied by feelings of anxiety. These symptoms may seem diverse but they indicate autonomic dystonia or dysfunction of the autonomic nervous system.


2. Symptoms do not appear in parts of the school that do not have this technology (Wi-Fi-free portables) and they do not appear in homes that do not have wireless technology.
3. We know that the heart is sensitive to and can be adversely affected by the same frequency used for Wi-Fi (2.4 GHz) at levels a fraction of federal guidelines (less than 1%) and at levels that have been recorded in two Ontario schools with Wi-Fi technology.
4. The incidence of sudden cardiac arrests (SCA) among young athletes is increasing and doctors don't know why. In one small Ontario community, the number of students experiencing SCA is disturbingly high. Whether WiFi and nearby cell phone antennas exacerbate SCA needs to be investigated further before students are subjected to these fields.

In conclusion it is unwise to install wireless technology (WiFi) in schools. We do not know what the long-term effects of low-level microwave radiation are on students and teachers. The safety of this technology on children has not been tested and I would advise that you follow the precautionary principle that states the following:

"In order to protect the environment, the precautionary approach shall be widely applied by States according to their capabilities. Where there are threats of serious or irreversible damage, lack of full scientific certainty shall not be used as a reason for postponing cost-effective measures to prevent environmental degradation."
(Rio Conference 1992).

The principle implies that we have a social responsibility to protect the public from exposure to harm, when scientific investigations have found a plausible risk. That "plausible risk" exists for microwave radiation at very low levels. These protections can be relaxed only if further scientific findings emerge that provide sound evidence that no harm will result. In some legal systems the application of the precautionary principle has been made a statutory requirement.

Sincerely,

A handwritten signature in black ink, reading "Stephen T. Sinatra". The signature is fluid and cursive, with the first name "Stephen" and last name "Sinatra" clearly legible.

Stephen T. Sinatra, M.D., F.A.C.C., F.A.C.N., C.N.S



Karolinska Institutet
Department of Neuroscience
Experimental Dermatology Unit

Stockholm, July 24, 2014

Mr. Thomas McKean, President, Board of Trustees
Mr. James Shay, President-Elect, Board of Trustees
Fay School
48 Main Street
Southborough, MA 01772

Ladies and Gentlemen,

It has been brought to my attention that children in your school are physically being impacted by radiation from WiFi antennas, and that some of the student's reactions have been severe. I was concerned to learn this. It is unwise to chronically expose children to this type of radiation, as their bodies are more sensitive than adults and the radiation has been shown to impair not just physiological functioning but cognitive function and learning.

Radiation of the kind emitted by WiFi transmitters impacts attention, memory, perception, learning capacity, energy, emotions and social skills. There is also diminished reaction time, decreased motor function, increased distraction, hyperactivity, and inability to focus on complex and long-term tasks. In some situations, children experience cardiac difficulties. In one Canadian school district, incidence of cardiac arrest in children was 40x the expected rate, and defibrillators have had to be placed at each school. Online time, particularly multi-tasking in young children, has been linked with a chronically distracted view of the world preventing learning critical social, emotional and relational skills. There is a physiological as well as psychological addiction taking place. I am sure, that as stewards of the lives of the children in your charge, you would not wish any of these outcomes.

Given the large and growing body of science indicating biological and health effects from the radiation emitted by antennas, it would be most imprudent at this time to permit wireless antennas on—or inside—your property. Understand the FCC exposure guidelines only protect against the acute power density, or acute thermal, effects, and they do nothing to protect against the other aspects of the radiation's risk, such the frequencies, amplitude, pulsing, intensity, polarity and biologically disruptive information content. Thus, until the FCC establishes guidelines for the non-thermal effects, any reliance by your school on current FCC guidelines, based solely on *thermal effects* would necessarily be incomplete. I urge a school of your caliber to be a leader on this issue, and appreciate that two wrongs do not make a right.

I enclose for your review the transcript of the Seletun Scientific Statement laying out the key concerns on this topic. If I can be of further help, please, do not hesitate to be in touch.

Yours truly,

Olle Johansson, Associate Professor
The Experimental Dermatology Unit,
Department of Neuroscience,
Karolinska Institute, 171 77 Stockholm, Sweden

Mailing address
Experimental Dermatology Unit
Department of Neuroscience
Karolinska Institutet
171 77 Stockholm
Sweden

Visiting address
Retziuslaboratoriet
Retzius väg 8
Solna

Direct	468-52 48 70 58
Switchboard	468-52 48 64 00
Fax	468-30 39 04
Fax (KI)	468-31 11 01

From: jmm@berkeley.edu

To: john.sterritt@lausd.net, monica.garcia@lausd.net,
marguerite.lamotte@lausd.net, tamar.galatzan@lausd.net,
steve.zimmer@lausd.net, sarah.bradshaw@lausd.net, nury.martinez@lausd.net,
richard.vladovic@lausd.net, enrique.boullt@lausd.net, pta31dist@aol.com,
ronald.chandler@lausd.net, lhc8767@lausd.net, bcohen@lausd.net,
superintendent@lausd.net, john.deasy@lausd.net, tim.delia@lausd.net,
senglish@advanceproj.com, wfletcher@utla.net, smfolsom@aol.com,
bforrester@utla.net, mark.hovatter@lausd.net, Daniel.hwang@lausd.net,
ainouye@utla.net, michelle.king@lausd.net, dlyell@utla.net,
yolanda.pujol@lausd.net, Irojas@lausd.net, azayas@SEIU99.org

CC: cheemf@lists.healthandenvironment.org

Sent: 2/8/2013 2:21:54 P.M. Pacific Standard Time

Subj: [cheemf] Adoption of Wi-Fi in Los Angeles USD classrooms

TO: Los Angeles Unified School District (LAUSD)

FROM: Joel M. Moskowitz, Ph.D.

Director, Center for Family and Community Health

School of Public Health

University of California, Berkeley

RE: Adoption of Wi-Fi in Classrooms

DATE: February 8, 2013

Based upon my review of the research of the health effects associated with exposure to radiofrequency (RF) electromagnetic radiation (EMR), especially microwave radiation, I feel compelled to register my concern that adoption of Wi-Fi in LAUSD classrooms is likely to put at risk the health of many students and employees in the District.

In December, Dr. Gayle Nicoll of URS Corporation asked me to serve as an expert reviewer for a report that URS prepared for the LAUSD regarding the adoption of Wi-Fi in classrooms. Since Ms. Nicoll could not assure me that URS has no conflicts of interest, I turned down her request and sent her references to recent studies about Wi-Fi radiation. I cc:ed Board members and key staff as I was concerned about the health risks of unnecessarily subjecting 660,000 children to 13,000 hours of Wi-Fi microwave radiation during their K-12 school years.

Although I have not seen the URS report, I imagine it is based on the FCC's outmoded 1996 safety standards which only protect the public from the **thermal risk of RF EMR exposure** (i.e., from heating of tissue). For the past three years, in numerous media interviews I have been calling on the FCC to strengthen its standards and testing procedures to protect the public and workers from the low-intensity, **non-thermal risks of RF EMR exposure** that have been reported in hundreds, if not thousands, of research studies. These include increased risk of neurological and cardiovascular problems, sperm damage and male infertility, reproductive health risks, and cancer.

The **precautionary principle** should be applied to this critical policy decision. This principle, developed at a U.N. environmental conference in 1992 states that in the absence of scientific consensus if an action has a suspected risk of causing harm, the burden of proof it is not harmful falls on those taking the action, and all reasonable measures to reduce the risk must be taken.

Internet access can be provided to students through wires or optical fiber without installing Wi-Fi in the classrooms.

For further information, please see my **Electromagnetic Radiation Safety web site** at <http://saferemr.blogspot.com> where I have archived news releases and links to recent reports by major scientific groups and political agencies.

Sincerely,

Joel M. Moskowitz, Ph.D.

=====
Joel M. Moskowitz, Ph.D.
Director
Center for Family and Community Health
The UC Berkeley Prevention Research Center
School of Public Health
University of California, Berkeley
50 University Hall
Berkeley, CA 94720-7360

Phone: 510-643-7314

E-mail: jmm@berkeley.edu

CFCH Web Site: <http://cfch.berkeley.edu>

EMR Safety Web Site: <http://saferemr.blogspot.com>

December 1, 2015

Montgomery County Schools
Carver Educational Services Center
850 Hungerford Drive
Rockville, MD 20850

Attention: Dr. Andrew Zuckerman, Chief Operating Officer
MCPS Board of Education Members

This letter of comment has been prepared after reviewing the *Montgomery County Public Schools Radiofrequency (RF) Summary Monitoring Report* dated July 2015 produced by AECOM Environment.

1) The instrument cited as being used for the peak measurements in section 7, a Narda SRM-3006, is not suitable to measure the very short (1 millisecond) spikes typically found in WiFi 802.11n communication. As stated on page 7-1, each data sweep takes 550 milliseconds, making the instrument unsuitable for reliably logging the short bursts typical in 802.11n WiFi communications. Palit et al conclude that 50% of the uplink traffic will be in bursts shorter than 2 milliseconds. The peak levels of those packets will not be reliably logged by a device with a 550 millisecond sweep time.

Palit et al, 2012. Anatomy of WiFi Access Traffic of Smartphones and Implications for Energy Saving Techniques. International Journal of Energy, Information and Communications, Vol. 3, Issue 1.

2) Even the average-level tests seem inconsistent with engineering reality. Figure 7.1 shows a background noise level mostly flat between 2.4GHz and 5.8Ghz. That noise (typically -70dBm) is generally consistent with the internal thermal noise in a quality wide-band measuring instrument. Two tiny peaks out of that noise are represented to be the "average electric field generated at one foot away from an AP in use at Beverly Farms Elementary School." Even with just the 802.11n beacon-frame idling, the peak field a foot away from an access point should be a million times higher than the levels of figure 7.1. Why do we just see a blip on the chart? Clearly some unusual 'averaging' has occurred, yet the parameters of that averaging, and the potential clinical implications of that averaging, are not noted in the annotation to the Figures. Further, Figure 7.2 shows a background noise level some 10dB higher than figure 7.1, something that would be very unusual in measurements at these Gigahertz frequencies.

3) The RF exposure estimates are additionally inadequate because, in reality, there is no way to meet the distancing that AECOM's report bases its measurements on for an individual student. In normal use, kids hover over devices. They hug them to the body. They put them in their laps at lunchtime, on the couch and in bed doing homework. It is entirely unrealistic to expect teachers and parents to guarantee that students always keep their Chromebooks at some arbitrary distance during use.

4) The report concludes with classroom RF measurement comparisons to an outdated 2007 BioInitiative Report recommendation of 0.1 uW/cm². (Section 7). Graphics need to be re-drawn with comparisons to the 2012 recommended BioInitiative level, and do so not only for a 12” spacing, but also for the one-inch distance measured from the Chromebook (Figure 7-3 and 7-4). Using an arbitrary 12” distance to report and compare to either the 2007 or 2012 BioInitiative recommendations will seriously underestimate RF exposures since students don’t always (or even typically) maintain a foot of distance. Their ‘leaning in’ and having to place their faces close to the device is common usage, and is unavoidable.

5) The methodology is not specific as to the number of operating devices and clustering of students at work – which is necessary to characterize exposures from a room full of operational wireless devices. Figure 2.1 shows multiple wireless devices connected to one wireless router. Measuring one or several Chromebooks rather than one Chromebook for each of the 25-35 students plus router isn't how a normal classroom operates. It **does not** produce RF measurements of a typical class using many wireless devices at once, so this is a fundamental flaw. It will underestimate RF exposures.

6) There is also a comment to be made here about the setup – how does this methodology reasonably reflect how smaller or younger children with short arms and torsos actually use tablets? What RF exposures they can expect to receive? The likely consequence to the measurements is greater exposure. Unless the students are using chopsticks instead of their fingers, or are using wired keyboards that increase the distance to the wireless device, RF exposures will be worse for the younger or smaller-stature students.

7) This Report appears to legitimize MCSD’s use of wireless in the classroom by asserting compliance with the 2007 BioInitiative Report recommendation, yet the report does not mention the significant revision of that threshold in the years between 2007 and 2012. Both BioInitiative Reports clearly state that their recommendations are interim and ‘that they may have to go lower.’ Recent studies of students reporting headache, irritability, concentration and behavior problems at levels as low as 0.003-0.006 uW/cm², indicate that neither BioInitiative Report threshold may be low enough to assure safety. As the co-editor of the BioInitiative Reports, and a founding member of the BioInitiative Working Group, the way in which our work has been invoked is not consistent with the findings of the BioInitiative Reports overall. The conclusions of this report cannot be said to give a positive assertion of safety because of the degree of uncertainty over whether the testing equipment was adequate (we believe it was not); the lack of comparison data; and the failure to measure RF exposures at realistic distances from the student(s).

8) Correct BioInitiative citations are:

BioInitiative Working Group, Cindy Sage and David O. Carpenter, Editors. BioInitiative Report: A Rationale for Biologically-based Public Exposure Standards for Electromagnetic Radiation at

www.bioinitiative.org, December 31, 2012.

BioInitiative Working Group, Cindy Sage and David O. Carpenter, Editors. BioInitiative Report: A Rationale for a Biologically-based Public Exposure Standard for Electromagnetic Fields (ELF and RF) at www.bioinitiative.org, August 31, 2007

CONCLUSION

The data in this report cannot therefore be used to infer safety, or lack of safety, of children in any of the tested locations.

Respectfully submitted,

Cindy Sage. MA
Sage Associates
Co-Editor, BioInitiative 2007 and 2012 Reports
sage@silcom.com

Prof. Trevor Marshall, PhD
Director, Autoimmunity Research Foundation,
Senior Member IEEE,
Founding chair (retired) IEEE EMBS (Buenaventura Chapter)
Fellow, European Association for Predictive, Preventive and Personalised Medicine (Brussels)
International Expert Council, Community of Practice: Preventative Medicine (Moscow)
trevor.m@trevormarshall.com



September 22, 2014

On behalf of the BioInitiative Working Group, we are writing to express our concern about the views expressed by CEOs from Google, Dell, Apple, Adobe, eBay, Facebook, the George Lucas Educational Foundation and others to the FCC supporting wireless technologies in schools.

Your letter to the FCC dated July 7, 2014 titled Education Superhighway, states:

*"Today, we are writing to you to urge swift bi-partisan action at your July 11, 2014 meeting to adopt the E-Rate modernization proposal set forth by Chairman Wheeler."
"By responsibly investing \$2 billion of unused funds and providing predictable ongoing support for Wi-Fi, the plan will make dramatic progress in bringing high-speed connectivity to our classrooms."*

No one denies that bringing high-speed connectivity to our classrooms is important. But it can be a wired connection and does not have to be WiFi. It does not reflect well on the ethics of your corporations to encourage the FCC to provide \$2 billion dollars for new wireless classroom infrastructure and devices for school children, knowing that wireless emissions have been classified as a Possible Human Carcinogen by the World Health Organization's International Agency for Research on Cancer (2011). To promote wireless technologies in schools is to deliberately and knowingly disregard current health warnings from international science and public health experts.

Saturating schools with wireless technology will likely create unnecessary liability for municipalities and result in a loss of public trust and confidence in the corporations that push their wireless products with a blind eye toward health concerns.

Epidemiological studies show links between radiofrequency radiation (RFR) exposure and cancers, neurological disorders, hormonal changes, symptoms of electrical hypersensitivity (EHS) and more. Laboratory studies show that RFR exposure increases risk of cancer, abnormal sperm, learning and memory deficits, and heart irregularities. Fetal exposures in both animal and human studies result in altered brain development in the young offspring, with disruption in learning, memory and behavior. The brain development of a fetus can be impaired by in-utero exposure to a pregnant woman. The evidence for these statements is based on hundreds of published, peer-reviewed scientific studies that report adverse effects at levels much lower than current FCC public safety limits. WiFi in schools, in contrast to wired internet connections, will increase risk of neurologic impairment and long-term risk of cancer in students. Corporations cannot avoid responsibility simply by asserting compliance with existing legal, but outdated and inadequate FCC public safety limits.

Today, corporations that deal with educational technology should be looking forward and helping school administrators and municipal leaders to access safe, wired solutions. Your corporations can reasonably foresee and offer alternatives to potentially hazardous exposures to wireless radiation by choosing to support wired educational technologies.



Thank you for your attention to this letter.

Cindy Sage, MA, Tel: (805) 969-0557 Email: sage@silcom.com
David O. Carpenter, MD, Tel: 518-525-2660 Email: dcarpenter@albany.edu
Co-Editors, BioInitiative 2012 Report
For the BioInitiative Working Group

Copies: CEOs signing Education Superhighway letter to the FCC
Federal Communications Commission
The White House, President Obama
US Secretary of Education Secretary Arne Duncan

Contributing Authors of the the 2007 and 2012 BioInitiative Working Groups

Jitendra Behari, PhD, India
Carlo V. Bellieni, MD, Italy
Igor Belyaev, Dr.Sc., Slovak Republic
Carl F. Blackman, PhD, USA
Martin Blank, PhD, USA
Michael Carlberg, MSc, Sweden
David O Carpenter, MD, USA
Zoreh Davanipour, DVM, PhD USA
Adamantia F. Fragopoulou, PhD, Greece
David Gee, Denmark
Yuri Grigoriev, MD, Russia
Kjell Hansson Mild, PhD, Sweden
Lennart Hardell, MD, PhD, Sweden
Martha Herbert, PhD, MD, USA
Paul Héroux, PhD, Canada
Michael Kundi, PhD, Austria
Henry Lai, PhD, USA
Ying Li, PhD, Canada
Abraham R. Liboff, PhD, USA
Lukas H. Margaritis, PhD, Greece
Henrietta Nittby, MD, PhD, Sweden
Gerd Oberfeld, MD, Austria
Bertil R. Persson, PhD, MD, Sweden
Iole Pinto, PhD, Italy
Paulraj Rajamani, PhD, India
Cindy Sage, MA, USA
Leif Salford, MD, PhD, Sweden
Eugene Sobel, PhD, USA
Amy Thomsen, MPH, MSPAS, USA



American Academy of Environmental Medicine

6505 E Central • Ste 296 • Wichita, KS 67206

Tel: (316) 684-5500 • Fax: (316) 684-5709

www.aaemonline.org

May 13, 2013

Executive Committee

President

Amy L. Dean, D.O., FAAEM
1955 Pauline Blvd Ste 100D
Ann Arbor, MI 48103

President-Elect

Janette Hope, M.D., FAAEM
304 W Los Olivos
Santa Barbara, CA 93105

Secretary

Jennifer Armstrong, M.D., FAAEM
3364 Carling Ave.
Ottawa, Ontario, Canada

Treasurer

Richard G. Jaeckle, M.D., FAAEM
8220 Walnut Hill Ln Ste 404
Dallas, TX 75231

Immediate Past President

A.L. Barrier, M.D., FAAO-HNS

Advisor

William J. Rea, M.D., FAAEM
Gary R. Oberg, M.D., FAAEM

Board of Directors

Craig Bass, M.D.
Robin Bernhoft, M.D., FAAEM
Martha Grout, M.D., MD(H)
W. Alan Ingram, M.D.
Derek Lang, D.O.
Allan D. Lieberman, M.D., FAAEM
Lisa Nagy, M.D.
Kalpana D. Patel, M.D., FAAEM

Continuing Medical Education

Chair

James W. Willoughby, II, D.O.
24 Main St.
Liberty, MO 64068

Co-Chair

Wm. Alan Ingram, M.D.
18015 Oak St Ste B
Omaha, NE 68130

Open Letter to the Superintendents
of the School Districts of the United States

The American Academy of Environmental Medicine (AAEM) strongly supports the use of wired Internet connections.

The AAEM comprises Medical Doctors, Osteopaths, and PhD researchers focusing on the effects of environmental agents on human health. For forty years the Academy has trained Physicians to treat the most difficult patients who are often overlooked by our medical system, because the cause of their illness, rather than being caused by an infection or traditionally understood cause, is related to more basic underlying causes such as chemical, toxic metal, food or radiation exposures.

In May 2011 the World Health Organization elevated exposure to wireless radiation, including WiFi, into the Class 2b list of Carcinogens.

There is consistent emerging science that shows people, especially children who are more vulnerable due to developing brains, and thinner skulls, are affected by the increasing exposure to wireless radiation. In September 2010, the Journal of the American Society for Reproductive Medicine-Fertility and Sterility, reported that only four hours of exposure to a standard laptop using WiFi caused DNA damage to human sperm.

In December 2012 the American Academy of Pediatrics- representing 60,000 pediatricians, wrote to Congress requesting it update the safety levels of microwave radiation exposure especially for children and pregnant women.

In a school setting, children are exposed to WiFi for an unprecedented period of time, for their entire childhood. Some of these signals will be much more powerful than is received at home, due to the need for the signals to go through walls, and serve multiple computers simultaneously. The school signals are dozens of times more powerful than the café and restaurant systems.

To install this system in your school district risks a widespread public health hazard that the medical system is not yet prepared to address. Statistics show that you can expect to see an immediate reaction in 3% and delayed effects in 30%, including teachers.

It is better to exercise caution and substitute with a safe alternate such as a wired connection, which is not classified as a possible Carcinogen. While more research is being conducted children must be protected. Wired technology is not only safer, it also stronger and more secure.

While the debate ensues about the dangers of WiFi, cell phone towers and cell phones, it is the doctors who must deal with the after affects. Until we can determine why some get sick and others do not, and some are debilitated for indeterminate amounts of time, we implore you to not take the risk, with the health of so many children who have entrusted you to keep them safe while at school.

Respectfully,

The Executive Committee of the American Academy of Environmental Medicine

Message to Schools and Colleges about Wireless Devices and Health

If wireless devices, such as Wi-Fi, are used in your schools and colleges, then the health of your students, your faculty, and your staff can be at risk. This is a difficult problem but an addressable one if you act.

Background: Wireless devices transmit information using radiofrequency/microwave radiation. The international biomedical research community has been studying the biological impact of such radiation for decades, but more intensely in recent years. Thousands of peer-reviewed studies published in biomedical research journals have contributed to our understanding of this impact. So many serious biological effects have been found that immediate responsive action is warranted. Further, these biological effects are occurring at levels of radiation far lower than earlier understood. Simply stated, a worldwide health crisis is emerging and is becoming a hallmark of the 21st Century. The international biomedical research community is trying to warn us; but we, in the USA, are not yet listening. I hope this message will help to change that.

As a scientist, I urge you to look into the **health impact of the radiofrequency/microwave radiation** produced by wireless devices. Examples of wireless devices of concern in our environment are Wi-Fi in all of its forms; cell phones and cell towers (especially those located on school grounds); cordless phones; wireless computers, whether desktop, laptop, or tablet versions; wireless baby monitors; wireless smart electricity meters; emerging wireless smart appliances; and microwave ovens (because they always leak radiation).

This crisis is the consequence of many factors. Here are some of them:

- All living things are bioelectrical in nature. That is why electrocardiograms and electroencephalograms work. They, of course, measure the tiny electrical signals that operate the heart and the brain. The critical tasks performed by these tiny electrical signals, and so many other electrical signals in all living things, can be disrupted by radiofrequency/microwave radiation.
- The levels of manmade radiofrequency/microwave radiation in our environment are increasing exponentially and already exceed, by many orders of magnitude, the levels at which all life on Earth evolved. Simply stated, we are drowning in a rising sea of manmade radiofrequency/microwave radiation.
- The invisible nature of radiofrequency/microwave radiation leaves the public and the decision-makers unaware of the rising levels of radiation around them.
- The genuine usefulness of wireless devices promotes denial of the risks.
- The intense advertising, the economic power, and the political power of profitable wireless industries enable them to dominate the public dialogue and to hold sway over government regulators and legislators.
- Current Federal standards for limiting the exposure of the public to radiofrequency/microwave radiation are outdated and overly permissive. Those standards are based on thermal heating alone. In effect, the Government claims that if you are not cooked too much by the radiation, then you are fine. Those Federal standards ignore the many biological effects that occur at much lower levels of radiation, leaving the public unprotected.
- Federal and state governments are advocating unlimited expansion of wireless technology, and are even co-funding such expansion and mandating the acceptance of wireless technology by the public. Such

actions reflect a widespread lack of understanding of, or willful blindness to, the underlying science and its consequences for public health.

- Some of the more serious consequences of exposure to radiofrequency/microwave radiation (such as DNA damage, cancer, and infertility) are especially nefarious because they give no early warning signs.
- Other consequences of exposure do give early warning signs (such as sleep disruption, headaches, fatigue, ringing in the ears, memory loss, dizziness, heart arrhythmia, and many others); but those signs are too often dismissed because they can have other causes as well, complicating identification of the true cause.
- The absence of routine training of physicians in the biological effects of radiofrequency/microwave radiation makes it difficult for physicians to identify the causes and to provide responsive guidance.
- Even aware individuals cannot control their exposure in any environment shared with others, because the radiation around them, much like second-hand smoke, is forced on them by unaware individuals. Only governments can fully solve this problem, but they are currently part of the problem. For now the public will have to protect itself, and that will require public education and action.

Fortunately, many of the services that wireless devices offer can be realized with much safer wired devices. The wired devices achieve connectivity with fiber-optic, coaxial, or Ethernet cables. The wired devices are faster, more reliable, and more cyber secure. They are, however, less mobile, often less convenient, and somewhat more expensive to install. But those drawbacks pale in comparison to the benefits of good health.

Simply stated, schools and colleges can protect their students, staff, and faculty from the health risks posed by wireless devices, including Wi-Fi, by converting to safe wired connectivity. If your institution lacks the resources to convert now, do consider shutting down your wireless devices anyway and converting as soon as you can. You can advance learning without leaving a trail of illness behind you, some of which can be lifelong.

As a suggested starting place for exploring the concerns about the radiation from wireless devices, I have appended an “Annotated List of References” and an “Annotated List of Videos”. Please view, especially, video (1) called “Wi-Fi in Schools, the Facts”, made in Australia, on page 6.

Regards,

Ronald M. Powell, Ph.D.
20316 Highland Hall Drive
Montgomery Village, MD 20886-4007
Telephone: 301-926-7568
Email: ronpowell@verizon.net

My background

I am a retired U.S. Government scientist (Ph.D., Applied Physics, Harvard University, 1975). During my Government career, I worked for the Executive Office of the President, the National Science Foundation, and the National Institute of Standards and Technology. For those organizations, respectively, I addressed Federal research and development program evaluation, energy policy research, and measurement development in support of the electronics and electrical-equipment industries and the biomedical research community. I currently interact with other scientists and with physicians around the world on the impact of the environment – including the radiofrequency/microwave environment – on human health.

ANNOTATED LIST OF REFERENCES

The international biomedical research community has conducted thousands of studies seeking to identify the biological effects of exposure to both low frequency and radiofrequency electromagnetic fields, extending into the microwave region. So many serious biological effects have been found from such fields, at levels earlier thought to be low enough to be safe, that immediate action is needed to alert and protect the public.

The most massive review of this biomedical literature is the 1479-page BioInitiative 2012 Report which considered about **1800** biomedical research publications, most issued in the previous five years. The BioInitiative 2012 Report was prepared by an international body of 29 experts, heavy in Ph.D.s and M.D.s, from 10 countries, including the USA which contributed the most experts (10). The review concludes that "The continued rollout of wireless technologies and devices puts global public health at risk from unrestricted wireless commerce unless new, and far lower[,] exposure limits and strong precautionary warnings for their use are implemented."

BioInitiative Working Group, Cindy Sage, M.A. and David O. Carpenter, M.D., Editors, BioInitiative Report: A Rationale for Biologically-based Public Exposure Standards for Electromagnetic Radiation, December 31, 2012

<http://www.bioinitiative.org>

A group of six doctors in Oregon, led by Paul Dart, M.D., released, in June 2013, a 74-page review of **279** biomedical research publications. This review makes the health case against "cell phones, base stations, Wi-Fi, Smart Meters and other RF [*radiofrequency*] or ELF [*extremely low frequency*] -emitting devices". The review notes that "The current levels of exposure need to be reduced rather than increased further. The FCC [*Federal Communications Commission*] must especially protect vulnerable groups in the population including children and teenagers, pregnant women, men of reproductive age, individuals with compromised immune systems, seniors, and workers." This review is posted on the website of the FCC at the link entitled "Health Effects of RF - Research Review (87)".

Biological and Health Effects of Microwave Radio Frequency Transmissions, A Review of the Research Literature, A Report to the Staff and Directors of the Eugene Water and Electric Board, June 4, 2013

<http://apps.fcc.gov/ecfs/comment/view?id=6017465430>

Michael Bevington, in 2013, published a book that summarizes the findings of **1828** international biomedical research publications. The book describes the symptoms caused by exposure to electromagnetic radiation, the many diseases associated with such exposure, and the relative risk levels associated with specific sources of electromagnetic radiation. The citations of papers include the PMID index numbers for easy location on the PubMed.gov website of the National Institutes of Health. This website provides the largest index to the biomedical research literature in the world.

Electromagnetic Sensitivity and Electromagnetic Hypersensitivity: A Summary by Michael Bevington
NEW EDITION: March 2013

<http://www.es-uk.info>

About 200 scientists from 39 countries around the world submitted an international appeal to the United Nations and to the World Health Organization in May 2015. These scientists seek improved protection of the public from harm from the radiation produced by many wireless sources, including "cellular and cordless phones and their base stations, Wi-Fi, broadcast antennas, smart meters, and baby monitors" among others.

Together, these scientists have published over 2000 peer-reviewed research papers on this subject.

<https://www.emfscientist.org/index.php/emf-scientist-appeal>

The International Agency for Research on Cancer, of the World Health Organization, has already classified radiofrequency electromagnetic fields as a Class 2B carcinogen ("possible carcinogen"), based primarily on the increased risk of brain cancer. That decision was made in 2011. Since then, the research supporting a higher classification of risk ("probable carcinogen", or even "known carcinogen") has continued to build.

http://www.iarc.fr/en/media-centre/pr/2011/pdfs/pr208_E.pdf

The American Academy of Environmental Medicine (AAEM), which trains physicians in preparation for Board Certification in Environmental Medicine, states: "The AAEM strongly supports the use of wired Internet connections, and encourages avoidance of radiofrequency such as from WiFi, cellular and mobile phones and towers, and 'smart meters'." AAEM further states that "The peer reviewed, scientific literature demonstrates the correlation between RF [*radiofrequency*] exposure and neurological, cardiac, and pulmonary disease as well as reproductive and developmental disorders, immune dysfunction, cancer and other health conditions. The evidence is irrefutable." The AAEM concludes: "To install WiFi in schools plus public spaces risks a widespread public health hazard that the medical system is not yet prepared to address."

AAEM, Wireless Radiofrequency Radiation in Schools, November 14, 2013

<http://www.aaemonline.org/pdf/WiredSchools.pdf>

The American Academy of Pediatrics (AAP), whose 60,000 doctors care for our children, supports the development of more restrictive standards for radiofrequency radiation exposure that would better protect the public, particularly the children. The AAP, in a letter to the Federal Communications Commission (FCC) and the Food and Drug Administration (FDA), dated August 29, 2013, states that "Children are not little adults and are disproportionately impacted by all environmental exposures, including cell phone radiation. Current FCC standards do not account for the unique vulnerability and use patterns specific to pregnant women and children. It is essential that any new standard for cell phones or other wireless devices be based on protecting the youngest and most vulnerable populations to ensure they are safeguarded throughout their lifetimes."

<http://apps.fcc.gov/ecfs/document/view?id=7520941318>

The U.S. Government bears a major responsibility for the exponential growth in the levels of radiation from wireless devices in the environment. In 1996, the U.S. Congress passed, and the President signed, the Telecommunications Act of 1996. Under pressure from the cell phone industries, this law included this provision: "No State or local government or instrumentality thereof may regulate the placement, construction, and modification of personal wireless service facilities [*cell towers*] on the basis of the environmental effects of radio frequency emissions to the extent that such facilities comply with the [*Federal Communications*] Commission's regulations concerning such emissions." Because the Federal Communications Commission's regulations on radiation exposure are so permissive, this provision prevents state and local governments from protecting their people from radiation from cell towers, based on health concerns.

Telecommunications Act of 1996

<https://transition.fcc.gov/Reports/tcom1996.pdf>

The Federal Communications Commission (FCC) has acted in partnership with the wireless industries by permitting wireless radiation levels far higher than the biomedical research literature indicates are necessary to protect human health. The success of the wireless industries in capturing the FCC, the committees in the U.S. Congress that oversee the FCC, and the Executive Branch is detailed in a new monograph from the Center for Ethics at Harvard University. As an example of that capture, the President recently appointed, as head of the FCC, the former head of the CTIA – The Wireless Association, which is the major lobbying organization for the wireless industry. This, of course, is the infamous "revolving door".

Norm Alster, Captured Agency: How the Federal Communications Commission is Dominated by the Industries It Presumably Regulates (2015)

<http://ethics.harvard.edu/news/new-e-books-edmond-j-safraresearch-lab>

Further, the U.S. Government's "American Recovery and Investment Act of 2009" provided funding that was used to motivate the installation of wireless smart meters (also called the "Advanced Metering Infrastructure" or "AMI") by offering cost sharing, in the form of grants, to the utilities that would adopt such meters.

https://www.smartgrid.gov/recovery_act/overview/smart_grid_investment_grant_program.html

Many states then extended the impact of the above Act by *mandating* the acceptance of wireless smart meters by the public. These meters contain microwave transmitters/receivers and are placed either on, or inside, every home and many businesses. A California court-ordered document indicates that each smart meter broadcasts bursts of radiation, on average about 10,000 times per day and up to a maximum of about 190,000 times per day. Such bursts flood neighborhoods with radiation, day and night, throughout the year.

http://emfsafetynetwork.org/wp-content/uploads/2011/11/PGERFDataOpt-outalternatives_11-1-11-3pm.pdf

Increasingly, the public is becoming aware of the threat that wireless radiation poses to health. The initial opposition focuses primarily on *mandated* sources of exposure, especially when the individuals exposed include the unborn and young children as they are among the most vulnerable. Thus, the strongest initial opposition is surfacing for cell towers, especially on school grounds; for Wi-Fi in schools and colleges; and for wireless smart meters placed on, or inside, homes and businesses. Most states now have opposition groups, and some states have even 10 or 20 such groups. These groups are pursuing relief through state regulatory bodies, through state legislatures, and through the courts. Below is a sampling of the hundreds of U.S. websites that reflect the nature and scope of the opposition to the unbridled expansion of wireless technology. Such websites seek to educate the public and decision-makers, and thus to promote responsive action, based on the underlying science.

The BabySafe Project

<http://www.babysafeproject.org/the-science/>

National Association for Children and Safe Technology

<http://www.nacst.org/>

Stop Smart Meter's listing of groups in the USA and other countries opposed to wireless smart meters

<http://stopsmartmeters.org/frequently-asked-questions/contacts-database/>

Smart Grid Awareness, a Website by SkyVision Solutions, Consumer Protection Advocate

<http://smartgridawareness.org>

ANNOTATED LIST OF VIDEOS

There are hundreds of videos on the Internet that address the impact of wireless radiation on health. Here are just a few that provide an especially good introduction to this topic. An Internet search will surface many more.

(1) An introduction to the health risks posed by Wi-Fi in schools

Wi-Fi in Schools, the Facts (September 9, 2013) (18 minutes)

Produced by Wi-Fi in Schools Australia.

<https://www.youtube.com/watch?v=QQryZbXlqXI&feature=youtu.be>

(2) Wide ranging overview of the impact of electromagnetic radiation on human health, particularly at microwave frequencies, with a special emphasis on children and the school environment

Electromagnetic Radiation Health for Children 2014 (70 minutes)

Presented by Dr. Erica Mallery-Blythe, a UK physician.

<https://www.youtube.com/watch?v=sNFdZVeXw7M>

(3) Documentary on the wireless industry's efforts to suppress public awareness of the health effects of wireless radiation

Microwaves, Science & Lies (2014) (90 minutes)

Produced by Jean Heches and Nancy de Meritens of France.

<https://vimeo.com/ondemand/17755/89417454>

(4) Samples of video testimony by individuals harmed by the radiation from wireless devices

Cell Phones Cause Cancer (October 17, 2012) (9 minutes)

Presented by Jimmy Gonzalez, Esq.

<https://www.youtube.com/watch?v=DII0VJd0IA8>

Woman suffers acute radiation exposure from a bank of smart meters (January 21, 2015) (3 minutes).

Produced by Maryland Smart Meter Awareness.

<https://www.youtube.com/watch?v=F9QZuWPw6Y0&feature=youtu.be>

Man experiences adverse health effects from exposure to a smart meter (March 7, 2013) (3 minutes).

Presented by Garic Schoen of Gaithersburg, MD.

Produced by Maryland Smart Meter Awareness.

<http://marylandsmartmeterawareness.org/smart-meter-news/maryland-ms-resident-testimony-to-economic-matters-committee-re-hb1038-on-march-14-2013/>

Individuals with high sensitivity to the radiation from wireless devices search for increasingly rare safe electromagnetic environments.

Searching for a Golden Cage (May 8, 2014) (13 minutes)

Produced by Nadav Neuhaus.

<http://time.com/golden-cage/>

Patrons:

Prof. Declan Kennedy
Prof. Vyvyan Howard
Prof. Risteard Mulcahy



Chairperson:

Juliet Duff
Glenville, Co. Cork

Hon. Secretary:

Dr. Philip Michael
Bandon, Co. Cork

Treasurer:

Dr. Elizabeth Cullen
Thomastown,
Kilcullen,
Co. Kildare
Tel: 045-485215

Website: www.ideaireland.org

Affiliated to: ISDE -
International Society of
Doctors for the Environment.
www.isde.org
and members of HEAL -
Health and Environment Alliance
www.env-health.org
and HCWH -
Health Care Without Harm.
www.noharm.org

Linked officially with WHO -
World Health Organisation

Affiliated to International
Physicians for the Prevention
of Nuclear War - IPPNW
(Nobel Prize Winner 1985)

Charity No. 14368

7th January, 2013

Dear Principal,

The Irish Doctors Environmental Association (IDEA) has very serious concerns in relation to the ubiquitous use of Wi-Fi in Irish schools, and alerts you to the warnings of many leading international scientists and medical doctors who believe Wi-Fi is harmful to health, especially children's health.

<http://wifiinschools.org.uk/resources/safeschools2012.pdf>

Wi-Fi is an unregulated technology and there is absolutely no evidence that it is safe.

Since May 31st, 2011, radiofrequency electromagnetic fields (as in Wi-Fi) have been classified by the World Health Organisation as 'possibly carcinogenic' to humans. The IDEA unequivocally supports the Council of Europe, The European Environmental Agency and The International Commission for Electromagnetic Safety (ICEMS) in urging the adoption of the Precautionary Principle to protect human health.

Warnings by Scientists and Doctors:

<http://www.iemfa.org/index.php/appeals>

The Precautionary Principle has already been adopted by a number of Governments and agencies internationally.

Governments & organisations banning and warning against Wi-Fi:

http://www.cellphonetaskforce.org/?page_id128

While we fully support the promotion of technology in education we urge you to use wired technologies for your own safety and that of your pupils and staff. The tragedy of avoidable illness is only superseded by the knowledge that it could have been avoided.

Yours sincerely

Elizabeth Cullen M.B. B.Ch. B.A.O. M.Sc. Ph. D

045-485215

Philip Michael M.B. B.Ch. B.A.O. D.C.H. MICGP

023 8844697

Komitéen for Strålebeskyttelse
c/o Advokatfirma Christian Harlang
Nytorg 5, 1.sal
DK-1450 København K
Denmark



PO Box 33
Maple Grove Village Postal
Outlet
Oakville, ON
Canada
L6J 7P5

April 9, 2014

Via email: rec@harlanglaw.dk

Dear members of The Committee on Radiation Protection/Komitéen for Strålebeskyttelse:

My name is Frank Clegg and I am the CEO of Canadians for Safe Technology, C4ST, a volunteer based, national organization which promotes the safe use of wireless technology.

In my previous role as President of Microsoft Canada, I witnessed the incredible benefits that technology can provide. I also witnessed the potential harmful effects if technology is not implemented safely. Though wireless technologies afford schools various advantages, this solution cannot overshadow the evidence which demonstrates cause for concern. I request that you consider the following important facts.

The Canadian Teachers' Federation (CTF) is a national alliance of provincial and territorial teacher organizations that represent nearly 200,000 elementary and secondary school teachers across Canada. In their submission to the public consultation of the Royal Society of Canada, Oct. 28, 2013, they submitted the following recommendations. (Safety Code 6 is Health Canada's guideline regarding the limits of radiation from wireless devices).

Recommendations...

... That Safety Code 6 include a recommendation for prudent use of Wi-Fi whenever possible including the recommendation to limit consistent exposure in schools by turning off wireless access points when not in use. ...

That Safety Code 6 exposure thresholds be based upon both thermal and biological effects of exposure to Wi-Fi.

... That the Expert Panel recommend an education program regarding the relative safety of Wi-Fi exposure and that appropriate resources be developed to educate the public regarding ways to avoid potential exposure risks of Wi-Fi access points and devices.

As reported by CBC News on Aug. 17, 2013, <http://www.cbc.ca/news/canada/toronto/story/2013/08/17/toronto-cell-phone-ban.html> "The Elementary Teachers' Federation of Ontario has updated its policy position on the student use of personal electronic devices, preferring for them to be turned off and put away unless a teacher says otherwise. That policy, which was amended at the union's annual general meeting, informs ETFO in its discussions with the government and school boards on related issues. A portion of that policy now states that such devices, which include cellphones, should "be stored and turned off during the instructional day unless their use is directly authorized by staff." In a separate resolution, ETFO voted to study the effects of non-ionizing electromagnetic radiation, the potentially harmful radiation emitted by cellphones. A report is due on the matter in February."

In a letter to the Peel Region, April 22, 2013, The American Academy of Environmental Medicine stated "To install this widespread wireless internet access system in Peel District schools risks a widespread public health hazard that the medical system is not yet prepared to address. Statistics show that you can expect to see an immediate reaction in 3% and delayed effects in 30%, including teachers."

In 2012, the BC Confederation of Parent Advisory Councils passed resolution 18 which states: "BCCPAC call on Boards of Education to cease to install Wi-Fi and other wireless networks in schools where other networking technology is feasible."
<http://www.bccpac.bc.ca/resolutions/wi-fi-classrooms-committee-report>

In May 2011, the World Health Organization (WHO) announced that the radiation emitted from wireless devices, including Wi-Fi, is a Class 2B carcinogen, which falls into the same category as lead and DDT.

You may already be aware that some schools and libraries in France and Switzerland have already removed Wi-Fi due to the suspected harmful health effects.

The Council of Europe, which includes 47 countries, adopted resolution 1815 which suggests in member countries "give preference to wired Internet connections, and strictly regulate(s) the use of mobile phones by schoolchildren on school premises."

The European Parliament (EU) resolutions 2008/2211(INI) & 2007/2252(INI,) state: "wireless technology (mobile phones, Wi-Fi / WiMAX, Bluetooth, DECT landline telephones) emits EMFs that may have adverse effects on human health... particularly to young people whose brains are still developing... **the limits on exposure to electromagnetic fields which have been set for the general public are obsolete.**" (emphasis in original)

Other countries such as Israel, Russia, Switzerland, Frankfurt, Bavaria, and Salzburg have followed suit making the difficult decision to use hard wired connections as well. Recently, France passed a law recommending hard wired technology in schools.

The Austrian Medical Chamber shares that "WiFi may lead to concentration difficulties and memory problems in certain individuals." The Austrian Medical Association recommends Wi-Fi free school environments.

The International Society of Doctors for the Environment (ISDE) and Irish Doctors Environmental Association (IDEA) advises to "Avoid Wi-Fi in home or work if possible, particularly in schools or hospitals. Use wired technology whenever possible" sharing that: "Because of the potentially increased risks for the fetus, infants and young children due to their thinner more permeable skulls and developing systems, particularly the immune and neurological systems, based on the precautionary principal and on the mounting evidence for harm at the sub-cellular level, we recommend that EMR exposure should be kept to a minimum."

The American Academy of Pediatrics (AAP) - 60,000 Pediatricians and Pediatric Surgeons calls for caution as well stating that "The differences in bone density and the amount of fluid in a child's brain compared to an adult's brain could allow children to absorb greater quantities of RF energy deeper into their brains than adults... the current exposure limits may not reflect the latest research on RF energy" and lends support to removing Wi-Fi from schools as well.

As stewards of the public trust, I urge you to ensure the safest possible learning environment for the students in your care and to set an example for school districts by removing Wi-Fi and adopting "Best Practices" which limit the use of other wireless technologies.

Sincerely,



Frank Clegg
CEO,
Canadians for Safe Technology (C4ST)
frank@c4st.org

cc: Susanne Hansen, sh.klodskov@gmail.com



28 February 2011

Chairman and Trustees
Kawartha Pine Ridge District School Board
Education Centre
1994 Fisher Drive
Peterborough, Ontario K9J7A1

Dear Sirs/Madams:

This is concerning potential adverse health effects associated with exposure to radiofrequency (RF) radiation, specifically that from wireless routers. I am a public health physician who has been involved in issues related to electromagnetic fields (EMFs) for a number of years. I served as the Executive Secretary for the New York Powerline Project in the 1980s, a program of research which showed that children living in homes with elevated magnetic fields coming from powerlines suffered from an elevated risk of developing leukemia. I have edited two books on effects of EMFs, including RF radiation. I served as the co-editor of the Bioinitiative Report (www.bioinitiative.org), a comprehensive review of the literature on this subject. The public health chapter from this report was subsequently published in a peer reviewed journal, and that is attached. Also I testified before the President's Cancer Panel on this subject in 2009, and a publication coming from that testimony is also attached. Thus this is a subject which I know well, and one on which I take a public health approach that has as a fundamental principle the need to protect against risk of disease even when one does not have all the information that would be desirable.

There is clear and strong evidence that intensive use of cell phones increases the risk of brain cancer, tumors of the auditory nerve and cancer of the parotid gland, the salivary gland in the cheek by the ear. The evidence for this conclusion is detailed in the attached publications. WiFi uses similar radiofrequency radiation (1.8 to 5.0 GHz), although the intensity of exposure in the immediate environment is much lower than what one gets from holding a cell phone close to your head. The difference between a cell phone and a WiFi environment, however, is that while the cell phone is used only intermittently a WiFi environment is continuous. In addition WiFi transmitters are indoors, where people (and in this case, children) may be very close to them. There is evidence from Scandinavian studies of cell phone usage that children who use cell phones are about five times more likely to develop brain cancer than if use starts as an adult. Thus it is especially important to protect children.

To my knowledge there has not been any health investigation of individuals living or working in WiFi environments as compared to others who are not. However, because the radiation is the same as those for cell phones, there is every reason to assume that the health effects would be the same, varying only in relation to the total dose of radiation. Wired facilities do not generate any RF radiation. While there is not specific proof that WiFi increases risk of cancer, there is certainly no evidence that it is safe. I urge you to not put WiFi in any school. Children should not be put at increased risk of developing cancer.

Yours sincerely,

David O. Carpenter, M.D.
Director, Institute for Health and the Environment
University at Albany

Dr., CEO Andrew Zuckerman
Montgomery County Schools
Carver Educational Services Center
850 Hungerford Drive
Rockville, MD 20850
U.S.A

13th December 2015

PhD Mikko Ahonen, Tampere, Finland
MD Lena Hedendal, Luleå, Sweden
MSc. Tarmo Koppel, Tallinn, Estonia

1. Regarding: Measurements related problems in the MCPS Wi-Fi Report

We have analysed the measurement report and would like to note the following:

- In the **Comparison-table 2.2.** the MCPS provides only average values, no peak values. In cell phone technologies (like GSM) the difference between average and peak value is 2-fold. **In Wireless local area technologies like Wi-Fi, the difference between average value and peak value is up to 100-fold** (Ferro & Potorti, 2005). Note that in the table 2.2. by the MCPS only average values are presented. Later you provide **in the chapter 7.2.2 Maximum, Instantaneous Power Density, which needs attention since these levels occasionally exceeded in your school measurements allowable EMC-levels (EN60601-1 → 3 V/m) for medical instruments** (Robinson *et al.*, 2003).

- **Almost all MCPS measurements were done in the near field of the devices under 3 wavelengths.** The wavelength for 2,4 GHz is 12,5 cm and for 5 GHz is 6 cm. That means that the near field will be <37,5 cm for 2,4 GHz and <18 cm for 5 GHz. In order to assess power density exposure in near field one needs to measure both electric and magnetic field components.

- The MCPS has not provided **information about Wi-Fi technology, namely it's beacon signal.** This signal, officially **SSID (Service Set Identifier)**, is created by the access point (AP) by sending constantly SSID 10 times in a second, at 10 Hz (Ferro and Poporti, 2005). **Mobile industry has patented technology to avoid this constant SSID sending for health reasons** (Swisscom, 2004). This SSID sending at 10 Hz is an additional risk-factor and it should be mentioned. Our brain operates in alpha, beta and gamma bands. This Wi-Fi beacon overlaps the alpha band. Low-frequency EMFs (including low-frequency pulses) have an effect on evoked potentials of the brain (Carrubba *et al.*, 2008).

- Because of the risk of this 10 Hz Beacon signal of Wi-Fi, The European Academy for Environmental Medicine has assigned very strict precautionary RF-levels for Wi-Fi (Belyaev et al., 2015). Please, pay attention to Wi-Fi RF power density peak-levels in the next picture.

RF source Max Peak/Peak Hold	Daytime exposure	Nighttime exposure	Sensitive populations ¹⁾
Radio broadcast (FM)	10,000 $\mu\text{W}/\text{m}^2$	1000 $\mu\text{W}/\text{m}^2$	100 $\mu\text{W}/\text{m}^2$
TETRA	1000 $\mu\text{W}/\text{m}^2$	100 $\mu\text{W}/\text{m}^2$	10 $\mu\text{W}/\text{m}^2$
DVB-T	1000 $\mu\text{W}/\text{m}^2$	100 $\mu\text{W}/\text{m}^2$	10 $\mu\text{W}/\text{m}^2$
GSM (2G) 900/1800 MHz	100 $\mu\text{W}/\text{m}^2$	10 $\mu\text{W}/\text{m}^2$	1 $\mu\text{W}/\text{m}^2$
DECT (cordless phone)	100 $\mu\text{W}/\text{m}^2$	10 $\mu\text{W}/\text{m}^2$	1 $\mu\text{W}/\text{m}^2$
UMTS (3G)	100 $\mu\text{W}/\text{m}^2$	10 $\mu\text{W}/\text{m}^2$	1 $\mu\text{W}/\text{m}^2$
LTE (4G)	100 $\mu\text{W}/\text{m}^2$	10 $\mu\text{W}/\text{m}^2$	1 $\mu\text{W}/\text{m}^2$
GPRS (2.5G) with PTCCH [*]	10 $\mu\text{W}/\text{m}^2$	1 $\mu\text{W}/\text{m}^2$	0.1 $\mu\text{W}/\text{m}^2$
(8.33 Hz pulsing)			
DAB+	10 $\mu\text{W}/\text{m}^2$	1 $\mu\text{W}/\text{m}^2$	0.1 $\mu\text{W}/\text{m}^2$
(10.4 Hz pulsing)			
Wi-Fi 2.4/5.6 GHz (10 Hz pulsing)	10 $\mu\text{W}/\text{m}^2$	1 $\mu\text{W}/\text{m}^2$	0.1 $\mu\text{W}/\text{m}^2$

Picture. Precautionary levels for RF-radiation. **For Wi-Fi less than 10 $\mu\text{W}/\text{m}^2$ (peak value), which is 0,001 $\mu\text{W}/\text{cm}^2$ (peak value).** By the European Academy for Environmental Medicine (Belyaev *et al.*, 2015, p. 356)

- We would like to draw attention to long-term exposure related health risks.

Radiofrequency radiation from Wi-Fi devices causes fertility problems as shown by several in vivo and in vitro studies (see for example Atasoy *et al.*, 2013, Avendaño *et al.*, 2012, Dasdag *et al.*, 2015a, Shokri *et al.*, 2015).

Additionally, **RF-radiation from Wi-Fi access points (AP) causes oxidative stress in cells which leads to several disorders** (see for example Nazıroğlu *et al.*, 2012, Aynali *et al.*, 2013, Salah *et al.*, 2013). The overall detrimental impact of RF radiation induced oxidative stress is summarised in the review of Yakymenko *et al.* (2015).

2. Regarding: The IARC classification of RF-EMF as Group 2B, i.e., ‘possibly’ carcinogenic to humans and the MCPS Report’s inaccurate interpretation

The classification of radiofrequency electromagnetic fields (RF-EMF) as Group 2B, i.e., ‘possibly’ carcinogenic to humans, was made by 30 scientists from 14 countries at a meeting 2011 for the International Agency for Research on Cancer (IARC), World Health Organization (IARC 2011, Baan et al. 2012). **The working group mainly based their classification on one cohort study (Schüz et al., 2006) and five case-control studies (Muscat et al., 2000, Inskip et al., 2001, Auvinen et al., 2002, The Interphone study group, 2010, Hardell et al., 2011).**

They also reviewed more than 40 studies that assessed the carcinogenicity of RF-EMF in rodents, including seven 2-year cancer bioassays and also many studies with endpoints relevant to mechanisms of carcinogenesis, including genotoxicity, effects on immune function, gene and protein expression, cell signaling, oxidative stress, and apoptosis (Baan et al., 2011).

The referred INTERPHONE study (The Interphone study group, 2010), in the MCPS radiation report, was one of the case-control studies. **The Interphone study was a multicentre study of mobile phone use and brain tumours, including malignant tumours in the brain as glioma and benign tumours as acoustic neuroma and meningioma.** The pooled analysis included 2708 glioma cases and 2972 controls (participation rates 64% and 53%, respectively). In the Interphone study a regular user of mobile phones had an average of at least one call per week for a period of ≥ 6 months.

This very low user group was compared to several other groups of low users compared to nowadays more extensive use of mobile phones. The highest group of users, ≥ 1640 hours was divided in three sub groups depending on how many years they had used a mobile phone. For the shortest time span on 1-4 years only 23 of the glioma cases and 8 of the controls had used their mobile phones for more than 1640 hours. If any of these 23 persons with a brain cancer or any of the 8 controls had used their mobile phones for only one year they would have used it at least in average for four and a half hours a day during a year. If they instead had talked in their mobile phones during four years it would be for an average of a little more than an hour a day.

For the group of users between 5 and 9 years, 84 cases and 73 controls, the use per day would be at least between 54 minutes and 30 minutes. **For the long user group of 10 years or more, 93 cases and 73 controls, they talked in their mobile phones for 27 minutes a day or less for more than 10 years of use.**

For the main part of cases their use of mobile phones had been for a lot less than four hours a day. Today when most people use only their mobile phone and landline phones both at home and at work are becoming scarce, an amount of 4 hours or more wireless telephone use / day for salesman, telephone operators and so on is not uncommon.

In the Interphone study there was an statistical significant increased risk for a malignant brain tumour of 1.4 times (odds ratio, OR, 1.4, 95% CI 1.03-1.89) only for the highest user group of a total on more than 1640 hours.

Hardell et al. (2011) in Sweden found that **cases who had used a mobile phone for more than 1 year had an increased risk for glioma of 1.3 (OR 1.3, 95% CI 1.1-1.6).**

The risk increased with increasing time since first use and with total call time, reaching 3.2 times (OR 3.2, CI 2.0-5.1) for more than 2000 hours of use. Use of the mobile phone on the same side of the head as the tumour was associated with higher risk.

Since 2011 several other studies have been published which are strengthening the possible association between RF-EMF and cancer. Using the Bradford Hill viewpoints for evaluating strengths of evidence of the risk for brain tumours associated with use of mobile and cordless phones the classification should be upgraded to group 1 carcinogen, i.e., “the agent is carcinogenic to humans” (Hardell & Carlberg, 2013).

New case-control studies have verified Hardell's studies (Coureau et al., 2014) and up to 20 years of mobile phone use have found even higher risk for brain tumours (Hardell & Carlberg, 2015).

A newly published study has found a tumor promotion effect on mice from exposure to radiofrequency electromagnetic fields below exposure limits for humans (Lerchl *et al.*, 2015). RF-EMFs do not cause direct DNA damage. On the contrary **numerous studies have shown generation of reactive oxygen species (ROS) that can cause oxidative damage of DNA. This is a well-known mechanism in carcinogenesis for many agents.** The broad biological potential of ROS and other free radicals makes radiofrequency radiation a potentially hazardous factor for human health, not only cancer risk but also other health effects (Yakymenko *et al.*, 2015).

The IARC classification of RF-EMF as Group 2B, possibly carcinogenic to humans, doesn't only include exposure from mobile phones near the ear. **The classification includes all sources of RF-EMFs.** The exposure from mobile phone base stations, Wi-Fi access points, smart phones, laptops and tablets can be long term, sometimes around the clock both at home and at school. **This constant exposure to lower levels of exposure may be as deleterious to health as higher exposure during short time** (Fragopoulou et al., 2012, Dasdag et al., 2015b). **This risk may be accentuated for children because their probable longer use of wireless devices** (Morgan et al., 2014). **Children are also growing and have more immature cells which can be more sensible to RF-EMF** (Markova et al., 2010)

In conclusion, long term health effects from RF EMFs are still under investigation and a significant amount of troublesome scientific evidence has surfaced. By using wireless technologies at close range, long term health risks cannot be excluded. Therefore, we recommend schools to use wired technologies.

Respectfully submitted

Sincerely,



Mr. Mikko Ahonen, PhD
Research manager of Finland, Institute of Environmental Health and Safety,
Tallinn, Estonia & Partner, Sustainable Mobile Inc, Tampere, Finland.
Piiskusalmentie 4, 33450 Siivikkala, Finland.
E-mail: mikko.ahonen@tutanota.com.



Mrs. Lena Hedendahl, MD
General Practitioner
Östra Skolgatan 12, 972 53 Luleå, Sweden
E-mail: lenahedendahl@telia.com



Mr. Tarmo Koppel, MSc., PhD Candidate
Department of Work Environment and Safety, Tallinn University of Technology,
Ehitajate tee 5, 19086 Tallinn, Estonia,
E-mail: tarmo.koppel@ttu.ee

REFERENCES:

- Atasoy HI, Gunal MY, Atasoy P, Elgun S, Bugdayci G. Immunohistopathologic demonstration of deleterious effects on growing rat testes of radiofrequency waves emitted from conventional Wi-Fi devices. *J Pediatr Urol*. 2013;9(2):223–9.
- Auvinen A, Hietanen M, Luukkonen R, Koskela RS. Brain tumors and salivary gland cancers among cellular telephone users. *Epidemiology* 2002; 13: 356–59
- Avendaño C, Mata A, Sanchez Sarmiento CA, Doncel GF. Use of laptop computers connected to internet through Wi-Fi decreases human sperm motility and increases sperm DNA fragmentation. *Fertility and Sterility*. 2012;97(1):39–45.e2.
- Aynali G, Nazıroğlu M, Çelik Ö, Doğan M, Yarıktaş M, Yasan H. Modulation of wireless (2.45 GHz)-induced oxidative toxicity in laryngotracheal mucosa of rat by melatonin. *Eur Arch Otorhinolaryngol*. 2013;270(5):1695–700.
- Baan R, Grosse Y, Lauby Secretan B, El Ghissassi F, Bouvard V, Benbrahim-Tallaa L, et al. Carcinogenicity of radiofrequency electromagnetic fields. *Lancet Oncol*. 2011;12(7):624–6.
- Belyaev I, Dean A, Eger H, Hubmann G, Jandrisovits R, Johansson O, et al. EUROPAEM EMF Guideline 2015 for the prevention, diagnosis and treatment of EMF-related health problems and illnesses. *Rev Environ Health*. 2015;30(4):337–71. Available at: <http://tinyurl.com/EUROPAEM-EMF>. Accessed on December 12, 2015.
- Carrubba S, Marino AA. The Effects of Low-Frequency Environmental-Strength Electromagnetic Fields on Brain Electrical Activity: A Critical Review of the Literature. *Electromagnetic Biology and Medicine*. 2008;1;27(2):83–101.
- Coureau G, Bouvier G, Lebailly P, Fabbro-Peray P, Gruber A, Leffondre K et al. Mobile phone use and brain tumours in the CERENAT, case-control study. *Occup Environ Med* 2014;71(7):514–22.
- Dasdag S, Taş M, Akdag MZ, Yegin K. Effect of long-term exposure of 2.4 GHz radiofrequency radiation emitted from Wi-Fi equipment on testes functions. *Electromagnetic Biology and Medicine*. 2015a;2;34(1):37–42.
- Dasdag S, Akdag MZ, Erdal ME, Erdal N, Ay OI, Ay ME, et al. Effects of 2.4 GHz radiofrequency radiation emitted from Wi-Fi equipment on microRNA expression in brain tissue. *Int J Radiat Biol* 2015b;91(7):555–61.
- Ferro E, Potorti F. Bluetooth and Wi-Fi wireless protocols: a survey and a comparison. *IEEE Wireless Communications*. 2005;12(1):12–26.
- Fragopoulou AF, Samara A, Antonelou MH, Xanthopoulou A, Papadopoulou A, Vougas K, et al. Brain proteome response following whole body exposure of mice to mobile phone or wireless DECT base radiation. *Electromagn Biol Med*. 2012;31(4)250–74.

Hardell L, Carlberg M, Hansson Mild K. Pooled analysis of case-control studies on malignant brain tumours and the use of mobile and cordless phones including living and deceased subjects. *Int J Oncol* 2011; 38: 1465–74.

Hardell L, Carlberg M. Using the Hill viewpoints from 1965 for evaluating strengths of evidence of the risk for brain tumours associated with use of mobile and cordless phones. *Rev Environ Health*. 2013;28:97-106.

Hardell L, Carlberg M. Mobile phone and cordless phone use and the risk for glioma – Analysis of pooled case-control studies in Sweden, 1997–2003 and 2007–2009. *Pathophysiology*. 2015;22(1):1-13.

Gandhi OP, Morgan LL, de Salles AA, Han Y-Y, Herberman RB, Davis DL. Exposure limits: the underestimation of absorbed cell phone radiation, especially in children. *Electromagnetic Biology and Medicine*. 2012;31(1):34–51.

IARC. World Health Organization. International Agency for Research on Cancer. Non-ionizing radiation, part 2: radiofrequency electromagnetic fields. IARC Monographs Volume 102. Lyon: IARC Press, 2013. Available at: <http://monographs.iarc.fr/ENG/Monographs/vol102/mono102.pdf>. Accessed on August 12, 2015.

Inskip PD, Tarone RE, Hatch EE, et al. Cellular-telephone use and brain tumors. *N Engl J Med* 2001; 344: 79–86.

Lerchl A, Klose M, Grote K, Wilhelm AF, Spathmann O, Fiedler T et al. Tumor promotion by exposure to radiofrequency electromagnetic fields below exposure limits for humans. *Biochem Biophys Res Commun*. 2015;459:585-90

Markova E, Malmgren L, Belyaev IY. Microwaves from mobile phones inhibit 53BP1 focus formation in human stem cells stronger than in differentiated cells: Possible mechanistic link to cancer risk. *Environ Health Perspect*. 2010;118(3):394-9.

Morgan LL, Kesari S, Davis DL. Why children absorb more microwave radiation than adults: The consequences. *J Microscopy Ultrastructure* 2. 2014;2:197–204.

Muscat JE, Malkin MG, Thompson S, et al. Handheld cellular telephone use and risk of brain cancer. *JAMA* 2000; 284: 3001–07

Nazıroğlu M, Çelik Ö, Özgül C, Çiğ B, Doğan S, Bal R, et al. Melatonin modulates wireless (2.45 GHz)-induced oxidative injury through TRPM2 and voltage gated Ca(2+) channels in brain and dorsal root ganglion in rat. *Physiol Behav*. 2012;105(3):683–92.

Salah MB, Abdelmelek H, Abderraba M. Effects of olive leave extract on metabolic disorders and oxidative stress induced by 2.45 GHz WIFI signals. *Environ Toxicol Pharmacol*. 2013;36(3):826–34.

Shokri S, Soltani A, Kazemi M, Sardari D, Mofrad FB. Effects of Wi-Fi (2.45 GHz) Exposure on Apoptosis, Sperm Parameters and Testicular Histomorphometry in Rats: A Time Course Study. *Cell J*. 2015;17(2):322–31.

Schüz J, Jacobsen R, Olsen JH, Boice JD Jr, McLaughlin JK, Johansen C. Cellular telephone use and cancer risk: update of a nationwide Danish cohort. *J Natl Cancer Inst* 2006;98: 1707–13.

Swisscom. Patent. Reducing electrosmog in wireless networks. 2004. Available at: <http://tinyurl.com/Wi-Fi-patent>. Accessed on December 12, 2015.

The Interphone study group. Brain tumour risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Int J Epidemiol*. 2010;39:675-94.

Yakymenko I, Tsybulin O, Sidorik E, Henshel D, Kyrylenko O, Kyrylenko S. Oxidative mechanisms of biological activity of low-intensity radiofrequency radiation. *Electromagn Biol Med*. 2015;19:1-16.

24 March 2014

Open letter by British medical doctors: Health and safety of Wi-Fi and mobile phones

We wish to highlight our concern over the safety of exposure to microwave radiation from wireless technology, particularly for vulnerable groups like children, pregnant women, the elderly and those with compromised health.

There is growing concern that chronic (long-term) exposure to radiofrequency/microwave radiation from wireless technologies causes damage, particularly genetic damage, cognitive damage, cancer and decreased fertility. There is now substantial evidence of a link between mobile phone use and brain cancer. This was recognised by the International Agency for Research on Cancer (IARC)'s 30-strong panel of scientists, which in 2011 classed radiofrequency radiation as "possibly carcinogenic".

Additionally, doctors are encountering a significant and growing number of people presenting with **a range of acute (short-term) symptoms from wireless radiation, including headaches, palpitations, rashes, fatigue, sleep disturbance, allergies and memory and concentration problems.**

International medical agencies have recognised the evidence of harm (see appended list) but these rulings may take many years to be reflected in public health policy. This controversy is a common characteristic of scientific understanding when environmental exposures are new.

New technologies and substances often come with scientific conflict, which can continue for several decades before consensus is achieved. Commercial pressures often delay the acceptance of health risks, even when scientific evidence is compelling. In the case of tobacco, asbestos, x-rays and leaded petrol, for example, it took many decades before damage was established and accepted by health agencies and, during those decades, millions of people suffered ill health and death as a result of the delay. Now, despite evidence of harm, wireless technology is being rolled out widely.

We urge health agencies and the public to act immediately to reduce exposure to radiofrequency/ microwave radiation. This is especially important for children, who are physiologically more vulnerable to this exposure, and for whom adults have a safeguarding responsibility. **Children's health should be put ahead of convenience and commercial benefits. Children should not use mobile phones except in an emergency, and WiFi should be replaced with wired alternatives in schools and other settings where children spend considerable time.**

Yours faithfully,

Dr Elizabeth Evans MA (Cantab), MBBS (Lond), DRCOG – Medical Doctor
Dr Andrew Tresidder MRCGP (1989), MBBS (Lond) – Medical Doctor
Dr Erica Mallery Blythe BM – Medical Doctor
Dr Elizabeth Cullen MBBCh BAO MSc PhD – Medical Doctor
Dr Philip Michael MBBCh BAO DCH MICGP – Medical Doctor
Dr Shideh Pouria MBBS, BSc, MRCP – Medical Doctor
Dr Rodney Adeniyi-Jones LRCP&SI, MRCP – Medical Doctor
Dr Jenny Goodman MA, MBChB – Ecological Physician

Dr Damian Downing MBBS, MSB – President BSEM
Dr Elena Toma MD - Psychiatrist
Dr Joan Kinder MA, MBBChir(Cantab), MRCPCH – retired Consultant Paediatrician
Dr Sarah Myhill MBBS – General Practitioner (GP)
Dr Dee Marshall MBBS, MFHom – Medical Doctor
Dr Charles Forsyth MBBS, FFHom – Medical Doctor
Dr Zac Cox BDS - Dentist

BCM SSITA London WC1N 3XX
www.ssita.org.uk

Appendix – International Rulings

1. In 2011 the **World Health Organization’s scientific panel, the International Agency for Research on Cancer (IARC)**, reviewed all the evidence on carcinogenesis (cancer-causing) and categorised electromagnetic radiation from mobile phones and Wi-Fi as **Possibly Carcinogenic (Class 2B)**.

See http://www.iarc.fr/en/media-centre/pr/2011/pdfs/pr208_E.pdf

2. **The Council of Europe has called for member states to take measures to reduce exposure to electromagnetic fields and give preference to wired internet connections for children, particularly in schools and classrooms.**

The Parliamentary Assembly stated that “the Assembly regrets that, despite calls for the respect of the precautionary principle and despite all the recommendations, declarations and a number of statutory and legislative advances, there is still a lack of reaction to known or emerging environmental and health risks and virtually systematic delays in adopting and implementing effective preventive measures. Waiting for high levels of scientific and clinical proof before taking action to prevent well-known risks can lead to very high health and economic costs, as was the case with asbestos, leaded petrol and tobacco.”

See <http://assembly.coe.int/mainf.asp?link=/documents/adoptedtext/ta11/eres1815.htm>

3. **The BioInitiative Report**, updated in 2012 by 29 scientists, states that **biological effects are clearly established and occur at very low levels of exposure to electromagnetic fields and radiofrequency radiation** from just minutes of exposure to mobile phone masts (cell towers), Wi-Fi, and wireless utility ‘smart’ meters.

See <http://www.bioinitiative.org/conclusions>

4. **The American Academy of Environmental Medicine** stated in a 2012 Position Paper that “**Multiple studies correlate RF exposure with diseases such as cancer, neurological disease, reproductive disorders, immune dysfunction, and electromagnetic hypersensitivity.**”

See http://aaemonline.org/emf_rf_position.html

6. **International Society of Doctors for the environment (ISDE) and Irish Doctors’ Environmental Association (IDEA)** state that “**there is sufficient scientific evidence to warrant more stringent controls** on the level and distribution of electromagnetic radiation [EMR]. The joint statement and recommendations are part of a call by medical and scientific experts for safe technologies in schools.”

See <http://www.env-health.org/news/members-news/article/isde-idea-statement-on>

5. **The Safe Schools Report 2012** lists statements by **other doctors and medical associations** raising concerns over children’s exposure to electromagnetic fields from Wi-Fi and other wireless technology.

See <http://wifiinschools.org.uk/resources/safeschools2012.pdf>



Dr. Magda Havas, B.Sc., Ph.D.

Environmental & Resource Studies, Trent University, Peterborough, ON, Canada
phone: (705) 748-1011 x7882 **fax:** (705) 748-1569 **email:** mhavas@trentu.ca

July 10, 2009.

Open Letter to Parents, Teachers, & School Boards Regarding Wi-Fi Networks in Schools and Cell Phone Antennas near School Property

I am a scientist who does research on the health effects of electromagnetic radiation and I am becoming increasingly concerned that a growing number of schools are installing WiFi networks and are making their school grounds available for cell phone antennas.

You will be told by both the federal government (Federal Communication Commission in the US; Health Canada and Industry Canada in Canada) as well as by the Wi-Fi provider that this technology is **safe** provided that exposures to radio frequency radiation remain below federal guidelines.

This information is **outdated** and **incorrect** based on the growing number of scientific publications that are reporting adverse health and biological effects below our “short-term, thermal-based” guidelines (see www.bioinitiative.org) and the growing number of scientific and medical organizations that are asking for stricter guidelines to be enforced.

For these reasons it is irresponsible to introduce Wi-Fi microwave radiation into a school environment where young children and school employees spend hours each day.

FACT:

- 1. GUIDELINES: Guidelines for microwave radiation (which is what is used in Wi-Fi) range 5 orders of magnitude in countries around the world.** The lowest guidelines are in Salzburg Austria and now in Liechtenstein. The guideline in these countries is 0.1 microW/cm². See short video (<http://videos.next-up.org/SfTv/Liechtenstein/AdoptsTheStandardOf06VmBioInitiative/09112008.html>). In Switzerland the guideline is 1 and in both Canada and the US it is 1000 microW/cm²!

Why do Canada and the US have guidelines that are so much higher than other countries? Our guidelines are based on a short-term (6-minute in Canada and 30-minute in US) heating effect. It is assumed that if this radiation does not heat your tissue it is “safe”. This is NOT correct. Effects are documented at intensities well below those that are able to heat body tissue. See attached report: *Analysis of Health and Environmental Effects of Proposed San Francisco Earthlink Wi-Fi Network* (2007). These biological effects include increased permeability of the blood brain barrier, increased calcium flux, increase in cancer and DNA breaks, induced stress proteins, and nerve damage. Exposure to this energy is associated with altered white blood cells in school children; childhood leukemia; impaired motor function, reaction time, and memory; headaches, dizziness, fatigue, weakness, and insomnia.

- 2. ELECTRO-HYPER-SENSITIVITY:** A growing population is adversely affected by these electromagnetic frequencies. The illness is referred to as “electro-hyper-sensitivity” (EHS) and is recognized as a disability in Sweden. The World Health Organization defines EHS as:

“... a phenomenon where individuals experience adverse health effects while using or being in the vicinity of devices emanating electric, magnetic, or electromagnetic fields (EMFs). . . EHS is a real and sometimes a debilitating problem for the affected persons, while the level of EMF in their neighborhood is no greater than is encountered in normal living environments. Their exposures are generally several orders of magnitude under the limits in internationally accepted standards.”

Health Canada acknowledges in their Safety Code 6 guideline that some people are more sensitive to this form of

energy but they have yet to address this by revising their guidelines.

Symptoms of EHS include sleep disturbance, fatigue, pain, nausea, skin disorders, problems with eyes and ears (tinnitus), dizziness, etc. It is estimated that 3% of the population are severely affected and another 35% have moderate symptoms. Prolonged exposure may be related to sensitivity and for this reason it is imperative that children's exposure to microwave radiation (Wi-Fi and mobile phones) be minimized as much as possible.

3. **CHILDREN'S SENSITIVITY:** Children are more sensitive to environmental contaminants and that includes microwave radiation. The Stewart Report (2000) recommended that children not use cell phones except for emergencies. The cell phone exposes your head to microwave radiation. A wireless computer (Wi-Fi) exposes your entire upper body and if you have the computer on your lap it exposes your reproductive organs as well. Certainly this is not desirable, especially for younger children and teenagers. For this reason we need to discourage the use of wireless technology by children, especially in elementary schools. That does not mean that students cannot go on the Internet. It simply means that access to the Internet needs to be through wires rather than through the air (wireless, Wi-Fi).
4. **REMOVAL OF WI-FI:** Most people do not want to live near either cell phone antennas or Wi-Fi antennas because of health concerns. Yet when Wi-Fi (wireless routers) are used inside buildings it is similar to the antenna being inside the building rather than outside and is potentially much worse with respect to exposure since you are closer to the source of emission.

Libraries in France are removing Wi-Fi because of concern from both the scientific community and their employees and patrons.

The Vancouver School Board (VSB) passed a resolution in January 2005 that prohibits construction of cellular antennas within 1000 feet (305 m) from school property.

Palm Beach, Florida, Los Angeles, California, and New Zealand have all prohibited cell phone base stations and antennas near schools due to safety concerns. The decision not to place cell antennas near schools is based on the likelihood that children are more susceptible to this form of radiation. **Clearly if we do not want antennas "near" schools, we certainly do not want antennas "inside" schools!** The safest route is to have wired internet access rather than wireless. While this is the more costly alternative in the short-term it is the least costly alternative in the long run if we factor in the cost of ill health of both teachers and students.

5. **ADVISORIES:** Advisories to limit cell phone use have been issued by the various countries and organizations including the UK (2000), Germany (2007), France, Russia, India, Belgium (2008) as well as the Toronto Board of Health and the Pittsburgh Cancer Institute (July 2008). While these advisories relate to cell phone use, they apply to Wi-Fi exposure as well since both use microwave radiation. If anything, Wi-Fi computers expose more of the body to this radiation than do cell phones.
6. **PRECAUTIONARY PRINCIPLE:** Even those who do not "accept" the science showing adverse biological effects of microwave exposure should recognize the need to be careful with the health of children. For this reason we have the Precautionary Principle, which states:

In order to protect the environment, the precautionary approach shall be widely applied by States according to their capability. Where there are threats of serious or irreversible damage, lack of full scientific certainty shall not be used as a reason for postponing cost effective measures to prevent environmental degradation.

In this case "States" refers to the School Board and those who make decisions about the health of children.

The two most important environments in a child's life are the home (especially the bedroom) and the school. For this reason it is imperative that these environments remain as safe as possible. **If we are to err, please let us err on the side of caution.**

Respectfully submitted,
Dr. Magda Havas,
Associate Professor
Trent University
July 10, 2009

Shallow Minds: How the Internet and Wi-Fi in Schools Can Affect Learning

By Cindy Lee Russell, MD
VP-Community Health, Santa Clara County Medical Association

Most of us cannot live without our computers, text messaging, e-mail, and immediate access to the vast cloud of information, especially kids and teenagers who have grown up in the age of the Internet. In fact, more schools are integrating computers at younger ages, even in kindergarten. Forty-nine states are phasing out cursive handwriting altogether. What effects does it have, however, on learning, brain development, cognition, and brain health? Studies have shown some interesting ways that technology is rewiring and shaping our brain, which may not be “all good.”

A growing body of scientific evidence suggests that the Internet, with its distractions and interruptions, is turning us into scattered, superficial thinkers. What does that portend for our kids?

Multitasking and Internet Addiction

Nicholas Carr explains, in his book “The Shallows,” that we are changing the way we process information. “Dozens of studies by psychologists, neurobiologists, educators, and Web designers point to the same conclusion: When we go online, we enter an environment that promotes cursory reading, hurried and distracted thinking, and superficial learning....The Net delivers precisely the kind of sensory and cognitive stimuli-repetitive, intensive, interactive, addictive, that have been shown to result in strong and rapid alterations in brain circuits and functions.”

Researchers from Stanford, in 2009, gave a battery of cognitive tests to a group of heavy and light media Internet multitaskers. They found that the heavy multitaskers were much more easily distracted by “irrelevant environmental stimuli” and had less control over their working memory. In addition, they were much less able to focus on a particular task. Professor Clifford Nass, who led the research, stated intensive multitaskers are “suckers for irrelevancy. Everything distracts them.” (5)

“Teaching is a human experience. Technology is a distraction when we need literacy, numeracy, and critical thinking.” Paul Thomas, author and associate professor of education at Furman University

Law School Professors Ban Laptops in Classrooms

Several years ago, professors who were irritated with students surfing the Web and hiding behind laptop screens began banning the use of the Internet or laptops in the classroom. Laptops have been banned in classes at Harvard Law School, Yale, George Washington University, University of Virginia, and South Texas College of Law, to mention a few. (4)(15) A 2006 study by Carrie Fried backed up the policies, demonstrating that students who used laptops in

class spent considerable time multitasking. They more importantly found that the level of laptop use was negatively related to several measures of student learning. (3)

A 2012 survey by Elon University, the Pew Internet, and American Life Project asked over 1,000 leaders in the U.S. their thoughts about cognition in our millennial generation. They were asked to consider how the Internet and its environment are changing, for better or worse. Overall, the survey found that multitasking is the new norm and that hyper-connectivity may be leading to a lack of patience and concentration. The “always on” ethos may be encouraging a culture of expectation and instant gratification.

Brain Maturation, Learning, Memory, and Intelligence

The maturation of intelligence requires quiet, deep thought, and time. Established research findings in cognitive science leads to the conclusion that laptop use, especially with Wi-Fi access, could interfere with learning.

The hippocampus, which lies under the cortex, is intimately involved in long-term memory storage. Initial experiences are stored and stabilized in the hippocampus and then later transferred to the cortex. Removal of the hippocampus does not affect long-term memories, but prevents new memories from forming.

Learning depends on the ability to transfer information from our working memory to long-term memory and weave this into other acquired knowledge. There is a bottleneck in the passage of working memory to long-term memory. We have a limited ability as humans to capture and process information. The Internet provides too many choices and too much information at once. Excess distracting information creates “overload,” preventing long-term memorization and important information is lost. No one disagrees that we need to protect our memories. As author Nicholas Carr highlights, personal memory is not just for the individual to function, but it shapes and sustains our collective cultural memory.

Brain Drain:

Adverse Neurologic and Health Effects of Wireless Microwave Communications

A growing body of peer reviewed research is showing neurologic damage to fetal brain and other systems from Wi-Fi and other microwave wireless sources. In a prior article, “Why-Fi: Is Wireless Communication Hazardous to Your Health?” in the Sept/Oct 2010 SCCMA *Bulletin*, the full range of effects of EMF from our cell phones and wireless devices was discussed. New basic science research in the last three years is confirming these findings. Initially, the Bioinitiative report of 2007 reviewed the biological effects of low level EMF. It found that there was clear evidence of adverse effects to living systems at current environmental exposures and at doses well below the threshold of the International Commission of Non-Ionizing Radiation Protection (ICNIRP) safety guidelines. Current microwave safety limits are based solely on the heating of tissue and do not take into account research showing negative biological effects on DNA, cancer, protein synthesis, skin tissue changes, sperm motility and viability, cognitive functioning, and disruption of the blood brain barrier.

Current Research on Cognition and Wireless Communication

Fetal Radiofrequency Radiation Exposure From 800-1900 MHz-Rated Cellular Telephones Affects Neurodevelopment and Behavior in Mice. *Scientific Reports*. March 2012.

Aldad et al noted that neurobehavioral disorders are increasingly prevalent in children with 3%-7% of school-aged children diagnosed with attention deficit hyperactivity disorder (ADHD). The etiology is unclear, however, an association between prenatal cellular telephone use and hyperactivity in children has been postulated by others. To test this, he exposed pregnant mice to cell phone radiation throughout gestation (days 1-17), with a sham cell phone control group. He found that the exposed group had dose responsive impaired neurologic transmission in the prefrontal cortex and that the mice exposed in utero were hyperactive and had impaired memory. He concluded “that these behavioral changes were due to altered neuronal developmental programming.”(3)

Microwave Radiation Induced Oxidative Stress, Cognitive Impairment, and Inflammation in Brain of Fischer Rats. Megha. 2012.

Megha evaluated the intensity of oxidative stress, cognitive impairment, and brain inflammation in rats exposed to typical cell phone microwave radiation. They were subjected to 900 and 1,800 MHz EMF for two hours a day, for 30 days. They state, “Significant impairment in cognitive function and induction of oxidative stress in brain tissues of microwave exposed rats were observed, in comparison with sham exposed groups... Results of the present study indicated that increased oxidative stress due to microwave exposure may contribute to cognitive impairment and inflammation in brain.”

Effect of Low Level Microwave Radiation Exposure on Cognitive Function and Oxidative Stress in Rats. Deshmukh. 2013.

The author highlights the exponential increase in wireless communication devices we are exposed to. He evaluated the effects of cell phone radiation on oxidation in tissues, in addition to cognition in rats. They subjected rats to 900 MHz EMF for two hours per day, five days a week, for 30 days, with an unexposed control group. “Results showed significant impairment in cognitive function and increase in oxidative stress, as evidenced by the increase in levels of MDA (a marker of lipid peroxidation) and protein carbonyl (a marker of protein oxidation) and unaltered GSH content in blood. Thus, the study demonstrated that low level MW radiation had significant effect on cognitive function and was also capable of leading to oxidative stress.”

The Internet Can Damage Teenage Brains

A large radiologic study from China, published July 2011, looked at structural brain changes in Internet-addicted teenagers. It is estimated that 24 million teenagers are addicted to the Internet in China. The researchers found a consistent atrophy of grey matter in parts of the brain and shrinkage of the surface of the brain in those addicted to the Internet. The effects were worse the longer the addiction. In addition, the study revealed changes in white matter of the brain, which

function to transmit messages in the brain to the grey matter. They concluded these structural abnormalities were most likely associated with functional impairments in cognitive control.

“It strikes me as a terrible shame that our society requires photos of brains shrinking in order to take seriously the common-sense assumption that long hours in front of screens is not good for our children’s health. Dr Aric Sigman, Fellow of the Royal Society of Medicine

WHO Classifies EMF as a Carcinogen

In 2011, The WHO/International Agency for Research on Cancer (IARC) classified radiofrequency electromagnetic fields as “possibly carcinogenic to humans (Group 2B), based on an increased risk for glioma, a malignant type of brain cancer¹, associated with wireless phone use.”

France Bans Wi-Fi in Schools, But Replaces With Ethernet

The French National Assembly, March 2013, passed an amendment to ban Wi-Fi in their schools until it’s proven “safe for human consumption.” They instead agreed to install far safer, wired Ethernet cable connections.

The Council of Europe has called for a ban on Wi-Fi use in schools and also recommends a wired alternative.

In Austria, the Austrian Medical Society has also issued a policy statement asking for a ban of Wi-Fi in schools.

The U.K. has a useful frequently-updated website on Wi-Fi in schools, which provides much scientific research. <http://www.wifinschools.org.uk/> Still the controversy persists.

The Cost of a Virtual World

There are a host of concerns with classroom technology, and the virtual world it creates, that have not been explored in the rush to “modernize” education and prevent our kids from becoming “computer illiterate,” despite the fact that computers are designed for ease of use. These issues range from distraction in the classroom, impairment of cognitive development and long-term memory, deficiency in learning social skills, Internet addiction, cyber bullying, access to inappropriate content, eye fatigue, and security risks to online learning networks. In addition, the sheer cost of computers and continuous upgrades is likely to break many school budgets. We have not mentioned the issue of toxic e-waste, another growing public health problem.

Common Sense

We will not get rid of the Internet or computers. We should not ignore, however, the enlarging body of science that points to real threats to public health and, especially, our children’s safety and well-being. The best approach is precautionary. Reduce the risk by reducing the microwave emissions. It is our obligation as physicians and parents to protect our children. They are the

future and our legacy.

1. Remove wireless devices (white boards and routers) in schools in favor of wired connections and fiberoptic.
2. If there is Wi-Fi, then give teachers the authority to turn it off when not in use or if they feel it is not necessary.
3. Ban cell towers near or on schools.
4. Limit screen time on computers.
5. Limit or ban cell phone use in the class.
6. Limit or ban cell phone use at home.
7. Do not allow laptops to be placed on laps.
8. Undertake independent scientific studies on Wi-Fi and computer use that look at acute and long-term health effects.
9. Train teachers how to recognize symptoms of EMF reactions.
10. Conduct meetings with parents and teachers to address this issue in each school.

References

1. **The Shallows: What the Internet Is Doing to Our Brains.** Nicholas Carr. 2010.
2. **Generation Y: The Internet's effects on cognition and education.**
www.Triplehelixblog.com
3. **In-class laptop use and its effects on student learning.** Carrie B. Fried. Sept 2006.
http://www.mcla.edu/Academics/uploads/textWidget/3424.00018/documents/laptop_use_in_the_classroom.pdf
4. **Banning Laptops in the Classroom: Is It Worth the Hassles?** Kevin Yamamoto.
<http://intra.albanylaw.edu/cr/insttech/pdfs/laptopban.pdf>
5. **Cognitive control in media multitasker.** C. Nass.
<http://www.pnas.org/content/early/2009/08/21/0903620106.abstract>
6. **Fetal radiofrequency radiation exposure from 800-1,900 MHz-rated cellular telephones affects neurodevelopment and behavior in mice.** 2012. Aldad.
<http://www.ncbi.nlm.nih.gov/pubmed/22428084>
7. **Effect of low level microwave radiation exposure on cognitive function and oxidative stress in rats.** Deshmukh PS, 2013 April, Indian J Biochem Biophy.
<http://www.ncbi.nlm.nih.gov/pubmed/23720885>
8. **Microwave radiation induced oxidative stress, cognitive impairment and inflammation in brain of Fischer rats.** Megha. Indian J Exp Biol. 2012, Dec;50(12):889-96.
<http://www.ncbi.nlm.nih.gov/pubmed/?term=microwave+radiation+induced+oxidative+stress+fischer+rats+Megha>
9. **Prevention of mobile phone induced skin tissue changes by melatonin in rat: an experimental study.** Ozguner FToxicol Ind Health. 2004 Sep;20(6-10):133-9.
<http://www.ncbi.nlm.nih.gov/pubmed/15941010>
10. **The effects of long-term exposure of magnetic field via 900-MHz GSM radiation on some biochemical parameters and brain histology in rats.** Celikozlu SD. Electromagn Biol Med. 2012 Dec;31(4):344-55.

- <http://www.ncbi.nlm.nih.gov/pubmed/22676902>
11. **France Bans Wi-Fi From Schools—We Should All Do the Same.**
<http://www.francesfox.com/france-bans-wifi-school/>
 12. **High Wired: Does Addictive Internet Use Restructure the Brain?** Scientific American. June 17, 2011. <https://www.scientificamerican.com/article.cfm?id=does-addictive-internet-use-restructure-brain>
 13. **Too much Internet use can damage teenagers' brain.** Mail Online. 18 July 2011. <http://www.dailymail.co.uk/sciencetech/article-2015196/Too-internet-use-damage-teenagers-brains.html>
 14. **Wi Fi in Schools U.K.** <http://www.wifiinschools.org.uk/>
 15. **Wi Fi in Schools Australia.** <http://www.wifi-in-schools-australia.org/p/worldwide.html>
 16. **Wide Web of diversions gets laptops evicted from lecture halls.**
<http://www.washingtonpost.com/wp-dyn/content/article/2010/03/08/AR2010030804915.html>
 17. **A Silicon Valley School That Doesn't Compute.**
<http://www.nytimes.com/2011/10/23/technology/at-waldorf-school-in-silicon-valley-technology-can-wait.html?pagewanted=all>
 18. **Effects of the exposure to mobile phones on male reproduction: a review of the literature.** <http://www.ncbi.nlm.nih.gov/pubmed/21799142>
 19. **Mobile phone radiation induces reactive oxygen species production and DNA damage in human spermatozoa in vitro.**
<http://www.ncbi.nlm.nih.gov/pubmed/19649291>
 20. **Evidence for mobile phone radiation exposure effects on reproductive pattern of male rats: role of ROS.** <http://www.ncbi.nlm.nih.gov/pubmed/22897402>
 21. **Effects of radiofrequency electromagnetic waves (RF-EMW) from cellular phones on human ejaculated semen: an in vitro pilot study.**
<http://www.ncbi.nlm.nih.gov/pubmed/18804757>
 22. **Rats Exposed to Cell Phone Microwaves Suffer Long-Term Memory Loss, According to New Study by University of Washington Researcher.**
<http://www.sciencedaily.com/releases/1999/12/991202070403.htm>
 23. **Spatial memory and learning performance and its relationship to protein synthesis of Swiss albino mice exposed to 10 GHz microwaves.**
<http://www.ncbi.nlm.nih.gov/pubmed/23952535>
 24. **Alterations of visual reaction time and short term memory in military radar personnel.** <http://www.ncbi.nlm.nih.gov/pubmed/23785684>
 25. **Relationship between cognition function and hippocampus structure after long-term microwave exposure.** <http://www.ncbi.nlm.nih.gov/pubmed/22998825>
 26. **Impairment of long-term potentiation induction is essential for the disruption of spatial memory after microwave exposure.**
<http://www.ncbi.nlm.nih.gov/pubmed/23786183>
 27. **Influence of microwave radiation on synaptic structure and function of hippocampus in Wistar rats.** <http://www.ncbi.nlm.nih.gov/pubmed/17535652>
 28. **A aquaporin 4 expression and effects in rat hippocampus after microwave radiation.** <http://www.ncbi.nlm.nih.gov/pubmed/20137298>
 29. **Relationship between millimeter wave irradiation in pregnant mice and c-Fos protein expression in hippocampus and learning and memory functions in their**

- offsprings.** <http://www.ncbi.nlm.nih.gov/pubmed/16405774>
30. **Effects of 7 Hz-modulated 450 MHz electromagnetic radiation on human performance in visual memory tasks.**
<http://www.ncbi.nlm.nih.gov/pubmed/12465659>
31. **Data Security Is a Classroom Worry, Too.**
http://www.nytimes.com/2013/06/23/business/data-security-is-a-classroom-worry-too.html?pagewanted=all&_r=0

New Jersey Education Association

JUSTFORMEMBERS

Sign in for member information and resources.

[Member Sign In](#)

Minimize health risks from electronic devices

Published in the September 2016 NJEA Review

by Adrienne Markowitz and Eileen Senn

Desktops, laptops, tablets, eBook readers, printers, projectors, smart boards, smart TVs, cellphones, cordless phones and wireless networks (WiFi) have become ubiquitous in schools. At their best, they are powerful tools for education. At their worst, they threaten the physical and mental health of teachers, paraeducators, secretaries, librarians and other school staff members and students who spend numerous hours using the devices.

Physical health risks from electronic devices include pain and tingling from repetitive strain injuries to the hands and wrists; pain in the neck, shoulders and back; dry, burning, itchy eyes, blurred vision and headaches; altered sleep patterns and next-day fatigue from exposure to blue screen light; distracted driving; and various health problems from exposure to radiation.

Mental health risks arise from stress due to raised expectations for multitasking, productivity and proficiency with devices; dealing with malfunctioning devices; student and colleague distraction from and addiction to devices; and intrusion of devices into nonwork time.

WiFi devices emit radiation

Radio frequency (RF) electromagnetic frequency (EMF) radiation is sent and/or received by the antennae of phones, routers and other wireless devices. RF radiation is capable of causing cancer, reproductive, neurological and ocular effects. The amount of radiation exposure received depends on the amount of time exposed and distance from the source. Radiation levels fall off exponentially with distance from antennae. If you double the distance, the radiation is four times less. If you triple the distance, it is nine times less, and so on. Children and developing fetuses are particularly at risk because their bodies are still growing. People with implanted medical devices are at risk for device interference.

Hazards and solutions

The most straightforward ways to minimize health risks are to use electronic devices in moderation and to maximize your distance from them. There are also specific solutions to specific hazards listed below.

Local associations should work with their UniServ field representative to negotiate solutions that are in the control of district administrators such as providing training and ergonomic equipment and hard-wiring devices. Individuals should take steps within their control, such as:

For repetitive strain injuries

- Use voice control/speech recognition.
- Use ergonomic alternatives to traditional mice and keyboards.
- Use as many fingers as possible when typing and both thumbs when texting.

For neck, shoulder and back pain

- Ensure an ergonomic workstation.
- When using a hand-held device, support it and the forearms.
- Avoid bending the head down or jutting it forward.
- Take frequent, short breaks from the device.
- Ensure good posture and change positions frequently.
- Stand and do stretching exercises.

For eye pain, blurred vision and headaches

- Use sufficient, but not excessive, lighting.
- Use assistive technology built into Apple, Android and Windows devices.
- Enlarge and darken the cursor and pointer.
- Enlarge the font; magnify the text.
- Use text-to-speech instead of reading.
- Use special computer glasses.
- Relax the eyes on a minibreak.

For altered sleep patterns and next-day fatigue

- Stop using devices at least one hour before bedtime.

For distracted driving

- Use hands-free devices, preferably speakerphones.
- Pull over and park.
- Let someone else drive.

For radiation exposure

- Keep devices away from the body and bedroom.
- Carry phones in briefcases, etc., not on the body.
- Put devices on desks, not laps.
- Hard wire all devices that connect to the internet.
- Hard wire all fixed devices such as printers, projectors and boards.
- Use hard-wired phones instead of cell or cordless phones.
- Text rather than call.
- Keep conversations short or talk in person.
- Put devices in airplane mode, which suspends EMF transmission by the device, thereby disabling Bluetooth, GPS, phone calls, and WiFi.
- Use speaker phone or ear buds instead of holding the phone next your head.
- Take off Bluetooth devices when not using them.

For stress

- Training in device use, assistive technology.
- Easy access to user manuals.
- Easily available technical support.

Cell phones and cancer

The National Toxicology Program (NTP) is conducting the largest set of laboratory rodent studies to date on cellphone RF radiation. The studies cost \$25 million and are designed to mimic human exposure. They are based on the cellphone

frequencies and modulations currently in use in the United States. The NTP studies are designed to look at effects in all parts of the body.

On May 27, 2016, NTP released a report with partial results of the studies. They found increased occurrence of rare brain tumors called gliomas and increases in nerve tumors called schwannoma of the heart in male rats. The released results are partial because more rat studies and all of the mouse studies will be forthcoming by 2017. The cells that became cancerous in the rats were the same types of cells as those that have been reported to develop into tumors in human cellphone users.

The EMF produced by cellphones was classified as possibly carcinogenic to humans by the World Health Organization in 2011. They found that long-term use of a cell phone might lead to two different types of tumors, gliomas and acoustic neuroma, a tumor of the auditory nerve.

For more information

- **“Job stress: Is it killing you?”** NJEA Review, May 2012.
- **“As schools lift bans on cell phones, educators weigh pros and cons.”** Kinjo Kiema, NEA Today, Feb. 23, 2015.
- **Be kind to your eyes**, NJEA Review, September 2012.
- **Computer workstations eTool, Occupational Safety and Health Administration** (OSHA).
- **“Stretching Exercises at Your Desk, 12 Simple Tips,”** WebMD.
- **“Cell phone facts and tips,”** Grassroots Environmental Education.
- **“Radiofrequency and microwave radiation,”** Occupational Safety and Health Administration (OSHA).
- **“Report of Partial Findings from the National Toxicology Program (NTP) Carcinogenesis Studies of Cell Phone Radiofrequency Radiation in Hsd: Sprague Dawley SD Rats (Whole Body Exposure).”**
- **“Low EMF Best Practices,”** Collaborative for High Performance Schools (CHPS), 2014.
- Microsoft Accessibility Center: www.microsoft.com/enable
- Apple Accessibility Center: www.apple.com/accessibility
- Google/Android Accessibility Center: www.google.com/accessibility/products-features.html

Adrienne Markowitz holds a Master of Science in Industrial Hygiene from Hunter College, City University of New York. Eileen Senn holds a Master of Science in Occupational Health from Temple University in Philadelphia. They are consultants with the New Jersey Work Environment Council, which is a frequent partner with NJEA on school health and safety concerns.

Adrienne Markowitz holds a Master of Science in Industrial Hygiene from Hunter College, City University of New York. Eileen Senn holds a Master of Science in Occupational Health from Temple University in Philadelphia. They are consultants with the New Jersey Work Environment Council, which is a frequent partner with NJEA on school health and safety concerns.

- View this article in **PDF**
- View this article optimized for **mobile phones**
- View this article optimized for **tablets**



Resolution 1815 (2011)¹

Final version

The potential dangers of electromagnetic fields and their effect on the environment

Parliamentary Assembly

1. The Parliamentary Assembly has repeatedly stressed the importance of states' commitment to preserving the environment and environmental health, as set out in many charters, conventions, declarations and protocols since the United Nations Conference on the Human Environment and the Stockholm Declaration (Stockholm, 1972). The Assembly refers to its past work in this field, namely [Recommendation 1863 \(2009\)](#) on environment and health: better prevention of environment-related health hazards, [Recommendation 1947 \(2010\)](#) on noise and light pollution, and more generally, [Recommendation 1885 \(2009\)](#) on drafting an additional protocol to the European Convention on Human Rights concerning the right to a healthy environment and [Recommendation 1430 \(1999\)](#) on access to information, public participation in environmental decision-making and access to justice – implementation of the Århus Convention.
2. The potential health effects of the very low frequency of electromagnetic fields surrounding power lines and electrical devices are the subject of ongoing research and a significant amount of public debate. According to the World Health Organization, electromagnetic fields of all frequencies represent one of the most common and fastest growing environmental influences, about which anxiety and speculation are spreading. All populations are now exposed in varying degrees to electromagnetic fields, the levels of which will continue to increase as technology advances.
3. Mobile telephony has become commonplace around the world. This wireless technology relies upon an extensive network of fixed antennae, or base stations, relaying information with radio-frequency signals. Over 1.4 million base stations exist worldwide and the number is increasing significantly with the introduction of third generation technology. Other wireless networks that allow high-speed Internet access and services, such as wireless local area networks, are also increasingly common in homes, offices and many public areas (airports, schools, residential and urban areas). As the number of base stations and local wireless networks increases, so does the radio-frequency exposure of the population.
4. While electrical and electromagnetic fields in certain frequency bands have wholly beneficial effects which are applied in medicine, other non-ionising frequencies, whether from extremely low frequencies, power lines or certain high frequency waves used in the fields of radar, telecommunications and mobile telephony, appear to have more or less potentially harmful, non-thermal, biological effects on plants, insects and animals as well as the human body, even when exposed to levels that are below the official threshold values.
5. As regards standards or threshold values for emissions of electromagnetic fields of all types and frequencies, the Assembly strongly recommends that the ALARA (as low as reasonably achievable) principle is applied, covering both the so-called thermal effects and the athermic or biological effects of electromagnetic emissions or radiation. Moreover, the precautionary principle should be applied when scientific evaluation does not allow the risk to be determined with sufficient certainty. Given the context of growing exposure of the population, in particular that of vulnerable groups such as young people and children, there could be extremely high human and economic costs if early warnings are neglected.

1. Text adopted by the Standing Committee, acting on behalf of the Assembly, on 27 May 2011 (see [Doc. 12608](#), report of the Committee on the Environment, Agriculture and Local and Regional Affairs, rapporteur: Mr Huss).



6. The Assembly regrets that, despite calls for the respect of the precautionary principle and despite all the recommendations, declarations and a number of statutory and legislative advances, there is still a lack of reaction to known or emerging environmental and health risks and virtually systematic delays in adopting and implementing effective preventive measures. Waiting for high levels of scientific and clinical proof before taking action to prevent well-known risks can lead to very high health and economic costs, as was the case with asbestos, leaded petrol and tobacco.

7. Moreover, the Assembly notes that the problem of electromagnetic fields or waves and their potential consequences for the environment and health has clear parallels with other current issues, such as the licensing of medication, chemicals, pesticides, heavy metals or genetically modified organisms. It therefore highlights that the issue of independence and credibility of scientific expertise is crucial to accomplish a transparent and balanced assessment of potential negative impacts on the environment and human health.

8. In light of the above considerations, the Assembly recommends that the member states of the Council of Europe:

8.1. in general terms:

8.1.1. take all reasonable measures to reduce exposure to electromagnetic fields, especially to radio frequencies from mobile phones, and particularly the exposure to children and young people who seem to be most at risk from head tumours;

8.1.2. reconsider the scientific basis for the present standards on exposure to electromagnetic fields set by the International Commission on Non-Ionising Radiation Protection, which have serious limitations, and apply ALARA principles, covering both thermal effects and the athermic or biological effects of electromagnetic emissions or radiation;

8.1.3. put in place information and awareness-raising campaigns on the risks of potentially harmful long-term biological effects on the environment and on human health, especially targeting children, teenagers and young people of reproductive age;

8.1.4. pay particular attention to “electrosensitive” people who suffer from a syndrome of intolerance to electromagnetic fields and introduce special measures to protect them, including the creation of wave-free areas not covered by the wireless network;

8.1.5. in order to reduce costs, save energy, and protect the environment and human health, step up research on new types of antenna, mobile phone and DECT-type device, and encourage research to develop telecommunication based on other technologies which are just as efficient but whose effects are less negative on the environment and health;

8.2. concerning the private use of mobile phones, DECT wireless phones, WiFi, WLAN and WIMAX for computers and other wireless devices such as baby monitors:

8.2.1. set preventive thresholds for levels of long-term exposure to microwaves in all indoor areas, in accordance with the precautionary principle, not exceeding 0.6 volts per metre, and in the medium term to reduce it to 0.2 volts per metre;

8.2.2. undertake appropriate risk-assessment procedures for all new types of device prior to licensing;

8.2.3. introduce clear labelling indicating the presence of microwaves or electromagnetic fields, the transmitting power or the specific absorption rate (SAR) of the device and any health risks connected with its use;

8.2.4. raise awareness on potential health risks of DECT wireless telephones, baby monitors and other domestic appliances which emit continuous pulse waves, if all electrical equipment is left permanently on standby, and recommend the use of wired, fixed telephones at home or, failing that, models which do not permanently emit pulse waves;

8.3. concerning the protection of children:

8.3.1. develop within different ministries (education, environment and health) targeted information campaigns aimed at teachers, parents and children to alert them to the specific risks of early, ill-considered and prolonged use of mobiles and other devices emitting microwaves;

8.3.2. for children in general, and particularly in schools and classrooms, give preference to wired Internet connections, and strictly regulate the use of mobile phones by schoolchildren on school premises;

- 8.4. concerning the planning of electric power lines and relay antenna base stations:
 - 8.4.1. introduce town planning measures to keep high-voltage power lines and other electric installations at a safe distance from dwellings;
 - 8.4.2. apply strict safety standards for the health impact of electrical systems in new dwellings;
 - 8.4.3. reduce threshold values for relay antennae in accordance with the ALARA principle and install systems for comprehensive and continuous monitoring of all antennae;
 - 8.4.4. determine the sites of any new GSM, UMTS, WiFi or WIMAX antennae not solely according to the operators' interests but in consultation with local and regional government authorities, local residents and associations of concerned citizens;
- 8.5. concerning risk assessment and precautions:
 - 8.5.1. make risk assessment more prevention oriented;
 - 8.5.2. improve risk-assessment standards and quality by creating a standard risk scale, making the indication of the risk level mandatory, commissioning several risk hypotheses to be studied and considering compatibility with real-life conditions;
 - 8.5.3. pay heed to and protect "early warning" scientists;
 - 8.5.4. formulate a human-rights-oriented definition of the precautionary and ALARA principles;
 - 8.5.5. increase public funding of independent research, in particular through grants from industry and taxation of products that are the subject of public research studies to evaluate health risks;
 - 8.5.6. create independent commissions for the allocation of public funds;
 - 8.5.7. make the transparency of lobby groups mandatory;
 - 8.5.8. promote pluralist and contradictory debates between all stakeholders, including civil society (Århus Convention).

How to Protect Your Family

1. Do not hold the phone directly against your head or body

Use speakerphone, other hands-free device, or another device that meets EHT guidance.

2. Beware of a weak signal

Your phone works harder and emits more radiation into you when the signal is weak or blocked.

3. Protect children, the pregnant abdomen, and men who want to become fathers

Children absorb twice as much radiation as adults. Sperm are especially vulnerable.

4. Do not sleep with your cell phone

Phones do not emit radiation when off or in airplane mode.

5. Corded landlines are swifter, safer, and more secure

Cordless phones also emit microwave radiation.

6. Do not text while moving e.g. driving, biking, blading, walking, skiing.

Milliseconds can make the difference between life or death.

Brought to you by:

The Global Campaign for Safer Cell Phones provides open-source resources that educate and motivate health professionals, teachers, parents, and students about simple and safer ways to use cell phones. Our multi-media toolkit can be downloaded at EHTrust.org.

Prepared by physicians, biomedical scientists, and neurosurgeons: including Kevin O'Neill, FRCS (SN); Ronald B. Herberman, MD; Charles Teo, MD; David Servan-Schreiber, MD, PhD; Annie J. Sasco, MD, DrPh; Stephen Sinatra, MD; John West, MD; Richard Stein, MD, PhD; Devra Lee Davis, PhD, MPH

Find out more about how to reduce your risk and get the latest news on our website EHTrust.org and #PracticeSafeTech.

Doctors' Advice to Patients & Their Families

Cell Phones & Health: Simple Precautions Make Sense



This pamphlet reviews precautionary advice by governments to reduce your exposure to cell phone radiation.

**ENVIRONMENTAL
HEALTH TRUST**

Cell phones are two-way microwave radios that were never properly tested for safety.

Antennas for cell phones are continually searching for signals from towers to send and receive information. The body or brain absorbs about half the radiation emitted from a phone at any time.

Children absorb more radiation than adults



Courtesy Om P. Gandhi and Green America


People who use cell phones more than 30 minutes per day (average) for 10 years have a doubled or higher risk of brain cancer. Teenagers who start to use cell phones develop 4 to 5 times more brain cancer in less than a decade. Links to studies can be found at EHTrust.org.

WORLD HEALTH ORGANIZATION (2011)

On May 31, 2011, the World Health Organization/International Agency for Research on Cancer classified the electromagnetic radiation from cell phones as possibly carcinogenic to humans based on an increased risk for glioma, a malignant type of brain cancer, associated with use of mobile phones.

Can Cell Phones Cause Cancer and Disease?


A number of well-designed studies indicate that cell phone use increases cancer risk. Cell phones change the brain. After 50 minutes of holding a cell phone on or next to the ear, the brain of a healthy adult shows significant changes in glucose metabolism—the brain's fuel—at precisely those areas reached by cell phone radiation. The long-term impact of these changes is unknown. Recent research suggests that cell phones, cordless phones, and other wireless devices can also disturb heart rate.



Cell Phones Lower Sperm Count

Men who use cell phones 2 to 4 hours per day have 30% lower sperm count than non-users; daily use over 4 hours results in 40% lower sperm count. Studies have shown that microwave radiation can damage sperm.

Agarwal, *Fertility and Sterility*, 2008



HEALTH CANADA (2011)

“Health Canada reminds cell phone users that they can take practical measures to reduce RF exposure. The department also encourages parents to reduce their children's RF exposure from cell phones since children are typically more sensitive to a variety of environmental agents.”

How to Protect Your Family

Choose wired over wireless

- Get a corded landline phone and forward cell phones to it.
- Use corded plug-in (non-wireless) connections for printer, mouse, internet, gaming, and routers, etc.—with all wireless functions OFF.

Safeguard sleep

- Stop use of and power off computers, tablets and phones one hour (or longer) before bedtime.
- Do not sleep with a cell phone, tablet or laptop.

Maximize distance from self and others

- Radiation decreases as you move away from the source.
- Locate devices away from laps, bras and pockets, as the radiation can damage sperm and may be linked to breast cancer.
- Use air tube headset or use speaker phone at maximum distance from head and body.

Protect children

- Children are particularly vulnerable and should not use cell phones except in an emergency.
- If children are using a phone or other device for work or play, select all 3 of the following settings to turn off the radiation from cell phones and wireless devices: Airplane mode ON, and Wi-Fi OFF, and Bluetooth OFF.
- Don't use a mobile device while a child is on your lap or in your arms.

Power off in vehicles

- Mobile devices distract drivers, emit more intense radiation during travel, and metal surroundings create radiation hot-zones.

more & news
EHTrust.org

Doctors' Advice to Patients & Families

Wireless & Health: Simple Precautions Make Sense



This pamphlet reviews
precautionary advice
by Doctors and governments to
reduce exposure to wireless radiation.

 **ENVIRONMENTAL
HEALTH TRUST**

Brought to you by: Physicians, biomedical scientists, and neurosurgeons—including Charles Teo, MD; Martha Herbert, MD, PhD; Anthony B. Miller, MD; Süleyman Kaplan, PhD; Annie J. Sasco, MD, PhD; Stephen T. Sinatra, MD; Dr. Erica Mallery-Blythe, BMBS; Devra Lee Davis, PhD, MPH

Wireless Radiation Exposures

Wireless devices—such as tablets, laptops, game consoles, baby monitors, and wifi routers—all emit the same type of wireless microwave radiation as cell phones. Children, pregnant women, and men hoping to father healthy children are more vulnerable to this exposure.

The Child Brain and Skull Absorb up to Ten Times More Invisible Wireless Radiation than the Adult

(Gandhi et al., 2012)



Cell phone radiation absorbed into brain, skull, and eyes of 6-year-old

(Fernandez et al., 2015)


New anatomically based research shows radiation absorption.

Increased Cancer Risk

Wireless radiofrequency radiation was classified as a Class 2B “Possible Human Carcinogen” by the World Health Organization’s International Agency for Research on Cancer in 2011. Evidence *has increased* since 2011, indicating that cell phone and wireless radiation should be classified as a “probable carcinogen” because of increased brain cancer (Davis et al., 2015).


Those exposed at younger ages show up to eight times more brain cancer risk.

Radiation Absorption from Tablet into a 6-Year-Old’s Brain (in this image: white & yellow areas are the highest absorption)



Federal University of Rio Grande do Sul, UFRGS Ferreira and de Salles (2015)

Wireless Exposure Damages Brains



Animals prenatally exposed to cell phone radiation develop damaged and fewer brain cells.

How Wireless Impacts Children’s Health


Accumulating research shows that wireless microwave radiation affects fetal brain development, the immune system and reproductive function. *Even very low doses* of wireless radiation affect brain metabolism and electrical activity.

Yale University researchers found that when pregnant mice were exposed to radiation from a simulated operating phone, their offspring had impaired memory, increased hyperactivity and altered brains. Several research studies show wireless impairs brain development at various ages.

The Reproductive System

Numerous studies indicate that wireless exposures decrease sperm quantity and quality (including altered DNA) and damage testes and ovaries.

Testes Damaged by Wi-Fi Equipment Radiation



Long-term Wi-Fi exposure decreases testes volume

(Dasdag et al., 2015)

Government Protections Worldwide

Over 20 countries—including France, Belgium, Israel and the European Union—have taken policy actions to “reduce exposure to children,” addressing cell phones, laptops, cell towers and Wi-Fi in schools in order to protect public health.

detailed references available at EHTrust.org