

# **Circumstances of Opioid-Related Overdose Deaths in Maryland Region of Injury: Baltimore City**

2020 State Unintentional Drug Overdose Reporting System (SUDORS)

Please note: The Maryland Vital Statistics Administration provides official death counts and rates for the State of Maryland. They can be viewed online at <https://health.maryland.gov/vsa/Pages/reports.aspx>. The State Unintentional Drug Overdose Reporting System (SUDORS) is designed to supplement vital statistics data by providing information about the circumstances of overdose death. Please refer to the Vital Statistics Administration for case counts and rates.

Prepared on February 27, 2023

### What is SUDORS?

The State Unintentional Drug Overdose Reporting System (SUDORS) is an enhanced surveillance system collecting over 600 variables about the circumstances of fatal overdose in Maryland. The Maryland SUDORS program works closely with the Office of the Chief Medical Examiner, the Vital Statistics Administration, and multiple law enforcement agencies in Maryland to access, review, and systematically document information about overdose death in Maryland. There is an 8-13 month lag in availability of data due to federal requirements for SUDORS data abstraction.

### What will you find here?

This data packet includes a select set of variables from the 2020 Maryland SUDORS data set. These include: substances contributing to the cause of death, mental health, substance use, place of overdose death, recent release from an institution, presence of potential bystanders, previous overdose, treatment for pain, prescribed buprenorphine or methadone, naloxone administration, route of administration, usual industry, and usual occupation.

### How can SUDORS data be used?

The information in this data packet can be used to inform local prevention and response activities around partner engagement, identification of priority populations, provision of services, and evaluation of existing programs. Please refer to the OD2A Toolkit for additional information.

### What if you need more information?

If you need information not provided in this report, please send an email with your contact information, a description of needed data and/or a request for a variable list, and the subject line "SUDORS data needed" to [mdh.mvdrs@maryland.gov](mailto:mdh.mvdrs@maryland.gov).

### KEY FINDINGS:

#### Baltimore City

- 96% (961) of fatal overdoses were opioid-related; for consistency with previous reports, those opioid-related deaths are the focus of this report.
- About 1 in 6 decedents (18%) were in treatment for mental health or substance use at the time of overdose death.
- More than half (52%) of decedents died in their own home.
- 1 in 12 decedents (8%) had recently been released from an institution, such as a hospital, jail or prison, detention facility, or supervised residential facility.
- 3 out of 5 decedents (60%) were in the vicinity of a potential bystander during or shortly before the fatal overdose.
- 1 in 25 decedents (4%) were treated for pain at the time of the fatal overdose.
- 1 in 7 decedents (14%) were administered naloxone.
- Just over 1 in 5 decedents (22%) had evidence of injection drug use.
- About 1 in 5 decedents (19%) worked in the construction industry.

***See pages 3-15 for additional information!***

## Table of Contents

---

**Important Note on Data Sources:** Circumstance data is limited to information documented in the law enforcement and/or medical examiner files; these are likely underestimated as death investigators might have limited information. Information on mental health and substance use history comes solely from these sources and not medical records or other treatment records.

### Maryland Overdose Deaths by Substance

Figure 1: Substance(s) Contributing to Cause of Death.....	4
--	---

### Prevalence of Mental Health, Substance Abuse, and Other Addiction


Figure 2: Overall.....	5
Figure 3: by Sex.....	5
Figure 4: by Age (Years).....	6
Figure 5: by Race/Ethnicity.....	6
Figure 6: by Education Level.....	7
Figure 7: by Marital Status.....	7
Figure 8: by Urban-Rural Classification of County of Residence.....	8
Figure 9: by Service in the U.S. Armed Forces.....	9

### Additional Information

Figure 10: Place of Overdose Death.....	9
Figure 11: Recent Release From Institution.....	10
Figure 12: Presence of Potential Bystanders, Previous Overdose, Treatment for Pain, and Prescribed Buprenorphine or Methadone.....	11
Figure 13: Naloxone Administration.....	12
Figure 14: Route of Administration.....	13
Table 1: Usual Industry of Opioid Overdose Decedents.....	14
Table 2: Usual Occupation of Opioid Overdose Decedents.....	15

**Circumstances of Opioid-Related Overdose Deaths in Maryland,  
Baltimore City**

Figure 1: Substance(s) Contributing to Cause of Death, 2020



<sup>1</sup>Opioid includes: illicitly manufactured fentanyl, heroin, prescription opioids, and any other opioids.

<sup>2</sup>Stimulant includes: cocaine, eutylone, methamphetamine, and any other stimulants.

<sup>3</sup>Other includes substances such as: antidepressants, antipsychotics, benzodiazepines, muscle relaxants, PCP, and others.


\*Per Centers for Disease Control and Prevention (CDC) requirement: data suppressed due to fewer than 11 cases.

**In 2020, 96% of overdose deaths were opioid related.**


The remaining analysis is limited to opioid-related fatal overdose, and consequently excludes:

- Stimulant only deaths (n=33)
- Other only deaths (n<11)

**Figure 2: Prevalence of Mental Health, Substance Abuse, and Other Addiction<sup>1</sup>, 2020**



**Figure 3: Prevalence of Mental Health, Substance Abuse, and Other Addiction<sup>1</sup> by Sex, 2020**



<sup>1</sup>Health history data may be incomplete. See note on Table of Contents (page 3).

## Circumstances of Opioid-Related Overdose Deaths in Maryland, Baltimore City

Figure 4: Prevalence of Mental Health, Substance Abuse, and Other Addiction<sup>1</sup> by Age (Years), 2020





Figure 5: Prevalence of Mental Health, Substance Abuse, and Other Addiction<sup>1</sup> by Race/Ethnicity, 2020



<sup>1</sup>Health history data may be incomplete. See note on Table of Contents (page 3).

<sup>2</sup>All other includes: Asian/Pacific Islander, American Indian/Alaska Native, unknown/unspecified, and two or more races.

\*Per Centers for Disease Control and Prevention (CDC) requirement: data suppressed due to fewer than 11 cases.

## Circumstances of Opioid-Related Overdose Deaths in Maryland, Baltimore City

Figure 6: Prevalence of Mental Health, Substance Abuse, and Other Addiction<sup>1</sup> by Education Level, 2020






Figure 7: Prevalence of Mental Health, Substance Abuse, and Other Addiction<sup>1</sup> by Marital Status, 2020



<sup>1</sup>Health history data may be incomplete. See note on Table of Contents (page 3).

\*Per Centers for Disease Control and Prevention (CDC) requirement: data suppressed due to fewer than 11 cases.

Figure 8: Prevalence of Mental Health, Substance Abuse, and Other Addiction<sup>1</sup> by Urban-Rural<sup>2</sup> Classification of County of Residence, 2020




<sup>1</sup>Health history data may be incomplete. See note on Table of Contents (page 3).

<sup>2</sup>Large metro: counties in central or fringe metropolitan statistical areas of 1 million or more population; small or medium metro: counties in metropolitan statistical areas of populations less than 999,999; rural: counties in micropolitan statistical areas and nonmetropolitan counties of less than 49,999 (defined by the 2013 NCHS Urban-Rural Classification Scheme for Counties).


\*Per Centers for Disease Control and Prevention (CDC) requirement: data suppressed due to fewer than 11 cases.



**Figure 9: Prevalence of Mental Health, Substance Abuse, and Other Addiction<sup>1</sup> by Service in the U.S. Armed Forces, 2020**



**Figure 10: Place of Overdose Death, 2020**




<sup>1</sup>Health history data may be incomplete. See note on Table of Contents (page 3).

<sup>2</sup>All other includes: commercial establishment, hospice care, jail/prison, long-term care facility, natural area, and nursing home.

\*Per Centers for Disease Control and Prevention (CDC) requirement: data suppressed due to fewer than 11 cases.


Figure 11: Recent Release From Institution<sup>1</sup>, 2020



<sup>1</sup>Recent release from an institution is defined as deaths that occurred within a month of the decedent being released from or admitted to an institutional setting. The decedent is considered institutionalized if they spent one or more nights in the institution.

<sup>2</sup>All other includes: long term residential health facility (e.g., nursing home), other psychiatric treatment, psychiatric hospital, supervised residential facilities not related to alcohol or substance abuse treatment, and unknown type of institution.

Figure 12: Presence of Potential Bystanders, Previous Overdose, Treatment for Pain, and Prescribed Buprenorphine or Methadone, 2020




<sup>1</sup>A potential bystander is defined as a person aged  $\geq 11$  years who was physically nearby either during or shortly preceding a drug overdose and potentially had an opportunity to intervene or respond to the overdose. This includes any persons in the same structure (e.g., same room or same building, but different room) as the decedent during that time. For example, the family member of an opioid overdose decedent who was in another room during the fatal incident would be considered a potential bystander if that person might have had an opportunity to provide life-saving measures such as naloxone administration, if adequate

<sup>2</sup>Prior overdose is defined by as a drug overdose related to ANY substance in which one of the following occurred: treatment in an emergency department of other medical center, medical services responded but the person refused to be transported to the hospital, or naloxone was administered by a layperson and medical treatment was not sought.

<sup>3</sup>Treated for pain is defined as a situation in which a decedent was receiving any type of treatment for acute and/or chronic pain at the time of the fatal overdose, including prescription opioid pain relievers.

<sup>4</sup>Prescribed buprenorphine or methadone is defined as a situation in which there is any evidence, including witness reports, prescription history, and/or scene evidence that suggests that the decedent was prescribed methadone or buprenorphine at the time of the fatal overdose for either medication-assisted treatment or pain relief.

Figure 13: Naloxone Administration, 2020



<sup>1</sup>Naloxone administration is defined as EITHER a situation in which a decedent was administered naloxone for their fatal opioid overdose by an EMS responder, law enforcement officer, firefighter, or health care worker, OR a situation in which naloxone was *reportedly* administered by a layperson and there was evidence that naloxone was *actually* administered.


<sup>2</sup>First responder includes: law enforcement, emergency medical services, firefighters, or other trained professional.

<sup>3</sup>Hospital includes: emergency department, inpatient hospital setting, or a critical care center.

<sup>4</sup>Lay-person includes: as a person using drugs/alcohol with the decedent, intimate partner, friend, family member, roommate, or a bystander with no relationship to the decedent.

Note: Counts may exceed the total number of cases due to multiple administration.

Figure 14: Route of Administration, 2020



<sup>1</sup>All other includes: sublingual and transdermal.

Note: Counts may exceed the total number of cases due to multiple findings at the scene.

**Circumstances of Opioid-Related Overdose Deaths in Maryland,  
Baltimore City**

**Table 1: Usual Industry<sup>1</sup> of Opioid Overdose Decedents, 2020**

<b>Usual Industry Sector<sup>1</sup></b>	<b>n</b>	<b>%</b>
Construction	170	18.7%
Accommodation and food services	69	7.6%
Other services (except public administration)	58	6.4%
Health care and social assistance	56	3.2%
Administrative and support and waste management and remediation services	52	5.7%
Retail trade	51	5.6%
Transportation and warehousing	49	5.4%
Not in workforce (homemaker, student, volunteer, unable to work)	47	5.2%
Manufacturing	41	4.5%
Public administration	23	2.5%
Arts, entertainment, and recreation	18	2.0%
Professional, scientific, and technical services	11	1.2%
All other industries <sup>2</sup>	38	4.0%
Missing, unknown, and not enough information	225	24.8%
<b>Total</b>	<b>908</b>	<b>100.0%</b>

<sup>1</sup>Major industry sectors defined using the 2012 North American Industry

<sup>2</sup>All other industries includes: finance and insurance; educational services; wholesale trade; real estate and rental and leasing; Military; utilities; agriculture, forestry, fishing and hunting; information.

\*Analysis limited to decedents of workforce age, 18-65 years.

<b>Most Common Usual Industry, by Sex</b>	
<b>Among males:</b>	<b>Among females:</b>
1. Construction (25.5%)	1. Healthcare and social assistance (13.0%)
2. Transportation and warehousing (6.3%)	2. Accommodation and food services (11.8%)
3. Accommodation and food services (6.0%)	3. Not in workforce (11.4%)

<b>Most Common Usual Industry, by Race and Ethnicity</b>	
<b>Among White, Non-Hispanics:</b>	<b>Among Black, Non-Hispanics:</b>
1. Construction (24.8%)	1. Construction (14.3%)
2. Accommodation and food services (8.6%)	2. Healthcare and social assistance (7.7%)
3. Not in workforce (7.0%)	3. Accommodation and food services (7.3%)
<b>Among Hispanics:</b>	
1. Construction (44.4%)	

<b>Most Common Usual Industry, by Age Group</b>	
<b>Among 25-44 year-olds:</b>	<b>Among 45+ year-olds:</b>
1. Construction (16.5%)	1. Construction (19.8%)
2. Accommodation and food services (8.6%)	2. Accommodation and food services (6.7%)
3. Other services (except public administration) (7.9%)	3. Healthcare and social assistance (6.5%)

**Circumstances of Opioid-Related Overdose Deaths in Maryland,  
Baltimore City**

**Table 2: Usual Occupation<sup>1</sup> of Opioid Overdose Decedents, 2020**

<b>Usual Major Occupation<sup>1</sup></b>	<b>n</b>	<b>%</b>
Construction and extraction	164	18.1%
Transportation and material moving	113	12.4%
Food preparation and serving related	61	6.7%
Building and grounds cleaning and maintenance	55	6.1%
Not in workforce (homemaker, student, volunteer, unable to work)	48	5.3%
Personal care and service	38	4.2%
Sales and related	33	3.6%
Installation, maintenance, and repair	29	3.2%
Production	25	2.8%
Office and administrative support	25	2.8%
Management	20	2.2%
Protective service	15	1.7%
Healthcare support	13	1.4%
Arts, design, entertainment, sports, and media	12	1.3%
Community and social service	12	1.3%
All other occupations <sup>2</sup>	31	3.4%
Missing, unknown, and not enough information	214	23.6%
<b>Total</b>	<b>908</b>	<b>100.0%</b>

<sup>1</sup>Usual occupation defined using the 2010 Standard Occupational Classification (SOC) system major groups.

<sup>2</sup>All other occupations includes: healthcare practitioners and technical occupations; architecture and engineering occupations; education, training, and library; business and financial operations; Military; life, physical, and social science; computer and mathematics; farming, fishing, and forestry; legal occupations

\*Analysis limited to decedents of workforce age, 18-65 years.

<b>Most Common Usual Occupation, by Sex</b>	
<b>Among males:</b>	<b>Among females:</b>
1. Construction and extraction (24.8%)	1. Not in workforce (11.4%)
2. Transportation and material moving (14.2%)	2. Food preparation and serving related (10.6%)
3. Building and grounds cleaning and maintenance (5.6%)	3. Transportation and material moving (7.7%)

<b>Most Common Usual Occupation, by Race and Ethnicity</b>	
<b>Among White, Non-Hispanics:</b>	<b>Among Black, Non-Hispanics:</b>
1. Construction and extraction (23.8%)	1. Transportation and material moving (15.7%)
2. Food preparation and serving related (8.0%)	2. Construction and extraction (13.6%)
3. Transportation and material moving (7.3%)	3. Food preparation and serving related (6.3%)
<b>Among Hispanics:</b>	
1. Construction and extraction (44.4%)	

<b>Most Common Usual Occupation, by Age Group</b>	
<b>Among 18-44 year-olds:</b>	<b>Among 45+ year-olds:</b>
1. Construction and extraction (16.4%)	1. Construction and extraction (19.0%)
2. Transportation and material moving (11.0%)	2. Transportation and material moving (13.3%)
3. Food preparation and serving related (8.7%)	3. Building and grounds cleaning and maintenance (6.5%)