

Expedited Partner Therapy in Maryland for Chlamydia, Gonorrhea and Trichomoniasis

RECOMMENDED PARTNER TREATMENT REGIMENS*

Chlamydia Only	□ Doxycycline (Vibramycin) 100 mg orally 2 times/day for 7 days NOTE: Substitute azithromycin (Zithromax) 1 gram orally in a single dose for partners who may be pregnant, or unlikely to adhere to a 7-day regimen
Gonorrhea Only	☐ Cefixime (Suprax) 800 mg orally in a single dose
Gonorrhea/Chlamydia Coinfection or Chlamydia Not Ruled Out	 □ Cefixime (Suprax) 800 mg orally in a single dose PLUS □ Doxycycline (Vibramycin) 100 mg orally 2 times/day for 7 days NOTE: Substitute azithromycin (Zithromax) 1 gram orally in a single dose for doxycycline for partners who may be pregnant, or unlikely to adhere to a 7-day regimen
Trichomoniasis	 Metronidazole (Flagyl) 500 mg orally 2 times/day for 7 days (Male partners) Metronidazole (Flagyl) 2 g orally in a single dose (Female partners) Tinidazole (Tindamax) 2 g orally in a single dose (Alternative regimen for all non-pregnant partners)

*Based on CDC 2021 STI Treatment Guidelines. https://www.cdc.gov/std/treatment-guidelines/default.htm) Last reviewed June, 2024.