

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey Cecil County Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN9: Percentage of students who rode with a driver who had been drinking alcohol (in a car or other vehicle, one or more times during the 30 days before the survey)	13.3 (11.4-15.4)	1,647	24.5 (18.2-32.1)	224		●	
QN10: Percentage of students who drove a car or other vehicle when they had been drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	4.3 (2.9-6.3)	726	14.5 (8.4-24.0)	100		●	
QN11: Percentage of students who texted or e-mailed while driving a car or other vehicle (on at least 1 day during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	28.3 (23.4-33.8)	734	31.1 (22.5-41.2)	100			●
QN13: Percentage of students who carried a weapon on school property (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)	4.2 (3.3-5.5)	1,651	11.4 (7.5-16.9)	224		●	
QN14: Percentage of students who carried a gun (not counting the days when they carried a gun only for hunting or for a sport such as target shooting, on at least 1 day during the 12 months before the survey)	3.5 (2.7-4.6)	1,648	6.8 (3.9-11.7)	225			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey
Cecil County
Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN16: Percentage of students who were threatened or injured with a weapon on school property (such as a gun, knife, or club, one or more times during the 12 months before the survey)	5.2 (4.2-6.5)	1,654	18.8 (14.2-24.5)	226		●	
QN18: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)	6.2 (5.1-7.5)	1,654	17.9 (12.9-24.2)	226		●	
QN19: Percentage of students who were ever physically forced to have sexual intercourse (when they did not want to)	5.8 (4.2-8.0)	1,640	19.3 (14.6-25.1)	222		●	
QN20: Percentage of students who experienced sexual violence (being forced by anyone to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey)	7.1 (5.7-8.8)	1,620	23.8 (17.8-31.2)	218		●	
QN21: Percentage of students who experienced sexual dating violence (being forced by someone they were dating or going out with to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)	4.0 (2.9-5.6)	1,011	14.7 (9.7-21.8)	158		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey
Cecil County
Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN22: Percentage of students who experienced physical dating violence (being physically hurt on purpose by someone they were dating or going out with [counting such things as being hit, slammed into something, or injured with an object or weapon] one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)	6.9 (5.5-8.8)	1,047	19.0 (13.4-26.2)	163		●	
QN23: Percentage of students who were bullied on school property (ever during the 12 months before the survey)	18.2 (15.9-20.6)	1,642	37.4 (30.6-44.9)	224		●	
QN24: Percentage of students who were electronically bullied (counting being bullied through texting, Instagram, Facebook, or other social media, ever during the 12 months before the survey)	12.5 (10.7-14.6)	1,643	29.2 (23.4-35.8)	218		●	
QN25: Percentage of students who felt sad or hopeless (almost every day for >=2 weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)	24.2 (21.6-27.0)	1,629	66.7 (60.2-72.7)	218		●	
QN26: Percentage of students who seriously considered attempting suicide (ever during the 12 months before the survey)	11.8 (10.1-13.9)	1,641	45.9 (38.8-53.2)	223		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey
Cecil County
Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN27: Percentage of students who made a plan about how they would attempt suicide (during the 12 months before the survey)	9.1 (7.7-10.8)	1,639	43.1 (36.4-50.0)	220		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey Cecil County Risk Behaviors and Sexual Identity Report

Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN32: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)	8.2 (6.6-10.2)	1,652	20.0 (14.9-26.2)	226		●	
QNFRCIG: Percentage of students who currently smoked cigarettes frequently (on 20 or more days during the 30 days before the survey)	1.9 (1.3-2.7)	1,652	5.2 (3.0-8.7)	226		●	
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)	1.6 (1.1-2.5)	1,652	3.8 (2.0-7.0)	226			●
QN34: Percentage of students who ever used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo])	41.3 (38.2-44.5)	1,618	58.1 (50.4-65.3)	220		●	
QN35: Percentage of students who currently used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo], on at least 1 day during the 30 days before the survey)	18.1 (15.4-21.2)	1,527	25.3 (19.3-32.3)	193		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey Cecil County Risk Behaviors and Sexual Identity Report

Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN36: Percentage of students who usually got their own electronic vapor products by buying them in a store (such as a convenience store, supermarket, discount store, gas station, or vape store, during the 30 days before the survey, among students who currently used electronic vapor products and who were aged <18 years)	6.0 (3.5-10.1)	252	8.9 (3.1-23.0)	44			●
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, dip, snus, or dissolvable tobacco products [such as Redman, Levi Garrett, Beechnut, Skoal, Skoal Bandits, Copenhagen, Camel Snus, Marlboro Snus, General Snus, Ariva, Stonewall, or Camel Orbs], not counting any electronic vapor products, on at least 1 day during the 30 days before the survey)	5.4 (4.3-6.7)	1,647	7.6 (4.6-12.3)	222			●
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars, on at least 1 day during the 30 days before the survey)	9.1 (7.6-10.9)	1,647	14.7 (10.4-20.4)	223		●	
QNTB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)	12.8 (10.8-15.0)	1,646	25.3 (19.7-31.8)	224		●	
QNTB3: Percentage of students who currently smoked cigarettes or cigars or used smokeless tobacco (on at least 1 day during the 30 days before the survey)	14.5 (12.5-16.8)	1,645	25.6 (20.0-32.2)	224		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey
Cecil County
Risk Behaviors and Sexual Identity Report

Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNTB4: Percentage of students who currently smoked cigarettes or cigars or used smokeless tobacco or electronic vapor products (on at least 1 day during the 30 days before the survey)	24.3 (21.4-27.5)	1,558	37.6 (30.8-44.8)	205		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey Cecil County Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN40: Percentage of students who ever drank alcohol (at least one drink of alcohol, on at least 1 day during their life)	57.8 (54.4-61.1)	1,572	77.2 (69.1-83.6)	209		●	
QN41: Percentage of students who had their first drink of alcohol before age 13 years (other than a few sips)	14.1 (12.4-16.1)	1,555	30.9 (23.7-39.2)	212		●	
QN42: Percentage of students who currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)	29.8 (26.7-33.2)	1,582	44.3 (36.9-52.0)	207		●	
QN44: Percentage of students who currently were binge drinking (had four or more drinks of alcohol in a row for female students or five or more drinks of alcohol in a row for male students, within a couple of hours, on at least 1 day during the 30 days before the survey)	16.8 (14.3-19.7)	1,605	26.2 (20.7-32.6)	214		●	
QN46: Percentage of students who ever used marijuana (one or more times during their life)	32.1 (28.8-35.6)	1,604	51.1 (43.5-58.6)	217		●	
QN47: Percentage of students who tried marijuana for the first time before age 13 years	6.4 (5.2-7.9)	1,613	14.6 (10.4-20.2)	216		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey
Cecil County
Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual†	Gay, Lesbian, or Bisexual More Likely Than Heterosexual†	No Difference†
	% 95% CI*	N	% 95% CI	N			
QN48: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)	19.0 (16.4-21.9)	1,620	31.5 (24.8-39.0)	217		●	
QN49: Percentage of students who ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)	2.7 (1.9-3.7)	1,642	10.0 (6.5-14.9)	223		●	
QN51: Percentage of students who ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)	1.3 (0.8-2.0)	1,654	5.9 (3.3-10.3)	226		●	
QN52: Percentage of students who ever used methamphetamines (also called "speed," "crystal," "crank," or "ice," one or more times during their life)	1.5 (1.0-2.3)	1,650	8.2 (5.1-12.9)	227		●	
QN53: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)	3.1 (2.3-4.2)	1,645	11.8 (7.7-17.6)	227		●	
QN54: Percentage of students who ever used synthetic marijuana (also called "K2," "Spice," "fake weed," "King Kong," "Yucatan Fire," "Skunk," or "Moon Rocks," one or more times during their life)	4.7 (3.7-5.9)	1,653	16.8 (12.1-22.8)	227		●	

*Confidence interval.

†Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey
Cecil County
Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN56: Percentage of students who ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their	10.3 (8.8-12.1)	1,654	28.7 (23.1-35.1)	225		●	
QN57: Percentage of students who ever injected any illegal drug (used a needle to inject any illegal drug into their body, one or more times during their life)	1.6 (1.0-2.7)	1,648	11.6 (6.2-20.9)	225		●	
QN58: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)	14.6 (12.7-16.7)	1,651	23.6 (18.4-29.6)	221		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey
Cecil County
Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN59: Percentage of students who ever had sexual intercourse	37.7 (33.6-42.0)	1,503	52.7 (43.8-61.5)	194		●	
QN60: Percentage of students who had sexual intercourse for the first time before age 13 years	2.8 (2.0-4.0)	1,495	7.3 (4.1-12.9)	192		●	
QN61: Percentage of students who had sexual intercourse with four or more persons during their life	7.3 (5.9-8.9)	1,504	10.1 (6.0-16.3)	191			●
QN62: Percentage of students who were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)	26.6 (23.4-30.1)	1,501	38.0 (30.3-46.4)	191		●	
QN63: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	19.0 (14.9-23.9)	387	17.0 (10.0-27.3)	72			●
QN64: Percentage of students who did not use a condom during last sexual intercourse (among students who were currently sexually active)	44.4 (39.1-49.9)	382	62.9 (50.8-73.5)	70		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey
Cecil County
Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN65: Percentage of students who did not use birth control pills before last sexual intercourse (to prevent pregnancy among students who were currently sexually active)	70.5 (65.0-75.4)	372	78.4 (67.5-86.4)	69			●
QNIUDIMP: Percentage of students who did not use an IUD (e.g., Mirena or ParaGard) or implant (e.g., Implanon or Nexplanon) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	98.9 (97.0-99.6)	372	100.0 (.-)	69			●
QNSHPARG: Percentage of students who did not use a shot (e.g., Depo-Provera), patch (e.g., OrthoEvra), or birth control ring (e.g., NuvaRing) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	91.2 (87.2-94.0)	372	97.8 (91.1-99.5)	69		●	
QNOTHHPL: Percentage of students who did not use birth control pills; an IUD or implant; or a shot, patch, or birth control ring (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	60.5 (54.6-66.2)	372	76.2 (65.8-84.1)	69		●	
QNDUALBC: Percentage of students who did not use both a condom during and birth control pills; an IUD or implant; or a shot, patch, or birth control ring before last sexual intercourse (to prevent STD and pregnancy among students who were currently sexually active)	87.3 (83.3-90.5)	371	95.2 (85.9-98.5)	69		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey
Cecil County
Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy during last sexual intercourse (among students who were currently sexually active)	9.3 (6.4-13.3)	372	27.2 (17.3-40.0)	69		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey
Cecil County
Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual†	Gay, Lesbian, or Bisexual More Likely Than Heterosexual†	No Difference‡
	% 95% CI*	N	% 95% CI	N			
QNOBESE: Percentage of students who had obesity (>= 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts)¶	15.6 (13.4-18.1)	1,409	17.5 (12.1-24.6)	188			●
QNOWT: Percentage of students who were overweight (>= 85th percentile but <95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts)¶	16.3 (14.1-18.7)	1,409	19.3 (14.3-25.4)	188			●
QN68: Percentage of students who described themselves as slightly or very overweight	30.4 (28.3-32.7)	1,630	36.2 (29.2-43.8)	221			●
QN70: Percentage of students who did not drink fruit juice (100% fruit juices one or more times during the 7 days before the survey)	32.9 (30.5-35.5)	1,640	35.5 (28.1-43.7)	224			●
QN71: Percentage of students who did not eat fruit (one or more times during the 7 days before the survey)	14.4 (12.5-16.5)	1,638	19.1 (13.5-26.4)	224			●
QNFR0: Percentage of students who did not eat fruit or drink 100% fruit juices (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)	8.8 (7.3-10.5)	1,634	12.0 (8.0-17.7)	224			●

*Confidence interval.

†Based on t-test analyses, p<0.05.

¶Based on reference data from the 2000 CDC Growth Charts. In 2017, new, slightly different ranges were used to calculate biologically implausible responses to height and weight questions.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey Cecil County Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN72: Percentage of students who did not eat green salad (one or more times during the 7 days before the survey)	44.0 (41.2-46.8)	1,629	44.0 (37.4-50.9)	223			●
QN73: Percentage of students who did not eat potatoes (one or more times during the 7 days before the survey)	28.3 (26.0-30.7)	1,640	31.5 (26.0-37.6)	224			●
QN74: Percentage of students who did not eat carrots (one or more times during the 7 days before the survey)	50.4 (47.6-53.1)	1,636	51.4 (44.2-58.6)	224			●
QN75: Percentage of students who did not eat other vegetables (one or more times during the 7 days before the survey)	17.8 (15.8-20.1)	1,616	21.8 (15.3-30.1)	219			●
QNVEG0: Percentage of students who did not eat vegetables (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	6.3 (5.1-7.8)	1,601	7.4 (4.5-11.9)	218			●
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	17.5 (15.5-19.7)	1,635	21.1 (16.1-27.1)	221			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey

Cecil County

Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	11.2 (9.4-13.3)	1,635	14.3 (10.0-20.0)	221			●
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)	5.7 (4.5-7.3)	1,635	9.2 (5.7-14.5)	221			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey
Cecil County
Risk Behaviors and Sexual Identity Report

Physical Activity

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN79: Percentage of students who were not physically active at least 60 minutes per day on 5 or more days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	62.2 (59.4-65.0)	1,629	78.7 (72.0-84.1)	221		●	
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	16.2 (13.6-19.3)	1,629	23.7 (17.9-30.7)	221		●	
QNPA7DAY: Percentage of students who were not physically active at least 60 minutes per day on all 7 days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	81.7 (79.8-83.5)	1,629	89.6 (84.5-93.1)	221		●	
QN80: Percentage of students who watched television 3 or more hours per day (on an average school day)	23.0 (21.2-25.0)	1,622	25.7 (20.3-31.9)	222			●
QN81: Percentage of students who played video or computer games or used a computer 3 or more hours per day (counting time spent on things such as Xbox, PlayStation, an iPad or other tablet, a smartphone, texting, YouTube, Instagram, Facebook, or other social media, for something that was not school work, on an average school day)	39.2 (36.3-42.1)	1,625	55.6 (47.9-62.9)	217		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey
Cecil County
Risk Behaviors and Sexual Identity Report

Physical Activity

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN82: Percentage of students who did not attend physical education (PE) classes on 1 or more days (in an average week when they were in school)	54.4 (49.8-59.1)	1,605	60.1 (52.0-67.6)	216			●
QNDLYPE: Percentage of students who did not attend physical education classes on all 5 days (in an average week when they were in school)	95.5 (94.1-96.5)	1,605	98.0 (94.1-99.4)	216		●	
QN84: Percentage of students who had a concussion from playing a sport or being physically active (one or more times during the 12 months before the survey)	15.1 (13.3-17.1)	1,622	20.0 (14.8-26.4)	221			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey
Cecil County
Risk Behaviors and Sexual Identity Report

Other	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
Health Risk Behavior							
QN85: Percentage of students who were never tested for human immunodeficiency virus (HIV) (not including tests done when donating blood)	89.0 (86.9-90.8)	1,624	81.3 (75.0-86.3)	219	●		
QN86: Percentage of students who did not see a dentist (for a check-up, exam, teeth cleaning, or other dental work, during the 12 months before the survey)	18.9 (16.8-21.2)	1,624	33.9 (26.4-42.2)	221		●	
QNNODNT: Percentage of students who never saw a dentist (for a check-up, exam, teeth cleaning, or other dental work)	1.1 (0.7-1.9)	1,624	4.7 (2.4-8.8)	221		●	
QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma	25.9 (23.7-28.1)	1,616	40.8 (33.9-48.1)	220		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey
Cecil County
Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual†	Gay, Lesbian, or Bisexual More Likely Than Heterosexual†	No Difference†
	% 95% CI*	N	% 95% CI	N			
QN91: Percentage of students who missed school one or more days for any reason, with or without permission (during the 30 days before the survey)	60.4 (57.5-63.3)	1,625	75.5 (68.5-81.4)	220		●	
QN92: Percentage of students who missed school one or more days for any reason, without permission (during the 30 days before the survey)	15.6 (13.4-18.2)	1,624	23.4 (18.0-30.0)	220		●	
QN93: Percentage of students who talked on a cell phone while driving (on at least 1 day during the 30 days before the survey, among students who drove a car or other vehicle)	29.9 (25.1-35.1)	780	31.7 (22.3-43.0)	103			●
QN94: Percentage of students who who smoked a whole cigarette before age 13 years (for the first time)	6.2 (5.0-7.8)	1,627	18.0 (13.1-24.3)	226		●	
QN95: Percentage of students who usually obtained their own cigarettes by buying them in a store or gas station (during the 30 days before the survey among students who smoked cigarettes during the 30 days before the survey and who were aged <18 years)	9.6 (5.3-16.9)	103	17.8 (7.5-36.7)	34			●

*Confidence interval.

†Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey Cecil County Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN96: Percentage of students who were not asked to show proof of age (among students who bought or tried to buy cigarettes at a store during the 30 days before the survey)	54.0 (36.0-71.0)	33	-	21			
QN97: Percentage of students who did not have someone refuse to sell them cigarettes because of their age (among students who tried to buy cigarettes during the 30 days before the survey)	74.1 (63.7-82.3)	79	64.1 (45.2-79.4)	35			●
QN98: Percentage of students who smoked tobacco in a hookah, narghile, or other type of waterpipe (on at least 1 day during the 30 days before the survey)	3.0 (2.3-3.9)	1,611	10.0 (6.3-15.5)	217		●	
QN99: Percentage of students who usually use a kind of flavoring other than tobacco flavor with an electronic vapor product (among students who have ever used an electronic vapor product)	98.6 (96.9-99.4)	565	94.3 (85.8-97.9)	108			●
QN100: Percentage of students who have ever used an electronic vapor product to smoke marijuana, THC or hash oil, or THC wax	8.7 (7.1-10.7)	1,548	17.7 (13.1-23.3)	210		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

- = Fewer than 30 students in the denominator (subgroup).

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey
Cecil County
Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN101: Percentage of students who think they definitely or probably will smoke a cigarette, cigar, cigarillo, or little cigar in the next year	16.3 (14.0-18.9)	1,651	27.7 (22.2-34.0)	223		●	
QN102: Percentage of students who used flavored tobacco products (flavored cigars, cigarillos, or little cigars, flavored smokeless tobacco products, or both, excluding menthol cigarettes, during the 30 days before the survey)	8.6 (7.2-10.2)	1,627	13.5 (9.2-19.4)	215			●
QN103: Percentage of students who tried or used tobacco products for the first time (during the 12 months before the survey)	8.5 (7.3-9.9)	1,596	8.5 (5.4-13.2)	208			●
QN104: Percentage of students who did not completely quit using all tobacco products (during the 12 months before the survey, among students who used any tobacco products during the 12 months before the survey)	65.0 (59.4-70.2)	314	71.6 (58.5-81.8)	67			●
QN105: Percentage of students who live with someone who now smokes cigarettes or cigars	38.9 (36.1-41.7)	1,644	50.1 (42.6-57.5)	216		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey Cecil County Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN106: Percentage of students who report that the rule about smoking inside their home is that there are no rules about smoking, that smoking is allowed in some places or at sometimes, or that smoking is allowed anywhere, inside their home (not counting decks, garages, or porches)	16.9 (15.1-18.8)	1,645	38.8 (31.8-46.3)	222		●	
QN107: Percentage of students who were in the same room with someone who was smoking (on at least one day during the 7 days before the survey)	33.2 (30.3-36.2)	1,651	54.9 (46.7-62.9)	220		●	
QN109: Percentage of students who are transgender	1.1 (0.7-2.0)	1,642	15.0 (10.6-20.7)	222		●	
QN110: Percentage of students who used an indoor tanning device (such as a sunlamp, sunbed, or tanning booth [not including getting a spray-on tan], one or more times during the 12 months before the survey)	7.4 (6.1-9.0)	1,621	10.4 (6.9-15.3)	219			●
QN111: Percentage of students who do not participate in any extracurricular activities at school (such as sports, band, drama, clubs, or student government)	35.4 (31.8-39.1)	1,608	40.9 (33.5-48.7)	215			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey Cecil County Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN112: Percentage of students who were not taught or who were not sure if they were taught in any of their classes about the dangers of tobacco use (during the last school year)	32.8 (29.0-37.0)	1,606	34.8 (27.2-43.3)	214			●
QN113: Percentage of students who say definitely yes or probably yes that young people who smoke have more friends	26.2 (23.8-28.8)	1,605	30.1 (23.9-37.1)	215			●
QN114: Percentage of students who say definitely yes or probably yes that smoking makes young people look cool or fit in	14.8 (13.0-16.9)	1,598	23.4 (16.6-31.7)	214		●	
QN115: Percentage of students who strongly or somewhat approve or who neither approve nor disapprove of someone their age having one or two drinks of an alcoholic beverage nearly every day	37.2 (34.3-40.2)	1,581	49.1 (41.6-56.7)	209		●	
QN116: Percentage of students who think people are at no risk or slight risk of harming themselves (physically and in other ways) if they have five or more drinks of alcohol once or twice a week (such as beer, wine, or liquor)	22.0 (19.3-24.9)	1,583	23.3 (17.9-29.8)	210			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2016 YOUTH RISK BEHAVIOR SURVEY RESULTS

Maryland High School Survey
Cecil County
Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN117: Percentage of students who do not have an adult outside of school they can talk to about things that are important to them	10.3 (8.9-11.9)	1,575	27.3 (21.2-34.3)	206		●	
QN118: Percentage of students who would not feel comfortable seeking help from adults besides their parents if they had an important question affecting their life	16.6 (14.6-18.8)	1,594	30.1 (23.6-37.5)	208		●	
QN120: Percentage of students who strongly disagree, disagree, or are not sure that their teachers really care about them and give them a lot of encouragement	49.7 (46.5-53.0)	1,590	53.8 (45.6-61.8)	211			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.