



Dental First Aid



For more information on oral health, visit health.maryland.gov/oral-health.

Non-Discrimination Statement

The Maryland Department of Health (MDH) complies with applicable federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, or disability in its health programs and activities.

English

Help is available in your language: 410-767-5300 (TTY: 1-800-735-2258).
These services are available for free.

Español/Spanish

Hay ayuda disponible en su idioma: 410-767-5300 (TTY: 1-800-735-2258).
Estos servicios están disponibles gratis.

Chinese

中文
用您的语言为您提供帮助: 410-767-5300
(TTY: 1-800-735-2258) 这些服务都是免费的



Fluoride and Water Testing Laboratories

County	Laboratory	Testing Details
Baltimore	Enviro-Chem Laboratories, Inc. 410-472-1112	Please call for water sample requirements. Lab supplies container. Cost: \$20 per sample.
Baltimore	Martel Laboratories JDS, Inc. 410-825-7790	Please call for water sample requirements. Lab supplies container. Cost: \$25 per sample.
Baltimore	Phase Separation Science, Inc. 410-747-8770	Please call for water sample requirements. Lab supplies container. Cost: \$35 per sample. Results in 5 business days.
Baltimore City	Microbac Laboratories Inc. Baltimore Division 410-633-1800	Please call for water sample requirements. Lab supplies container. Cost: \$20 per sample. Results in 7 business days.
Dorchester	Environmental Health Division (Dorchester County Health Department) 410-228-1167	Please call for water sample requirements. Lab supplies container. Cost: \$50 per sample. Results in 4-6 weeks.
Frederick	Fredericktowne Labs, Inc. 301-293-3340	Please call for water sample requirements. Lab supplies container. Cost: \$25 per sample. Results in 3-5 business days.
Frederick	Catoctin Labs, Inc. 301-663-5323	Please call for water sample requirements. Use standard 250 mL bottle for collection. Cost: \$20 and collection fees per sample. Results in 5 business days.
Queen Anne's	Chesapeake Environmental Laboratory, Inc. 410-643-0800	Please call for water sample requirements. Lab supplies container. Cost: \$20 per sample. Results in 5 business days.
Queen Anne's	Water Testing Aardvark Laboratory 410-643-7711	Please call for water sample requirements. Lab supplies container. Cost: \$35 per sample. Results in 7-10 business days.
Queen Anne's	Environmental Health Division (Queen Anne's County Local Health Department) 410-748-2281	Please call for water sample requirements. Lab supplies container. Cost: \$20 per sample. Results in 5 business days.
Washington	Washington County Division of Environmental Management Laboratory 240-313-2604	Please call for water sample requirements. Lab supplies container. Cost: \$12 per sample. Results in 2 weeks.
Wicomico	Eastern Shore Regional Lab 410-219-9005	Please call for water sample requirements. Lab supplies container. Cost: \$95 per sample; no cost for doctors. Results in 28 days.

What is Fluoride? Approximately 70 years of evidence supports the effectiveness and safety of fluoride in community water systems and its ability to prevent, reduce, or even reverse the onset and development of tooth decay. Through continuous research, drinking optimally fluoridated water has been scientifically proven to be safe and effective.

Fluoride is a natural element found in rocks and soil, in fresh water, and in ocean water. Waters in and around the United States have natural fluoride levels that range from 0.1 to more than 12 parts per million (ppm). It is known that in order to prevent tooth decay, the optimal fluoride level for a community water system is 0.7ppm. When the natural fluoride level in a community water system is below 0.7 ppm, cities and towns add supplemental fluoride to their community water systems as a cost efficient way to help reduce tooth decay.

What are the Benefits of Fluoridated Water?

- Fluoridated water:
- ◆ Prevents tooth decay.
 - ◆ Strengthens tooth enamel.
 - ◆ Kills germs that cause tooth decay.
 - ◆ Is a cost effective means of improving community health.

The addition of supplemental fluoride to drinking water can be compared to the addition of supplemental Vitamin D to milk.

- ◆ Both fluoride and vitamin D occur naturally and provide significant health benefits when present in a person’s diet at sufficient levels.
- ◆ Sufficient levels of Vitamin D in a person’s diet can prevent rickets in children and bone disease in adults. Sufficient levels of fluoride in a person’s drinking water can reduce tooth decay and strengthen tooth enamel.

97% of the Maryland population on community water systems, or a total of 5,060,379 million people, had access to optimally fluoridated water in 2012 according to CDC data. Maryland ranks 4th in the nation for community water fluoridation and surpasses the Healthy People 2020 target of 79.6%.

Having Water Tested for Fluoride

Consult your medical or dental provider before making any decisions about dietary fluoride supplements (vitamins) which come in the form of liquid drops or tablets. If you do not know the concentration of fluoride in your community water supply, it is important to contact your local health department before a medical or dental provider prescribes fluoride supplements for children ages 6 months - 16 years of age. If you are on a private well, you should have it tested before taking any fluoride supplements.

Dietary Fluoride Supplementation Schedule

Age	<0.3 ppm F	0.3 to 0.6 ppm F	>0.6 ppm F
Birth to 6 months	0	0	0
6 mo to 3 years	0.25 mg	0	0
3 to 6 years	0.50 mg	0.25 mg	0
6 to 16 years	1.00 mg	0.50 mg	0

Approved by the American Dental Association Council on Scientific Affairs—2010

To Learn More, Check Out:

Centers for Disease Control and Prevention _ www.cdc.gov
American Dental Association _____ www.ada.org
National Institutes of Health _____ www.nidcr.nih.gov
Campaign for Dental Health _____ www.ilikemyteeth.org
Children’s Dental Health Project _ [www.cdhp.org/
resources/309-fluoride-
fluoridation](http://www.cdhp.org/resources/309-fluoride-fluoridation)

This flipchart is designed to aid school nurses, school health aides, pre-school and school teachers, pre-school and school administrators, daycare coordinators, healthcare providers, and other health and administrative personnel in the effective management of dental emergencies and basic oral health principles, prevention, and care. Although these first aid procedures should provide temporary relief and assistance, they are not intended to permanently resolve a dental problem.

A local dentist or physician should be consulted as soon as possible in most situations. If a child does not have a dentist, please contact:

- ◆ Your local health department (see tab for Local Dental Services/Oral Health Resource Guide)
- ◆ Maryland Healthy Smiles Program (Medicaid eligible)_____ 1-855-934-9812 (program members)
www.member.MDhealthysmiles.com
1-844-275-8753 (dental providers)
www.provider.MDhealthysmiles.com
- ◆ Maryland State Dental Association_____ 410-964-2880
www.msda.com
- ◆ Maryland Department of Health, Office of Oral Health_____ 410-767-5300
health.maryland.gov/oralhealth

Proper toothbrushing is not difficult. Just follow these steps:



Brush with Fluoride Toothpaste

2 minutes

+2 times a day

4 a Healthy Smile



Place the toothbrush against the gum line at a 45-degree angle. Use a small circular motion to brush your teeth.



Brush back and forth on the chewing surfaces of your teeth holding the brush flat.



Brush the back of your teeth!



Brush your tongue! This will freshen your breath and remove germs.

Brush for at least two minutes, two times a day.

Be sure to brush at bedtime to remove germs that can cause decay while you are sleeping.

LOSS OF PRIMARY (BABY) TOOTH

If a primary tooth (baby tooth) is knocked out and traumatically lost:

- ◆ DO NOT attempt to replace the tooth in the child’s mouth.
- ◆ Control any bleeding and check for other injuries. If bleeding doesn’t stop, contact a dentist or physician immediately.
- ◆ Notify the child’s parent or caregiver.
- ◆ Find the tooth and send it with the parent or caregiver to the dentist.
- ◆ If the injury site has been contaminated with soil, check the child’s records for date of the most recent tetanus shot.

IF A PERMANENT TOOTH IS KNOCKED OUT AND IS INTACT, TIME IS CRITICAL!

LOSS OF PERMANENT TOOTH

Contact the parent or caregiver and arrange to have the child taken to a dentist IMMEDIATELY. Many times the tooth can be successfully reimplanted and saved if accomplished within ONE HOUR. Look at the accident area for the tooth that was knocked out.

If found and dirty (prior to going to a dentist):

- ◆ Gently rinse the tooth under tap water (remember to plug sink!) DO NOT CLEAN OR SCRUB TOOTH! Hold tooth by the crown (top part of the tooth) and gently tease tooth back into socket. Be careful not to put the tooth in backwards!
- ◆ Child and/or parent or caregiver should hold tooth in socket while being transported to a dentist (the child may do this by biting on a clean gauze or cloth).

If the tooth cannot be placed back into the socket:

- ◆ Place tooth in a glass of cold milk or saline (whichever is available).
- ◆ Avoid storage in water.
- ◆ If the wound site has been contaminated by dirt or soil, record this information and send it with the injured child to aid the dentist in determining the necessity for a tetanus shot.



Joanna Douglass, BDS, DDS

Empty socket of an avulsed permanent central incisor



Photo credit: Washington Interdisciplinary Children’s Oral Health Promotion (ICOHP)

DENTAL TRAUMA

In all occasions of oral or dental trauma, serious head injury should be ruled out.

SIGNS AND SYMPTOMS OF CONCUSSION MAY INCLUDE:
NAUSEA/VOMITING, DIZZINESS/HEADACHE, DILATED PUPILS, COLD/CLAMMY SKIN.

POSSIBLE JAW DISLOCATION OR FRACTURE

In all occasions of oral or dental trauma, serious head injury should be ruled out. Signs and symptoms of concussion may include: nausea/vomiting, dizziness/headache, dilated pupils, cold/clammy skin.

Note time of injury and other injuries that may be present.

If a jaw fracture or dislocation is suspected, do the following:

- ◆ Contact the parent or caregiver immediately and arrange for the child to be taken to an oral surgeon or hospital emergency room;
- ◆ Elevate the head and apply a cold compress;
- ◆ Stabilize the patient as well as possible;
- ◆ Monitor vital signs (breathing, pulse) and consciousness and record all observations. If any loss of consciousness or difficulty breathing occurs, call your institutional emergency telephone number immediately!

BROKEN TOOTH

- Control any bleeding;
- Avoid further trauma to the area if possible;
- Rinse dirt from injured area with warm water;
- Apply a cold compress on the cheek next to the injured tooth to reduce swelling;
- Observe broken end of tooth for bleeding;
- Find broken tooth fragments if possible. Keep tooth piece in gauze moistened with water;
- Make an immediate referral to a dentist.

DISPLACED TOOTH

- If the tooth has been pushed up into the socket or gum by trauma, control bleeding and be supportive.
- Immediate referral to a dentist.
- The tooth may discolor long after the original trauma and will need attention by a dentist at that time.

DO NOT ATTEMPT TO PULL THE TOOTH INTO POSITION.



Larger “chip” that involves the enamel and dentin



Tooth fracture involving the root



Intruded lateral incisor in a 3 year old

Photo credit: Washington Interdisciplinary Children’s Oral Health Promotion (ICOHP)

TOOTHBRUSHING

Plaque contains a sticky layer of germs which can lead to tooth decay and gum disease. Brush with toothpaste containing fluoride for at least two minutes, twice a day (after breakfast and before bed). This keeps the teeth and gums healthy.


Brushing Tips:

- ◆ Always use a soft-bristle toothbrush and always use toothpaste with fluoride. Use the correct amount of toothpaste according to age. (see chart)
- ◆ Replace your toothbrush at least every 3 months.
- ◆ Never share toothbrushes, it spreads germs.
- ◆ Children at high risk for tooth decay should use toothpaste that has fluoride as soon as the first tooth appears.
- ◆ Children younger than age 8 need supervision to make sure they do not use too much toothpaste and that they brush their teeth thoroughly.

How Much
Toothpaste with
Fluoride?



Under age 3: Smear about the size of a grain of rice



Age 3 and older: Pea-sized amount

Spit, do not rinse after brushing.

TEETHING

Babies begin teething around 5 months of age. Local discomfort is common with tooth eruption and can be associated with cold, fever or diarrhea.

Strategies to ease the pain and discomfort of teething

Keep it safe: Make sure liquid-filled teething toys are made of durable materials that the baby can not chew a hole into. Use plastic teething toys that do not contain bisphenol A (BPA). Find teething toys that do not have loose pieces that could break off in baby’s mouth and cause choking.

Cool it: Give the baby a refrigerated clean wet washcloth, pacifier, or teething ring to chew on. Cold helps the pain of sore gums. Teething rings should NOT be put in the freezer.

Freeze it: Unlike teething rings, certain frozen foods such as sliced peaches, can help ease teething pain.

Massage it: Gently rub the baby’s gums with a clean finger for 2 minutes. Many babies find the pressure soothing.

Don’t use it: Oral health and medical providers do not recommend using teething gels and liquids on babies’ gums.



FLOSSING

Flossing removes plaque between teeth and above and below the gum line - areas a toothbrush cannot reach.

By combining; brushing with toothpaste that contains fluoride and flossing at the same time everyday (and with the appropriate diet), you can thoroughly prevent the germ-containing plaque from causing cavities and gum disease.

Children under the age of 8 should not floss without an adult’s help. A dentist or dental hygienist can show you how to floss your child’s teeth.

TOOTH EXTRACTION COMPLICATIONS

The child has been instructed by the dentist:

Not to rinse or swish for 24 hours after an extraction, as this could wash out the blood clot forming at the extraction site. Normal drinking is permissible. Instruct the child to follow these steps:

- ◆ Not to use straws for 24 hours; the suction may dislodge the blood clot.
- ◆ Not to eat excessively cold or hot foods for 24 hours because this could dislodge the blood clot.
- ◆ Not to smoke since smoking could delay tissue healing.
- ◆ Not to spit or play with the extraction site (especially with the tongue).

If the bleeding is determined to be more than oozing (has a bright red color) or is alarming the child, the following is recommended:

- ◆ Fold a 2” x 2” sterile gauze pad and place on the extraction site (wearing gloves), having the child bite down on it for about 30 minutes. Replace soaked 2” x 2” gauze pads as necessary.
- ◆ If bleeding cannot be controlled within an hour or is extensive, advise the parent or caregiver to immediately call the child’s dentist’s office or take the child to a hospital emergency room.



ORTHODONTIC (BRACES) OR OTHER APPLIANCES

If a wire or appliance becomes loose or broken and cannot be removed easily, contact the parent or caregiver to take the child to a dentist (preferably to their orthodontist) immediately.

If a broken appliance can be removed easily, take the following steps:

- ◆ A blunt item (tongue depressor) may be used to gently bend the wire so that it is no longer irritating oral tissue.
- ◆ If cheek or gum is pierced by a wire, ease the tissue off the wire if possible. Cover the wire with cotton gauze, cotton balls, or wax and contact the parent or caregiver for referral to the orthodontist.
- ◆ If the protruding wire cannot be bent, simply cover the end of it with a piece of cotton gauze, a cotton ball or wax so that it is no longer causing irritation.
- ◆ DO NOT REMOVE THE WIRE. Contact the parent or caregiver for referral to the child’s orthodontist.
- ◆ Do not attempt to remove the wire if it is broken off and/or embedded in the cheeks, gums or tongue.

Note: Most children with braces have orthodontic wax and know how to apply it.

GUM INFLAMMATION

A blow (trauma) to the mouth can cause the gum tissue to swell and bleed. Evaluate the child for signs and symptoms of a concussion and additional injuries if trauma caused the bleeding.

A cold compress may be applied to the area from the outside of the cheek to help control swelling.

If appropriate, while wearing gloves, apply direct pressure with a sterile 2” x 2” gauze to the injured gum or cheek to control the bleeding and contact the child’s parent or caregiver according to institutional policy.



Inflamed or irritated gum tissue and/or gingivitis can be caused by:

- ◆ **Poor oral hygiene** - this can be corrected by daily removal of plaque by brushing and flossing with toothpaste that contains fluoride.
- ◆ **Puberty** (hormones associated with adolescence)
- ◆ **Pregnancy** (increased hormone levels)
- ◆ **Smoking** (affects gum tissue)

Sudden bleeding or swelling of the gums may be the result of food or foreign body impaction. Locate and, if possible, remove the debris with a soft bristled toothbrush. You may need to wear gloves (non-latex).

In general, red, swollen gums should be rinsed thoroughly with a warm salt water solution (containing one dissolved teaspoon of salt in an 8 oz. glass of warm water) for 15-30 seconds and spit out completely. The child’s parent or caregiver should be contacted for referral to a dentist if the bleeding cannot be controlled.

OBJECTS WEDGED BETWEEN TEETH

If an object becomes wedged between teeth:

- ◆ The child may use dental floss to remove the object, if age appropriate.
- ◆ Remember to assist the child in guiding the floss gently between teeth, to prevent injury to gum tissue.
- ◆ Do not try to remove the object with a sharp or pointed tool/instrument. This may result in injury.
- ◆ If unsuccessful, contact the parent or caregiver for referral to a dentist.

Flossing is as easy as brushing once you know how.

Follow these steps:



Wrap about 18 inches of floss around the middle fingers.

Slide the floss between the teeth. (Do not snap the floss into the gums but guide the floss gently just below the gumline).

As the floss reaches the gum line, make a C-shape around the tooth until you feel pressure against your tooth.

Gently scrape the tooth surface with the floss.

Move the floss as it becomes soiled and repeat the process for each tooth. Floss every day.

If assisting a child to floss, you should wear gloves (non-latex).

DENTAL FIRST AID SUPPLIES

The following should be present in your first aid kit for use in dental emergencies:

- Disposable non-latex gloves (should be worn at all times)
- Mask
- Eyewear
- Cotton swabs
- Sterile gauze squares 2” x 2” or pads
- Toothbrushes
- Dental floss
- Ice pack or wet frozen washcloth
- Saline or salt water
- Flashlight
- Tongue blade or dental mouth mirror



Medications cannot be used in most schools without a written approved protocol specific to each drug and signed by a physician or dentist and written permission from a parent or caregiver.

When examining the mouth and surrounding structures, ALWAYS wash your hands (before and after) and ALWAYS wear non-latex gloves!

TOOTH ERUPTION, SHEDDING, AND TOOTH PAIN



Erupting Primary (baby) Teeth and Teeth Pain

Local discomfort is common with tooth eruption and can be associated with cold, fever or diarrhea.

See tab for **Teething and Flossing** for strategies to help with infant teething.

Eruption sites (locations where teeth are about to appear) should be hard and blanched (white). If an eruption site is soft, it may indicate an eruption cyst. In this instance, contact the parent or caregiver for referral to a dentist.

Shedding (losing) Primary (baby) Teeth and Teeth Pain

- ◆ Pain associated with losing primary teeth is normal.
- ◆ Encourage the parent to remove the tooth at home.
- ◆ Determine if pain is due to incomplete shedding of the primary tooth. If so, refer the child to a dentist.
- ◆ Advise the child to avoid the site while eating.

Erupting Permanent Teeth and Teeth Pain

No treatment is necessary unless the child is experiencing prolonged pain (more than 24 hours). This may be caused by inflammation around an impacted or partially impacted tooth.

Primary (baby) Teeth

Upper Teeth		Erupt	Shed
Central incisor		8-12 mos.	6-7 yrs.
Lateral incisor		9-13 mos.	7-8 yrs.
Canine (cuspid)		16-22 mos.	10-12 yrs.
First molar		13-19 mos.	9-11 yrs.
Second molar		25-33 mos.	10-12 yrs.
Lower Teeth		Erupt	Shed
Second molar		23-31 mos.	10-12 yrs.
First molar		14-18 mos.	9-11 yrs.
Canine (cuspid)		17-23 mos.	9-12 yrs.
Lateral incisor		10-16 mos.	7-8 yrs.
Central incisor		6-10 mos.	6-7 yrs.

Used with permission of the Arkansas Office of Oral Health

Permanent Teeth

Upper Teeth		Erupt
Central incisor		7-8 yrs.
Lateral incisor		8-9 yrs.
Canine (cuspid)		11-12 yrs.
First premolar (first bicuspid)		10-11 yrs.
Second premolar (second bicuspid)		10-12 yrs.
First molar		6-7 yrs.
Second molar		12-13 yrs.
Third molar (wisdom tooth)		17-21 yrs.
Lower Teeth		Erupt
Third molar (wisdom tooth)		17-21 yrs.
Second molar		11-13 yrs.
First molar		6-7 yrs.
Second premolar (second bicuspid)		11-12 yrs.
First premolar (first bicuspid)		10-12 yrs.
Canine (cuspid)		9-10 yrs.
Lateral incisor		7-8 yrs.
Central incisor		6-7 yrs.

Used with permission of the Arkansas Office of Oral Health

CHILD ABUSE / DENTAL NEGLECT

Tooth decay is the most common chronic disease in childhood, five times more common than asthma. Tooth decay is an infection. Left untreated, tooth decay can cause pain, malnutrition, difficulty concentrating and learning, and poor overall health.

The Code of Maryland Regulations (COMAR) requires health practitioners, educators, human service workers and police officers to report any suspected child abuse and neglect to local department of social services or to a local law enforcement agency. Dental neglect is a form of child abuse not often considered though it may have profound repercussions on the life of a child.

Dental neglect is the willful failure of a parent or guardian to seek and follow through with treatment necessary to ensure a level of oral health essential for adequate function and freedom from pain and infection.

Source: American Academy of Pediatric Dentistry

Some Signs of Dental Neglect

- ◆ Child is unable to eat normally;
- ◆ Child does not smile with his or her teeth;
- ◆ Child demonstrates changes in behavior (for example, puts hand in front of mouth, failure to thrive);
- ◆ Crying, in constant pain;
- ◆ Teeth appear rotted, grossly discolored; and
- ◆ Perpetually bad mouth odor.

TO REPORT NEGLECT

Maryland Child Protective Services


1-800-332-6347

District of Columbia Family Services Administration

202-671-7233

Delaware Child Protective Services

1-800-292-9582



Facial swelling (cellulitis) related to untreated dental decay.



Photo credit: mchoralhealth.org
Abscesses are present on upper gum, teeth have severe decay.



Photo credit: midatlanticpanda.org
View shows teeth decayed to the gumline.

TOOTHACHE AND SWELLING

The following steps should be taken for a child with suspected toothache or facial swelling:


- ◆ Clean the area around the sore tooth thoroughly by rinsing the mouth with warm salt water.
- ◆ Check the child’s temperature and observe for respiratory distress.
- ◆ Administer appropriate over-the-counter pain medication if needed.



DO NOT PLACE ASPIRIN ON THE GUM OR ACHING TOOTH!

- ◆ If the child’s face is swollen, apply a cold compress or ice. Alternate 10 minutes on and 5 minutes off.
- ◆ If the child has a localized abscess or gum boil (with or without pain), avoid heat.
- ◆ Refer to a dentist.

NOTE: Your local health department, the Maryland Office of Oral Health, or the Maryland State Dental Association can be contacted for names of dentists who accept emergency patients. In addition, area dental schools, and other community dental clinics can be contacted for emergency care. (See **Local Dental Services** in this flip chart or Oral Health Resource Guide on the web health.maryland.gov/oral-health for contact information.)



ORAL ULCERS

Oral ulcers can be the result of:

- ◆ Aphthous lesion/canker sore (on softer mouth tissues)
- ◆ Cold sore/fever blister or traumatic lesion (on harder mouth tissues)

The following steps should be taken for ulcers:

- ◆ Always wear non-latex gloves if touching lesions. Before and after inspecting lesions and removing gloves, wash hands thoroughly.
- ◆ Observe the location, type and severity of lesions.
- ◆ Take the child’s temperature. Administer appropriate mild over-the-counter pain medication (if needed and possible).
- ◆ Tell the child to avoid spicy, hard, crunchy foods.
- ◆ Apply ice to the area to provide temporary relief.
- ◆ Contact the child’s parent or caregiver for referral to a physician or dentist if fever and/or lesions persist.



Cold sore/fever blister
Photo credit: www.healthtap.com




Aphthous lesion/canker sore
Photo credit: www.fitnesshealthpros.com

Oral ulcers can be a symptom of:

- Herpes virus
- Mumps
- German measles
- Impetigo
- Chicken pox
- Streptococcal infection
- Stress
- Other diseases or disorders

Nutrition and Oral Health



The germs that cause tooth decay feed on sugar and produce acid. The acid attacks enamel, the protective outer layer of a tooth. Once the enamel is worn down, a cavity is formed. Unless it is treated by a dentist, the cavity will grow until it destroys the tooth.

Every sip or bite of a sugary food or drink starts a new acid attack. Acid attacks last 20-40 minutes. Frequent sipping and snacking result in a continuous acid attack.

Sodas, sports drinks, sweet tea and juice contain sugar and acid which can erode tooth enamel. Even diet or “sugar free” soft drinks have acids that can harm teeth. Sodas have no nutritional value, only empty calories!



More and more people are becoming concerned about sugary drinks — the empty calories, the long-term impact on kids' tastes, the potential for obesity and diabetes.

The Better Beverage Finder website helps families make better choices. The site allows users to search for beverage choices any mother can love and find where to get them in Howard County. If you don’t live in Howard County, you can discover the best choices to look for in your community.

Visit the Resources section at the website for free downloadable campaign posters. Help spread the news about better beverages!

www.betterbeveragefinder.org



HOW MUCH SUGAR?

The SMALLER the acid number, the STRONGER the acid!	Acid*	Sugar**
	Low = Bad	Per 12 ounce Serving (1 can)
Pure Water	7.00	0.0 tsp.
Barq’s Root Beer	4.61	10.7 tsp.
Diet 7UP	3.67	0.0 tsp.
Sprite	3.42	9.0 tsp.
Diet Coke	3.39	0.0 tsp.
Mountain Dew	3.22	11.0 tsp.
Fresca	3.20	0.0 tsp.
Orange Slice	3.12	11.9 tsp.
Diet Pepsi	3.05	0.0 tsp.
Nestea	3.04	5.0 tsp.
Gatorade	2.95	3.3 tsp.
Dr. Pepper	2.92	9.5 tsp.
Hawaiian Fruit Punch	2.82	10.2 tsp.
Orange Minute Maid Soda	2.80	11.2 tsp.
Coke Classic	2.53	9.3 tsp.
Pepsi	2.49	9.8 tsp.
Battery acid	1.00 (ouch!)	0.0 tsp.

*Laboratory tests, University of Minnesota School of Dentistry, 2000.

**USDA: 4.2 grams = 1 teaspoons of sugar



INJURY PREVENTION

Preventing oral injuries is important for many reasons. Injured primary (baby) teeth can turn brown or black, be painful, become infected, or have to be removed. Primary teeth keep space for permanent teeth; injuries to a child’s primary teeth can damage the permanent teeth. Injuries to primary or permanent teeth can also affect a child’s speech, nutrition, self-esteem, and overall health.

Dental injuries are the most common type of injuries to the face. Half of these injuries can be prevented.



In the United States, five million teeth are knocked out each year - mainly front teeth. 33 percent of all five-year-olds have injured their primary teeth. 60 percent of facial injuries occur during sports practice. The cost to repair a broken tooth is more expensive than a mouth guard.

Wearing a Mouth Guard Helps Protect Teeth, Lips and Tongue from Trauma.

Types of Mouth Guards	Oral Tissue Injury
<p>Stock Mouth Guard</p> <p>Inexpensive, pre-formed, ready to wear, can be bought at many sporting goods stores.</p> <p>Often do not fit very well, may be bulky, may make breathing and talking difficult.</p>	<p>In all cases of trauma, rule out serious head injury.</p> <ul style="list-style-type: none">◆ Have the child rinse and spit repeatedly with warm water.◆ Apply firm but gentle pressure (wearing non-latex gloves after washing hands) to the bleeding area with 2"x2" sterile gauze for at least 5 minutes. If injury site has been contaminated with soil, check the child’s records for date of most recent tetanus shot.◆ Vigorous bleeding may be expected initially.◆ Remove foreign body if easily visible and accessible.◆ Carefully check for broken/fractured teeth and avulsed (knocked out) teeth. Also look for tooth fragments in mouth, lip and cheek.◆ If the tooth has been shoved into the socket, contact a dentist immediately.◆ If swelling or bruising is present, apply ice, alternating 10 minutes on and 5 minutes off (lip injury).◆ Notify parent or caregiver.◆ If bleeding still persists after 15 minutes or cannot be controlled by simple pressure, or if the injury is severe, the child should be taken to the hospital emergency room, according to constitutional policy.
<p>Boil and Bite Mouth Guard</p> <p>Most commonly used, can be bought at many sporting goods stores.</p> <p>Should be softened in water, inserted and allowed to adapt to the shape of the mouth.</p>	
<p>Custom-Fitted Mouth Guard</p> <p>Made by a dentist, best at preventing injury.</p> <p>Most expensive type of mouth guard.</p>	

ORAL PIERCING COMPLICATIONS

There are many potential complications from piercing in and around the mouth. Most adolescents are unaware of the complications of intraoral piercing.

<p>Possible Complications</p> <ul style="list-style-type: none">Tooth fracture or injuryStud aspirationAllergic reactionNerve damageSpeech impedimentGingival (gum) recessionInfection	 <p>Radiograph shows root recession</p> <p>Photo credit: Washington Interdisciplinary Children’s Oral Health Promotion (ICOHP)</p>	 <p>Gum recession</p> <p>The inner surface of the lip bolt</p> <p>Photo credit: Washington Interdisciplinary Children’s Oral Health Promotion (ICOHP)</p>
---	--	--

Management of Complications

If the adolescent presents with inflammation around a piercing, refer the child to a dentist or physician who should:

- ◆ Remove the jewelry;
- ◆ Perform local debridement;
- ◆ Start antibiotics, if needed;
- ◆ Provide close follow-up.

Complication of oral infection may include sepsis and airway obstruction.

Reference: Smiles for Life, A National Oral Health Curriculum. Course 4: Acute Dental Problems “Oral Piercing Complications.” www.smilesforlifeoralhealth.org, 01/13/2015.

Dental Schools

Howard University Dental School, Washington, DC

New Patients.....202-806-0008

Pediatric Clinic.....202-806-0307

University of Maryland Dental School, Baltimore, MD

Pediatric Clinic.....410-706-4213

Adult Clinic410-706-7063

Clinical Dental Programs Operated by

Local Health Departments and Federally Qualified Health Centers

Allegany County

Allegany County Health Dept.

301-759-5030

Anne Arundel County

Anne Arundel County Health Dept.

Annapolis Health Center

410-222-7138

N. County Health Service Center

410-222-6861

Chase Brexton Health Services

410-837-2050

Baltimore City

Baltimore City Health Dept.

Druid Dental Clinic

410-396-0840

Eastern Dental Clinic

443-984-3548

Chase Brexton Health Services

410-837-2050

Family Health Centers of Baltimore

410-354-2000

Healthcare for the Homeless

410-837-5533

Park West Health Center

410-542-7800

Total Health Care

410-383-8300

University of MD Rehabilitation & Orthopaedic Institute

UM Rehab Dentistry

410-448-6290

Cleft Palate Clinic

410-448-6383

University of Maryland School of Dentistry

General Dentistry (Pre-Doctoral Clinic)

410-706-7063

Advanced General Dentistry (Post-Doctoral Clinic)

410-706-2940

Urgent Care (Emergency) Dentistry

410-706-2716

Pediatric Dentistry

410-706-4213

Orthodontics

410-706-7803

Endodontics

410-706-2860

Faculty Practice

410-706-7961

Periodontics

410-706-0896

Prosthodontics

410-706-7952

University of MD Medical System

Dept. of Oral and Maxillofacial Surgery

410-328-5566

Baltimore County

Baltimore County Health Dept.

410-887-2780

Chase Brexton Health Services

410-837-2050

Calvert County

Calvert Community Dental Care

410-535-8402

Caroline County

Choptank Community Health System

Federsburg Dental Center

410-754-7583

Goldsboro Family Dental Center

410-482-2224

Carroll County

Access Carroll

410-871-1478

Carroll County Health Dept.

410-876-4928

Cecil County

West Cecil Health Center

410-378-9696

Charles County

Charles County Health Dept.

301-609-6886

Health Partners, Inc.

301-645-3556

Dorchester County

Choptank Community Health System

Cambridge Dental Center

410-228-9381

Frederick County

Frederick County Health Dept.

301-600-1041

Frederick Health Dental Clinic.

.....240-566-7005

Garrett County

Garrett County Health Dept.

401-334-7660

Harford County

Harford County Health Dept.

443-922-7670

Howard County

Chase Brexton Health Services

410-884-7831

Kent County

refer to Caroline County listings

Montgomery County

CCI Health & Wellness Services

Gaithersburg Dental Clinic

240-720-0510

Rockville Dental Clinic

301-340-7525

Silver Spring Dental Clinic, Fenton St.

301-585-1250

Silver Spring Dental Clinic, Lamberton Dr.

301-364-9390

Montgomery County Health Dept.

Colesville Adult Dental Clinic

240-777-4471

Fenton Street Dental Clinic

240-777-3135

Germantown Health Center

240-777-3290

Metro Court Adult Dental Clinic

240-773-0304

Dennis Avenue Health Center

240-777-1875

Prince George’s County

CCI Health & Wellness Services

240-624-2323

Greater Baden Medical Services

Brandywine

301-888-2233

Oxon Hill

301-686-1665

Prince George’s County Health Dept.

301-583-5900

Deamonte Driver Dental Project

301-593-5900

Queen Anne’s County

refer to Caroline County listings

Somerset County

Chesapeake Health Care Dental

Princess Anne Clinic

410-651-5151

Salisbury Clinic

443-944-9600

St. Mary’s County

St. Mary’s County Health Dept.

301-475-4316

Talbot County

refer to Caroline County listings

Washington County

Family Healthcare of Hagerstown

301-745-3777

Wicomico County

Wicomico County Health Dept.

410-334-3401

Worcester County

Worcester County Health Dept.

410-641-0240

Download the

Maryland Oral Health Resource Guide

(health.maryland.gov/oral-health) for

a complete listing of dental services

provided by local health departments

and other public dental programs.

Maryland Oral Health

Resource Guide

2020 Edition

Office of Oral Health
Cancer and Chronic Disease Bureau
Prevention and Health Promotion Administration
Maryland Department of Health

