

CODING FOR
Pediatric
Preventive
Care **2021**



**Bright
Futures™**

prevention and health
promotion for infants,
children, adolescents,
and their families™

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



Coding for Pediatric Preventive Care, 2021

This resource contains comprehensive listings of codes that may not be used by your practice on a regular basis. We recommend that you identify the codes most relevant to your practice and include those on your encounter form or billing sheet.

Following are the *Current Procedural Terminology (CPT®)*, Healthcare Common Procedure Coding System (HCPCS) Level II, and *International Classification of Diseases, 10th Revision, Clinical Modification (ICD-10-CM)* codes most commonly reported by pediatricians in providing preventive care services. The pediatrician, not the staff, is ultimately responsible for the appropriate codes to report.

SYMBOL DESCRIPTION

- A bullet at the beginning of a code means it is a new code for the current year.
- + A plus sign means the code is an add-on code.

The recommendations in this publication do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

© 2021 American Academy of Pediatrics

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means—electronic, mechanical, photocopying, recording, or otherwise—without prior written permission from the publisher.

Printed in the United States of America

Current Procedural Terminology (CPT®) 5-digit codes, nomenclature, and other data are copyright 2020 American Medical Association (AMA). All rights reserved. No fee schedules, basic units, relative values, or related listings are included in *CPT*. The AMA assumes no liability for the data contained herein.

The Bright Futures/American Academy of Pediatrics “Recommendations for Preventive Pediatric Health Care,” also known as the “periodicity schedule,” is a schedule of screenings and assessments recommended at each well-child visit from infancy through adolescence. The following services and codes coincide with this schedule. For more details on the periodicity schedule, see www.aap.org/periodicityschedule.

PREVENTIVE MEDICINE SERVICE CODES

Services included under these codes include measurements (eg, length/height, head circumference, weight, body mass index, blood pressure) and age- and gender-appropriate examination and history (initial or interval).

- ▶ Preventive medicine service codes are not time-based; therefore, time spent during the visit is not relevant in selecting the appropriate preventive medicine service code.
- ▶ If an illness or abnormality is discovered, or a preexisting problem is addressed, in the process of performing the preventive medicine service, and if the illness, abnormality, or problem *is significant enough to require additional work* to perform the key components of a problem-oriented evaluation and management (E/M) service (ie, history, physical examination, medical decision-making, counseling/care coordination, or a combination of those), the appropriate office or other outpatient service code (**99202–99215**) should be reported in addition to the preventive medicine service code. Append modifier **25** to the office or other outpatient service code (eg, **99392** and **99213 25**).
- ▶ An *insignificant or trivial illness*, abnormality, or problem encountered in the process of performing the preventive medicine service should not be separately reported.
- ▶ The comprehensive nature of the preventive medicine service codes reflects an age- and gender-appropriate history and physical examination and is not synonymous with the comprehensive examination required for some other E/M codes (eg, **99204**, **99205**, **99215**).
- ▶ Immunizations and ancillary studies involving laboratory, radiology, or other procedures, or screening tests (eg, vision, developmental, hearing) identified with a specific *CPT* code, are reported and paid for separately from the preventive medicine service code.
- ▶ For more information on coding during the COVID-19 pandemic (including telemedicine and telehealth), refer to the Coding at the AAP website (www.aap.org/coding) and its page dedicated to this coding.

Preventive Medicine Services: New Patients

- ▶ Initial comprehensive preventive medicine E/M of an individual includes an age- and gender-appropriate history; physical examination; counseling, anticipatory guidance, or risk factor reduction interventions; and the ordering of laboratory or diagnostic procedures.
- ▶ A *new patient* is defined as one who has not received any professional face-to-face services rendered by physicians and other qualified health care professionals (QHPs) who may report E/M services and reported by a specific *CPT* code(s) from a physician/other QHP, or another physician/other QHP of the exact same specialty and subspecialty who belongs to the same group practice, within the past 3 years.

| <i>CPT</i> Codes | <i>ICD-10-CM</i> Codes |
|--|---|
| 99381 Infant (younger than 1 year) | Z00.110 Health supervision for newborn under 8 days old or Z00.111 Health supervision for newborn 8 to 28 days old or Z00.121 Routine child health exam <i>with abnormal findings</i> or Z00.129 Routine child health exam <i>without abnormal findings</i> |
| 99382 Early childhood (age 1–4 years) | Z00.121 |
| 99383 Late childhood (age 5–11 years) | Z00.129 |
| 99384 Adolescent (age 12–17 years) | |
| 99385 18 years or older | Z00.00 General adult medical exam <i>without abnormal findings</i> Z00.01 General adult medical exam <i>with abnormal findings</i> |

Preventive Medicine Services: Established Patients

Periodic comprehensive preventive medicine reevaluation and management of an individual includes an age- and gender-appropriate history; physical examination; counseling, anticipatory guidance, or risk factor reduction interventions; and the ordering of laboratory or diagnostic procedures.

| CPT Codes | ICD-10-CM Codes |
|--|--|
| 99391 Infant (younger than 1 year) | Z00.110 Health supervision for newborn under 8 days old or |
| | Z00.111 Health supervision for newborn 8 to 28 days old or |
| | Z00.121 Routine child health exam <i>with abnormal findings</i> or |
| | Z00.129 Routine child health exam <i>without abnormal findings</i> |
| 99392 Early childhood (age 1–4 years) | Z00.121 |
| 99393 Late childhood (age 5–11 years) | Z00.129 |
| 99394 Adolescent (age 12–17 years) | |
| 99395 18 years or older | Z00.00 General adult medical exam <i>without abnormal findings</i> |
| | Z00.01 General adult medical exam <i>with abnormal findings</i> |

Preventive Medicine Services: With And Without Abnormal Findings

The use of an *ICD-10-CM* code for *with abnormal findings* (eg, **Z00.121**) does not mean that an additional E/M service must or can be used.

Abnormal findings can be trivial or incidental issues that do not require additional work, but the condition is still documented or listed as contributory. Examples of abnormal findings include abnormal screening results, new acute problem, or unstable or worsening chronic condition.

A stable chronic condition (whether addressed or not) would *not* warrant the use of an abnormal findings code. You can link an abnormal findings *ICD-10-CM* code to a screening if the screen is normal; the abnormality will be identified with the appropriate *ICD-10-CM* code so the payer will be aware.

COUNSELING, RISK FACTOR REDUCTION, AND BEHAVIOR CHANGE INTERVENTION CODES

- ▶ Used to report services provided for the purpose of promoting health and preventing illness or injury.
- ▶ They are distinct from other E/M services that may be reported separately when performed. However, one exception is you cannot report counseling codes (**99401–99404**) in addition to preventive medicine service codes (**99381–99385** and **99391–99395**).
- ▶ Counseling will vary with age and address such issues as family dynamics, diet and exercise, sexual practices, injury prevention, dental health, and diagnostic or laboratory test results available at the time of the encounter.
- ▶ Codes are time-based, where the appropriate code is selected according to the approximate time spent providing the service. Codes may be reported when the midpoint for that time has passed. For example, once 8 minutes are documented, one may report **99401**.
- ▶ Extent of counseling or risk factor reduction intervention must be documented in the patient chart to qualify the service based on time.
- ▶ Counseling or interventions are used for persons without a specific illness for which the counseling might otherwise be used as part of treatment.
- ▶ Cannot be reported with patients who have symptoms or established illness.
- ▶ For counseling individual patients with symptoms or established illness, report an office or other outpatient service code (**99202–99215**) instead.
- ▶ For counseling groups of patients with symptoms or established illness, report **99078** (physician educational services rendered to patients in a group setting) instead.

Preventive Medicine, Counseling

CPT Codes

- 99401** Preventive medicine counseling or risk factor reduction intervention(s) provided to an individual; approximately 15 minutes
- 99402** approximately 30 minutes
- 99403** approximately 45 minutes
- 99404** approximately 60 minutes
- 99411** Preventive medicine counseling or risk factor reduction intervention(s) provided to individuals in a group setting; approximately 30 minutes
- 99412** approximately 60 minutes

ICD-10-CM Codes for Preventive Counseling

- ▶ The diagnosis codes reported for preventive counseling will vary depending on the reason for the encounter.
- ▶ Remember that the patient cannot have symptoms or established illness; therefore, the diagnosis codes reported cannot reflect symptoms or illnesses.
- ▶ Examples of some possible diagnosis codes include

- Z28.3** Underimmunized status (Also include an additional code, eg, **Z28.82** [caregiver refusal].)
- Z71.3** Dietary surveillance and counseling
- Z71.82** Exercise counseling
- Z71.84** Encounter for health counseling related to travel
- Z71.89** Other specified counseling
- Z71.9** Counseling, unspecified

Behavior Change Interventions, Individual

- ▶ Used only when counseling a patient (not parent) on smoking cessation (**99406, 99407**).
- ▶ If counseling a patient's parent or guardian on smoking cessation, do not report these codes (**99406, 99407**) under the patient; instead, refer to preventive medicine counseling codes (**99401–99404**) if the patient

is not currently experiencing adverse effects (eg, illness), or include under the problem-related E/M service if patient is present for a sick visit (**99202–99215**).

- ▶ Codes **99406–99409** may be reported in addition to the preventive medicine service codes.

CPT Codes

- 99406** Smoking and tobacco use cessation counseling visit; intermediate, greater than 3 minutes up to 10 minutes
- 99407** intensive, greater than 10 minutes
- 99408** Alcohol or substance (other than tobacco) abuse structured screening (eg, Alcohol Use Disorder Identification Test [AUDIT], Drug Abuse Screening Test [DAST]) and brief intervention (SBI) services; 15 to 30 minutes
- 99409** greater than 30 minutes

ICD-10-CM Codes for Risk Factor Reduction and Behavior Change Interventions

- F10.10** Alcohol abuse, uncomplicated
- F11.10** Opioid abuse, uncomplicated
- F12.10** Cannabis abuse, uncomplicated
- F13.10** Sedative, hypnotic or anxiolytic abuse, uncomplicated
- F13.90** Sedative, hypnotic, or anxiolytic use, unspecified, uncomplicated
- F15.90** Other stimulant use, unspecified, uncomplicated
- F16.90** Hallucinogen use, unspecified, uncomplicated
- F17.290** Nicotine dependence, other tobacco products (*Includes* Electronic nicotine delivery systems [ENDS]/vaping products)
- Z71.41** Alcohol abuse counseling and surveillance of alcoholic
- Z71.51** Drug abuse counseling and surveillance of drug abuser
- Z71.6** Tobacco abuse counseling
- Z87.891** Personal history of nicotine dependence
- Z91.89** Other specified personal risk factors, presenting as hazards to health not elsewhere classified

OTHER PREVENTIVE MEDICINE SERVICES

Oral Health

CPT Code

99188 Application of topical fluoride varnish by a physician or other qualified health care professional

Refer to page 15 for the definition of QHP.

ICD-10-CM Codes

Z00.121 Routine child health exam *with abnormal findings*

Z00.129 Routine child health exam *without abnormal findings*

Z29.3 Encounter for prophylactic fluoride administration

Z91.841 Risk for dental caries, low

Z91.842 Risk for dental caries, moderate

Z91.843 Risk for dental caries, high

Z91.849 Unspecified risk for dental caries

Pelvic Examination

- ▶ Preventive medicine service codes (**99381–99385** and **99391–99395**) include a pelvic examination as part of the age- and gender-appropriate examination.
- ▶ If the patient is having a problem, the physician can report an office or other outpatient E/M service code (**99212–99215**) for the visit and attach modifier **25**, which identifies that the problem-oriented pelvic visit is a separately identifiable E/M service by the same physician on the same date of service.
- ▶ Link the appropriate *ICD-10-CM* code for the well-child or well-adult examination with abnormal findings (**Z00.121** or **Z00.01**) to the preventive medicine service code, but link a different diagnosis code (eg, **N89.8** [vaginal discharge], **N94.4** [primary dysmenorrhea]) to the office or other outpatient E/M service code (eg, **99212**).
- ▶ Anticipatory or periodic contraceptive management is not a “problem” and is therefore included in the preventive medicine service code; however, if contraception creates a problem (eg, breakthrough bleeding, vomiting), the service can be reported separately with an office or other outpatient service code.

ICD-10-CM Codes

- Z01.411** Gynecological exam *with abnormal findings*
- Z01.419** Gynecological exam *without abnormal findings*
- Z11.51** Screening for human papillomavirus (HPV)
- Z12.72** Screening for malignant neoplasm of vagina
- Z30.011** Initial prescription of contraceptive pills
- Z30.012** Prescription of emergency contraception
- Z30.013** Initial prescription of injectable contraceptive
- Z30.014** Initial prescription of intrauterine contraceptive device (IUD)
- Z30.015** Encounter for initial prescription of vaginal ring hormonal contraceptive
- Z30.016** Encounter for initial prescription of transdermal patch hormonal contraceptive device
- Z30.017** Encounter for initial prescription of implantable subdermal contraceptive
- Z30.018** Encounter for initial prescription of other contraceptives
- Z30.02** Counseling and instruction in natural family planning to avoid pregnancy
- Z30.09** General counseling and advice on contraception
- Z30.40** Surveillance of contraceptives, unspecified
- Z30.41** Surveillance of contraceptive pills
- Z30.42** Surveillance of injectable contraceptive
- Z30.430** Insertion of IUD
- Z30.431** Routine checking of IUD
- Z30.432** Removal of IUD
- Z30.433** Removal and reinsertion of IUD
- Z30.44** Encounter for surveillance of vaginal ring hormonal contraceptive device
- Z30.45** Encounter for surveillance of transdermal patch hormonal contraceptive device
- Z30.46** Encounter for surveillance of implantable subdermal contraceptive
- Z30.49** Surveillance of other contraceptives

Health Risk Assessments

CPT Codes

- 96160** Administration of patient-focused health risk assessment instrument (eg, health hazard appraisal) with scoring and documentation, per standardized instrument (eg, CRAFFT)
- 96161** Administration of caregiver-focused health risk assessment instrument (eg, depression inventory) for the benefit of the patient, with scoring and documentation, per standardized instrument

NOTE: Code **96161** can be reported for a postpartum screening administered to a mother as part of a routine newborn check but billed under the baby's name. Link to *ICD-10-CM* code **Z00.121** or **Z00.129** for normal screening results during a routine well-baby examination. Do *not* report *ICD-10-CM* code **Z13.31** or **Z13.32** under the baby, as those are *only for the maternal record*.

- ▶ Used to report administration of *standardized* health risk assessment instruments on the patient (**96160**) or a primary caregiver (eg, parent) on behalf of the patient (**96161**). Code **96161** requires that the questions and answers relate to the primary caregiver's health and behaviors, not the patient's.
- ▶ *CPT* defines standardization as, "Standardized instruments are validated tests that are administered and scored in a consistent or 'standard' manner consistent with their validation."

Unlisted Preventive Medicine Service

CPT Code

- 99429** Unlisted preventive medicine service

Report code **99429** only when a more specific preventive medicine service code does not exist.

SCREENING CODES

Developmental/Autism Screening and Emotional/Behavioral Assessment

| CPT Codes | ICD-10-CM Codes |
|---|--|
| 96110 Developmental screening, per instrument, scoring and documentation | Z13.41 Encounter for autism screening Z13.42 Encounter for screening for global developmental delays (milestones) |
| 96127 Brief emotional/behavioral assessment (eg, depression inventory) with scoring and documentation, per standardized instrument | Z13.31 Encounter for screening for depression |

- ▶ Used to report administration of standardized developmental/autism screening instruments (**96110**) or behavioral/emotional assessments (**96127**). See page 10 for the definition of *standardized*.
- ▶ Often reported when performed in the context of preventive medicine services but may also be reported when screening or assessment is performed with other E/M services (eg, acute illness or follow-up office visits).
- ▶ Clinical staff (eg, registered nurse) typically administers and scores the completed instrument, while the physician incorporates the interpretation component into the accompanying E/M service.
- ▶ When a standardized screening or assessment is administered along with any E/M service (eg, preventive medicine service), both services should be reported, and modifier **25** (significant, separately identifiable E/M service by the same physician on the same day of the procedure or other service) may need to be appended to the E/M code to show the E/M service was distinct and necessary at the same visit.
- ▶ For more information on reporting **96110** and **96127** instruments, refer to https://www.aap.org/en-us/Documents/coding_factsheet_developmentalscreeningtestingandEmotionalBehavioraassessment.pdf.

Hearing Screening

| CPT Codes | ICD-10-CM Codes |
|---|---|
| 92551 Screening test, pure tone, air only | Z00.121 Routine child health exam <i>with abnormal findings</i> Z00.129 Routine child health exam <i>without abnormal findings</i> |
| 92552 Pure tone audiometry (threshold), air only | |
| 92567 Tympanometry (impedance testing) | |

- ▶ Requires use of calibrated electronic equipment; tests using other methods (eg, whispered voice, tuning fork) are not reported separately.
- ▶ Includes testing of both ears; append modifier **52** when a test is applied to only one ear.
- ▶ For newborn hearing screenings for young patients, including those patients who are nonverbal or have developmental delays, other hearing assessment methods may be more appropriate.

| CPT Codes | ICD-10-CM Codes |
|--|---|
| 92558 Evoked otoacoustic emissions, screening (qualitative measurement of distortion product or transient evoked otoacoustic emissions), automated analysis | Z00.110 Health supervision for newborn under 8 days old or Z00.111 Health supervision for newborn 8 to 28 days old or Z00.121 Routine child health exam <i>with abnormal findings</i> Z00.129 Routine child health exam <i>without abnormal findings</i> P09 Abnormal findings on neonatal screening |
| 92650 Auditory evoked potentials screening of auditory potential with broadband stimuli; automated analysis | |
| 92558 Evoked otoacoustic emissions, screening (qualitative measurement of distortion product or transient evoked otoacoustic emissions), automated analysis | |
| 92587 Distortion product evoked otoacoustic emissions; limited evaluation (to confirm the presence or absence of hearing disorder, 3-6 frequencies) or transient evoked otoacoustic emissions, with interpretation and report | |
| 92588 comprehensive diagnostic evaluation, with interpretation and report | |

- ▶ Codes **Z01.10** (encounter for examination of ears and hearing without abnormal findings) and **Z01.118** (encounter for examination of ears and hearing with other abnormal findings) are reported only when a patient presents for an encounter specific to ears and hearing, not for a routine well-child examination at which a hearing screening is performed.
- ▶ Failed hearing screenings will most likely result in a follow-up office visit (eg, **99212–99215**). Code **Z01.110** (encounter for hearing examination following failed hearing screening) is reported when a specific disorder cannot be identified or when the follow-up hearing screening findings are normal. You can also report **Z01.118** (encounter for examination of ears and hearing with other abnormal findings) and include the code for the abnormal findings (eg, **R94.120** [abnormal auditory function study]).

Vision Screening

| CPT Codes | ICD-10-CM Codes |
|---|---|
| 99173 Screening test of visual acuity quantitative, bilateral | Z00.121 Routine child health exam <i>with abnormal findings</i> |
| 99174 Instrument-based ocular screening (eg, photoscreening, automated-refraction), bilateral, with remote-analysis and report | Z00.129 Routine child health exam <i>without abnormal findings</i> |
| 99177 Instrument-based ocular screening (eg, photoscreening, automated-refraction), bilateral, with on-site analysis | |

Z01.00 and **Z01.01** (examination of eyes and vision with and without abnormal findings) are reported only for routine examination of eyes and vision, not when a vision screening is done during a routine well-child examination.

- ▶ To report code **99173**, you must employ graduate visual acuity stimuli that allow a quantitative estimate of visual acuity (eg, Snellen chart).
- ▶ Codes **99174** and **99177** are reported for instrument-based ocular screening for esotropia, exotropia, anisometropia, cataracts, ptosis, hyperopia, and myopia.

- ▶ Code **99177** is reported in lieu of **99174** when the screening instrument provides you with immediate pass or fail results.
- ▶ When acuity (**99173**) or instrument-based ocular screening (eg, **99174**) is measured as part of a general ophthalmologic service or an E/M service of the eye (eg, for an eye-related problem or symptom), it is considered part of the diagnostic examination of the office or other outpatient service code (**99202–99215**) and is not reported separately.
- ▶ Other identifiable services unrelated to the screening test provided at the same time are reported separately (eg, preventive medicine services).
- ▶ Failed vision screenings will most likely result in a follow-up office visit (eg, **99212–99215**). Report the follow-up screening with **Z01.020** (encounter for examination of eyes and vision following failed vision screening without abnormal findings) if normal results or **Z01.021** (encounter for examination of eyes and vision following failed vision screening with abnormal findings) if abnormal results. If abnormal, link to the diagnosis code for the reason for the failure (eg, **H52.1-** [myopia]); when a specific disorder cannot be identified, report **R94.118** (abnormal results of other function studies of eye).

IMMUNIZATIONS

Immunization Administration (IA)

Pediatric IA Codes

90460 Immunization administration (IA) through 18 years of age via any route of administration, with counseling by physician or other qualified health care professional; first or only component of each vaccine or toxoid administered

+90461 each additional vaccine or toxoid component administered (List separately in addition to code for primary procedure.)

Report **90461** in conjunction with **90460**.

- ▶ *Component* refers to all antigens in a vaccine that prevent diseases caused by 1 organism. Multivalent antigens or multiple serotypes of antigens against a single organism are considered a single component of vaccines. Combination vaccines are vaccines that contain multiple vaccine components. Conjugates or adjuvants contained in vaccines are not considered to be component parts of the vaccine, as defined previously.
- ▶ A QHP is an individual who by education, training, licensure/regulation, facility credentialing (when applicable), and payer policy is able to perform a professional service within his or her scope of practice and to independently report a professional service. These professionals are distinct from *clinical staff*. A *clinical staff member* is a person who works under the supervision of a physician or other QHP and who is allowed by law, regulation, facility, and payer policy to perform or assist in the performance of specified professional services but does not individually report any professional services.
- ▶ Code **90460** is used to report the first or only component in a single vaccine given during an encounter. You can report **90460** more than once during a single office encounter. Code **90461** is considered an add-on code to **90460** (hence the + symbol next to it). This means that the provider will use **90461** in addition to **90460** if more than 1 component is contained within a single vaccine administered. CPT codes **90460** and **90461** are reported regardless of route of administration.

- ▶ Pediatric IA codes (**90460**, **90461**) are reported only when both of the following requirements are met:
 1. The patient must be 18 years or younger.
 2. The physician or other QHP must perform face-to-face vaccine counseling associated with the administration.

NOTE: The clinical staff can do the actual administration of the vaccine.

- ▶ If *both* of these requirements are not met, report a non–age-specific IA code (**90471–90474**) instead.

Non–age-specific IA Codes

- ▶ Report a *CPT* code for both the administration and product and an *ICD-10-CM* code for each vaccine administered during a patient encounter.

90471 IA (includes percutaneous, intradermal, subcutaneous, or intramuscular injections); one vaccine (single or combination vaccine/toxoid)

Do not report **90471** in conjunction with **90473**.

+90472 each additional vaccine (single or combination vaccine/toxoid) (List separately to code for primary procedure.)

Use **90472** in conjunction with **90460**, **90471**, or **90473**.

90473 IA (includes intranasal or oral administration); one vaccine (single or combination vaccine/toxoid)

Do not report **90473** in conjunction with **90471**.

+90474 each additional vaccine (single or combination vaccine/toxoid) (List separately to code for primary procedure.)

Use **90474** in conjunction with **90460**, **90471**, or **90473**.

- ▶ Codes **90471** and **90473** are used to code for the first immunization given during a single office visit. Codes **90472** and **90474** are considered *add-on* codes (hence the **+** symbol next to them) to **90460**, **90471**, and **90473**. This means that the provider will use **90472** or **90474** in addition to **90460**, **90471**, or **90473** if more than 1 vaccine is administered during a visit. There can be only 1 first administration during a given visit. (See vignettes 3, 4, and 5 on pages 21–23.)

- ▶ If during a single encounter for a patient 18 years or younger, a physician or other QHP only counsels on some of the vaccines, report code **90460** (and **90461** when applicable) for those counseled on and defer to codes **90472** or **90474**, as appropriate, for those that are not counseled on.
- ▶ The following vignettes may help illustrate the correct use of the administration codes (see pages 25–28 for a full list of vaccine product codes):

NOTE: The coding vignettes are for teaching purposes only and do not necessarily follow every payer’s reporting requirements.

Vignette 1

A 2-month-old established patient presents for her checkup. The following vaccines are ordered: Pentacel (diphtheria-tetanus-acellular pertussis [DTaP], *Haemophilus influenzae* type b [Hib], inactivated poliovirus [IPV]), pneumococcal, and rotavirus. The physician counsels the parents on all of them, consent is obtained and the nurse administers them all.

How are the appropriate codes for this service selected?

Step 1: Select appropriate E/M code.

99391 Preventive medicine service, established patient, infant (age younger than 1 year)

Step 2: Select appropriate vaccine product codes.

90698 DTaP-Hib-IPV (Pentacel) product

90670 Pneumococcal product

90680 Rotavirus vaccine, oral use

Step 3: Select appropriate IA codes by considering the following questions:

- ▶ Is the patient 18 years or younger?
- ▶ If the patient is younger than 18 years, did the physician or other QHP perform the face-to-face vaccine counseling, discussing the specific risks and benefits of the vaccines?

If the answer to both questions is yes, select a code from the pediatric IA code family (**90460**, **90461**). If the answer to one of the questions is no, select a code from the non–age-specific IA code family (**90471–90474**).

In this vignette, the answer to both questions is yes. Therefore, IA codes **90460** and **90461** will be reported.

Step 4: Select the appropriate *ICD-10-CM* diagnosis codes.

Diagnosis codes are used along with *CPT* codes to reflect the outcome of a visit. The *CPT* codes tell a carrier what was done, and *ICD-10-CM* codes tell a carrier why it was done.

The vaccine product *CPT* code and its corresponding IA *CPT* code are always linked to the same *ICD-10-CM* code. This is because the vaccine product and work that goes into administering that product are intended to provide prophylactic vaccination against a certain type of disease.

ICD-10-CM lists only a single code to describe an encounter in which a patient receives a vaccine. The code is **Z23**, and it is reported at any encounter when a vaccine is given, including routine well-child or adult examinations.

The diagnosis codes for the 3 vaccines and 3 IA codes used in this vignette are as follows:

| <i>CPT</i> Codes | | <i>ICD-10-CM</i> Codes |
|-------------------|---|------------------------|
| 99391 25 | Preventive medicine service, established patient, <1 year | Z00.129 |
| 90698 | DTaP-Hib-IPV (Pentacel) product | Z23 |
| 90670 | Pneumococcal product | Z23 |
| 90680 | Rotavirus vaccine, oral use | Z23 |
| 90460 (×3) | Pediatric IA (Pentacel, pneumococcal, rotavirus), first component | Z23 |
| 90461 (×4) | Pediatric IA (Pentacel), each additional component | Z23 |

Vignette 2

A 5-year-old established patient is at a physician's office for her annual well-child examination. The patient is scheduled to receive her first hepatitis A vaccine; her fifth DTaP vaccine; and the influenza vaccine. After distributing the Vaccine Information Statements and discussing the risks and benefits of immunizations with her parents, the physician administers the vaccines.

How are the appropriate codes for this service selected?

Step 1: Select appropriate E/M code.

99393 Preventive medicine service, established patient, age 5 to 11 years

Step 2: Select appropriate vaccine product codes.

90633 Hepatitis A vaccine, pediatric/adolescent dosage (2-dose schedule), for intramuscular use

90700 DTaP, for use in individuals younger than 7 years, for intramuscular use

90686 Influenza virus vaccine, quadrivalent (IIV4), split virus, preservative free, 0.5 mL dosage, for IM use

Step 3: Select appropriate IA codes by considering the following questions:

- ▶ Is the patient 18 years or younger?
- ▶ If the patient is younger than 18 years, did the physician or other qualified health care professional perform the face-to-face vaccine counseling, discussing the specific risks and benefits of the vaccines?

If the answer to both questions is yes, select a code from the pediatric IA code family (**90460**, **90461**). If the answer to one of the questions is no, select a code from the non-age-specific IA code family (**90471–90474**).

In this vignette, the answer to both questions is yes. Therefore, IA codes **90460** and **90461** will be reported.

Step 4: Select the appropriate *ICD-10-CM* diagnosis codes.

Diagnosis codes are used along with *CPT* codes to reflect the outcome of a visit. The *CPT* codes tell a carrier what was done, and *ICD-10-CM* codes tell a carrier why it was done.

The vaccine product *CPT* code and its corresponding IA *CPT* code are always linked to the same *ICD-10-CM* code. This is because the vaccine product and work that goes into administering that product are intended to provide prophylactic vaccination against a certain type of disease.

ICD-10-CM lists only a single code to describe an encounter in which a patient receives a vaccine. The code is **Z23**, and it is reported at any encounter when a vaccine is given, including routine well-child or adult examinations.

The diagnosis codes for the 3 vaccines and 3 IA codes used in this vignette are as follows:

| <i>CPT Codes</i> | | <i>ICD-10-CM Codes</i> |
|-------------------|---|------------------------|
| 99393 25 | Preventive medicine service, established patient, 5–11 years | Z00.129 |
| 90633 | Hepatitis A vaccine product | Z23 |
| 90460 | Pediatric IA (hepatitis A vaccine), first component | Z23 |
| 90700 | DTaP vaccine product | Z23 |
| 90460 | Pediatric IA (DTaP vaccine), first component | Z23 |
| 90461 (×2) | Pediatric IA (DTaP vaccine), each additional component | Z23 |
| 90686 | Influenza virus vaccine, quadrivalent, preservative free, 0.5 mL dosage | Z23 |
| 90460 | Pediatric IA (influenza vaccine), first component | Z23 |

Alternative Coding

| <i>CPT Codes</i> | | <i>ICD-10-CM Codes</i> |
|-------------------|---|------------------------|
| 99393 25 | Preventive medicine service, established patient, 5–11 years | Z00.129 |
| 90633 | Hepatitis A vaccine product | Z23 |
| 90700 | DTaP vaccine product | Z23 |
| 90686 | Influenza virus vaccine, quadrivalent, preservative free, 0.5 mL dosage | Z23 |
| 90460 (×3) | Pediatric IA (hepatitis A, DTaP, influenza vaccines), first component | Z23 |
| 90461 (×2) | Pediatric IA (DTaP vaccine), second and third components | Z23 |

NOTE: Most payers do not want multiple line items of codes **90460** or **90461**; therefore, follow the alternative coding.

Rationale

Because the patient is younger than 18 years and there is physician counseling, pediatric IA codes are reported (**90460** and **90461**). Each vaccine administered will be reported with its own **90460** (hepatitis A, DTaP, and influenza). The only vaccine with multiple components is DTaP. Because the first component (ie, diphtheria) was counted in **90460**, only the second and third components (ie, tetanus and acellular pertussis) are reported with **90461** with 2 units.

Vignette 3

A 19-year-old patient presents to the office to complete a college physical examination (in college the patient will be living in a dormitory). He is due for a tetanus-diphtheria-acellular pertussis (Tdap) booster, meningococcal vaccine, and intranasal influenza vaccine. The physician counsels the patient on each, and the nurse administers each.

| <i>CPT Codes</i> | | <i>ICD-10-CM Codes</i> |
|-------------------------|---|-------------------------------|
| 99395 25 | Preventive medicine service, established patient, 18–39 years | Z02.0 |
| 90715 | Tdap product | Z23 |
| 90471 | IA, first injection | Z23 |
| 90734 | Meningococcal conjugate vaccine (MenACWY-D or MenACWY-CRM) | Z23 |
| 90472 | IA, each additional injection | Z23 |
| 90672 | Influenza virus vaccine, quadrivalent, live (LAIV4), for intranasal use | Z23 |
| 90474 | IA, each additional oral or intranasal | Z23 |

Rationale

The patient is older than 18 years; therefore, despite physician counseling, pediatric IA codes cannot be reported. Instead, codes **90471** and **90474** must be used.

Vignette 4

A 17-year-old patient presents to the office for her annual checkup and to complete a college physical examination (in college the patient will be living in a dormitory). The patient is healthy and due for a Tdap booster, meningococcal vaccine, first HPV (9-valent) vaccine, and influenza vaccine. The physician counsels the patient only on the meningococcal and HPV vaccines, and the nurse administers each. The patient is asked to return in 4 to 6 weeks for her second HPV vaccine.

| <i>CPT Codes (First Visit Only)</i> | | <i>ICD-10-CM Codes (First Visit Only)</i> |
|---|---|---|
| 99395 25 | Preventive medicine service, established patient, 12–17 years | Z00.0 and Z02.0 |
| 90734 | Meningococcal (MCV4) product | Z23 |
| 90651 | HPV (9-valent) product | Z23 |
| 90460 (×2) | Pediatric IA (meningococcal and HPV), first component | Z23 |
| 90715 | Tdap product | Z23 |
| 90472 (×2) | IA, each additional injection (Tdap) | Z23 |
| 90686 | Influenza virus vaccine, quadrivalent, preservative free, 0.5 mL dosage | Z23 |

Rationale

Because the physician documents counseling only for the meningococcal and HPV vaccines, code **90460** can be reported only for those vaccines because the patient meets the age criteria. For the Tdap and influenza vaccines, defer to non-pediatric IA codes (**90471**, **90472**). In this case, however, a first vaccine code is already reported with code **90460**, so the additional IA code **90472** has to be reported. While *ICD-10-CM* does not provide official ages for the “adult” *ICD-10-CM* codes (**Z00.00** and **Z00.01**) in lieu of the well-child examination codes, many payers use age 17 years as the cutoff. Refer to specific payer policy for details.

Vignette 5

A 6-month-old patient presents to the office for her routine checkup and to receive vaccines. The patient is due for DTaP, pneumococcal, and hepatitis B vaccines. During the examination, the physician finds an upper respiratory infection and fever. The physician counsels the parent on the vaccines but decides to defer for 2 weeks. The physician completes the well-baby checkup on that day.

Two weeks later, the patient returns. The patient is afebrile and asymptomatic and is seen only by the nurse. The DTaP, pneumococcal, and hepatitis B vaccines are administered.

| <i>CPT Code (First Visit)</i> | | <i>ICD-10-CM Code (First Visit)</i> |
|---|---|--|
| 99391 | Preventive medicine service, established patient, <1 year | Z00.121 |
| An appropriate acute sick visit (eg, 99213) may be reported in addition with modifier 25 and linked to an appropriate <i>ICD-10-CM</i> code. | | |
| <i>CPT Codes (2 Weeks Later)</i> | | <i>ICD-10-CM Codes (2 Weeks Later)</i> |
| 90700 | DTaP product | Z23 |
| 90670 | Pneumococcal product | Z23 |
| 90744 | Hepatitis B vaccine product | Z23 |
| 90471 | IA (DTaP), first vaccine | Z23 |
| 90472 (×2) | IA (pneumococcal, hepatitis B), each additional vaccine | Z23 |

Rationale

If counseling occurs outside the IA service, there is no way to report it separately. Therefore, in this vignette, there is nothing separate to report during the well-baby visit, and when the patient returns and sees only the nurse, pediatric IA codes cannot be reported; defer to codes **90471–90474**. During the preventive medicine service, when an acute illness is detected, a code from **99212–99215** can be reported if the service is significant and separately identifiable. Code **9921x** is reported with modifier **25**. When the patient returns *only for vaccines*, an E/M service is not reported. The *ICD-10-CM* code will be reported for *with abnormal findings* (**Z00.121**) because an abnormality was identified during the encounter.

For more information on IA codes, refer to the Coding at the AAP website (www.aap.org/coding) and its page dedicated to vaccine coding.

How to Code When Immunizations Are Not Administered

ICD-CM-10 Codes

- ▶ For many reasons, immunizations are not given during routine preventive medicine services. Parents may refuse vaccines or defer them, a patient may be ill at the time and it is counteractive to administer, or the patient may already have had the disease or be immune.
- ▶ Because of tracking purposes and quality measures, it is important to report non-administration as part of the *ICD-10-CM* codes. The following *ICD-10-CM* codes were created to report why a vaccine is not given:

Vaccination not carried out due to

- Z28.01** Acute illness
- Z28.02** Chronic illness or condition
- Z28.03** Immunocompromised state
- Z28.04** Allergy to vaccine or component
- Z28.1** Religious reasons
- Z28.20** Unspecified reason
- Z28.21** Patient refusal
- Z28.81** Patient had disease being vaccinated against
- Z28.82** Caregiver refusal
- Z28.83** Vaccine was unavailable (eg, manufacturer delay)
- Z28.89** Other reason

Vignette

A 1-year-old presents for his routine well-child examination. He is scheduled to receive his first measles, mumps, rubella; hepatitis A; and varicella vaccines. Because he had a documented case of varicella when he was 9 months of age, the varicella vaccine is not given.

Report the following *ICD-10-CM* codes linked to the E/M service:

- Z23** Encounter for immunization
- Z28.81** Vaccination not carried out due to patient having had the disease being vaccinated against

VACCINES FOR CHILDREN PROGRAM

The rules for reporting vaccines for patients who qualify for the Vaccines for Children (VFC) program vary greatly. Some states require that the product code be submitted, while others require the IA codes. Some require the use of modifiers, while others do not. Currently, the VFC program does not recognize component-based vaccine counseling; therefore, you will not be paid for CPT code **90461**. The American Academy of Pediatrics continues to work on changing this so pediatric providers can be properly compensated for giving multiple-component vaccines. Also be sure to check with your individual state Medicaid plan for varying rules, including, but not limited to, being able to report code **99211** in addition to IA codes for vaccine-only encounters. Be sure to get these rules in writing.

Commonly Administered Pediatric Vaccines

| Product Code | Separately report the administration with codes 90460–90461 or 90471–90474. | Manufacturer | Brand | No. of Vaccine Components |
|--------------|--|--------------|---|---------------------------|
| 90702 | Diphtheria and tetanus toxoids (DT), adsorbed when administered to younger than seven years, for IM use | SP | Diphtheria and Tetanus Toxoids Adsorbed | 2 |
| 90700 | Diphtheria, tetanus toxoids, and acellular pertussis vaccine (DTaP), when administered to <7 years, for IM use | SP GSK | DAPTACEL INFANRIX | 3 |
| 90696 | Diphtheria, tetanus toxoids, and acellular pertussis vaccine and inactivated poliovirus vaccine (DTaP-IPV), when administered to children 4-6 years of age, for IM use | GSK SP | KINRIX Quadracel | 4 |
| 90697 | Diphtheria, tetanus toxoids, acellular pertussis vaccine, inactivated poliovirus vaccine, w PRP-OMP conjugate vaccine, and hepatitis B vaccine (DTaP-IPV-Hib-HepB), for IM use | Merck SP | VAXELIS | 6 |

Commonly Administered Pediatric Vaccines (continued)

| Product Code | Separately report the administration with codes 90460–90461 or 90471–90474. | Manufacturer | Brand | No. of Vaccine Components |
|--------------|---|--------------|-------------------------------|---------------------------|
| 90698 | Diphtheria, tetanus toxoids, acellular pertussis vaccine, <i>Haemophilus influenzae</i> type b, and inactivated poliovirus vaccine (DTaP-IPV/Hib), for IM use | SP | Pentacel | 5 |
| 90723 | Diphtheria, tetanus toxoids, acellular pertussis vaccine, Hepatitis B, and inactivated poliovirus vaccine (DTaP-Hep B-IPV), for IM use | GSK | PEDIARIX | 5 |
| 90633 | Hepatitis A vaccine (Hep A), pediatric/adolescent dosage, 2 dose, for IM use | GSK Merck | HAVRIX VAQTA | 1 |
| 90740 | Hepatitis B vaccine (Hep B), dialysis or immunosuppressed patient dosage, 3 dose, for IM use | Merck | RECOMBIVAX HB | 1 |
| 90743 | Hepatitis B vaccine (Hep B), adolescent, 2 dose, for IM use | Merck | RECOMBIVAX HB | 1 |
| 90744 | Hepatitis B vaccine (Hep B), pediatric/adolescent dosage, 3 dose, for IM use | Merck GSK | RECOMBIVAX HB ENERGIX-B | 1 |
| 90746 | Hepatitis B vaccine (Hep B), adult dosage, for IM use | Merck GSK | RECOMBIVAX HB ENERGIX-B | 1 |
| 90747 | Hepatitis B vaccine (Hep B), dialysis or immunosuppressed patient dosage, 4 dose, for IM use | GSK | ENERGIX-B | 1 |
| 90647 | <i>Haemophilus influenzae</i> type b vaccine (Hib), PRP-OMP conjugate, 3 dose, for IM use | Merck | PedvaxHIB | 1 |
| 90648 | <i>Haemophilus influenzae</i> type b vaccine (Hib), PRP-T conjugate, 4 dose, for IM use | SP GSK | ActHIB HIBERIX | 1 |
| 90651 | Human Papillomavirus vaccine types 6, 11, 16, 18, 31, 33, 45, 52, 58, nonavalent (HPV), 2 or 3 dose schedule, for IM use | Merck | GARDASIL 9 | 1 |

| Product Code | Separately report the administration with codes 90460–90461 or 90471–90474. | Manufacturer | Brand | No. of Vaccine Components |
|--------------|--|--------------|--------------------|---------------------------|
| 90707 | Measles, mumps, and rubella virus vaccine (MMR), live, for subcutaneous use | Merck | M-M-R II | 3 |
| 90710 | Measles, mumps, rubella, and varicella vaccine (MMRV), live, for subcutaneous use | Merck | ProQuad | 4 |
| 90619 | Meningococcal conjugate vaccine, serogroups A, C, W, Y, quadrivalent, tetanus toxoid carrier (MenACWY-TT), for IM use | SP | MenQuadfi | 1 |
| 90620 | Meningococcal recombinant protein and outer membrane vesicle vaccine, serogroup B (MenB-4C), 2 dose schedule, for IM use | GSK | Bexsero | 1 |
| 90621 | Meningococcal recombinant lipoprotein vaccine, serogroup B, 2 or 3 dose schedule, for IM use | Pfizer | Trumenba | 1 |
| 90734 | Meningococcal conjugate vaccine, serogroups A, C, W, Y, quadrivalent, diphtheria toxoid carrier (MenACWY-D) or CRM197 carrier (MenACWY-CRM), for IM use | SP GSK | Menactra Menveo | 1 |
| 90670 | Pneumococcal conjugate vaccine, 13 valent (PCV13), for IM use | Pfizer | PREVNAR 13 | 1 |
| 90732 | Pneumococcal polysaccharide vaccine, 23-valent (PPSV23), adult or immunosuppressed patient dosage, when administered to 2 years or older, for subcutaneous or IM use | Merck | PNEUMOVAX 23 | 1 |
| 90713 | Poliovirus vaccine (IPV), inactivated, for subcutaneous or IM use | SP | IPOL | 1 |
| 90680 | Rotavirus vaccine, pentavalent (RV5), 3 dose schedule, live, for oral use | Merck | RotaTeq | 1 |
| 90681 | Rotavirus vaccine, human, attenuated (RV1), 2 dose schedule, live, for oral use | GSK | ROTARIX | 1 |

Commonly Administered Pediatric Vaccines (continued)

| Product Code | Separately report the administration with codes 90460–90461 or 90471–90474. | Manufacturer | Brand | No. of Vaccine Components |
|--------------|---|-------------------------------|---|---------------------------|
| 90714 | Tetanus and diphtheria toxoids (Td) adsorbed, preservative free, when administered to seven years or older, for IM use | MBL SP | TDVAX TENIVAC | 2 |
| 90715 | Tetanus, diphtheria toxoids and acellular pertussis vaccine (Tdap), when administered to 7 years or older, for IM use | SP GSK | ADACEL BOOSTRIX | 3 |
| 90716 | Varicella virus vaccine (VAR), live, for subcutaneous use | Merck | VARIVAX | 1 |
| 90749 | Unlisted vaccine or toxoid | Please see <i>CPT</i> manual. | | |
| 90672 | Influenza virus vaccine, quad (LAIV), live, intranasal use | AstraZeneca | Flumist Quad | 1 |
| 90674 | Influenza virus vaccine, quad (ccIIIV4), derived from cell cultures, subunit, preservative and antibiotic free, 0.5 mL dosage, IM | Seqirus | Flucelvax | 1 |
| 90682 | Influenza virus vaccine, quad (RIV4), derived from recombinant DNA, HA protein only, preservative and antibiotic free, IM use | Seqirus | Flublok Quad | 1 |
| 90685 | Influenza virus vaccine, quad (IIIV4), split virus, preservative free, 0.25ml dose, for IM use | Seqirus GSK SP | Afluria Fluarix Fluzone Quad | 1 |
| 90686 | Influenza virus vaccine, quad (IIIV4), split virus, preservative free, 0.5ml dosage, for IM use | Seqirus GSK GSK SP | Afluria FLUARIX Quad FLULAVAL Fluzone Quad | 1 |
| 90687 | Influenza virus vaccine, quad (IIIV4), split virus, 0.25ml dosage, for IM use | Seqirus SP | Afluria Quad Fluzone Quad | 1 |
| 90688 | Influenza virus vaccine, quad (IIIV4), split virus, 0.5ml dosage, for IM use | Seqirus SP | Afluria Fluzone Quad | 1 |
| 90756 | Influenza virus vaccine, quad (ccIIIV4), derived from cell cultures, subunit, antibiotic free, 0.5mL dosage, for IM use | Seqirus | Flucelvax Quad | 1 |

Current at time of publication. Developed and maintained by the American Academy of Pediatrics. Updated periodically at https://www.aap.org/en-us/Documents/coding_vaccine_coding_table.pdf. For reporting purposes only. Any vaccine products still US Food and Drug Administration pending are not listed in this resource.

LABORATORY

Two different practice models surround the conducting of laboratory tests: blood is drawn in office and specimen is sent to an outside laboratory for analysis, or blood is drawn and laboratory tests are performed in the physician's practice. Never report the laboratory code for a laboratory test that the practice does not run in-house or is not financially responsible for and billed by the outside laboratory. In those cases, report only the blood draw and specimen handling, as appropriate.

Model 1: Blood is drawn in office and specimen is sent to an outside laboratory for analysis.

CPT Code

99000 Handling and/or conveyance of specimen for transfer from the physician's office to a laboratory

Venipuncture CPT Codes

36406 Venipuncture, younger than 3 years, necessitating physician's skill, not to be used for routine venipuncture

36410 Venipuncture, 3 years or older, necessitating physician's skill, for diagnostic or therapeutic purposes (not to be used for routine venipuncture)

36415 Collection of venous blood by venipuncture

36416 Collection of capillary blood specimen (eg, finger, heel, ear stick)

Venipuncture ICD-10-CM Codes

Link to *ICD-10-CM* codes for the well-child examination or for specific screening tests.

Model 2: Blood is drawn and laboratory tests are performed in the physician's practice.

Venipuncture CPT Codes

- 36406** Venipuncture, younger than 3 years, necessitating physician's skill, not to be used for routine venipuncture
- 36410** Venipuncture, 3 years or older, necessitating physician's skill, for diagnostic or therapeutic purposes (not to be used for routine venipuncture)
- 36415** Collection of venous blood by venipuncture
- 36416** Collection of capillary blood specimen (eg, finger, heel, ear stick)

Venipuncture ICD-10-CM Codes

Link to *ICD-10-CM* codes for the well-child examination or for specific screening tests.

Bilirubin CPT Codes

- 82247** Bilirubin, total
- 88720** Bilirubin, total, transcutaneous

Bilirubin ICD-10-CM Code

- Z13.228** Encounter for screening for other metabolic disorder

Dyslipidemia Screening CPT Codes

- 80061** Lipid panel (includes total cholesterol, high-density lipoprotein [HDL] cholesterol, and triglycerides)
- 82465** Cholesterol, serum, total
- 83718** Lipoprotein, direct measurement, high-density cholesterol (HDL cholesterol)
- 84478** Triglycerides

Dyslipidemia Screening ICD-10-CM Code

- Z13.220** Encounter for screening for lipid disorders

Anemia Screening CPT Code

- 85018** Blood count; hemoglobin

Anemia Screening ICD-10-CM Code

Z13.0 Encounter for screening for diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (eg, anemia)

Lead Screening CPT Code

83655 Lead

Lead Screening ICD-10-CM Code

Z13.88 Encounter for screening for disorder due to exposure to contaminants

Newborn Metabolic Screening HCPCS Code

NOTE: See Healthcare Common Procedure Coding System Codes section on page 34 for explanation of HCPCS codes.

S3620 Newborn metabolic screening panel, includes test kit, postage, and the laboratory tests specified by the state for inclusion in this panel (eg, galactose; hemoglobin, electrophoresis; hydroxyprogesterone, 17-D; phenylalanine [phenylketonuria (PKU)]; and thyroxine, total)

NOTE: Only report code **S3620** if you are billing for the actual running of the laboratory test or test kit. Otherwise only report the appropriate blood collection code (eg, **36416**).

Newborn Metabolic Screening ICD-10-CM Codes

Report the diagnosis codes for the state-specific newborn screening tests conducted. Examples include

Z13.0 Encounter for screening for diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (eg, anemia, sickle cell)

Z13.21 Encounter for screening for nutritional disorder

Z13.228 Encounter for screening for other metabolic disorders (eg, PKU, galactosemia)

Z13.29 Encounter for screening for other suspected endocrine disorder (eg, thyroid)

Papanicolaou Smear HCPCS Code

NOTE: See Healthcare Common Procedure Coding System Codes section on page 34 for explanation of HCPCS codes.

Q0091 Screening Papanicolaou smear; obtaining, preparing, and conveyance of cervical or vaginal smear to laboratory

Papanicolaou Smear CPT Code

Collection of a cervical specimen via a pelvic examination is included in the preventive medicine service code (**99381–99385** and **99391–99395**).

Papanicolaou Smear ICD-10-CM Codes

Z12.4 Encounter for screening for malignant neoplasm of cervix (excludes HPV)

Z12.72 Encounter for screening for malignant neoplasm of vagina

Z12.79 Encounter for screening for malignant neoplasm of other genitourinary organs

Z12.89 Encounter for screening for malignant neoplasms of other sites

Tuberculosis Testing (Mantoux/Purified Protein Derivative [PPD])

Administration of PPD Test

| <i>CPT Code</i> | <i>ICD-10-CM Code</i> |
|---|---|
| 86580 Skin test; tuberculosis, intradermal | Z11.1 Encounter for screening for respiratory tuberculosis |

NOTE: There is no separate administration code for the PPD test. Do not report one.

Reading of PPD Test

If patient returns to have a nurse read the test results, report

| <i>CPT Codes</i> | <i>ICD-10-CM Codes</i> |
|--|---|
| 99211 Office or other outpatient services (negative PPD outcome) | Z11.1 Encounter for screening for respiratory tuberculosis (<i>if test is negative</i>) |
| 99212–99215 Office or outpatient services (physician service for <i>positive</i> encounter) | R76.11 Nonspecific reaction to tuberculin skin tuberculosis (<i>if test is positive</i>) |

Sexually Transmitted Infection and HIV Screening CPT Codes

- 86701** Antibody; HIV-1
- 86703** Antibody; HIV-1 and HIV-2; single assay
- 87490** Infectious agent detection by nucleic acid (DNA or RNA);
Chlamydia trachomatis, direct probe technique
- 87491** Infectious agent detection by nucleic acid (DNA or RNA);
C trachomatis, amplified probe technique
- 87590** Infectious agent detection by nucleic acid (DNA or RNA);
Neisseria gonorrhoeae, direct probe technique
- 87591** Infectious agent detection by nucleic acid (DNA or RNA);
N gonorrhoeae, amplified probe technique
- 87810** Infectious agent detection by immunoassay with direct
optical observation; *C trachomatis*
- 87850** Infectious agent detection by immunoassay with direct
optical observation; *N gonorrhoeae*

Sexually Transmitted Infection and HIV Screening ICD-10-CM Codes

- Z11.3** Encounter for screening for infections with a predominantly
sexual mode of transmission (excludes HPV and HIV)
- Z11.8** Encounter for screening for other infectious and parasitic
diseases (eg, chlamydia)

HEALTHCARE COMMON PROCEDURE CODING SYSTEM CODES

- ▶ The HCPCS Level II codes are procedure codes used to report services and supplies not included in the *CPT* nomenclature.
- ▶ Like *CPT* codes, HCPCS Level II codes are part of the standard procedure code set under the Health Insurance Portability and Accountability Act of 1996.
- ▶ Certain payers may require that HCPCS codes be reported in lieu of or as a supplement to *CPT* codes.
- ▶ The HCPCS nomenclature contains many codes for reporting nonphysician provider patient education, which can be an integral service in the provision of pediatric preventive care.
- ▶ Examples of HCPCS Level II codes relevant to pediatric preventive care include

| | |
|--------------|--|
| S0302 | Completed Early and Periodic Screening, Diagnosis, and Treatment service (List in addition to code for appropriate E/M service.) |
| S0610 | Annual gynecologic examination; new patient |
| S0612 | Annual gynecologic examination; established patient |
| S0613 | Annual gynecologic examination, clinical breast examination without pelvic examination |
| S0622 | Routine examination for college, new or established patient (List separately in addition to appropriate E/M code.) |
| S9444 | Parenting classes, nonphysician provider, per session |
| S9445 | Patient education, not otherwise classified, nonphysician provider, individual, per session |
| S9446 | Patient education, not otherwise classified, nonphysician provider, group, per session |
| S9447 | Infant safety (including cardiopulmonary resuscitation) classes, nonphysician provider, per session |
| S9451 | Exercise classes, nonphysician provider, per session |
| S9452 | Nutrition classes, nonphysician provider, per session |
| S9454 | Stress management classes, nonphysician provider, per session |

Commonly Reported *ICD-10-CM* Codes for Pediatric Preventive Services

| <i>ICD-10-CM</i> Code | Descriptor | Special Coding Conventions |
|--|---|--|
| Encounter and Examination Codes | | |
| Z00.110 | Newborn check under 8 days old | Outpatient codes only |
| Z00.111 | Newborn check 8 to 28 days old | Outpatient codes only |
| Z00.121 | Routine child health examination <i>with abnormal findings</i> | First-listed <i>ICD-10-CM</i> code only. |
| Z00.129 | <i>without abnormal findings</i> | |
| Z00.00 | General adult medical examination <i>without abnormal findings</i> | First-listed <i>ICD-10-CM</i> code only. Typically used for patients 18 years and older (payer policy). |
| Z00.01 | <i>with abnormal findings</i> | |
| Z02.0 | Examination for admission to educational institution | Not required in addition to a Z00 code |
| Z02.4 | Examination for driving license | |
| Z02.5 | Examination for participation in sport | |
| Z01.110 | Hearing examination following failed hearing screening | First-listed <i>ICD-10-CM</i> code only. Do not report as a secondary code or in addition to a Z00 code. |
| Z23 | Immunizations | This is the only code in <i>ICD-10-CM</i> for vaccines. Link to both the product and administration <i>CPT</i> codes. |
| Z29.3 | Encounter for prophylactic fluoride administration | |

Commonly Reported ICD-10-CM Codes for Pediatric Preventive Services (continued)

| ICD-10-CM Code | Descriptor | Special Coding Conventions |
|--|---|----------------------------|
| <p>Screening Codes A screening code is not necessary if the screening is inherent to a routine examination, but it can be reported and oftentimes payers require it.</p> | | |
| Z11.1 | Respiratory tuberculosis | |
| Z11.3 | Infections with a predominantly sexual mode of transmission (<i>excludes</i> HPV and HIV) | |
| Z12.4 | Encounter for screening for malignant neoplasm of cervix (<i>excludes</i> HPV) | |
| Z12.79 | Malignant neoplasm of other genitourinary organs | |
| Z12.89 | Malignant neoplasms of other sites | |
| Z13.29 | Other suspected endocrine disorder | |
| Z13.1 | Diabetes mellitus | |
| Z13.228 | Other metabolic disorders (eg, inborn errors of metabolism, galactosemia, PKU) | |
| Z13.220 | Lipid disorders | |
| Z13.21 | Nutritional disorder | |
| Z13.228 | Other metabolic disorder | |
| Z13.29 | Other suspected endocrine disorder | |
| Z13.0 | Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (eg, anemia, sickle cell) | |
| Z13.31 | Encounter for screening for depression | |
| Z13.89 | Other disorders | |
| Z13.41 | Encounter for autism screening | |
| Z13.42 | Encounter for screening for global developmental delays (milestones) | |
| Z13.88 | Disorder due to exposure to contaminants (eg, lead) | |

| <i>ICD-10-CM</i> Code | Descriptor | Special Coding Conventions |
|------------------------------|--|--|
| Preventive Counseling | | |
| Z71.3 | Dietary surveillance and counseling | |
| Z71.82 | Exercise counseling | |
| Z71.84 | Health counseling related to travel | |
| Z71.89 | Other specified counseling | |
| Z71.9 | Counseling, unspecified | |
| Underimmunized Status | | |
| Z28.3 | Underimmunized status | A status code is informative and may affect the course of treatment and its outcome. Report when this is the case. |
| Vaccines Not Given | | |
| Z28.01 | Acute illness | |
| Z28.04 | Allergy to vaccine or components | |
| Z28.82 | Caregiver refusal | |
| Z28.02 | Chronic illness or condition | |
| Z28.03 | Immune compromised state | |
| Z28.21 | Patient refusal | |
| Z28.81 | Patient had disease being vaccinated for | |
| Z28.1 | Religious reasons | |
| Z28.89 | Other reason | |
| Z28.83 | Vaccine was unavailable (eg, manufacturer delay) | |
| Z28.20 | Unspecified reason | |

Healthcare Effectiveness Data and Information Set Measures Related to Pediatric Preventive Care

| Measure Topic | Measure | Coding Options |
|---|--|--|
| Child and Adolescent Well-Care Visits: Well-Child Visits in the First 15 Months of Life (W15) | At least 6 well-child examinations by 15 months of age | <i>ICD-10-CM</i> Z00.110, Z00.111, Z00.121, Z00.129 <i>CPT</i> 99381, 99382, 99391, 99392 |
| Child and Adolescent Well-Care Visits: Well-Child Visits in the Third, Fourth, Fifth, and Sixth Years of Life (W34) | One or more comprehensive well-child visits with a PCP (per year) | <i>ICD-10-CM</i> Z00.121, Z00.129 <i>CPT</i> 99382, 99392 |
| Child and Adolescent Well-Care Visits: Adolescent Well-Care Visits (AWC) | At least one annual comprehensive well-care encounter (per year) for adolescents and young adults aged 12–21 years | <i>ICD-10-CM</i> Z00.00, Z00.01, Z00.121, Z00.129 <i>CPT</i> 99384, 99385, 99394, 99395 |
| Lead Screening in Children (LSC) | By age 2 years, have had one or more capillary or venous lead blood tests for lead poisoning | <i>CPT</i> 83655 |
| Chlamydia Screening in Women (CHL) | Sexually active women aged 16–24 years who received at least one chlamydia test each year | <i>CPT</i> 87110, 87270, 87320, 87490–87492, 87810 |

| Measure Topic | Measure | Coding Options |
|---|---|--|
| Childhood Immunization Status (CIS) and Immunizations for Adolescents (IMA) | <p>By age 2 y, have DTaP (4 doses) IPV (3 doses) MMR (1 dose) Hib (3 doses) Hep B (3 doses) Varicella (1 dose) Pneumococcal (4 doses) Hep A (1 dose) Rotavirus (2–3 doses) Influenza (2 doses)</p> <p>By 13th birthday, have Meningococcal (1 dose) (Ages 11–13 y) Tdap (1 dose) (Ages 10–13 y) HPV (males/females) (2–3 doses) (Ages 9–13 y)</p> | Varies; refer to the Commonly Administered Pediatric Vaccines table on pages 25–28 for specific vaccine codes. |
| Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC) | For those aged 3–17 years who had an outpatient visit with a PCP during the measurement year and had evidence of BMI percentile documentation and counseling for nutrition and/or physical activity | <i>ICD-10-CM</i> Z68.51–Z68.54,^a Z71.3, Z02.5, Z71.82 <i>CPT</i> 3000F^a |

Abbreviations: BMI, body mass index; *CPT*, Current Procedural Terminology; DTaP, diphtheria, tetanus, acellular pertussis; Hep A, hepatitis A; Hep B, hepatitis B; Hib, *Haemophilus influenzae* type b; HPV, human papillomavirus; *ICD-10-CM*, International Classification of Diseases, 10th Revision, Clinical Modification; IPV, inactivated poliovirus; MMR, measles, mumps, rubella; PCP, primary care practitioner; Tdap, tetanus, diphtheria, acellular pertussis.

^a Body mass index codes should only be reported when there is a related condition (eg, obesity). Payers need to accept **3000F** in lieu of BMI *ICD-10-CM* codes for the BMI measure unless the patient has a related condition.



**Bright
Futures**[™]

prevention and health
promotion for infants,
children, adolescents,
and their families[™]

HRSA

Health Resources & Services Administration

This program is supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) as part of an award totaling \$5,000,000 with 10 percent financed with non-governmental sources. The contents are those of the author(s) and do not necessarily represent the official views of, nor an endorsement, by HRSA, HHS, or the U.S. Government. For more information, please visit HRSA.gov.

Also From Bright Futures!

Bright Futures Tool and Resource Kit, 2nd Edition

This online resource provides health care professionals with an organized and integrated compilation of current forms and materials needed to perform a well-child visit on one website. The new online platform allows easy access to the core tools for health supervision visits, including previsit questionnaires, visit documentation forms, parent and patient education handouts, and additional documentation forms and AAP-developed educational resources.

Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents, 4th Edition

This essential resource provides key background information and recommendations for themes critical to healthy child development along with well-child supervision standards for 31 age-based visits—from newborn through 21 years.

Additional Pediatric Coding Resources From the American Academy of Pediatrics

NEW EDITION!

Coding for Pediatrics 2021: A Manual for Pediatric Documentation and Payment, 26th Edition

Code it right the first time—and avoid delays and costly denials—with this widely used pediatric coding resource. This year's updated 26th edition includes all changes in *CPT* and *ICD-10-CM* codes—complete with expert guidance for their application.

NEW EDITION!

Pediatric ICD-10-CM 2021: A Manual for Provider-Based Coding, 6th Edition

This convenient resource condenses the vast *ICD-10-CM* code set into 500+ pages of pediatric-related codes and guidelines.

NEW EDITION!

AAP Pediatric Coding Newsletter™

This print and online monthly publication brings you proven coding strategies and how-to help with your toughest coding and payment challenges!

**To order these and other pediatric resources,
visit shop.aap.org.**

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



BFNC.2021.CPPCFEB
3-360/0221