

GUIDELINES FOR EMERGENCY CARE IN MARYLAND SCHOOLS

2015 EDITION
Second Maryland Edition

LIST OF CONTENTS

**Guidelines
for helping an
ill or injured
student when
the school
nurse is not
available**

- Amputations
- Anaphylaxis/Allergic Reactions (Part 1)
- Anaphylaxis/Allergic Reactions (Part 2)
- Asthma & Difficulty Breathing
- Behavioral Emergencies
- Bites
- Bleeding
- Blisters
- Bruises
- Burns
- Chest Pain
- CPR (Infant, Child, Adult & AEDs)
- Choking
- Child Abuse & Neglect
- Communicable Diseases Resources
- Cuts, Scratches & Scrapes
- Diabetes
- Diarrhea
- Drowning (Near)
- Ear Problems
- Electric Shock
- Eye Problems
- Fainting
- Fever
- Fractures & Sprains
- Frostbite
- Genitourinary Complaints
- Headache
- Head Injuries
- Hyperthermia (Heat) Emergencies
- Hypothermia (Cold) Emergencies
- Menstrual Difficulties
- Mouth & Jaw Injuries
- Neck & Back Pain
- Nose Problems
- Poisoning & Overdose
- Pregnancy
- Puncture Wounds
- Rape/Sexual Assault
- Rashes
- Seizures
- Shock
- Splinters
- Stabbing & Gunshot Injuries
- Stings
- Stomachaches & Pain
- Teeth Problems
- Tetanus Immunization
- Ticks
- Unresponsiveness
- Vomiting

Also Includes:

- Selected Resources for School Safety Planning & Emergency Preparedness



DEPARTMENT OF HEALTH

Robert R. Neall, Secretary

STATE DEPARTMENT OF EDUCATION

Dr. Karen Salmon, State Superintendent of Schools

EMERGENCY CARE GUIDELINES – 2ND EDITION

Update History Page

Please see the list below for updated pages to the 2015 edition of the Guidelines for Emergency Care in Maryland Schools.

Name of Page	Date of update
Bleeding	April 2019
Diabetes	May 2017
Teeth Problems	May 2019

GUIDELINES FOR EMERGENCY CARE IN MARYLAND SCHOOLS 2015 EDITION



Van T. Mitchell
Secretary



Kevin Seaman, MD, FACEP
Executive Director,
MIEMSS



Lillian M. Lowery, Ed.D
State Superintendent

This project was made possible with funding from a Centers for Disease Control Office of Public Health Emergency Preparedness (PHEP) Cooperative Agreement provided through the Maryland Department of Health and Mental Hygiene, Office of Preparedness and Response.

ACKNOWLEDGEMENTS

Special thanks go to the following organizations for the original development of these resources and updated editions:

Ohio Emergency Medical Services for Children Program, Ohio Department of Public Safety, Pennsylvania EMS for Children Program and Pennsylvania Department of Health. Permissions were obtained from the Ohio and Pennsylvania EMS for Children Programs for allowing reproductions of portions of this document to reflect Maryland specific law and regulations.

We would also like to acknowledge the following contributors:

Laurel A. Moody, RN, MS, MSN
Assistant Professor, Nursing Education
Stevenson University
Lead Editor

Cheryl Duncan De Pinto, MD, MPH, FAAP
Medical Director
Office of Population Health Improvement
Department of Health and Mental Hygiene

Cynthia Wright-Johnson, MSN, RN
Program Director
Emergency Medical Services for Children
Program at Maryland Institute for Emergency
Medical Services Systems

Alicia L. Mezu, MSN/Ed, BSN, BS, RN
Health Services Specialist
Maryland State Department of Education
Division of Student, Family and School Support
Student Services and Strategic Planning Branch

Allen R. Walker, MD, MBA, FAAP
Associate State EMS Medical Director for
Pediatrics at Maryland Institute for Emergency
Medical Services Systems

Mary Nasuta RN, MS, NCSN
Nurse Coordinator, Health Services
Harford County Public Schools

Keisha Major RN, BSN, MSN
Health Services Nurse Manager
Prince George's County Public Schools

Patricia Wince, RN, MSN
Supervisor, Health Services
St. Mary's County Public Schools

Special thanks to Stevenson University for supporting the leadership and participation of Laurel A. Moody for the completion of this revised guideline.

Sincere thanks to James W. Brown, Director of Education Services, Aleithea Warmack and Gail Kostas at Maryland Institute for Emergency Medical Services Systems for formatting the revised guidelines.

ABOUT THE GUIDELINES

The *Guidelines for Emergency Care in Maryland Schools* was originally developed in 2005 by the Guidelines Committee of the Maryland State School Health Council. It reflected input from Maryland Department of Health and Mental Hygiene, Maryland State Department of Education, Maryland Emergency Medical Systems for Children, as well as local health department and local school systems' school health services staff.

This updated edition, *Guidelines for Emergency Care in Maryland Schools, Second Maryland Edition* is the product of a careful review of previous content. It reflects changes in Maryland policy and statute and updated best practice recommendations for providing first aid and emergency care to students in Maryland schools, when the school nurse is not available. The *Guidelines* were adapted from similar documents in use in other states.

The *Guidelines* contain **recommended** procedures to serve as “*what to do in an emergency information*” for school staff with minimal training to guide decision making in an actual emergency. The algorithms contained in the *Guidelines* reflect established first aid and emergency response standards. It is not the intent of the *Guidelines* to supersede or make invalid any laws or rules established by a school system, a school board, or the state of Maryland.

Users of these *Guidelines* should review the “How to Use the Guidelines” section and familiarize themselves with the format of the document prior to an emergency.

It is strongly recommended that staff who are in a position to provide first aid to students, complete an approved first aid and CPR course. School staff should consult their school nurse or local school health services coordinator with questions about any of the recommendations. School-specific instructions may be added as needed, and in some cases is explicitly recommended within certain *Guidelines*.

This document can be downloaded and printed from the following websites:


http://pophealth.dhmh.maryland.gov/Documents/Guide_for_Emergency_Care_in_MD_Schools.pdf

http://marylandpublicschools.org/about/Documents/DSFSS/SSSP/SHS/SHSGuidelines/EmergencyCareGuidelines2015_08262015.pdf

www.miemss.org

HOW TO USE THE EMERGENCY GUIDELINES

- In an emergency, refer first to the guideline for treating the most severe symptoms (unresponsiveness, bleeding, etc.)
- Learn when EMS/9-1-1 (Emergency Medical Services) should be contacted. Copy the “When to Call 9-1-1 for EMS” page and post in key locations.
- The last page of the *Guidelines* contains important information about emergency numbers in your area. Please complete this information as soon as you receive the guidelines, as you will need to have this information ready in an emergency situation.
- The *Guidelines* are arranged in alphabetical order for quick access.
- A colored flow chart format is used to guide you easily through all steps and symptoms from beginning to ending. See the **Key to Shapes and Colors**.
- Take some time to familiarize yourself with the **Emergency Procedures for Injury or Illness**. These procedures give a general overview of the recommended steps in an emergency situation and the safeguards that should be taken.
- In addition, information has been provided about **Infection Control, Planning for Students with Special Needs, School Safety Planning and Emergency Preparedness**.



WHEN TO CALL 9-1-1 FOR EMERGENCY MEDICAL SERVICES (EMS)

Call EMS / 9-1-1 if:

- The person is unresponsive, semi-responsive, or unusually confused
- The person's airway is blocked
- The person is not breathing
- The person is having difficulty breathing, has shortness of breath or is choking
- The person has no pulse when checked by a trained person
- The person has bleeding that won't stop
- The person is coughing up or vomiting blood
- The person has been poisoned
- The person has a seizure for the first time or a seizure that lasts more than five minutes
- The person has injuries to the neck or back
- The person has sudden, severe pain anywhere in the body
- The person's condition is life-threatening (for example: amputations) or other injuries that may leave the person permanently disabled unless he/she receives immediate care (for example: severe eye injuries)
- The person's condition could worsen or become life-threatening on the way to the hospital
- Moving the person could cause further injury (for example: neck injury)
- The person needs the skills or equipment of paramedics or emergency medical technicians
- Distance or traffic conditions would cause a delay in getting the person to the hospital



EMERGENCY PROCEDURES FOR INJURY OR ILLNESS

1. Remain calm and assess the situation. Be sure the situation is safe for you to approach. The following dangers will require caution: live electrical wires, gas leaks, building damage, fire or smoke, traffic, or violence.
2. A responsible adult should stay at the scene and give help until the person designated to handle emergencies arrives.
3. Send word to the person designated to handle emergencies. This person will take charge of the emergency and render any further first aid needed.
4. Do **NOT** give medications unless there has been prior approval by the student's parent or legal guardian and health care provider **OR** according to local school system policy or the student's emergency/action plan.
5. Do **NOT** move a severely injured or ill student unless absolutely necessary for immediate safety. If EMS/9-1-1 states moving is necessary, follow guidelines in **NECK AND BACK PAIN** section.
6. The responsible school authority or a designated school staff should notify the parent or legal guardian of the emergency as soon as possible.
7. If the parent or legal guardian cannot be reached, notify an emergency contact or the parent or legal guardian substitute listed on the student's Emergency Contact card / form. Arrange for transportation of the student by EMS/9-1-1, if necessary.
8. A responsible adult should stay with the injured / ill student.
9. Fill out a report for all injuries or illnesses requiring above procedures as required by local school system policy.

POST-CRISIS INTERVENTION FOLLOWING SERIOUS INJURY OR DEATH

- Discuss with counseling staff or critical incident stress management team.
- Determine level of intervention for staff and students.
- Designate private rooms for private counseling/ defusing.
- Escort affected students, siblings, close friends, and other highly stressed individuals to counselors/critical incident stress management team.
- Assess stress level of staff. Recommend counseling to all staff.
- Follow-up with students and staff who receive counseling.
- Designate staff person(s) to attend funeral.
- Allow for changes in normal routines or schedules to address injury or death.

PLANNING FOR STUDENTS WITH SPECIAL NEEDS

Some students in your school may have special emergency care needs due to health conditions, physical abilities, or communication challenges and need to be included in emergency and disaster planning.

HEALTH CONDITIONS:

Some students may have special conditions that put them at risk for life-threatening emergencies, including but not limited to:

- Seizures
- Diabetes
- Asthma or other breathing difficulties
- Life-threatening or severe allergic reactions
- Technology-dependent or medically fragile conditions

Your school nurse or other school health professional, along with the student's parent or legal guardian and health care provider should develop individual emergency/action plans for these students when they are enrolled. These emergency/action plans should be made available to appropriate staff at all times.

In the event of an emergency situation, refer to the student's emergency/action Plan.

The American College of Emergency Physicians and the American Academy of Pediatrics developed an *Emergency Information Form for Children (EIF) with Special Needs* that is included on the next pages. It can also be downloaded from <http://www.aap.org>. This form provides standardized information that can be used to prepare the caregivers and health care system for emergencies in children with special health care needs. The EIF will ensure a child's complicated medical history is concisely summarized and available when needed most - when the child has an emergency health problem when neither parent nor physician is immediately available*.

PHYSICAL ABILITIES:

Other students in your school may have special emergency needs due to their physical abilities. For example, students who are:

- In wheelchairs
- Temporarily on crutches/walking casts
- Unable or have difficulty walking up or down stairs

These students will need special arrangements in the event of a school-wide emergency (e.g., fire, tornado, evacuation, etc.). A plan should be developed and a responsible person should be designated to assist these students to safety. All staff should be aware of this plan.

* The emergency/action plan should also contain provisions to ensure availability of medications during an emergency such as lockdowns and school evacuations.

PLANNING FOR STUDENTS WITH SPECIAL NEEDS (CONTINUED)

COMMUNICATION CHALLENGES:

Other students in your school may have sensory impairments or have difficulty understanding special instructions during an emergency. For example, students who have:

- Vision impairments
- Hearing impairments
- Processing disorders
- Limited English proficiency
- Behavior or developmental disorders
- Emotional or mental health issues

These students may need special communication considerations in the event of a school-wide emergency. All staff should be aware of plans to communicate information to these students.

Emergency Information Form for Children With Special Needs

Last name:



American Academy of Pediatrics



Date form completed
By Whom

Revised
Revised

Initials
Initials

Name:		Birth date:	Nickname:
Home Address:		Home/Work Phone:	
Parent/Guardian:	Emergency Contact Names & Relationship:		
Signature/Consent**:			
Primary Language:	Phone Number(s):		
Physicians:			
Primary care physician:		Emergency Phone:	
		Fax:	
Current Specialty physician: Specialty:		Emergency Phone:	
		Fax:	
Current Specialty physician: Specialty:		Emergency Phone:	
		Fax:	
Anticipated Primary ED:		Pharmacy:	
Anticipated Tertiary Care Center:			

Diagnoses/Past Procedures/Physical Exam:	
1.	Baseline physical findings:
2.	
3.	Baseline vital signs:
4.	
Synopsis:	
	Baseline neurological status:

*Consent for release of this form to health care providers

EMERGENCY
INFORMATION
FORM

Last name:

Diagnoses/Past Procedures/Physical Exam continued:	
Medications:	Significant baseline ancillary findings (lab, x-ray, ECG):
1.	
2.	
3.	
4.	Prostheses/Appliances/Advanced Technology Devices:
5.	
6.	

Management Data:	
Allergies: Medications/Foods to be avoided	and why:
1.	
2.	
3.	
Procedures to be avoided	and why:
1.	
2.	
3.	

Immunizations											
Dates						Dates					
DPT						Hep B					
OPV						Varicella					
MMR						TB status					
HIB						Other					

Antibiotic prophylaxis:

Indication:

Medication and dose:

Common Presenting Problems/Findings With Specific Suggested Managements		
Problem	Suggested Diagnostic Studies	Treatment Considerations

Comments on child, family, or other specific medical issues:	
Physician/Provider Signature:	Print Name:

© American College of Emergency Physicians and American Academy of Pediatrics. Permission to reprint granted with acknowledgement.

INFECTION CONTROL

To reduce the spread of infectious diseases (diseases that can be spread from one person to another), it is important to follow Standard Precautions. **Standard Precautions** is an approach to infection control that combines the major features of Universal Precautions (UP) and Body Substance Isolation (BSI) and are based on the principle that all blood, body fluids, secretions, excretions except sweat, non-intact skin, and mucous membranes may contain transmissible infectious agents. Standard Precautions include a group of infection prevention practices that apply to all patients, regardless of suspected or confirmed infection status, in any setting in which health care is delivered. These include: hand hygiene, use of gloves, gown, mask, eye protection, or face shield, depending on the anticipated exposure; and safe injection practices. The following list describes standard precautions:

- **Wash hands thoroughly** with running water and soap for at least 15 seconds:
 1. Before and after physical contact with any student (*even if gloves have been worn*)
 2. Before and after eating or handling food
 3. After cleaning
 4. After using the restroom
 5. After providing any first aid

Be sure to scrub between fingers, under fingernails and around the tops and palms of hands. If soap and water are not available, an alcohol-based waterless hand sanitizer may be used according to manufacturer's instructions.

- Wear disposable gloves when in contact with blood and other body fluids.
- Wear protective eyewear when body fluids may come in contact with eyes (e.g., squirting blood).
- Wipe up any blood or body fluid spills as soon as possible (*wear disposable gloves*). Double the trash in plastic bags and dispose of immediately. Clean the area with an appropriate cleaning solution.
- Send soiled clothing (i.e., clothing with blood, stool, or vomit) home with the student in a double-bagged plastic bag.
- Do not touch your mouth or eyes while giving any first aid.

Source: Siegel JD, Rhinehart E, Jackson M, Chiarello L, and the Healthcare Infection Control Practices Advisory Committee, 2007 Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings. <http://www.cdc.gov/ncidod/dhqp/pdf/isolation2007.pdf>

AMPUTATIONS

Always use Standard Precautions

Stump – The end of a limb left after amputation

Amputation – The removal of a limb or other appendage

Call EMS/9-1-1.



Control bleeding by applying direct pressure to stump. See “Bleeding.”

Elevate the injured extremity/stump.

Support the affected extremity for comfort. Treat person for shock. See “Shock.”

CARE OF AMPUTATED PART:

- Locate part if possible.
- Do not attempt to clean.
- Wrap in a dry sterile dressing.
- Place in a clean plastic bag.
- Place plastic bag on ice.
- **Do NOT place amputated part directly on ice or in water.**
- Transport amputated part with person or as soon as it is located.

Notify responsible school authority and parent or legal guardian.

ANAPHYLAXIS / ALLERGIC REACTIONS (PART 1)

Anaphylaxis is a serious, rapidly progressing, whole body allergic reaction that can be fatal if not treated immediately. It can occur in a person who has a hypersensitivity to foods, insect stings, medications, or other allergens. **Children may experience a delayed reaction up to 2 hours following the allergen exposure.** The risk of anaphylaxis and death from anaphylaxis is higher among persons with asthma. Students with life-threatening allergies or who are at risk for anaphylaxis should be known to appropriate school staff. Symptoms of anaphylaxis are contained in the algorithm that follows.

Epinephrine is the medication of choice to treat anaphylaxis. According to the Annotated Code of Maryland, Education Article, Section 7-426.2, every local school system shall have stock auto-injector epinephrine to respond to a life-threatening event for students with no known history of anaphylaxis. The law reads as follows:

“Each county board shall establish a policy for public schools within its jurisdiction to authorize the school nurse and other school personnel to administer auto-injectable epinephrine, if available, to a student who is determined to be or perceived to be in anaphylaxis, regardless of whether the student:

- (1) Has been identified as having an anaphylactic allergy, as defined in § 7-426.1 of this subtitle; or
- (2) Has a prescription for epinephrine as prescribed by an authorized licensed health care practitioner under the Health Occupations Article.”

In addition, staff in each school building should be trained to use the epinephrine auto-injector.


1. INSERT YOUR LOCAL SCHOOL SYSTEM POLICY AND PROTOCOL ON THE USE OF EPINEPHRINE AUTO-INJECTORS BEHIND THIS TAB FOR YOUR REFERENCE.

2. Complete the information below:


Names of persons trained to use epinephrine auto-injector

Contact Number


ANAPHYLAXIS / ALLERGIC REACTIONS (PART 2)




ASTHMA & DIFFICULTY BREATHING



BEHAVIORAL EMERGENCIES




BITES




BITES

BLEEDING




BLISTERS




BRUISES

If student comes to school with unexplained unusual or frequent bruising, consider the possibility of child abuse. See "Child Abuse" section.



BURNS



CHEST PAIN

A person with chest pain may:

- **Be awake**, able to talk but complains of severe chest pain
- **Complain of chest pain** or pressure located in the center of the chest
- **State that the pain** feels like pressure, fullness, squeezing, or heaviness in the chest
- **State that pain** travels to shoulders, neck, lower jaw or down arms
- **State that pain** lasts more than 3-5 minutes
- **States that pain** has stopped completely and returned a short time later



CALL EMS/9-1-1.

- Have person rest quietly.
- Place person in a position of comfort.
- Loosen any tight clothing.

Observe for these additional vague symptoms:

- Lightheadedness or "feeling dizzy"
- Sweating
- Nausea
- Shortness of breath
- Ache, heartburn, or indigestion
- Fainting or loss of responsiveness

Monitor airway, breathing and signs of circulation.
See "CPR and AED."

Send for CPR trained staff. If person stops breathing or becomes unresponsive, begin CPR.
See "CPR and AED."

Notify responsible school authority and parent or legal guardian or emergency contact for adults.

NOTES ON PERFORMING CPR & AED

The American Heart Association (AHA) issued new CPR guidelines for laypersons in 2010. A compression-to-ventilation ratio of 30:2 is one emphasized component of these guidelines. Code of Maryland Regulations (COMAR 13A.05.05.09) requires at least one person in each school to become trained in CPR and the use of AEDs. Also, COMAR 13A.05.05.11 requires annual in-service training plans that include training in first aid and CPR for school personnel.

AUTOMATIC EXTERNAL DEFIBRILLATORS (AEDs)

An automatic external defibrillator (AED) is a small electronic device that helps to restore a normal heart rhythm by delivering an electric shock to the heart after detecting a life-threatening irregular rhythm. AEDs are not substitutes for CPR but are designed to increase the effectiveness of basic life support when integrated into the CPR cycle.



AEDs are safe to use for **any infant, child or adult in cardiac arrest, according to the American Heart Association (AHA)**. * Some AEDs are capable of delivering a “child” energy dose through smaller child pads. Use child pads / child system for infants and children younger than age 8, if available. If child system is not available, use adult AED and pads. Do not use the child pads or “child” energy dose for adults in cardiac arrest. The location of AEDs should be known to all school personnel.

CHEST COMPRESSIONS FOR INFANT, CHILD, AND ADULT

The AHA is placing more emphasis on the use of effective chest compressions during CPR. CPR chest compressions produce blood flow from the heart to the vital organs. While performing chest compressions on any infant, child or adult, **Push Hard, Fast and Deep.**



BARRIER DEVICES

Barrier devices, to prevent the spread of infections from one person to another, can be used when performing rescue breathing. Several different types (e.g., face shields, pocket masks) exist. It is important to learn and practice using these devices in the presence of a trained CPR instructor before attempting to use them in an emergency situation. Rescue breathing technique may be affected by these devices.



* AHA Guidelines, 2010.

AUTOMATIC EXTERNAL DEFIBRILLATORS (AEDs) FOR ALL AGES (INFANT, CHILD, & ADULT)



CPR and AEDs are to be used when a person is unresponsive or when breathing or heart beat stops.

If your school has an AED, this guideline will refresh information provided in training courses as to incorporating AED use into CPR cycles.

1. Tap or gently shake the shoulder. Shout, "Are you OK?" If person is unresponsive, shout for help and **send one person to CALL EMS/9-1-1 and another person to get your school's AED if available.**
2. Follow primary steps for CPR (see "CPR" for appropriate age group – infant and over 1 year, including adults).
3. If available, set up the AED according to the manufacturer's instructions. Turn on the AED and follow the verbal instructions provided. Incorporate AED into CPR cycles according to instructions and training method.

IF CARDIAC ARREST OR COLLAPSE WAS WITNESSED:

1. Use the AED first if immediately available. If not, begin CPR.
2. Prepare AED to check heart rhythm and deliver 1 shock as necessary.
3. Begin 30 CPR chest compressions in about 20 seconds followed by 2 normal rescue breaths. See age-appropriate CPR guideline.
4. Complete 5 cycles of CPR (30 chest compressions in about 20 seconds to 2 breaths for a rate of 100 compressions per minute).
5. Prompt another AED rhythm check.
6. Rhythm checks should be performed after every 2 minutes (about 5 cycles) of CPR.
7. REPEAT CYCLES OF 2 MINUTES OF CPR TO 1 AED RHYTHM CHECK UNTIL VICTIM RESPONDS OR HELP ARRIVES.



IF CARDIAC ARREST OR COLLAPSE WAS NOT WITNESSED:

1. Start CPR first. See age appropriate CPR guideline. Continue for 5 cycles or about 2 minutes of 30 chest compressions in about 20 seconds to 2 breaths at a rate of at least 100 compressions per minute.
2. Prepare the AED to check the heart rhythm and deliver a shock as needed.
3. REPEAT CYCLES OF 2 MINUTES OF CPR TO 1 AED RHYTHM CHECK UNTIL VICTIM RESPONDS OR HELP ARRIVES.



CPR FOR INFANTS UNDER 1 YEAR

CPR is to be used when a student is unresponsive or when breathing or heart beat stops.

1. Gently shake infant. If no response, shout for help and send someone to **call EMS/9-1-1 and get your school's AED, if available.**
2. Turn the infant onto his/her back as a unit by supporting the head and neck.
3. Evaluate for signs of circulation, which include breathing, moving, or coughing.
4. If no signs of circulation exist, begin CPR, beginning with chest compressions at a rate of 100 compressions-per-minute. Remember to allow the chest to return to its normal position in between each compression. **Push hard, fast, and deep.**



Begin CPR:

1. Find finger position near center of breastbone just below the nipple line. (Make sure fingers are NOT over the very bottom of the breastbone.)
2. Compress chest hard and fast at rate of 30 compressions in about 20 seconds with 2 or 3 fingers about 1/3 to 1/2 the depth of the infant's chest.

Use equal compression and relaxation times. Limit interruptions in chest compressions.

3. If you have been trained to provide breathing, provide two (2) breaths with each breath lasting 1 second and watch for the chest to rise with each breath.
4. REPEAT CYCLES OF 30 COMPRESSIONS TO 2 BREATHS AT A RATE OF 100 COMPRESSIONS PER MINUTE UNTIL INFANT STARTS BREATHING EFFECTIVELY ON THEIR OWN OR HELP ARRIVES.



IF CHEST DOES NOT RISE WITH RESCUE BREATH (AIR DOES NOT GO IN):

Re-tilt head back. Try to give 2 breaths again.

CPR AGE 1 THRU ADULTS

CPR is to be used when a student is unresponsive or when breathing or heart beat stops.

1. Tap or gently shake the shoulder. Shout, "Are you OK?" If child is unresponsive, shout for help and send someone to **call EMS/9-1-1 and get your school's AED, if available.**
2. Turn the child onto his/her back as a unit by supporting the head and neck. If head or neck injury is suspected, **DO NOT BEND OR TURN NECK.**
3. Evaluate for signs of circulation, which include breathing, moving, or coughing.
4. If no signs of circulation exist, begin CPR, beginning with chest compressions at a rate of 100 compressions-per-minute. Remember to allow the chest to return to its normal position in between each compression. **Push hard, fast, and deep.**



Begin CPR:

1. Find hand position near center of breastbone just below the nipple line. (Make sure hand(s) are **NOT** over the very bottom of the breastbone.
2. Compress chest hard and fast at rate of 30 compressions in about 20 seconds with 1 or 2 hands* about 2 inches in depth.

Use equal compression and relaxation times. Limit interruptions in chest compressions.
3. If you have been trained to provide breathing, provide two (2) breaths with each breath lasting 1 second and watch for the chest to rise with each breath.
4. REPEAT CYCLES OF 30 COMPRESSIONS TO 2 BREATHS AT A RATE OF 100 COMPRESSIONS PER MINUTE UNTIL PATIENT STARTS BREATHING EFFECTIVELY ON THEIR OWN OR HELP ARRIVES.



* Hand positions for child CPR:

- **1 hand:** Use heel of 1 hand only
- **2 hands:** Use heel of 1 hand with second on top of first

CHOKING

Call EMS/9-1-1 after starting rescue efforts.

INFANTS UNDER 1 YEAR

Begin the following if the infant is choking and is unable to breathe. However, if the infant is coughing or crying, do **NOT** do any of the following, but call EMS, try to calm the child and watch for worsening of symptoms. If cough becomes ineffective (loss of sound), begin step 1 below.


1. Position the infant, with head slightly lower than chest, face down on your arm and support the head (support jaw; do **NOT** compress throat).
2. Give up to 5 back slaps with the heel of hand between infant's shoulder blades.
3. If object is not coughed up, position infant face up on your forearm with head slightly lower than rest of body.
4. With 2 or 3 fingers, give up to 5 chest thrusts near center of breastbone, just below the nipple line.
5. Open mouth and look. If foreign object is seen, sweep it out with the finger.
6. REPEAT STEPS 1-5 UNTIL OBJECT IS COUGHED UP OR INFANT STARTS TO BREATHE OR BECOMES UNCONSCIOUS.
7. Call EMS/9-1-1 after 2 minutes (5 cycles of 30 compressions to 2 rescue breaths) if not already called

IF INFANT BECOMES UNRESPONSIVE, BEGIN THE STEPS OF INFANT CPR.



CHILDREN OVER 1 YEAR OF AGE & ADULTS

Begin the following if the victim is choking and unable to breathe. Ask the victim: "Are you choking?" If the victim nods yes or can't respond, help is needed. However, if the victim is coughing, crying or speaking, do **NOT** do any of the following, but call EMS, try to calm him/her and watch for worsening of symptoms. If cough becomes ineffective (loss of sound) and victim cannot speak, begin step 1 below



1. Stand behind an adult, or stand or kneel behind child with arms encircling patient.
2. Place thumbside of fist against middle of abdomen just above the navel. (Do NOT place your hand over the very bottom of the breastbone. Grasp fist with other hand).
3. Give up to 5 quick inward and upward abdominal thrusts.
4. REPEAT STEPS 1-2 UNTIL OBJECT IS COUGHED UP AND THE CHILD STARTS TO BREATHE OR CHILD BECOMES UNCONSCIOUS.

IF CHILD OR ADULT BECOMES UNRESPONSIVE, PLACE ON BACK AND BEGIN THE STEPS OF CPR.

FOR OBESE OR PREGNANT PERSONS:

Stand behind person and place your arms under the armpits to encircle the chest. Press with quick backward thrusts.




CHILD ABUSE & NEGLECT

According to Family Article Title 5, Subtitle 7; COMAR 13A.12.05.01D; Education Article Section 6-202, child abuse and neglect includes child physical abuse, sexual abuse, human trafficking of youth, neglect, mental injury abuse and mental injury neglect. Child abuse and neglect is a complicated issue with many potential signs. Some signs are listed below. This is not a complete list:

- Depression, hostility, low-self-esteem, poor self-image
- Evidence of repeated injuries or unusual injuries
- Lack of explanation or unlikely explanation for an injury
- Pattern bruises or marks (e.g., burns in the shape of a cigarette or iron, bruises or welts in the shape of a hand)
- Unusual knowledge of sex, inappropriate touching or engaging in sexual play with other children
- Severe injury or illness without medical care
- Poor hygiene, underfed appearance

All school staff are required to report suspected child abuse and neglect (COMAR 13A.12.05.01). Failure to report may result in revocation of licensure or certification and loss of employment. Follow local school system policy on reporting Child Abuse and Neglect.

All communication should be done in a nonjudgmental and confidential manner



COMMUNICABLE DISEASES RESOURCES


The Maryland Department of Health and Mental Hygiene offers advice on the control of communicable disease.

More information can be found at:
[http://phpa.dhmh.maryland.gov/
IDEHASharedDocuments/guidelines/
CDSummary_FINAL_2011_Nov.pdf](http://phpa.dhmh.maryland.gov/IDEHASharedDocuments/guidelines/CDSummary_FINAL_2011_Nov.pdf)

Reportable Diseases
[http://phpa.dhmh.maryland.gov/SitePages/
reportable-diseases.aspx](http://phpa.dhmh.maryland.gov/SitePages/reportable-diseases.aspx)

Follow local school system
policy for reporting of
communicable diseases.

CUTS, SCRATCHES & SCRAPES



CUTS, SCRATCHES & SCRAPES


DIABETES

A student with diabetes should be known to appropriate school staff. An emergency/ action plan must be developed.


A student with diabetes may have the following symptoms:

- Irritability and feeling upset
- Change in personality
- Sweating and feeling “shaky”
- Loss of responsiveness
- Confusion or strange behavior
- Rapid, deep breathing


Refer to student’s emergency/action plan.



DIARRHEA



DROWNING (NEAR)




EAR PROBLEMS


DRAINAGE FROM EAR




EARACHE



OBJECT IN EAR CANAL




ELECTRIC SHOCK



EYE PROBLEMS


EYE INJURY




*Eye Problems
continued on next page*

EYE PROBLEMS (CONT.)


PARTICLE IN EYE



CHEMICALS IN EYE



FAINTING



FEVER

Fever is defined as a temperature >100.0° F orally; an oral temperature of 100.0° F is approximately equivalent to 101.0° F rectally or temporally (Temporal Artery Forehead Scan), or 99.5° F axillary (armpit).

Take student's temperature.
Note if student has fever.


Have the student lie down in a room
that affords privacy.

Give no medication, unless
previously authorized.

Contact responsible
school authority and
parent or legal guardian.
Follow local school
system policy for
students with fever.

FEVER

FRACTURES & SPRAINS



FROSTBITE

Frostbite can result in the same type of tissue damage as a burn. It is a serious condition and requires medical attention.

Exposure to cold even for short periods of time may cause "HYPOTHERMIA" in children (see "*Hypothermia*"). The nose, ears, chin, cheeks, fingers, and toes are the parts most often affected by frostbite.

Frostbitten skin may:

- Look discolored (flushed, grayish-yellow, pale)
- Feel cold to the touch
- Feel numb to the student

Deeply frostbitten skin may:

- Look white or waxy
- Feel firm or hard (frozen)

- Take the student to a warm place.
- Remove cold or wet clothing and give student warm, dry clothes.
- Protect cold part from further injury.
- **Do NOT rub or massage the cold part or apply heat such as a water bottle or hot running water.**
- Do not break open any blisters.
- Cover part loosely with nonstick, sterile dressings or dry blanket.

Does extremity/part:

- Look discolored – grayish, white or waxy?
- Feel firm/hard (frozen)?
- Have a loss of sensation?

YES

NO


CALL EMS/9-1-1.
Keep student warm and part covered.

Contact responsible authority and parent or legal guardian.

Contact responsible authority and parent or legal guardian.
Encourage medical care.

Keep student and part warm.

GENITOURINARY COMPLAINTS


Genitourinary complaints include trauma, injury, infection of the genitals (penis, testicles, vagina, vulva and surrounding area), itching, unusual color, or unusual odor.

Sexually transmitted infection and abuse must be ruled out regardless of person's age.


Physical examination should be left to medical professionals.

Complaints may include:


- Painful urination
- Blood or pus in urine
- Discharge from penis, vagina, or urethra
- Ulcers
- Swollen or painful inguinal (groin) lymph glands
- Swelling, redness, and tenderness of genitals
- Injury
- Trauma
- Rash




HEADACHE




HEAD INJURIES




HYPERTHERMIA (HEAT) EMERGENCIES




HYPOTHERMIA (COLD) EMERGENCIES




MENSTRUAL DIFFICULTIES



MOUTH & JAW INJURIES




NECK & BACK PAIN



NOSE PROBLEMS

NOSEBLEED

See "Head Injuries" section if you suspect a head injury other than a nosebleed or broken nose.



BROKEN NOSE


- Care for nose as in "Nosebleed" above.
- Contact responsible school authority and parent or legal guardian.
- **URGE MEDICAL CARE.**

*Nose Problems
continued on next page*


NOSE PROBLEMS

NOSE PROBLEMS (CONT.)

OBJECT IN NOSE




POISONING & OVERDOSE




PREGNANCY

Pregnant students should be known to appropriate school staff. Follow local school system communication policy regarding confidentiality. Under Health General, Sections 20-102 and 20-104, minors may consent to pregnancy related services. Consider issues of confidentiality when addressing pregnancy concerns. Follow local school system policy regarding confidential communication.


Pregnancy may be complicated by any of the following:




PUNCTURE WOUNDS



RAPE / SEXUAL ASSAULT



RASHES



SEIZURES

Seizures may be any of the following:

- Episodes of staring with loss of eye contact
- Staring involving twitching of the arm and leg muscles
- Generalized jerking movements of the arms and legs
- Unusual behavior for that person (e.g., running, belligerence, making strange sounds, etc.)
- If head injury is suspected, do not move the student.

A student with a history of seizures should be known to appropriate school staff. An emergency/action plan should be developed, containing a description of the onset, type, duration, and after effects of the seizures.

Refer to student's emergency/action plan.

- If student seems off balance, place him/her on the floor (on a mat) for observation and safety.
- **Do NOT restrain movements.**
- Move surrounding objects to avoid injury.
- **Do NOT place anything in between the teeth or give anything by mouth.**
- Keep airway clear by placing student on his/her side. A pillow should **NOT** be used.

Observe details of the seizure for parent/legal guardian, emergency personnel or physician. Note:

- Duration
- Kind of movement or behavior
- Body parts involved
- Loss of responsiveness, etc.

NO

- Is student having a seizure lasting longer than *5 minutes*?
- Is student having seizures following one another at short intervals?
- Is student *without a known history* of seizures having a seizure?
- Is student having any breathing difficulties after the seizure?

Seizures are often followed by sleep. The student may also be confused. This may last from 15 minutes to an hour or more. After the sleeping period, the student should be encouraged to participate in all normal class activities.


YES

Contact responsible school authority and parent or legal guardian.


CALL EMS/9-1-1.




SHOCK



SPLINTERS



STABBING & GUNSHOT INJURIES



STINGS

Most reactions to insect stings and bites cause mild local swelling, redness, itching or pain. However, some people experience life-threatening allergic reactions.

This type of reaction, anaphylaxis, is a serious, sudden, rapidly progressing whole body allergic reaction that can be fatal. According to the Annotated Code of Maryland, Education Article, Section 7-426.2, each county board shall establish a policy authorizing the school nurse and other school personnel to administer a stock epinephrine auto-injector to respond to anaphylaxis regardless of whether the student has a history of anaphylaxis or a prescription for epinephrine. See Anaphylaxis/Allergic Reaction section.

Students with a history of allergy to stings should be known to appropriate school staff. Adult(s) supervising students during normal activities should be aware of stings and should watch for signs of anaphylaxis, which may be delayed.

Does the student have any symptoms of a severe allergic reaction, which may include?

- Swelling of the back of the mouth / throat or tongue; feeling like the throat is closing; difficulty swallowing; hoarseness or change in quality of voice
- Coughing; wheezing; shortness of breath; difficulty breathing; noisy breathing; "air hunger" or gasping for air
- Dizzy / lightheaded; fainting; unresponsiveness
- Hives; generalized itching, tingling and / or swelling of face or extremities
- Nausea; abdominal pain or cramps; vomiting; diarrhea
- Uneasiness; agitation; panic; feeling of impending doom

- NO
- Remove stinger if present.
 - Wash area with soap and water.
 - Apply cold compress.

Contact parent or legal guardian.

- YES
- **Immediately** administer auto-injector epinephrine according to local school system policy for use of school stock auto-injector epinephrine, or call trained staff to administer.
 - Follow local school system policy for students with severe allergic reactions.
 - See "Anaphylaxis/Allergic Reaction" section.

CALL EMS/9-1-1.



Any student receiving epinephrine **must** be transported to the hospital.


Position child for comfort and offer reassurance while awaiting EMS.

Contact responsible school authority and parent or legal guardian.

STOMACHACHES & PAIN

Stomachaches/pain may have many causes, including:

- Illness
- Hunger
- Overeating
- Diarrhea
- Food poisoning
- Injury
- Menstrual difficulties
- Psychological issues
- Stress
- Constipation
- Gas pain
- Pregnancy



STOMACHACHES & PAIN

TEETH PROBLEMS

Refer to the “Dental First Aid for Children flip chart from the Office of Oral Health, Maryland Department of Health (MDH).

Flip chart:

<https://phpa.health.maryland.gov/oralhealth/Documents/FlipChart.pdf>

Resource Guides:

<https://phpa.health.maryland.gov/oralhealth/Documents/OralHealthResourceGuide.pdf> (English)

<https://phpa.health.maryland.gov/oralhealth/Documents/OralHealthResourceGuideSpanish.pdf> (Spanish)

TETANUS IMMUNIZATION

Protection against tetanus should be considered with any wound, even a minor one. After any wound, check the student's immunization record for tetanus and notify parent or legal guardian.

A **minor wound** may need a tetanus booster if it has been at least **10 years** since the last tetanus shot or if the student is **5 years old or younger**.

Other wounds such as those contaminated by dirt, feces, and saliva (or other body fluids); puncture wounds; amputations; and wounds resulting from crushing, burns, and frostbite may need a tetanus booster if it has been more than **5 years** since last tetanus shot.

The need for a tetanus immunization should be determined by a licensed health care provider.

TICKS

Students should be inspected for ticks after time in woods or brush. Ticks may carry serious infections and must be completely removed.

Do NOT handle ticks with bare hands.

Tick identification information:

<http://phpa.dhmh.maryland.gov/OIDEOR/CZVBD/SitePages/lyme-disease.aspx>

Refer to your school system policy regarding the removal of ticks.

Wear disposable gloves when exposed to blood and other body fluids.

Wash the tick area gently with soap and water before attempting removal.


- Using tweezers, grasp the tick as close to the skin surface as possible and pull upward with steady, even pressure.
- **Do NOT twist or jerk the tick as the mouth parts may break off.** It is important to remove the *ENTIRE* tick.
- Take care not to squeeze, crush or puncture the body of the tick as its fluids may carry infection.

- After removal, wash the tick area thoroughly with soap and water.
- Wash your hands.
- Apply a bandage.


Dispose of tick following local school system policy.

Contact responsible school authority and parent or legal guardian.

UNRESPONSIVENESS



VOMITING



SCHOOL SAFETY PLANNING AND EMERGENCY PREPAREDNESS

All school staff should be aware of the school's safety plan and be prepared for a variety of emergencies, disasters and hazards. The pages that follow contain important information and resources related to:

1. School safety planning
2. Emergency preparedness for kids and parents
3. Emergency kit checklist

It is important that the school safety and emergency plan be quickly accessible if needed.

Please insert your school's safety plan into this binder behind this tab.

EMERGENCY PREPAREDNESS RESOURCES

The following resources may be helpful when addressing emergency preparedness in schools.

General Information for the Community

Emergency Supply Kits:

<http://preparedness.dhmh.maryland.gov/Documents/Fact%20SheetEmergencyKit.pdf>

Community and Personal Preparedness (this page also includes some links to Emergency Preparedness for Kids, which may be especially relevant/useful for educators):

<http://preparedness.dhmh.maryland.gov/SitePages/Community%20And%20Personal%20Preparedness.aspx>

Emergency Planning and Other Resources for Schools

Emergency Planning Guidelines for Local School Systems and Schools, Maryland State Department of Education, 2013

1. See Appendix A-2 – Hazard Profile Key
2. See Appendix I on page 209 for a list of local and state Emergency Management contacts

<http://www.marylandpublicschools.org/w/EmergencyPlanningGuidelines2013.pdf>

Lesson plans for educators on emergency preparedness

<http://www.ready.gov/kids/educators>

National Center for School Crisis and Bereavement

<http://www.schoolcrisiscenter.org/index.html>

Maryland Center for School Safety

www.MCFSS.maryland.gov

Emergency Preparedness Resources for Kids and Parents

Be a Hero!

This newly redesigned website includes games and materials for kids and parents, as well as lesson plans about emergency preparedness for educators.

<http://www.ready.gov/kids>

Let's Get Ready! Planning Together for Emergencies

This page, featuring characters from Sesame Street, has fun videos and activities kids will love. Included is a guide for educators under the provider tab.

<http://www.sesamestreet.org/parents/topicsandactivities/toolkits/ready>

Ready Wrigley

The Ready Wrigley page from the CDC includes an interactive site, as well as coloring books on hurricanes and earthquakes.

<http://www.cdc.gov/phpr/readywrigley/>

EMERGENCY PREPAREDNESS RESOURCES (CONT.)

American Academy of Pediatrics

The AAP has a dedicated website for family and care givers that includes resources for emergencies, disaster preparedness and response to school violence.

<https://healthychildren.org/English/safety-prevention/at-home/Pages/Getting-Your-Family-Prepared-for-a-Disaster.aspx>

<https://healthychildren.org/English/safety-prevention/at-home/Pages/Family-Disaster-Supplies-List.aspx>

<https://healthychildren.org/English/news/Pages/AAP-Offers-Resources-to-Help-Parents,-Children-and-Others-Cope-in-the-Aftermath-of-School-Violence.aspx>

EMERGENCY SUPPLY KIT CHECKLIST



Emergency Supply Kit Checklist

Be prepared for any emergency. Assemble an emergency supply kit with items to take care of yourself, your family and your pets for three or more days. Keep your kit in sturdy and easy-to-carry backpacks or duffle bags. Depending on the situation, you may be told by authorities to shelter-in-place (stay inside) or evacuate with your kit. The following checklist will help you put your kit together.

WATER and FOOD

- one gallon of water per person, per day, plus water for pets
- ready-to-eat canned meats, fish, soups, beans, vegetables and fruits. *Choose foods that need little or no cooking.*
- salt, pepper, sugar, spices
- powdered milk, tea, instant coffee
- high-energy snacks: nuts, protein bars, trail mix, peanut butter
- comfort foods: granola, dried fruits, cookies, crackers, hard candy, cocoa
- foods for infants, individuals with special needs and pets
- paper cups, plates and plastic utensils
- camp cook kit or pans
- manual can opener
- aluminum foil, plastic wrap

Rotate the food in your supply kit regularly. Some foods should be used within six months, such as powdered milk, dried fruits and crackers. Other foods will keep for up to one year, such as canned soups and meats, fruits, vegetables and juices, peanut butter, jelly, hard candy and canned nuts. Foods that can be stored indefinitely (in air-tight containers away from heat) include vegetable oil, dried corn and wheat, baking powder, soybeans, instant coffee, tea and cocoa, salt, rice, bouillon products and dry pasta.

EMERGENCY SUPPLIES

- cash, traveler's checks, coins
- battery-operated radio, NOAA Weather Radio
- flashlight
- batteries
- cell phone
- face masks
- maps of your area and nearby states
- whistle
- extra set of house and car keys
- small fire extinguisher
- wrench or pliers to turn off utilities
- plastic garbage bags with twist ties
- toilet paper, towelettes
- household chlorine bleach (*to purify water*)
- matches in a waterproof container
- soap, detergent, alcohol-based hand sanitizer
- toothbrushes and toothpaste, dental floss, deodorant, shampoo, shaving supplies
- feminine supplies, condoms
- lip balm, sunscreen
- infant supplies (diapers, bottles, etc.)
- pet supplies (litter, flea collar, etc.)
- books, playing cards, board games

FIRST AID KIT

- first aid manual
- prescription drugs; a two week supply of every household member's vital medications
- nonprescription drugs: aspirin or other pain reliever, allergy medicine, anti-diarrhea medication, antacid, laxative, antibiotic ointment, vitamins, eye wash
- prescribed medical supplies, such as glucose and blood pressure monitoring equipment
- scissors, tweezers, magnifying glass
- sterile needle, safety razor blade
- thermometer
- insect repellent
- mirror
- sterile adhesive bandages (Band-Aids) in assorted sizes, gauze pads and roller bandages
- hypoallergenic adhesive tape
- several pairs of disposable gloves
- isopropyl alcohol, hydrogen peroxide
- antiseptic, antiseptic spray
- cold packs and heat packs

-more-

EMERGENCY SUPPLY KIT CHECKLIST (CONTINUED)

CLOTHING and BEDDING

- include at least one complete change of clothing and shoes per person
- long pants and long sleeve shirt
- sturdy shoes or work boots
- thermal underwear, regular underwear
- several pairs of socks
- warm hat and work gloves
- jacket or coat, rain gear, poncho
- sleeping bag
- blankets, space blankets, pillows
- towels, washcloths
- extra prescription glasses, sunglasses

IMPORTANT DOCUMENTS

Keep copies (not originals) in a waterproof, portable container or scan to a CD or USB drive

- bank account numbers (checking, savings)
- credit account numbers, with company names and contact information
- Social Security cards and records
- passports
- family records: birth, marriage and death certificates, divorce decree
- wills, living wills, advanced directives
- power of attorney papers
- medical records
- current medical and eyeglass prescriptions
- immunization records of family, pets
- all insurance policies (life, health, auto, home, hazard)
- deeds, mortgages, titles, rental agreement
- stocks and bonds, securities, investment statements
- bank loan agreements, other contracts
- motor vehicle titles, bill of sale, serial or VIN numbers, driver's licenses
- employment records
- recent tax returns
- records of valuable collections, appraisals
- school transcripts, diplomas
- safe deposit box location and extra key, inventory of contents
- original manuscripts, discs
- journals, diaries, genealogies
- inventory of household goods (including photographs)
- current photographs of family members, pets
- favorite photographs of family members, pets and events

Learn more at <http://preparedness.dhmh.maryland.gov> and www.facebook.com/MarylandOPR.

EMERGENCY PHONE NUMBERS

Complete this page as soon as possible and update as needed.

EMERGENCY MEDICAL SERVICES (EMS) INFORMATION

Know how to contact your EMS. Most areas use 9-1-1; others use a 7-digit phone number.

+ **EMERGENCY PHONE NUMBER: 9-1-1 OR** _____

+ Name of EMS agency _____

+ Their average emergency response time to your school _____

+ Directions to your school _____

+ Location of the school's AED(s) _____

BE PREPARED TO GIVE THE FOLLOWING INFORMATION & DO NOT HANG UP BEFORE THE EMERGENCY DISPATCHER HANGS UP:

- Name and school name _____
- School telephone number _____
- Address and easy directions _____
- Nature of emergency _____
- Exact location of injured person (e.g., behind building in parking lot) _____
- Help already given _____
- Ways to make it easier to find you (e.g., standing in front of building, red flag, etc.).

OTHER IMPORTANT PHONE NUMBERS

+ **School Nurse** _____

+ **Responsible School Authority** _____

+ **Poison Control Center** **1-800-222-1222**

+ **Fire Department** **9-1-1 or** _____

+ **Police** **9-1-1 or** _____

+ Hospital or Nearest Emergency Facility _____

+ County Children Services Agency _____

+ Rape Crisis Center _____

+ Suicide Hotline _____

+ Local Health Department _____

+ Taxi _____

+ Other medical services (e.g., dentists): _____