

## The BESURE Study 2004-2010

Danielle German, PhD, MPH
Assistant Scientist

Department of Health, Behavior and Society
Johns Hopkins Bloomberg School of Public Health





#### Overview

- HET, IDU, MSM: focus on last round
  - Overview of sample characteristics and types of data
  - Sample of analyses, focus on HIV
- Next steps



**HET: 2007 AND 2010** 



### **HET** overview

|                                   | Wave 1<br>July – Oct 2007 | Wave 2<br>Sept – Dec 2010 |
|-----------------------------------|---------------------------|---------------------------|
| Data collection method            | Venue – based             | Respondent Driven         |
| Total recruited *                 | 332                       | 338                       |
| HIV prevalence                    | 3.9%                      | 6.3%                      |
| Prevalence unrecognized infection | 83.3%                     | 62.5%                     |



<sup>\*</sup>Complete survey and HIV test

### Sample characteristics: HET1 and HET2

| Characteristic |                                                                       | Wave 1<br>(n=332)        | Wave 2<br>(n=386)        |
|----------------|-----------------------------------------------------------------------|--------------------------|--------------------------|
| Gender         | Male<br>Female                                                        | 54%<br>46%               | 48%<br>52%               |
| Race/Ethnicity | Black, not Hispanic * White, not Hispanic Other                       | 89%<br>4%<br>8%          | 95%<br>2%<br>3%          |
| Age            | 18-24<br>25-34<br>35-44 *<br>45 or older *                            | 24%<br>27%<br>33%<br>16% | 25%<br>26%<br>17%<br>32% |
| Education      | Less than high school High school grad or GED More than high school * | 37%<br>42%<br>20%        | 42%<br>44%<br>14%        |

### Sample characteristics: HET1 and HET2

| Characteristic                   |                                                      | Wave 1<br>(n=332) | Wave 2<br>(n=386) |
|----------------------------------|------------------------------------------------------|-------------------|-------------------|
| Employment                       | Full or part-time * Disabled for work * Unemployed * | 41%<br>11%<br>36% | 25%<br>16%<br>45% |
| Homelessness                     | Current * Past year *                                | 10%<br>24%        | 19%<br>39%        |
| Arrested in past year            | Yes                                                  | 29%               | 24%               |
| Health insurance                 | Yes *                                                | 48%               | 63%               |
| Healthcare provider in past year | Yes                                                  | 78%               | 74%               |
| Ever injection drug use          | Yes                                                  | 21%               | 22%               |
| Drug use in past year            | Injection drug use<br>Crack use                      | 13%<br>22%        | 9%<br>17%         |

### What counts as "sexual risk"? (HET1)

- Unprotected sex?
  - 84% report any unprotected vaginal or anal sex in past year
- Type of partner?
  - 72% report unprotected sex with only main partners
  - 38% report unprotected sex with casual or exchange partners

# Unprotected intercourse by STD history

|                                      | No STD | Any STD | <i>p</i> -value |
|--------------------------------------|--------|---------|-----------------|
| Comparing all groups                 |        |         |                 |
| No unprotected sex                   | 89%    | 11%     | 0.01            |
| With main partners only              | 93%    | 7%      |                 |
| With casual/exchange partners only   | 82%    | 18%     |                 |
| Comparing casual/exchange to main    |        |         |                 |
| With main partners only              | 93%    | 7%      | 0.003           |
| With casual/exchange partners only   | 82%    | 18%     |                 |
| Comparing main to no unprotected sex |        |         |                 |
| No unprotected sex                   | 89%    | 11%     | 0.37            |
| With main partners only              | 93%    | 7%      |                 |



### Research question

- Reported unprotected sex with casual or exchange partners (HET1): 38%
  - What are the associated socio-demographic characteristics?
  - What are the associated sexual behaviors?
  - What are other associated risks?



### Characteristics associated with unprotected sex with casual or exchange partners: Demographics

| Characteristic                     |                                  | Total #<br>(n=301)   | Unprotected % (n=115)    | Prevalence<br>Ratio             |
|------------------------------------|----------------------------------|----------------------|--------------------------|---------------------------------|
| Current homelessness               | Yes                              | 27                   | 63%                      | 1.8 *                           |
| High school education              | Yes                              | 192                  | 35%                      | -                               |
| Race/ethnicity                     | African-Amer                     | 271                  | 37%                      | -                               |
| Age                                | 18-24<br>25-33<br>34-41<br>42-50 | 80<br>77<br>71<br>73 | 21%<br>44%<br>48%<br>41% | Ref.<br>2.0 *<br>2.3 *<br>1.9 * |
| Gender                             | Male                             | 161                  | 44%                      | 1.4 *                           |
| Marital status                     | Married/living                   | 34                   | 32%                      | -                               |
| Living below poverty level         | Yes                              | 190                  | 37%                      | -                               |
| Healthcare past year               | Yes                              | 238                  | 35%                      | 0.69 *                          |
| Health insurance status            | No insurance                     | 156                  | 41%                      | -                               |
| Arrested/ booked past year *p<0.05 | Yes                              | 89                   | 49%                      | 1.5 *                           |

### Characteristics associated with unprotected sex with casual or exchange partners: Sex and HIV

| Characteristic                  |                                                                          | Total #<br>(n=301)   | Unprotected % (n=115)   | Prev.<br>Ratio                  |
|---------------------------------|--------------------------------------------------------------------------|----------------------|-------------------------|---------------------------------|
| Number of opposite sex partners | 1<br>2<br>3<br>4 or more                                                 | 99<br>70<br>47<br>85 | 9%<br>34%<br>47%<br>71% | Ref.<br>3.8 *<br>5.2 *<br>7.8 * |
| Past year, any sex with:        | Same sex partners Concurrent partners                                    | 20<br>178            | 65%<br>57%              | 1.8 *<br>5.0 *                  |
| Substance use in past year      | Injection drug use<br>Non-injection drug use<br>Binge drinking (past mo) | 36<br>173<br>100     | 58%<br>46%<br>48%       | 1.6 *<br>1.7 *<br>1.4 *         |
| Ever told had STD               | Yes                                                                      | 37                   | 60%                     | 1.7 *                           |
| HIV status                      | Positive                                                                 | 10                   | 50%                     | -                               |
| Ever HIV test                   | Yes                                                                      | 255                  | 37%                     | -                               |
| HIV test in past 2 years        | Yes                                                                      | 190                  | 36%                     | -                               |

### Characteristics associated with unprotected sex with casual or exchange partners: Multivariate

| Characteristic                  |                                  | Total #<br>(n=301)   | Unprot. %<br>(n=115)     | Prev.<br>Ratio                  | Adj.<br>P. R.                   |
|---------------------------------|----------------------------------|----------------------|--------------------------|---------------------------------|---------------------------------|
| Currently homeless              | Yes                              | 27                   | 63%                      | 1.8 *                           | 1.2 *                           |
| Age                             | 18-24<br>25-33<br>34-41<br>42-50 | 80<br>77<br>71<br>73 | 21%<br>44%<br>48%<br>41% | Ref.<br>2.0 *<br>2.3 *<br>1.9 * | Ref.<br>1.6 *<br>1.9 *<br>2.2 * |
| Number of opposite sex partners | 1<br>2<br>3<br>4 or more         | 99<br>70<br>47<br>85 | 9%<br>34%<br>47%<br>71%  | Ref.<br>3.8 *<br>5.2 *<br>7.8 * | Ref.<br>-<br>-<br>3.5 *         |
| Concurrent sex partners         | Yes                              | 178                  | 57%                      | 5.0 *                           | 2.1 *                           |
| Any binge drinking past month   | Yes                              | 100                  | 48                       | 1.4 *                           | 1.2 *                           |
| Ever told by physician had STD  | Yes                              | 37                   | 60%                      | 1.7 *                           | 1.2 *                           |

### Summary

- Limitations:
  - Cross-sectional, not causal
  - Voluntary enrollment
  - Self-report
  - Don't know if findings also true in HET2 sample (different characteristics)
- Prevention implications:
  - Associated with variety of potential risk behaviors
  - Homelessness and older age as co-factors

#### **IDU: 2006 AND 2009**



#### **IDU** overview

|                                    | Wave 1<br>Aug 2006 – Jan 2007 | Wave 2<br>Sept – Dec 2009 |
|------------------------------------|-------------------------------|---------------------------|
| Data collection method             | Respondent Driven             | Respondent Driven         |
| Total IDU in past year recruited * | 539                           | 507                       |
| HIV prevalence                     | 11.9%                         | 16.2%                     |
| Prevalence unrecognized infection  | 50%                           | 48%                       |



<sup>\*</sup>Complete survey and HIV test

### Sample characteristics: IDU1 and IDU2

| Characteristic         |                                                 | Wave 1<br>(n=539) | Wave 2<br>(n=507) |
|------------------------|-------------------------------------------------|-------------------|-------------------|
| Gender                 | Male<br>Female *                                | 61%<br>38%        | 73%<br>27%        |
| Race/Ethnicity         | Black, not Hispanic * White, not Hispanic Other | 54%<br>41%<br>6%  | 80%<br>16%<br>4%  |
| Age                    | 18-34<br>35-44<br>45 or older *                 | 27%<br>42%<br>31% | 14%<br>23%<br>64% |
| Education              | Less than high school<br>High school or more    | 48%<br>52%        | 43%<br>57%        |
| Homelessness past year | Yes                                             | 58%               | 54%               |
| Arrested in past year  | Yes                                             | 47%               | 44%               |
| Hepatitis C diagnosis  | Yes *                                           | 29%               | 50%               |

\* p<0.05

### Sample characteristics: IDU1 and IDU2

| Characteristic                                 |              | Wave 1<br>(n=539) | Wave 2<br>(n=507) |
|------------------------------------------------|--------------|-------------------|-------------------|
| Years since first IDU                          | 10 or more * | 61%               | 86%               |
| Daily IDU                                      | Yes *        | 79%               | 90%               |
| Crack use past year                            | Yes          | 46%               | 43%               |
| Used someone's needle past year                | Yes *        | 53%               | 28%               |
| Always used sterile needle past year           | Yes          | 24%               | 25%               |
| Used someone's cooker, cotton, water past year | Yes          | 64%               | 69%               |



### Research questions (IDU2)

- Characteristics associated with HIV infection
  - Total: 16.2%
  - RDS Adjusted: 13.2 (95% C.I.: 9.4, 16.8)
- Characteristics associated with unrecognized infection
  - Total: 8.3 (or 48% of HIV positive)
  - RDS Adjusted: 4.7 (2.7 7.8)



## Characteristics associated with HIV: socio-demographic (IDU2)

| Characteristic     |                                                   | Total #<br>(n=507) | HIV+ %<br>(n=82) | Odds Ratio<br>(RDS Adj.) |
|--------------------|---------------------------------------------------|--------------------|------------------|--------------------------|
| Gender             | Male<br>Female                                    | 370<br>136         | 18%<br>13%       | Ref<br>-                 |
| Race/Ethnicity     | Non-Hispanic White<br>Non-Hispanic Black<br>Other | 83<br>404<br>20    | 7%<br>18%<br>10% | Ref<br>-<br>-            |
| Age                | 18-34<br>35-44<br>> 45                            | 69<br>116<br>322   | 6%<br>15%<br>19% | Ref<br>-<br>2.5 *        |
| Education          | At least high school                              | 287                | 15%              | 0.3 *                    |
| Homeless past year | Yes                                               | 274                | 13%              | -                        |
| Arrested past year | Yes                                               | 223                | 14%              | -                        |
| Any STI past year  | Yes                                               | 33                 | 15%              | -                        |
| Diagnosed with HCV | Yes                                               | 255                | 18%              | 2.2 *                    |

# Characteristics associated with HIV: drug and sex behaviors (IDU2)

| Characteristic                        |                            | Total #<br>(n=507) | HIV+<br>%<br>(n=82) | Odds Ratio<br>(RDS Adj.) |
|---------------------------------------|----------------------------|--------------------|---------------------|--------------------------|
| Years since first injection           | Less than 10<br>10 or more | 70<br>437          | 4%<br>18%           | Ref.<br>10.9 *           |
| Daily injection drug use              | Yes                        | 454                | 17%                 | -                        |
| Crack use past year                   | Yes                        | 218                | 12%                 | -                        |
| Used anyone's needle past year        | Yes                        | 144                | 13%                 | 0.3 *                    |
| Share cooker, cotton, water past year | Yes                        | 351                | 16%                 | -                        |
| Number sex partners past year         | 0<br>1<br>2 or more        | 70<br>199<br>230   | 29%<br>18%<br>11%   | Ref<br>0.5 *<br>-        |
| Any unprotected sex past year         | Yes                        | 349                | 10%                 | 0.3 *                    |
| Last sex with HIV+ partner            | Yes                        | 12                 | 3%                  | 5.3 *                    |

# Characteristics associated with HIV (IDU2)

| Characteristic                |                        | Total #<br>(n=507) | HIV+ %<br>(n=82)  | Odds Ratio<br>(RDS Adj.) | Adj. Odds<br>Ratio |
|-------------------------------|------------------------|--------------------|-------------------|--------------------------|--------------------|
| Age                           | 18-34<br>35-44<br>> 45 | 69<br>116<br>322   | 6%<br>15%<br>19%  | Ref<br>-<br>2.5 *        | Ref<br>-<br>-      |
| Years since first IDU         | 10 or more             | 437                | 18%               | 10.9 *                   | 7.0 *              |
| Education                     | At least h.s.          | 287                | 15%               | 0.3 *                    | 0.2 *              |
| Diagnosed with HCV            | Yes                    | 255                | 18%               | 2.2 *                    | -                  |
| Used anyone's needle          | Yes                    | 144                | 13%               | 0.3 *                    | 0.4 *              |
| Number sex partners past year | 0<br>1<br>2 or more    | 70<br>199<br>230   | 29%<br>18%<br>11% | Ref<br>0.5 *<br>-        | Ref.<br>-<br>-     |
| Any unprotected sex           | Yes                    | 349                | 10%               | 0.3 *                    | 0.2 *              |
| Last sex with HIV+ partner    | Yes                    | 12                 | 3%                | 5.3 *                    | 6.6 *              |


## Characteristics associated with unrecognized HIV infection (IDU2)

- 48% of HIV-positive
- No difference:
  - Socio-demographic characteristics
  - Drug use patterns
  - Injection risk behaviors



### HIV testing and unrecognized HIV infection

- Among those unaware of HIV positive status (n=42)
  - Ever test for HIV: 72%
  - Never test for HIV: 28%
- Among those ever tested for HIV (n=34)
  - Tested in past year: 36%
  - Not tested in past year:64%



### Summary

- Limitations:
  - Cross-sectional, not causal
  - Voluntary enrollment
  - Self-report
  - Comparison to IDU1 (Different characteristics)
- Prevention implications:
  - Low HIV risk behaviors among HIV-positive IDUs
  - Continued need for testing
  - Need to counter "fear of being positive"
  - Increase "always sterile needle"
  - Persistently high rates of cooker, cotton, water sharing
  - High arrest and homelessness

### MSM: 2004/2005 AND 2008



#### MSM overview

|                                   | Wave 1<br>July – Oct 2007 | Wave 2<br>Sept – Dec 2010 |
|-----------------------------------|---------------------------|---------------------------|
| Data collection method            | Venue – based             | Venue – based             |
| Total MSM in past year *          | 645                       | 448                       |
| HIV prevalence                    | 37.7%                     | 37.5%                     |
| Prevalence unrecognized infection | 58.4%                     | 78.4%                     |



<sup>\*</sup>Complete survey and HIV test

## Sample characteristics: MSM1 and MSM2

| Characteristic       | Wave 1<br>(n=645)                                              | Wave 2<br>(n=448)        |                          |
|----------------------|----------------------------------------------------------------|--------------------------|--------------------------|
| Race/Ethnicity       | Black, not Hispanic * White, not Hispanic * Other              | 63%<br>31%<br>6%         | 71%<br>23%<br>5%         |
| Age                  | 18-24 * 25-34 35-44 * 45 or older                              | 24%<br>26%<br>31%<br>18% | 31%<br>30%<br>24%<br>15% |
| Education            | High school or less College or some college Graduate education | 49%<br>45%<br>6%         | 48%<br>45%<br>6%         |
| Sexual identity      | Homosexual Bisexual Heterosexual or other *                    | 63%<br>32%<br>5%         | 68%<br>30%<br>2%         |
| Current homelessness | Yes *                                                          | 3%                       | 6%                       |

<sup>\*</sup> *p*<0.05

## Sample characteristics: MSM1 and MSM2

| Characteristic                 | Wave 1<br>(n=645)                | Wave 2<br>(n=448)        |                          |
|--------------------------------|----------------------------------|--------------------------|--------------------------|
| Sex partners in past year      | Men only * Men and women *       | 67%<br>33%               | 76%<br>24%               |
| # male partners past year      | 1<br>2-3 *<br>4-8<br>9 or more * | 26%<br>28%<br>23%<br>23% | 24%<br>34%<br>26%<br>16% |
| Unprotected anal sex past year | Yes *                            | 53%                      | 36%                      |
| Ever IDU                       | Yes *                            | 17%                      | 6%                       |
| Non-injection drugs past year  | Yes *                            | 49%                      | 59%                      |
| Ever tested for HIV            | Yes                              | 87%                      | 90%                      |
| Health insurance               | None<br>Public<br>Private        | 36%<br>23%<br>41%        | 37%<br>19%<br>44%        |
| Doctor's visit past year       | Yes                              | 77%                      | 81%                      |

### Research questions (MSM2)

- Characteristics associated with HIV infection
  - Total: 37.5%
- Characteristics associated with unrecognized infection
  - Total: 74.4% of HIV positive
- Racial disparities?



## Characteristics associated with HIV infection (MSM2): Socio-demographic

| Characteristic  |                                                                | Total %<br>(n=448) | HIV+ %<br>(n=168) | Prevalence<br>Ratio |
|-----------------|----------------------------------------------------------------|--------------------|-------------------|---------------------|
| Race/Ethnicity  | Non-Hispanic white                                             | 31%                | 18%               | Ref                 |
|                 | Non-Hispanic Black                                             | 62%                | 45%               | 2.5 *               |
|                 | Other                                                          | 6%                 | 25%               | -                   |
| Age             | < 24                                                           | 24%                | 29%               | Ref                 |
|                 | 25-34                                                          | 26%                | 42%               | -                   |
|                 | 35-44                                                          | 30%                | 40%               | -                   |
|                 | > 45                                                           | 18%                | 43%               | -                   |
| Education       | Graduate education College or some college High school or less | 49%<br>45%<br>6%   | 29%<br>34%<br>42% | Ref.<br>-<br>-      |
| Homelessness    | Current                                                        | 3%                 | 41%               | -                   |
| Sexual identity | Homosexual                                                     | 63%                | 39%               | Ref                 |
|                 | Bisexual                                                       | 32%                | 35%               | -                   |
|                 | Heterosexual or other                                          | 5%                 | 20%               | -                   |

## Characteristics associated with HIV infection (MSM2): Sex and drug

| Characteristic                          |                                                            | Total %<br>(n=448)       | HIV+ %<br>(n=168)        | Prevalence<br>Ratio |
|-----------------------------------------|------------------------------------------------------------|--------------------------|--------------------------|---------------------|
| # of male sex<br>partners, past<br>year | 1<br>2-3<br>4-8<br>9 or more                               | 26%<br>28%<br>23%<br>23% | 34%<br>35%<br>44%<br>39% | Ref<br>-<br>-<br>-  |
| Sex partners in past year               | Men only<br>Men and women                                  | 67%<br>34%               | 40%<br>30%               | Ref<br>-            |
| Unprotected sex in past year            | None With main partners only With casual/exchange partners | 47%<br>25%<br>27%        | 35%<br>33%<br>52%        | Ref<br>-<br>1.5 *   |
| Substance use                           | Ever IDU<br>Non-injection drug use past year               | 17%<br>49%               | 30%<br>41%               | -                   |

## Characteristics associated with HIV infection (MSM2): Health services

| Characteristic                     |                                          | Total %<br>(n=448) | HIV+ %<br>(n=168) | Prevalence<br>Ratio |
|------------------------------------|------------------------------------------|--------------------|-------------------|---------------------|
| Ever tested for HIV                | Yes                                      | 86%                | 38%               | -                   |
| Ever told had STD                  | Yes                                      | 16%                | 51%               | 1.5 *               |
| Health insurance                   | No health insurance<br>Public<br>Private | 36%<br>23%<br>41%  | 40%<br>48%<br>31% | Ref.<br>-<br>-      |
| Visit doctor's office in past year | Yes                                      | 77%                | 39%               | -                   |



## Characteristics associated with HIV infection: Multivariate

| Characteristic               |                                                            | Total %<br>(n=448) | HIV+<br>%         | Prevalence<br>Ratio | Adjusted<br>Prevalence<br>Ratio |
|------------------------------|------------------------------------------------------------|--------------------|-------------------|---------------------|---------------------------------|
| Race/Ethnicity               | Non-Hispanic White<br>Non-Hispanic Black<br>Other          | 31%<br>62%<br>6%   | 18%<br>45%<br>25% | Ref<br>2.5 *<br>1.4 | Ref.<br>2.5 *<br>-              |
| Unprotected sex in past year | None With main partners only With casual/exchange partners | 47%<br>25%<br>27%  | 35%<br>33%<br>52% | Ref<br>0.9<br>1.5 * | Ref<br>-<br>1.4 *               |
| Ever told had STD            | Yes                                                        | 16%                | 51%               | 1.5 *               | 1.5 *                           |

# Unrecognized HIV infection (MSM2): Socio-demographic

| Characteristic  |                      | Total %<br>(n=448) | HIV+ %<br>(n=168) | Unrecog %<br>(n=125) | P.R.  |
|-----------------|----------------------|--------------------|-------------------|----------------------|-------|
| Race/Ethnicity  | Non-Hispanic White   | 31%                | 18%               | 47%                  | Ref   |
|                 | Non-Hispanic Black   | 62%                | 45%               | 77%                  | -     |
|                 | Other                | 6%                 | 25%               | 100%                 | 2.1 * |
| Age             | < 24                 | 24%                | 29%               | 85%                  | Ref   |
|                 | 25-34                | 26%                | 42%               | 75%                  | -     |
|                 | 35-44                | 30%                | 40%               | 79%                  | -     |
|                 | > 45                 | 18%                | 43%               | 50%                  | 0.6 * |
| Education       | Graduate education   | 49%                | 29%               | 50%                  | Ref.  |
|                 | College or some coll | 45%                | 34%               | 77%                  | -     |
|                 | High school or less  | 6%                 | 42%               | 76%                  | -     |
| Homelessness    | Current              | 3%                 | 41%               | 73%                  | -     |
| Sexual identity | Homosexual           | 63%                | 39%               | 72%                  | Ref   |
|                 | Bisexual             | 32%                | 35%               | 81%                  | -     |
|                 | Heterosexual/other   | 5%                 | 20%               | 50%                  | -     |

# Unrecognized HIV infection (MSM2): Sex and drug

| Characteristic                          |                                                            | Total %<br>(n=448)       | HIV+ %<br>(n=168)        | Unrecog %<br>(n=125)     | P.R.               |
|-----------------------------------------|------------------------------------------------------------|--------------------------|--------------------------|--------------------------|--------------------|
| # of male sex<br>partners, past<br>year | 1<br>2-3<br>4-8<br>9 or more                               | 26%<br>28%<br>23%<br>23% | 34%<br>35%<br>44%<br>39% | 76%<br>74%<br>76%<br>70% | Ref<br>-<br>-<br>- |
| Sex partners in past year               | Men only<br>Men and women                                  | 67%<br>34%               | 40%<br>30%               | 71%<br>88%               | Ref<br>1.2 *       |
| Unprotected sex in past year            | None With main partners only With casual/exchange partners | 47%<br>25%<br>27%        | 35%<br>33%<br>52%        | 73%<br>81%<br>74%        | Ref<br>-<br>-      |
| Substance use                           | Ever IDU<br>Non-injection drug use<br>in past year         | 17%<br>49%               | 30%<br>41%               | 74%<br>79%               | -                  |

## Unrecognized HIV infection (MSM2): Health services

| Characteristic                     |                                          | Total %<br>(n=448) | HIV+ %<br>(n=168) | Unrecog %<br>(n=125) | P.R.               |
|------------------------------------|------------------------------------------|--------------------|-------------------|----------------------|--------------------|
| Ever told had STD                  | Yes                                      | 16%                | 51%               | 66%                  | -                  |
| Health insurance                   | No health insurance<br>Public<br>Private | 36%<br>23%<br>41%  | 40%<br>48%<br>31% | 80%<br>73%<br>68%    | Ref.<br>0.9 *<br>- |
| Visit doctor's office in past year | Yes                                      | 77%                | 39%               | 71%                  | 0.8 *              |



## Characteristics associated with unrecognized HIV infection: Multivariate

| Characteristic        |                    | Total %<br>(n=448) | HIV+ % | Unrecog %<br>(n=125) | P.R.  | A.P.R. |
|-----------------------|--------------------|--------------------|--------|----------------------|-------|--------|
| Race/Ethnicity        | Non-Hispanic White | 31%                | 18%    | 47%                  | Ref   | Ref.   |
|                       | Non-Hispanic Black | 62%                | 45%    | 77%                  | 1.6   | -      |
|                       | Other              | 6%                 | 25%    | 100%                 | 2.1 * | -      |
| Age                   | < 24               | 24%                | 29%    | 85%                  | Ref   | Ref    |
|                       | 25-34              | 26%                | 42%    | 75%                  | 0.9   | -      |
|                       | 35-44              | 30%                | 40%    | 79%                  | 0.9   | -      |
|                       | > 45               | 18%                | 43%    | 50%                  | 0.6 * | 0.7 *  |
| Visit doctor's office | Yes                | 77%                | 39%    | 71%                  | 0.8 * | 0.8 *  |



### Summary

- Limitations:
  - Cross-sectional, not causal
  - Voluntary enrollment
  - Self-report
  - Comparison to MSM1 (different characteristics)
- Prevention implications:
  - Drastic need for innovative evidence-based prevention, especially among African-American men
  - Drastic need for targeted testing and link to care, especially among young men
  - Racial disparities not accounted for by behavior

### BESURE next steps

- Round 3: MSM3 2011
- BESURE Blog!



#### With thanks to:

- Study participants
- Community partners
- Frangiscos Sifakis
- Data team: Vivian Towe, Cathy Maulsby,
   Andrea Villante, Trang Nguyen, Beth Linas
- Data collection teams
- David Holtgrave, David Celentano, Carl Latkin
- DHMH, CDC

