

Maryland Department of Health
Center for STI Prevention

EXPEDITED PARTNER THERAPY
RECOMMENDED PARTNER TREATMENT REGIMENS*

Chlamydia Only	<input type="checkbox"/> Doxycycline (Vibramycin) 100 mg orally 2 times/day for 7 days <i>NOTE: Substitute azithromycin (Zithromax) 1 gram orally in a single dose for partners who may be pregnant, or unlikely to adhere to a 7-day regimen</i>
Gonorrhea Only	<input type="checkbox"/> Cefixime (Suprax) 800 mg orally in a single dose
Gonorrhea/Chlamydia Coinfection or Chlamydia Not Ruled Out	<input type="checkbox"/> Cefixime (Suprax) 800 mg orally in a single dose PLUS <input type="checkbox"/> Doxycycline (Vibramycin) 100 mg orally 2 times/day for 7 days <i>NOTE: Substitute azithromycin (Zithromax) 1 gram orally in a single dose for doxycycline for partners who may be pregnant, or unlikely to adhere to a 7-day regimen</i>
Trichomoniasis	<input type="checkbox"/> Metronidazole (Flagyl) 500 mg orally 2 times/day for 7 days (Presumed vaginal infection) <input type="checkbox"/> Metronidazole (Flagyl) 2 g orally in a single dose (Presumed penile infection) <input type="checkbox"/> Tinidazole (Tindamax) 2 g orally in a single dose (Alternative regimen for all non-pregnant partners)

*Based on CDC 2021 STI Treatment Guidelines
<https://www.cdc.gov/std/treatment-guidelines/default.htm>

Last revised February 2023
Document updated when pertinent changes are made.