

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN9: Percentage of students who never or rarely wore a seat belt when riding in a car driven by someone else	14.7 (13.0-16.6)	1,794	16.2 (12.4-20.9)	366			●
QN10: Percentage of students who rode one or more times during the past 30 days in a car or other vehicle driven by someone who had been drinking alcohol	20.9 (18.7-23.2)	1,791	28.6 (23.6-34.3)	363		●	
QN11: Among students who drove a car or other vehicle during the past 30 days, the percentage who drove when they had been drinking alcohol one or more times during the past 30 days	4.5 (2.9-6.9)	633	6.7 (3.3-13.1)	147			●
QN12: Among students who drove a car or other vehicle during the past 30 days, the percentage who texted or e-mailed while driving on one or more of the past 30 days	18.5 (15.0-22.5)	669	29.9 (22.1-39.1)	162		●	
QN13: Percentage of students who carried a weapon such as a gun, knife, or club on one or more of the past 30 days	16.7 (14.7-18.9)	1,755	21.0 (16.9-25.7)	340			●
QN15: Percentage of students who carried a weapon such as a gun, knife, or club on school property on one or more of the past 30 days	7.9 (6.7-9.2)	1,746	9.6 (6.5-14.0)	340			●
QN16: Percentage of students who did not go to school on one or more of the past 30 days because they felt they would be unsafe at school or on their way to or from school	9.5 (7.7-11.6)	1,802	16.4 (12.4-21.5)	373		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QN17: Percentage of students who had been threatened or injured with a weapon such as a gun, knife, or club on school property one or more times during the past 12 months	10.3 (9.0-11.9)	1,800	16.8 (13.0-21.5)	371		●	
QN20: Percentage of students who were in a physical fight on school property one or more times during the past 12 months	22.7 (19.7-26.1)	1,757	28.8 (24.0-34.3)	365		●	
QN21: Percentage of students who had ever been physically forced to have sexual intercourse when they did not want to	9.5 (8.2-11.0)	1,775	20.4 (16.0-25.7)	353		●	
QN22: Among students who dated or went out with someone during the past 12 months, the percentage who had been physically hurt on purpose by someone they were dating or going out with one or more times during the past 12 months	10.0 (8.3-12.0)	1,310	21.3 (17.2-26.0)	280		●	
QN23: Among students who dated or went out with someone during the past 12 months, the percentage who had been forced by someone they were dating or going out with to do sexual things that they did not want to one or more times during the past 12 months	7.0 (5.7-8.6)	1,297	15.2 (11.0-20.5)	277		●	
QN24: Percentage of students who had ever been bullied on school property during the past 12 months	11.9 (10.1-13.9)	1,777	23.1 (17.9-29.1)	349		●	
QN25: Percentage of students who had ever been electronically bullied during the past 12 months	7.4 (6.1-9.0)	1,782	15.3 (11.9-19.3)	342		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey
Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN26: Percentage of students who felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	24.9 (23.0-27.0)	1,737	43.5 (38.3-48.9)	331		●	
QN27: Percentage of students who seriously considered attempting suicide during the past 12 months	12.7 (11.2-14.3)	1,770	31.2 (25.8-37.1)	341		●	
QN28: Percentage of students who made a plan about how they would attempt suicide during the past 12 months	11.0 (9.5-12.7)	1,771	24.1 (19.7-29.0)	343		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey Risk Behaviors and Sexual Identity Report

Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN32: Percentage of students who smoked a whole cigarette for the first time before age 13 years	6.4 (5.2-8.0)	1,786	13.3 (9.7-18.0)	368		●	
QN33: Percentage of students who smoked cigarettes on one or more of the past 30 days	6.8 (5.3-8.6)	1,685	16.6 (12.4-22.0)	317		●	
QNFRCIG: Percentage of students who smoked cigarettes on 20 or more of the past 30 days	1.2 (0.6-2.2)	1,685	2.6 (1.1-6.0)	317			●
QN35: Among students who were less than 18 years of age and who reported current cigarette use, the percentage who usually got their own cigarettes by buying them in a store or gas station during the past 30 days	33.4 (23.5-45.0)	97	30.0 (18.7-44.4)	51			●
QN39: Percentage of students who used chewing tobacco, snuff, or dip on one or more of the past 30 days	4.8 (3.6-6.3)	1,783	15.9 (11.5-21.7)	366		●	
QN40: Percentage of students who smoked cigars, cigarillos, or little cigars on one or more of the past 30 days	11.1 (9.6-12.8)	1,768	31.0 (25.1-37.5)	362		●	
QNANYTOB: Percentage of students who smoked cigarettes or cigars or used chewing tobacco, snuff, or dip on one or more of the past 30 days	13.0 (11.4-14.9)	1,659	31.7 (25.5-38.7)	307		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN41: Percentage of students who had at least one drink of alcohol on one or more days during their life	55.1 (52.5-57.7)	1,681	74.7 (68.6-80.1)	323		●	
QN42: Percentage of students who had their first drink of alcohol other than a few sips before age 13 years	18.0 (16.0-20.2)	1,756	32.4 (27.8-37.5)	361		●	
QN43: Percentage of students who had at least one drink of alcohol on one or more of the past 30 days	23.6 (20.9-26.5)	1,650	39.5 (33.9-45.5)	299		●	
QN44: Percentage of students who had five or more drinks of alcohol in a row, that is, within a couple of hours, on one or more of the past 30 days	10.0 (8.5-11.8)	1,702	20.2 (16.4-24.8)	327		●	
QN47: Percentage of students who used marijuana one or more times during their life	41.1 (38.1-44.1)	1,675	61.7 (55.4-67.7)	310		●	
QN48: Percentage of students who tried marijuana for the first time before age 13 years	12.0 (10.3-13.9)	1,706	25.2 (19.8-31.6)	319		●	
QN49: Percentage of students who used marijuana one or more times during the past 30 days	20.5 (18.1-23.1)	1,726	36.3 (29.9-43.1)	322		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN50: Percentage of students who used any form of cocaine, including powder, crack, or freebase one or more times during their life	4.5 (3.5-5.7)	1,775	17.3 (12.5-23.4)	364		●	
QN51: Percentage of students who sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times during their life	8.0 (6.6-9.5)	1,769	26.2 (20.7-32.6)	362		●	
QN52: Percentage of students who used heroin one or more times during their life	4.7 (3.7-5.8)	1,755	16.8 (12.3-22.5)	357		●	
QN53: Percentage of students who used methamphetamines one or more times during their life	3.8 (3.0-4.9)	1,770	16.7 (11.9-22.8)	363		●	
QN54: Percentage of students who used ecstasy one or more times during their life	5.5 (4.5-6.7)	1,767	20.1 (14.3-27.4)	363		●	
QN55: Percentage of students who took steroid pills or shots without a doctor's prescription one or more times during their life	4.5 (3.5-5.7)	1,781	18.2 (13.3-24.4)	361		●	
QN56: Percentage of students who have taken a prescription drug (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax) without a doctor's prescription one or more times during their life	10.9 (9.4-12.7)	1,786	27.4 (21.4-34.3)	366		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey
Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN57: Percentage of students who used a needle to inject any illegal drug into their body one or more times during their life	3.8 (2.8-5.3)	1,766	16.5 (11.4-23.3)	354		●	
QN58: Percentage of students who were offered, sold, or given an illegal drug by someone on school property during the past 12 months	26.2 (23.6-29.0)	1,774	32.1 (26.3-38.5)	342			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN59: Percentage of students who ever had sexual intercourse	53.5 (49.9-57.1)	1,479	61.8 (54.1-68.9)	252		●	
QN60: Percentage of students who had sexual intercourse for the first time before age 13 years	14.8 (12.7-17.2)	1,482	15.3 (11.4-20.2)	259			●
QN61: Percentage of students who had sexual intercourse with four or more people during their life	22.2 (19.1-25.6)	1,478	19.0 (14.0-25.2)	255			●
QN62: Percentage of students who had sexual intercourse with one or more people during the past three months	36.6 (33.6-39.8)	1,473	44.4 (37.3-51.7)	256		●	
QN63: Among students who had sexual intercourse during the past three months, the percentage who drank alcohol or used drugs before last sexual intercourse	16.4 (12.9-20.6)	502	29.1 (20.7-39.2)	111		●	
QN64: Among students who had sexual intercourse during the past three months, the percentage who did not use a condom during last sexual intercourse	34.7 (30.7-39.0)	501	57.2 (44.8-68.8)	104		●	
QN65: Among students who had sexual intercourse during the past three months, the percentage who did not use birth control pills to prevent pregnancy before last sexual intercourse	89.6 (86.6-92.1)	497	89.8 (81.4-94.7)	103			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNOTHH: Among students who had sexual intercourse during the past three months, the percentage who did not use an IUD or implant or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	91.1 (87.2-93.8)	497	90.1 (80.8-95.2)	103			●
QNOTHHPL: Among students who had sexual intercourse during the past three months, the percentage who did not use birth control pills, an IUD or implant, or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	80.7 (76.8-84.1)	497	79.9 (69.6-87.4)	103			●
QNDUALBC: Among students who had sexual intercourse during the past three months, the percentage who did not use both a condom during last sexual intercourse and birth control pills, an IUD or implant, or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	91.3 (88.1-93.7)	494	94.8 (87.2-98.0)	99			●
QNBCNONE: Among students who had sexual intercourse during the past three months, the percentage who used no method of birth control to prevent pregnancy before last sexual intercourse	14.5 (11.3-18.4)	497	30.9 (22.4-40.8)	103		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QNOWT: Percentage of students who were overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex) [§]	16.7 (14.9-18.6)	1,686	21.5 (16.6-27.3)	330			●
QNOBESE: Percentage of students who were obese (i.e., at or above the 95th percentile for body mass index, by age and sex) [§]	13.3 (11.7-15.2)	1,686	16.3 (13.2-20.0)	330			●
QN66: Percentage of students who described themselves as slightly or very overweight	21.7 (19.6-23.9)	1,789	29.1 (24.0-34.8)	362		●	
QN71: Percentage of students who did not drink 100% fruit juices during the past seven days	23.5 (20.9-26.3)	1,747	28.2 (22.4-34.9)	356			●
QN72: Percentage of students who did not eat fruit during the past seven days	24.0 (21.6-26.6)	1,750	25.1 (19.8-31.3)	360			●
QN73: Percentage of students who did not eat green salad during the past seven days	51.1 (48.3-54.0)	1,734	45.9 (40.9-51.1)	364			●
QN74: Percentage of students who did not eat potatoes during the past seven days	45.1 (42.2-48.0)	1,726	40.4 (35.6-45.4)	359			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

[§]Based on reference data from the 2000 CDC Growth Charts.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

**Baltimore City, MD High School Survey
Risk Behaviors and Sexual Identity Report**

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN75: Percentage of students who did not eat carrots during the past seven days	68.9 (66.2-71.5)	1,731	64.7 (58.8-70.2)	357			●
QN76: Percentage of students who did not eat other vegetables during the past seven days	26.3 (24.0-28.6)	1,716	24.3 (19.2-30.3)	359			●
QNFRVG: Percentage of students who ate fruits or drank 100% fruit juices or ate vegetables less than five times per day during the past seven days	81.1 (79.2-83.0)	1,668	79.8 (74.4-84.3)	341			●
QNFRUIT: Percentage of students who ate fruits or drank 100% fruit juices less than two times per day during the past seven days	72.2 (69.7-74.5)	1,723	72.4 (66.2-77.8)	351			●
QNFR0: Percentage of students who ate fruits or drank 100% fruit juices 0 times per day during the past seven days	11.0 (9.7-12.6)	1,723	9.8 (7.0-13.5)	351			●
QNFR1: Percentage of students who ate fruits or drank 100% fruit juices less than one time per day during the past seven days	48.3 (45.7-50.8)	1,723	51.4 (45.3-57.5)	351			●
QNFR3: Percentage of students who ate fruits or drank 100% fruit juices less than three times per day during the past seven days	80.1 (78.2-82.0)	1,723	80.2 (74.2-85.0)	351			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

**Baltimore City, MD High School Survey
Risk Behaviors and Sexual Identity Report**

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNVEG: Percentage of students who ate vegetables less than three times per day during the past seven days	87.6 (85.7-89.3)	1,693	85.2 (80.3-89.1)	352			●
QNVEG0: Percentage of students who ate vegetables 0 times per day during the past seven days	13.2 (11.0-15.7)	1,693	8.3 (5.2-13.0)	352			●
QNVEG1: Percentage of students who ate vegetables less than one time per day during the past seven days	52.9 (50.1-55.7)	1,693	48.2 (42.5-53.9)	352			●
QNVEG2: Percentage of students who ate vegetables less than two times per day during the past seven days	80.5 (78.4-82.4)	1,693	74.5 (69.5-78.9)	352	●		
QNFRVG2: Percentage of students who ate fruits or drank 100% fruit juices less than two times per day and ate vegetables less than three times per day during the past seven days	91.4 (89.8-92.8)	1,668	91.2 (87.2-94.1)	341			●
QN77: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day during the past seven days	23.6 (21.3-26.0)	1,674	31.4 (26.4-36.9)	341		●	
QNSODA0: Percentage of students who drank a can, bottle, or glass of soda or pop during the past seven days	77.4 (74.7-80.0)	1,674	79.7 (74.2-84.3)	341			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day during the past seven days	17.7 (15.7-19.9)	1,674	22.8 (18.4-27.9)	341			●
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day during the past seven days	11.8 (9.7-14.2)	1,674	14.5 (10.8-19.2)	341			●
QN79: Percentage of students who did not eat breakfast on all of the past seven days	76.8 (74.3-79.2)	1,706	85.7 (81.5-89.2)	357		●	
QNNOBKFT: Percentage of students who ate breakfast on none of the past seven days	22.0 (19.7-24.5)	1,706	24.2 (19.4-29.7)	357			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey Risk Behaviors and Sexual Identity Report

Physical Activity

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN80: Percentage of students who were physically active for a total of at least 60 minutes per day less than five of the past seven days	68.0 (65.3-70.6)	1,687	78.3 (73.6-82.3)	351		●	
QNPA0DAY: Percentage of students who were physically active for a total of at least 60 minutes per day on 0 of the past seven days	26.9 (25.0-29.0)	1,687	26.8 (22.0-32.2)	351			●
QNPA7DAY: Percentage of students who were physically active for a total of at least 60 minutes per day on less than seven of the past seven days	80.8 (78.2-83.1)	1,687	86.9 (82.2-90.5)	351		●	
QN81: Percentage of students who watched three or more hours per day of TV on an average school day	43.7 (40.8-46.5)	1,669	35.6 (29.9-41.7)	352	●		
QN82: Percentage of students who played video or computer games or used a computer for something that was not school work three or more hours per day on an average school day	37.3 (34.9-39.9)	1,685	34.1 (27.6-41.2)	351			●
QN83: Percentage of students who did not attend physical education (PE) classes in an average week when they were in school	69.4 (66.3-72.4)	1,654	62.1 (55.7-68.1)	352	●		
QNDLYPE: Percentage of students who attended physical education (PE) classes less than daily in an average week when they were in school	85.1 (82.0-87.8)	1,654	90.1 (85.7-93.2)	352		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey
Risk Behaviors and Sexual Identity Report

Other

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN85: Percentage of students who were not sure if they had ever been taught or who had not ever been taught in school about AIDS or HIV infection	21.3 (18.7-24.1)	1,632	31.7 (25.9-38.1)	332		●	
QN86: Percentage of students who had ever been told by a doctor or nurse that they had asthma	35.1 (32.4-38.0)	1,624	36.1 (30.7-41.8)	329			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN87: Percentage of students who tried or used tobacco products (for example, cigarettes, cigars, cigarillos, Black and Milds, little cigars, or smokeless tobacco) for the first time during the past 12 months	14.1 (12.4-16.0)	1,653	28.1 (23.1-33.8)	317		●	
QN89: Percentage of students who used some flavored tobacco product during the past 30 days	11.9 (10.1-14.1)	1,662	28.3 (22.5-35.0)	311		●	
QN90: Percentage of students who used only flavored cigars, cigarillos, little cigars, or only flavored smokeless tobacco products, or both flavored cigar products and flavored smokeless tobacco products during the past 30 days	11.5 (9.7-13.6)	1,643	26.6 (21.3-32.7)	294		●	
QN91: Percentage of students who have smoked a 100 or more cigarettes (5 or more packs) during their life	1.6 (0.9-2.8)	1,689	2.0 (0.9-4.2)	346			●
QN92: Among students who bought or tried to buy cigarettes in a store during past 30 days, the percentage who were not asked to show proof of age	66.2 (56.4-74.8)	93	40.3 (28.2-53.7)	57	●		
QN94: Percentage of students who usually smoked menthol cigarettes during the past 30 days	2.2 (1.4-3.3)	1,640	3.6 (2.1-6.2)	318			●
QN95: Percentage of students who usually smoked cigarillos or small cigars, blunts, little cigars that look like brown cigarettes, or some other type of cigar during the past 30 days	9.8 (8.3-11.5)	1,723	22.5 (17.7-28.2)	335		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN96: Percentage of students who live with someone who now smokes cigarettes or cigars	45.9 (42.6-49.3)	1,761	51.5 (45.1-57.8)	333			●
QN97: Percentage of students who describe the rule about smoking inside the home where they live is that it is always allowed inside their home or allowed only at some times or in some places	26.5 (23.6-29.5)	1,756	43.6 (35.9-51.5)	346		●	
QN98: Percentage of students who were in the same room with someone who was smoking cigarettes on one or more of the past 7 days	32.9 (29.9-36.1)	1,792	49.6 (44.0-55.2)	367		●	
QN99: Among students who drank alcohol during the past 30 days and who have a usual type of alcohol that they drink, the percentage who reported liquor, such as vodka, rum, scotch, bourbon, or whisky, as their type of alcohol they usually drank during the past 30 days	44.2 (38.6-50.0)	463	40.1 (31.3-49.7)	154			●
QN100: Percentage of students who took a prescription drug without a doctor's prescription one or more times during the past 30 days	7.7 (6.5-9.2)	1,784	22.8 (16.8-30.3)	364		●	
QN102: Percentage of students who spend less than 20 minutes actually exercising or playing sports during an average physical (PE) class or who do not take PE	69.1 (65.6-72.5)	1,656	78.8 (74.4-82.6)	347		●	
QN103: Percentage of students who described their health in general as good, fair, or poor	48.2 (45.5-50.9)	1,649	55.1 (48.5-61.5)	349		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN104: Percentage of students who have had asthma symptoms during the past 12 months	20.4 (18.2-22.8)	1,627	30.1 (24.0-37.1)	336		●	
QN105: Percentage of students who used an indoor tanning device such as a sunlamp, sunbed, or tanning booth one or more times during the past 12 months	8.3 (7.0-9.7)	1,636	25.3 (19.6-32.0)	349		●	
QN106: Percentage of students who say definitely yes or probably yes that young people who smoke have more friends	57.5 (54.7-60.3)	1,591	63.0 (56.8-68.9)	337			●
QN107: Percentage of students who say definitely yes or probably yes that smoking makes young people look cool or fit in	33.5 (31.2-35.8)	1,614	39.1 (33.4-45.1)	344			●
QN108: Percentage of students who say definitely not or probably not that smokers have shorter lives than non-smokers	15.8 (13.7-18.0)	1,610	23.6 (18.7-29.3)	350		●	
QN109: Percentage of students who were not taught or who were not sure if they were taught in any of their classes about the dangers of tobacco use during the last school year	37.9 (34.2-41.7)	1,589	44.2 (39.1-49.4)	333		●	
QN110: Percentage of students who responded that their parents would feel it would be a little wrong, not at all wrong, or not sure for them to drink beer, wine, or hard liquor (such as vodka, whiskey, or gin) at least once or twice a month	21.1 (18.5-23.8)	1,600	32.1 (27.4-37.2)	346		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Baltimore City, MD High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN111: Percentage of students who think people are at no risk or slight risk of harming themselves (physically or in other ways) if they have one or two drinks of alcohol (beer, wine, or liquor) nearly every day	40.2 (37.1-43.4)	1,596	47.1 (40.1-54.2)	337			●
QN112: Percentage of students who think people are at no risk or slight risk of harming themselves (physically or in other ways) if they have five or more drinks of alcohol (beer, wine, or liquor) once or twice a week	32.7 (29.7-35.8)	1,575	38.7 (32.8-45.0)	331			●
QN113: Percentage of students who do not have an adult outside of school they can talk to about things that are important to them	15.7 (13.2-18.6)	1,554	19.3 (15.1-24.4)	301			●
QN114: Percentage of students who would feel comfortable seeking help from 0 adults besides their parents if they had an important question affecting their life	23.7 (21.6-25.9)	1,592	19.8 (15.8-24.5)	340			●
QN116: Percentage of students who strongly disagree, disagree, or are not sure that their teachers really care about them and give them a lot of encouragement	39.2 (35.8-42.8)	1,597	40.9 (34.2-48.0)	336			●
QN117: Percentage of students who do not participate in any extracurricular activities at school such as sports, band, drama, clubs, or student government	40.6 (37.4-44.0)	1,553	44.8 (38.7-51.1)	309			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.