

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN9: Percentage of students who never or rarely wore a seat belt when riding in a car driven by someone else	13.1 (10.8-15.8)	936	36.2 (24.4-50.0)	92		●	
QN10: Percentage of students who rode one or more times during the past 30 days in a car or other vehicle driven by someone who had been drinking alcohol	22.6 (19.3-26.3)	937	34.1 (24.2-45.7)	90		●	
QN11: Among students who drove a car or other vehicle during the past 30 days, the percentage who drove when they had been drinking alcohol one or more times during the past 30 days	13.5 (10.0-17.8)	507	41.2 (29.1-54.3)	42		●	
QN12: Among students who drove a car or other vehicle during the past 30 days, the percentage who texted or e-mailed while driving on one or more of the past 30 days	42.9 (37.5-48.4)	517	65.8 (53.3-76.3)	47		●	
QN13: Percentage of students who carried a weapon such as a gun, knife, or club on one or more of the past 30 days	20.6 (17.3-24.4)	925	37.9 (25.5-52.1)	89		●	
QN15: Percentage of students who carried a weapon such as a gun, knife, or club on school property on one or more of the past 30 days	4.4 (3.2-6.2)	924	28.8 (15.5-47.3)	90		●	
QN16: Percentage of students who did not go to school on one or more of the past 30 days because they felt they would be unsafe at school or on their way to or from school	8.4 (6.7-10.5)	935	22.1 (15.1-31.3)	94		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QN17: Percentage of students who had been threatened or injured with a weapon such as a gun, knife, or club on school property one or more times during the past 12 months	6.8 (5.3-8.6)	937	29.3 (20.2-40.5)	93		●	
QN20: Percentage of students who were in a physical fight on school property one or more times during the past 12 months	11.2 (9.2-13.7)	927	41.8 (31.5-52.9)	94		●	
QN21: Percentage of students who had ever been physically forced to have sexual intercourse when they did not want to	8.5 (6.7-10.7)	933	31.0 (22.9-40.3)	90		●	
QN22: Among students who dated or went out with someone during the past 12 months, the percentage who had been physically hurt on purpose by someone they were dating or going out with one or more times during the past 12 months	9.0 (6.9-11.5)	696	25.7 (17.2-36.6)	72		●	
QN23: Among students who dated or went out with someone during the past 12 months, the percentage who had been forced by someone they were dating or going out with to do sexual things that they did not want to one or more times during the past 12 months	9.6 (7.2-12.7)	694	23.8 (15.4-34.9)	73		●	
QN24: Percentage of students who had ever been bullied on school property during the past 12 months	19.9 (17.3-22.7)	932	54.0 (45.2-62.6)	87		●	
QN25: Percentage of students who had ever been electronically bullied during the past 12 months	14.9 (12.8-17.3)	934	26.6 (18.9-36.1)	86		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey
Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN26: Percentage of students who felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	25.3 (22.1-28.7)	926	44.4 (34.8-54.5)	89		●	
QN27: Percentage of students who seriously considered attempting suicide during the past 12 months	13.8 (11.4-16.6)	934	33.7 (26.0-42.3)	87		●	
QN28: Percentage of students who made a plan about how they would attempt suicide during the past 12 months	10.9 (9.0-13.1)	932	27.8 (17.3-41.4)	87		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey Risk Behaviors and Sexual Identity Report

Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN32: Percentage of students who smoked a whole cigarette for the first time before age 13 years	12.0 (9.9-14.5)	925	33.2 (22.4-46.3)	95		●	
QN33: Percentage of students who smoked cigarettes on one or more of the past 30 days	18.1 (14.7-22.0)	886	50.4 (39.6-61.2)	86		●	
QNFRDIG: Percentage of students who smoked cigarettes on 20 or more of the past 30 days	9.0 (6.9-11.6)	886	25.9 (17.3-36.8)	86		●	
QN35: Among students who were less than 18 years of age and who reported current cigarette use, the percentage who usually got their own cigarettes by buying them in a store or gas station during the past 30 days	15.4 (10.8-21.6)	126	11.2 (4.3-26.0)	37			●
QN39: Percentage of students who used chewing tobacco, snuff, or dip on one or more of the past 30 days	10.1 (7.8-12.9)	935	33.2 (19.9-49.8)	93		●	
QN40: Percentage of students who smoked cigars, cigarillos, or little cigars on one or more of the past 30 days	10.4 (8.1-13.3)	931	35.3 (23.7-48.8)	93		●	
QNANYTOB: Percentage of students who smoked cigarettes or cigars or used chewing tobacco, snuff, or dip on one or more of the past 30 days	20.8 (17.3-24.8)	879	56.6 (44.4-68.1)	84		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey
Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN41: Percentage of students who had at least one drink of alcohol on one or more days during their life	65.1 (60.7-69.2)	915	77.1 (67.0-84.8)	82		●	
QN42: Percentage of students who had their first drink of alcohol other than a few sips before age 13 years	23.4 (20.5-26.7)	920	43.6 (33.8-54.1)	91		●	
QN43: Percentage of students who had at least one drink of alcohol on one or more of the past 30 days	35.0 (30.5-39.9)	904	51.8 (41.3-62.2)	81		●	
QN44: Percentage of students who had five or more drinks of alcohol in a row, that is, within a couple of hours, on one or more of the past 30 days	22.9 (19.2-27.0)	910	37.1 (28.0-47.2)	82		●	
QN47: Percentage of students who used marijuana one or more times during their life	36.7 (32.3-41.3)	911	59.5 (49.8-68.4)	79		●	
QN48: Percentage of students who tried marijuana for the first time before age 13 years	9.5 (7.4-12.0)	913	25.7 (17.1-36.6)	80		●	
QN49: Percentage of students who used marijuana one or more times during the past 30 days	19.5 (16.1-23.5)	921	34.6 (24.9-45.8)	82		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN50: Percentage of students who used any form of cocaine, including powder, crack, or freebase one or more times during their life	5.8 (4.4-7.8)	933	35.6 (21.9-52.1)	92		●	
QN51: Percentage of students who sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times during their life	7.7 (5.8-10.2)	935	41.9 (30.4-54.3)	91		●	
QN52: Percentage of students who used heroin one or more times during their life	3.9 (2.5-5.8)	931	31.5 (22.3-42.4)	92		●	
QN53: Percentage of students who used methamphetamines one or more times during their life	4.3 (2.9-6.5)	934	26.0 (17.9-36.3)	93		●	
QN54: Percentage of students who used ecstasy one or more times during their life	7.4 (5.4-9.9)	934	34.5 (22.6-48.7)	92		●	
QN55: Percentage of students who took steroid pills or shots without a doctor's prescription one or more times during their life	5.3 (3.7-7.4)	932	38.4 (27.2-51.0)	92		●	
QN56: Percentage of students who have taken a prescription drug (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax) without a doctor's prescription one or more times during their life	13.6 (11.0-16.7)	935	44.3 (32.5-56.8)	94		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey
Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN57: Percentage of students who used a needle to inject any illegal drug into their body one or more times during their life	2.7 (1.7-4.2)	927	25.8 (18.1-35.3)	86		●	
QN58: Percentage of students who were offered, sold, or given an illegal drug by someone on school property during the past 12 months	23.8 (21.0-26.9)	933	37.3 (25.0-51.6)	82		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN59: Percentage of students who ever had sexual intercourse	52.7 (47.0-58.4)	844	62.5 (51.1-72.6)	57			●
QN60: Percentage of students who had sexual intercourse for the first time before age 13 years	6.8 (5.1-9.0)	846	19.3 (11.1-31.4)	59		●	
QN61: Percentage of students who had sexual intercourse with four or more people during their life	15.7 (12.7-19.3)	844	33.8 (24.8-44.2)	58		●	
QN62: Percentage of students who had sexual intercourse with one or more people during the past three months	38.3 (33.6-43.3)	846	48.4 (36.4-60.6)	60			●
QN63: Among students who had sexual intercourse during the past three months, the percentage who drank alcohol or used drugs before last sexual intercourse	24.0 (19.5-29.1)	322	-	27			
QN64: Among students who had sexual intercourse during the past three months, the percentage who did not use a condom during last sexual intercourse	36.3 (30.2-42.8)	320	-	25			
QN65: Among students who had sexual intercourse during the past three months, the percentage who did not use birth control pills to prevent pregnancy before last sexual intercourse	70.8 (64.4-76.6)	314	-	27			

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

- = Fewer than 30 students in this subgroup.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey
Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNOTHH: Among students who had sexual intercourse during the past three months, the percentage who did not use an IUD or implant or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	88.2 (84.0-91.4)	314	-	27			
QNOTHHPL: Among students who had sexual intercourse during the past three months, the percentage who did not use birth control pills, an IUD or implant, or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	59.1 (51.9-65.9)	314	-	27			
QNDUALBC: Among students who had sexual intercourse during the past three months, the percentage who did not use both a condom during last sexual intercourse and birth control pills, an IUD or implant, or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	82.4 (76.4-87.1)	312	-	25			
QNBCNONE: Among students who had sexual intercourse during the past three months, the percentage who used no method of birth control to prevent pregnancy before last sexual intercourse	6.0 (3.9-9.2)	314	-	27			

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

- = Fewer than 30 students in this subgroup.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QNOWT: Percentage of students who were overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex) [§]	16.1 (13.9-18.5)	886	19.3 (12.5-28.7)	78			●
QNOBESE: Percentage of students who were obese (i.e., at or above the 95th percentile for body mass index, by age and sex) [§]	14.3 (11.8-17.3)	886	13.7 (7.9-22.8)	78			●
QN66: Percentage of students who described themselves as slightly or very overweight	32.0 (28.5-35.8)	933	33.6 (23.2-45.9)	94			●
QN71: Percentage of students who did not drink 100% fruit juices during the past seven days	25.3 (22.7-28.1)	927	32.1 (23.4-42.3)	93			●
QN72: Percentage of students who did not eat fruit during the past seven days	18.2 (15.5-21.2)	932	28.9 (17.6-43.5)	93			●
QN73: Percentage of students who did not eat green salad during the past seven days	41.4 (37.8-45.0)	931	38.6 (28.6-49.7)	94			●
QN74: Percentage of students who did not eat potatoes during the past seven days	33.1 (30.2-36.2)	929	30.8 (22.4-40.7)	90			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

[§]Based on reference data from the 2000 CDC Growth Charts.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey
Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN75: Percentage of students who did not eat carrots during the past seven days	55.4 (51.8-59.0)	932	47.5 (36.0-59.4)	90			●
QN76: Percentage of students who did not eat other vegetables during the past seven days	17.6 (14.9-20.8)	932	28.1 (20.5-37.3)	91		●	
QNFRVG: Percentage of students who ate fruits or drank 100% fruit juices or ate vegetables less than five times per day during the past seven days	83.7 (81.1-86.0)	910	78.4 (68.7-85.7)	86			●
QNFRUIT: Percentage of students who ate fruits or drank 100% fruit juices less than two times per day during the past seven days	74.1 (70.7-77.2)	923	75.5 (65.5-83.3)	91			●
QNFR0: Percentage of students who ate fruits or drank 100% fruit juices 0 times per day during the past seven days	7.1 (5.6-9.0)	923	15.1 (7.1-29.2)	91			●
QNFR1: Percentage of students who ate fruits or drank 100% fruit juices less than one time per day during the past seven days	46.8 (43.2-50.6)	923	53.2 (44.0-62.2)	91			●
QNFR3: Percentage of students who ate fruits or drank 100% fruit juices less than three times per day during the past seven days	83.8 (81.3-86.0)	923	85.1 (76.1-91.2)	91			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey
Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNVEG: Percentage of students who ate vegetables less than three times per day during the past seven days	88.0 (85.8-89.8)	921	81.0 (71.4-87.9)	88			●
QNVEG0: Percentage of students who ate vegetables 0 times per day during the past seven days	7.0 (5.6-8.8)	921	12.5 (7.2-20.9)	88			●
QNVEG1: Percentage of students who ate vegetables less than one time per day during the past seven days	38.1 (35.1-41.3)	921	39.6 (28.9-51.3)	88			●
QNVEG2: Percentage of students who ate vegetables less than two times per day during the past seven days	76.2 (73.8-78.4)	921	67.7 (57.6-76.4)	88			●
QNFRVG2: Percentage of students who ate fruits or drank 100% fruit juices less than two times per day and ate vegetables less than three times per day during the past seven days	91.7 (89.9-93.3)	910	91.6 (83.6-95.9)	86			●
QN77: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day during the past seven days	20.4 (17.8-23.2)	913	31.3 (22.6-41.6)	90		●	
QNSODA0: Percentage of students who drank a can, bottle, or glass of soda or pop during the past seven days	73.8 (69.9-77.4)	913	73.0 (63.4-80.9)	90			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey
Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day during the past seven days	14.3 (11.7-17.4)	913	18.8 (12.0-28.2)	90			●
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day during the past seven days	8.6 (6.7-11.0)	913	16.7 (10.3-25.8)	90			●
QN79: Percentage of students who did not eat breakfast on all of the past seven days	64.9 (61.7-68.0)	928	77.1 (67.1-84.8)	93		●	
QNNOBKFT: Percentage of students who ate breakfast on none of the past seven days	15.8 (13.3-18.8)	928	28.6 (20.3-38.8)	93		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey
Risk Behaviors and Sexual Identity Report

Physical Activity

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN80: Percentage of students who were physically active for a total of at least 60 minutes per day less than five of the past seven days	56.6 (52.6-60.6)	924	77.7 (67.5-85.4)	93		●	
QNPA0DAY: Percentage of students who were physically active for a total of at least 60 minutes per day on 0 of the past seven days	19.0 (16.2-22.0)	924	32.9 (24.1-43.1)	93		●	
QNPA7DAY: Percentage of students who were physically active for a total of at least 60 minutes per day on less than seven of the past seven days	74.5 (71.2-77.6)	924	88.0 (80.1-93.1)	93		●	
QN81: Percentage of students who watched three or more hours per day of TV on an average school day	31.4 (28.5-34.4)	924	28.5 (19.4-39.8)	90			●
QN82: Percentage of students who played video or computer games or used a computer for something that was not school work three or more hours per day on an average school day	34.0 (30.6-37.6)	923	41.5 (29.4-54.7)	93			●
QN83: Percentage of students who did not attend physical education (PE) classes in an average week when they were in school	67.1 (61.8-72.0)	918	56.4 (45.9-66.3)	92	●		
QNDLYPE: Percentage of students who attended physical education (PE) classes less than daily in an average week when they were in school	73.8 (68.9-78.2)	918	84.0 (74.8-90.3)	92		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey
Risk Behaviors and Sexual Identity Report

Other

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN85: Percentage of students who were not sure if they had ever been taught or who had not ever been taught in school about AIDS or HIV infection	12.0 (9.4-15.1)	921	23.8 (15.7-34.5)	83		●	
QN86: Percentage of students who had ever been told by a doctor or nurse that they had asthma	28.4 (25.3-31.7)	920	35.0 (24.3-47.4)	83			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QN87: Percentage of students who tried or used tobacco products (for example, cigarettes, cigars, cigarillos, Black and Milds, little cigars, or smokeless tobacco) for the first time during the past 12 months	21.0 (17.8-24.7)	879	43.9 (32.9-55.4)	87		●	
QN89: Percentage of students who used some flavored tobacco product during the past 30 days	12.6 (9.9-15.8)	880	39.7 (28.0-52.7)	81		●	
QN90: Percentage of students who used only flavored cigars, cigarillos, little cigars, or only flavored smokeless tobacco products, or both flavored cigar products and flavored smokeless tobacco products during the past 30 days	12.5 (9.9-15.7)	879	38.5 (27.4-50.9)	80		●	
QN91: Percentage of students who have smoked a 100 or more cigarettes (5 or more packs) during their life	11.6 (9.3-14.3)	918	25.1 (18.1-33.6)	94		●	
QN92: Among students who bought or tried to buy cigarettes in a store during past 30 days, the percentage who were not asked to show proof of age	49.6 (39.4-59.9)	82	-	22			
QN94: Percentage of students who usually smoked menthol cigarettes during the past 30 days	12.4 (9.5-16.0)	899	27.1 (17.8-39.1)	84		●	
QN95: Percentage of students who usually smoked cigarillos or small cigars, blunts, little cigars that look like brown cigarettes, or some other type of cigar during the past 30 days	8.8 (6.8-11.4)	907	23.8 (15.3-35.1)	85		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

- = Fewer than 30 students in this subgroup.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN96: Percentage of students who live with someone who now smokes cigarettes or cigars	49.3 (45.6-53.0)	921	60.9 (48.9-71.7)	85		●	
QN97: Percentage of students who describe the rule about smoking inside the home where they live is that it is always allowed inside their home or allowed only at some times or in some places	23.9 (21.3-26.7)	922	42.3 (32.7-52.6)	85		●	
QN98: Percentage of students who were in the same room with someone who was smoking cigarettes on one or more of the past 7 days	45.9 (42.3-49.5)	935	61.7 (52.9-69.8)	95		●	
QN99: Among students who drank alcohol during the past 30 days and who have a usual type of alcohol that they drink, the percentage who reported liquor, such as vodka, rum, scotch, bourbon, or whisky, as their type of alcohol they usually drank during the past 30 days	43.3 (37.4-49.5)	310	48.5 (31.7-65.6)	43			●
QN100: Percentage of students who took a prescription drug without a doctor's prescription one or more times during the past 30 days	7.7 (5.6-10.4)	935	47.5 (35.4-60.0)	92		●	
QN102: Percentage of students who spend less than 20 minutes actually exercising or playing sports during an average physical (PE) class or who do not take PE	65.4 (60.6-70.0)	924	67.2 (56.5-76.4)	90			●
QN103: Percentage of students who described their health in general as good, fair, or poor	47.8 (43.7-51.8)	926	52.6 (42.0-63.0)	90			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN104: Percentage of students who have had asthma symptoms during the past 12 months	17.5 (15.0-20.4)	921	31.8 (24.1-40.6)	83		●	
QN105: Percentage of students who used an indoor tanning device such as a sunlamp, sunbed, or tanning booth one or more times during the past 12 months	12.8 (10.3-15.7)	925	32.2 (20.3-46.9)	94		●	
QN106: Percentage of students who say definitely yes or probably yes that young people who smoke have more friends	33.0 (29.7-36.5)	919	48.4 (38.9-58.1)	88		●	
QN107: Percentage of students who say definitely yes or probably yes that smoking makes young people look cool or fit in	16.5 (13.7-19.8)	920	33.0 (21.3-47.4)	90		●	
QN108: Percentage of students who say definitely not or probably not that smokers have shorter lives than non-smokers	8.4 (7.0-10.0)	923	20.5 (13.3-30.3)	87		●	
QN109: Percentage of students who were not taught or who were not sure if they were taught in any of their classes about the dangers of tobacco use during the last school year	22.5 (19.5-25.8)	912	34.5 (22.0-49.5)	82			●
QN110: Percentage of students who responded that their parents would feel it would be a little wrong, not at all wrong, or not sure for them to drink beer, wine, or hard liquor (such as vodka, whiskey, or gin) at least once or twice a month	31.9 (28.4-35.7)	921	39.2 (28.0-51.6)	88			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Caroline County, MD High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN111: Percentage of students who think people are at no risk or slight risk of harming themselves (physically or in other ways) if they have one or two drinks of alcohol (beer, wine, or liquor) nearly every day	39.0 (35.0-43.2)	910	45.6 (33.2-58.6)	87			●
QN112: Percentage of students who think people are at no risk or slight risk of harming themselves (physically or in other ways) if they have five or more drinks of alcohol (beer, wine, or liquor) once or twice a week	26.2 (22.7-30.1)	913	44.7 (33.7-56.2)	86		●	
QN113: Percentage of students who do not have an adult outside of school they can talk to about things that are important to them	14.0 (11.6-16.8)	902	38.0 (27.5-49.7)	83		●	
QN114: Percentage of students who would feel comfortable seeking help from 0 adults besides their parents if they had an important question affecting their life	19.6 (17.1-22.4)	920	26.7 (17.8-38.0)	89			●
QN116: Percentage of students who strongly disagree, disagree, or are not sure that their teachers really care about them and give them a lot of encouragement	42.7 (39.1-46.4)	918	52.4 (41.5-63.2)	88			●
QN117: Percentage of students who do not participate in any extracurricular activities at school such as sports, band, drama, clubs, or student government	33.3 (28.9-38.0)	908	50.1 (39.6-60.6)	77		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.