

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Charles County, MD High School Survey Risk Behaviors and Sexual Identity Report

#### Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN9: Percentage of students who never or rarely wore a seat belt when riding in a car driven by someone else	8.4 (7.2-9.8)	2,438	16.1 (12.1-21.0)	292		●	
QN10: Percentage of students who rode one or more times during the past 30 days in a car or other vehicle driven by someone who had been drinking alcohol	19.1 (17.3-21.0)	2,426	29.7 (23.8-36.3)	288		●	
QN11: Among students who drove a car or other vehicle during the past 30 days, the percentage who drove when they had been drinking alcohol one or more times during the past 30 days	7.8 (6.3-9.7)	1,069	14.9 (9.4-22.7)	121			●
QN12: Among students who drove a car or other vehicle during the past 30 days, the percentage who texted or e-mailed while driving on one or more of the past 30 days	32.1 (28.8-35.6)	1,100	40.5 (32.0-49.5)	129			●
QN13: Percentage of students who carried a weapon such as a gun, knife, or club on one or more of the past 30 days	16.9 (15.2-18.7)	2,414	24.1 (19.0-30.1)	275		●	
QN15: Percentage of students who carried a weapon such as a gun, knife, or club on school property on one or more of the past 30 days	3.6 (2.8-4.6)	2,397	11.3 (7.6-16.3)	277		●	
QN16: Percentage of students who did not go to school on one or more of the past 30 days because they felt they would be unsafe at school or on their way to or from school	6.8 (5.7-8.1)	2,441	14.6 (10.9-19.3)	294		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Charles County, MD High School Survey Risk Behaviors and Sexual Identity Report

#### Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>‡</sup>
	% 95% CI*	N	% 95% CI	N			
QN17: Percentage of students who had been threatened or injured with a weapon such as a gun, knife, or club on school property one or more times during the past 12 months	8.1 (7.0-9.4)	2,441	19.0 (14.5-24.4)	292		●	
QN20: Percentage of students who were in a physical fight on school property one or more times during the past 12 months	12.9 (11.4-14.6)	2,390	23.8 (18.7-29.8)	289		●	
QN21: Percentage of students who had ever been physically forced to have sexual intercourse when they did not want to	8.6 (7.4-10.1)	2,421	24.4 (19.6-29.9)	284		●	
QN22: Among students who dated or went out with someone during the past 12 months, the percentage who had been physically hurt on purpose by someone they were dating or going out with one or more times during the past 12 months	9.2 (7.7-11.0)	1,669	27.5 (21.6-34.3)	230		●	
QN23: Among students who dated or went out with someone during the past 12 months, the percentage who had been forced by someone they were dating or going out with to do sexual things that they did not want to one or more times during the past 12 months	8.4 (7.1-9.9)	1,657	24.8 (18.9-31.9)	232		●	
QN24: Percentage of students who had ever been bullied on school property during the past 12 months	17.8 (16.2-19.5)	2,431	31.5 (26.0-37.6)	286		●	
QN25: Percentage of students who had ever been electronically bullied during the past 12 months	12.2 (10.8-13.8)	2,428	22.9 (18.5-27.9)	285		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Charles County, MD High School Survey  
Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN26: Percentage of students who felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	24.7 (23.0-26.4)	2,392	44.7 (38.5-51.1)	279		●	
QN27: Percentage of students who seriously considered attempting suicide during the past 12 months	13.2 (11.8-14.6)	2,418	35.2 (29.2-41.7)	288		●	
QN28: Percentage of students who made a plan about how they would attempt suicide during the past 12 months	9.6 (8.4-11.0)	2,425	28.2 (23.0-34.1)	284		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Charles County, MD High School Survey  
Risk Behaviors and Sexual Identity Report

Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN32: Percentage of students who smoked a whole cigarette for the first time before age 13 years	6.1 (5.0-7.3)	2,416	17.8 (13.6-22.9)	292		●	
QN33: Percentage of students who smoked cigarettes on one or more of the past 30 days	10.0 (8.6-11.7)	2,291	29.1 (23.4-35.6)	261		●	
QNFRCIG: Percentage of students who smoked cigarettes on 20 or more of the past 30 days	3.0 (2.3-4.0)	2,291	10.1 (7.0-14.4)	261		●	
QN35: Among students who were less than 18 years of age and who reported current cigarette use, the percentage who usually got their own cigarettes by buying them in a store or gas station during the past 30 days	17.9 (12.2-25.4)	178	19.6 (12.0-30.4)	68			●
QN39: Percentage of students who used chewing tobacco, snuff, or dip on one or more of the past 30 days	5.1 (4.1-6.3)	2,427	14.1 (10.3-19.0)	292		●	
QN40: Percentage of students who smoked cigars, cigarillos, or little cigars on one or more of the past 30 days	11.0 (9.6-12.6)	2,417	30.3 (25.0-36.3)	289		●	
QNANYTOB: Percentage of students who smoked cigarettes or cigars or used chewing tobacco, snuff, or dip on one or more of the past 30 days	14.5 (12.8-16.3)	2,265	40.0 (33.7-46.7)	258		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Charles County, MD High School Survey  
Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN41: Percentage of students who had at least one drink of alcohol on one or more days during their life	59.1 (56.3-61.9)	2,348	81.3 (75.1-86.1)	263		●	
QN42: Percentage of students who had their first drink of alcohol other than a few sips before age 13 years	18.8 (17.0-20.7)	2,390	38.7 (33.1-44.6)	290		●	
QN43: Percentage of students who had at least one drink of alcohol on one or more of the past 30 days	28.6 (26.4-30.9)	2,311	53.4 (46.5-60.1)	257		●	
QN44: Percentage of students who had five or more drinks of alcohol in a row, that is, within a couple of hours, on one or more of the past 30 days	15.0 (13.2-16.9)	2,335	30.7 (25.0-37.1)	270		●	
QN47: Percentage of students who used marijuana one or more times during their life	33.6 (30.9-36.3)	2,357	57.9 (50.8-64.7)	261		●	
QN48: Percentage of students who tried marijuana for the first time before age 13 years	7.2 (6.2-8.5)	2,372	20.7 (16.2-26.0)	265		●	
QN49: Percentage of students who used marijuana one or more times during the past 30 days	17.8 (15.8-19.9)	2,374	40.0 (33.8-46.4)	267		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Charles County, MD High School Survey  
Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN50: Percentage of students who used any form of cocaine, including powder, crack, or freebase one or more times during their life	4.3 (3.5-5.2)	2,422	17.4 (13.1-22.7)	291		●	
QN51: Percentage of students who sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times during their life	7.8 (6.7-9.0)	2,415	28.4 (23.1-34.4)	292		●	
QN52: Percentage of students who used heroin one or more times during their life	2.2 (1.6-2.9)	2,397	16.6 (12.0-22.4)	289		●	
QN53: Percentage of students who used methamphetamines one or more times during their life	2.6 (2.0-3.4)	2,425	15.5 (11.9-20.0)	292		●	
QN54: Percentage of students who used ecstasy one or more times during their life	5.9 (4.9-7.1)	2,415	28.6 (23.0-34.8)	291		●	
QN55: Percentage of students who took steroid pills or shots without a doctor's prescription one or more times during their life	2.9 (2.3-3.8)	2,415	17.2 (12.8-22.8)	286		●	
QN56: Percentage of students who have taken a prescription drug (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax) without a doctor's prescription one or more times during their life	11.8 (10.4-13.4)	2,427	32.7 (27.4-38.6)	292		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Charles County, MD High School Survey  
Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN57: Percentage of students who used a needle to inject any illegal drug into their body one or more times during their life	2.3 (1.7-3.0)	2,412	10.8 (7.5-15.2)	279		●	
QN58: Percentage of students who were offered, sold, or given an illegal drug by someone on school property during the past 12 months	26.8 (24.8-28.9)	2,420	39.5 (33.3-46.0)	279		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Charles County, MD High School Survey Risk Behaviors and Sexual Identity Report

#### Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN59: Percentage of students who ever had sexual intercourse	43.6 (40.3-46.9)	2,161	65.2 (58.0-71.8)	222		●	
QN60: Percentage of students who had sexual intercourse for the first time before age 13 years	6.0 (4.9-7.3)	2,163	13.7 (9.7-18.9)	229		●	
QN61: Percentage of students who had sexual intercourse with four or more people during their life	12.8 (11.2-14.6)	2,159	27.3 (21.5-34.1)	228		●	
QN62: Percentage of students who had sexual intercourse with one or more people during the past three months	28.3 (25.4-31.5)	2,164	46.2 (39.5-53.1)	227		●	
QN63: Among students who had sexual intercourse during the past three months, the percentage who drank alcohol or used drugs before last sexual intercourse	18.9 (15.8-22.6)	575	36.6 (27.4-46.9)	102		●	
QN64: Among students who had sexual intercourse during the past three months, the percentage who did not use a condom during last sexual intercourse	30.9 (27.3-34.7)	566	55.2 (44.7-65.2)	99		●	
QN65: Among students who had sexual intercourse during the past three months, the percentage who did not use birth control pills to prevent pregnancy before last sexual intercourse	85.1 (81.6-88.1)	561	84.6 (76.0-90.6)	101			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.



2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Charles County, MD High School Survey  
Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QNOTHH: Among students who had sexual intercourse during the past three months, the percentage who did not use an IUD or implant or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	96.0 (93.9-97.4)	561	97.2 (91.0-99.2)	101			●
QNOTHHPL: Among students who had sexual intercourse during the past three months, the percentage who did not use birth control pills, an IUD or implant, or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	81.2 (77.4-84.4)	561	81.8 (72.6-88.5)	101			●
QNDUALBC: Among students who had sexual intercourse during the past three months, the percentage who did not use both a condom during last sexual intercourse and birth control pills, an IUD or implant, or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	91.4 (88.5-93.7)	553	92.8 (84.7-96.8)	98			●
QNBCNONE: Among students who had sexual intercourse during the past three months, the percentage who used no method of birth control to prevent pregnancy before last sexual intercourse	12.6 (9.4-16.7)	561	31.4 (22.8-41.5)	101		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Charles County, MD High School Survey Risk Behaviors and Sexual Identity Report

#### Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>‡</sup>
	% 95% CI*	N	% 95% CI	N			
QNOWT: Percentage of students who were overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex) <sup>§</sup>	16.9 (15.2-18.6)	2,313	20.2 (14.9-26.6)	264			●
QNOBESE: Percentage of students who were obese (i.e., at or above the 95th percentile for body mass index, by age and sex) <sup>§</sup>	11.5 (10.0-13.2)	2,313	14.3 (10.3-19.5)	264			●
QN66: Percentage of students who described themselves as slightly or very overweight	27.1 (25.2-29.2)	2,420	33.4 (27.7-39.8)	290		●	
QN71: Percentage of students who did not drink 100% fruit juices during the past seven days	22.4 (20.4-24.6)	2,381	19.5 (15.2-24.7)	286			●
QN72: Percentage of students who did not eat fruit during the past seven days	19.9 (18.1-22.0)	2,381	26.6 (21.5-32.4)	286		●	
QN73: Percentage of students who did not eat green salad during the past seven days	43.8 (41.5-46.2)	2,373	42.7 (37.1-48.4)	285			●
QN74: Percentage of students who did not eat potatoes during the past seven days	36.3 (34.4-38.2)	2,358	40.2 (34.4-46.3)	279			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

<sup>§</sup>Based on reference data from the 2000 CDC Growth Charts.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Charles County, MD High School Survey Risk Behaviors and Sexual Identity Report

#### Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>‡</sup>
	% 95% CI*	N	% 95% CI	N			
QN75: Percentage of students who did not eat carrots during the past seven days	60.3 (58.1-62.4)	2,368	61.4 (55.4-67.0)	281			●
QN76: Percentage of students who did not eat other vegetables during the past seven days	17.4 (15.5-19.4)	2,362	20.6 (15.8-26.3)	280			●
QNFRVG: Percentage of students who ate fruits or drank 100% fruit juices or ate vegetables less than five times per day during the past seven days	82.0 (80.3-83.6)	2,308	78.4 (72.9-83.1)	271			●
QNFRUIT: Percentage of students who ate fruits or drank 100% fruit juices less than two times per day during the past seven days	71.8 (69.9-73.7)	2,356	68.6 (61.9-74.5)	282			●
QNFR0: Percentage of students who ate fruits or drank 100% fruit juices 0 times per day during the past seven days	7.9 (6.6-9.6)	2,356	9.3 (6.5-13.3)	282			●
QNFR1: Percentage of students who ate fruits or drank 100% fruit juices less than one time per day during the past seven days	45.5 (43.2-47.8)	2,356	45.2 (38.4-52.1)	282			●
QNFR3: Percentage of students who ate fruits or drank 100% fruit juices less than three times per day during the past seven days	80.4 (78.6-82.1)	2,356	79.2 (73.4-83.9)	282			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Charles County, MD High School Survey Risk Behaviors and Sexual Identity Report

#### Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QNVEG: Percentage of students who ate vegetables less than three times per day during the past seven days	89.9 (88.3-91.2)	2,334	81.5 (76.1-85.9)	274	●		
QNVEG0: Percentage of students who ate vegetables 0 times per day during the past seven days	7.3 (6.2-8.7)	2,334	9.4 (6.4-13.7)	274			●
QNVEG1: Percentage of students who ate vegetables less than one time per day during the past seven days	42.0 (39.7-44.3)	2,334	37.8 (31.8-44.1)	274			●
QNVEG2: Percentage of students who ate vegetables less than two times per day during the past seven days	77.7 (75.7-79.7)	2,334	72.3 (66.7-77.3)	274			●
QNFRVG2: Percentage of students who ate fruits or drank 100% fruit juices less than two times per day and ate vegetables less than three times per day during the past seven days	92.7 (91.4-93.9)	2,308	89.6 (84.9-93.0)	271			●
QN77: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day during the past seven days	17.3 (15.3-19.5)	2,301	25.5 (20.4-31.4)	272		●	
QNSODA0: Percentage of students who drank a can, bottle, or glass of soda or pop during the past seven days	70.9 (68.8-72.8)	2,301	75.6 (69.7-80.8)	272			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Charles County, MD High School Survey  
Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>‡</sup>
	% 95% CI*	N	% 95% CI	N			
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day during the past seven days	11.9 (10.3-13.8)	2,301	17.9 (13.6-23.2)	272		●	
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day during the past seven days	6.8 (5.7-8.2)	2,301	11.6 (8.3-16.1)	272		●	
QN79: Percentage of students who did not eat breakfast on all of the past seven days	70.0 (67.8-72.1)	2,340	82.0 (76.8-86.3)	279		●	
QNNOBKFT: Percentage of students who ate breakfast on none of the past seven days	14.6 (12.9-16.4)	2,340	24.6 (19.6-30.4)	279		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Charles County, MD High School Survey  
Risk Behaviors and Sexual Identity Report

Physical Activity

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>‡</sup>
	% 95% CI*	N	% 95% CI	N			
QN80: Percentage of students who were physically active for a total of at least 60 minutes per day less than five of the past seven days	61.3 (59.1-63.5)	2,337	78.7 (73.3-83.3)	277		●	
QNPA0DAY: Percentage of students who were physically active for a total of at least 60 minutes per day on 0 of the past seven days	19.6 (17.7-21.6)	2,337	29.3 (23.6-35.7)	277		●	
QNPA7DAY: Percentage of students who were physically active for a total of at least 60 minutes per day on less than seven of the past seven days	77.8 (75.9-79.6)	2,337	88.9 (84.3-92.2)	277		●	
QN81: Percentage of students who watched three or more hours per day of TV on an average school day	42.9 (40.2-45.6)	2,301	35.2 (29.5-41.4)	274	●		
QN82: Percentage of students who played video or computer games or used a computer for something that was not school work three or more hours per day on an average school day	38.8 (36.6-41.0)	2,330	43.0 (36.9-49.4)	276			●
QN83: Percentage of students who did not attend physical education (PE) classes in an average week when they were in school	60.4 (57.1-63.7)	2,303	62.8 (55.8-69.3)	266			●
QNDLYPE: Percentage of students who attended physical education (PE) classes less than daily in an average week when they were in school	77.8 (75.0-80.3)	2,303	87.1 (81.7-91.0)	266		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Charles County, MD High School Survey  
Risk Behaviors and Sexual Identity Report

Other

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN85: Percentage of students who were not sure if they had ever been taught or who had not ever been taught in school about AIDS or HIV infection	15.2 (13.7-17.0)	2,295	20.1 (14.8-26.8)	263			●
QN86: Percentage of students who had ever been told by a doctor or nurse that they had asthma	24.6 (22.8-26.5)	2,289	31.6 (25.8-38.0)	266		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Charles County, MD High School Survey Risk Behaviors and Sexual Identity Report

#### Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN87: Percentage of students who tried or used tobacco products (for example, cigarettes, cigars, cigarillos, Black and Milds, little cigars, or smokeless tobacco) for the first time during the past 12 months	17.4 (15.7-19.2)	2,326	34.4 (28.1-41.3)	261		●	
QN89: Percentage of students who used some flavored tobacco product during the past 30 days	12.5 (10.9-14.4)	2,288	30.9 (24.8-37.8)	254		●	
QN90: Percentage of students who used only flavored cigars, cigarillos, little cigars, or only flavored smokeless tobacco products, or both flavored cigar products and flavored smokeless tobacco products during the past 30 days	12.7 (11.0-14.5)	2,273	31.8 (25.5-38.8)	253		●	
QN91: Percentage of students who have smoked a 100 or more cigarettes (5 or more packs) during their life	4.7 (3.8-5.8)	2,375	8.4 (5.8-12.1)	282		●	
QN92: Among students who bought or tried to buy cigarettes in a store during past 30 days, the percentage who were not asked to show proof of age	48.1 (39.2-57.1)	134	56.5 (39.8-71.9)	47			●
QN94: Percentage of students who usually smoked menthol cigarettes during the past 30 days	4.8 (3.9-5.9)	2,333	10.5 (7.2-15.2)	270		●	
QN95: Percentage of students who usually smoked cigarillos or small cigars, blunts, little cigars that look like brown cigarettes, or some other type of cigar during the past 30 days	9.9 (8.7-11.4)	2,371	27.3 (22.1-33.3)	275		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.



## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Charles County, MD High School Survey Risk Behaviors and Sexual Identity Report

#### Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN96: Percentage of students who live with someone who now smokes cigarettes or cigars	38.5 (36.3-40.8)	2,411	46.2 (39.7-52.8)	271		●	
QN97: Percentage of students who describe the rule about smoking inside the home where they live is that it is always allowed inside their home or allowed only at some times or in some places	16.7 (15.1-18.4)	2,406	33.8 (28.4-39.6)	282		●	
QN98: Percentage of students who were in the same room with someone who was smoking cigarettes on one or more of the past 7 days	30.0 (28.1-31.9)	2,430	42.9 (37.0-49.1)	294		●	
QN99: Among students who drank alcohol during the past 30 days and who have a usual type of alcohol that they drink, the percentage who reported liquor, such as vodka, rum, scotch, bourbon, or whisky, as their type of alcohol they usually drank during the past 30 days	56.2 (52.3-60.0)	670	58.0 (49.7-66.0)	140			●
QN100: Percentage of students who took a prescription drug without a doctor's prescription one or more times during the past 30 days	6.6 (5.6-7.8)	2,419	26.9 (22.1-32.4)	292		●	
QN102: Percentage of students who spend less than 20 minutes actually exercising or playing sports during an average physical (PE) class or who do not take PE	62.5 (59.3-65.5)	2,283	76.2 (70.2-81.3)	273		●	
QN103: Percentage of students who described their health in general as good, fair, or poor	50.1 (47.7-52.5)	2,308	63.4 (56.9-69.5)	275		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Charles County, MD High School Survey Risk Behaviors and Sexual Identity Report

#### Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN104: Percentage of students who have had asthma symptoms during the past 12 months	15.8 (14.3-17.5)	2,287	26.7 (21.0-33.2)	266		●	
QN105: Percentage of students who used an indoor tanning device such as a sunlamp, sunbed, or tanning booth one or more times during the past 12 months	6.6 (5.4-7.9)	2,291	20.6 (15.6-26.7)	273		●	
QN106: Percentage of students who say definitely yes or probably yes that young people who smoke have more friends	41.7 (39.3-44.3)	2,251	56.6 (50.2-62.7)	262		●	
QN107: Percentage of students who say definitely yes or probably yes that smoking makes young people look cool or fit in	23.8 (21.6-26.2)	2,261	37.9 (31.3-45.0)	261		●	
QN108: Percentage of students who say definitely not or probably not that smokers have shorter lives than non-smokers	12.1 (10.4-14.0)	2,266	22.5 (17.8-28.0)	266		●	
QN109: Percentage of students who were not taught or who were not sure if they were taught in any of their classes about the dangers of tobacco use during the last school year	29.1 (26.1-32.2)	2,241	40.6 (34.6-47.0)	260		●	
QN110: Percentage of students who responded that their parents would feel it would be a little wrong, not at all wrong, or not sure for them to drink beer, wine, or hard liquor (such as vodka, whiskey, or gin) at least once or twice a month	28.1 (25.9-30.3)	2,254	41.6 (35.3-48.2)	269		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Charles County, MD High School Survey Risk Behaviors and Sexual Identity Report

#### Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual <sup>†</sup>	Gay, Lesbian, or Bisexual More Likely Than Heterosexual <sup>†</sup>	No Difference <sup>‡</sup>
	% 95% CI*	N	% 95% CI	N			
QN111: Percentage of students who think people are at no risk or slight risk of harming themselves (physically or in other ways) if they have one or two drinks of alcohol (beer, wine, or liquor) nearly every day	38.0 (35.5-40.6)	2,232	43.7 (37.6-49.9)	266			●
QN112: Percentage of students who think people are at no risk or slight risk of harming themselves (physically or in other ways) if they have five or more drinks of alcohol (beer, wine, or liquor) once or twice a week	26.5 (23.8-29.3)	2,218	31.4 (25.6-37.9)	261			●
QN113: Percentage of students who do not have an adult outside of school they can talk to about things that are important to them	13.7 (12.2-15.2)	2,196	30.9 (25.6-36.9)	256		●	
QN114: Percentage of students who would feel comfortable seeking help from 0 adults besides their parents if they had an important question affecting their life	20.9 (19.2-22.6)	2,230	30.4 (24.9-36.5)	269		●	
QN116: Percentage of students who strongly disagree, disagree, or are not sure that their teachers really care about them and give them a lot of encouragement	50.5 (48.0-53.1)	2,232	54.0 (47.4-60.4)	267			●
QN117: Percentage of students who do not participate in any extracurricular activities at school such as sports, band, drama, clubs, or student government	30.8 (28.0-33.8)	2,198	42.6 (35.8-49.6)	253		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.