

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN9: Percentage of students who never or rarely wore a seat belt when riding in a car driven by someone else	6.2 (4.9-7.7)	1,814	14.1 (9.1-21.2)	175		●	
QN10: Percentage of students who rode one or more times during the past 30 days in a car or other vehicle driven by someone who had been drinking alcohol	17.2 (15.4-19.3)	1,809	25.8 (18.9-34.2)	173		●	
QN11: Among students who drove a car or other vehicle during the past 30 days, the percentage who drove when they had been drinking alcohol one or more times during the past 30 days	6.7 (5.0-8.8)	861	12.9 (6.4-24.4)	74			●
QN12: Among students who drove a car or other vehicle during the past 30 days, the percentage who texted or e-mailed while driving on one or more of the past 30 days	31.0 (26.9-35.4)	878	35.5 (24.6-48.2)	79			●
QN13: Percentage of students who carried a weapon such as a gun, knife, or club on one or more of the past 30 days	18.5 (16.2-21.0)	1,804	21.6 (15.5-29.3)	170			●
QN15: Percentage of students who carried a weapon such as a gun, knife, or club on school property on one or more of the past 30 days	2.3 (1.7-3.2)	1,804	10.1 (6.1-16.4)	170		●	
QN16: Percentage of students who did not go to school on one or more of the past 30 days because they felt they would be unsafe at school or on their way to or from school	4.6 (3.6-5.9)	1,816	16.8 (11.9-23.2)	177		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QN17: Percentage of students who had been threatened or injured with a weapon such as a gun, knife, or club on school property one or more times during the past 12 months	5.9 (4.7-7.3)	1,817	15.9 (10.3-23.8)	176		●	
QN20: Percentage of students who were in a physical fight on school property one or more times during the past 12 months	9.6 (8.2-11.3)	1,792	18.6 (12.7-26.3)	176		●	
QN21: Percentage of students who had ever been physically forced to have sexual intercourse when they did not want to	6.4 (5.2-7.9)	1,801	18.9 (13.2-26.4)	170		●	
QN22: Among students who dated or went out with someone during the past 12 months, the percentage who had been physically hurt on purpose by someone they were dating or going out with one or more times during the past 12 months	7.2 (5.9-8.8)	1,255	22.8 (16.1-31.3)	134		●	
QN23: Among students who dated or went out with someone during the past 12 months, the percentage who had been forced by someone they were dating or going out with to do sexual things that they did not want to one or more times during the past 12 months	9.3 (7.7-11.2)	1,253	30.9 (23.0-40.0)	133		●	
QN24: Percentage of students who had ever been bullied on school property during the past 12 months	16.8 (14.8-19.1)	1,808	40.8 (33.0-49.0)	171		●	
QN25: Percentage of students who had ever been electronically bullied during the past 12 months	11.1 (9.6-12.9)	1,808	29.6 (22.7-37.6)	170		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey
Risk Behaviors and Sexual Identity Report

Unintentional Injuries/Violence

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN26: Percentage of students who felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	20.5 (18.5-22.6)	1,790	55.5 (48.0-62.7)	171		●	
QN27: Percentage of students who seriously considered attempting suicide during the past 12 months	13.0 (11.3-14.9)	1,801	42.0 (32.8-51.8)	168		●	
QN28: Percentage of students who made a plan about how they would attempt suicide during the past 12 months	9.5 (8.1-11.2)	1,798	37.9 (29.9-46.7)	173		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey Risk Behaviors and Sexual Identity Report

Tobacco Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN32: Percentage of students who smoked a whole cigarette for the first time before age 13 years	7.5 (6.2-9.0)	1,805	21.6 (15.6-29.1)	172		●	
QN33: Percentage of students who smoked cigarettes on one or more of the past 30 days	13.2 (11.2-15.5)	1,729	31.6 (23.4-41.1)	154		●	
QNFRDIG: Percentage of students who smoked cigarettes on 20 or more of the past 30 days	4.9 (3.8-6.5)	1,729	8.5 (4.7-15.0)	154			●
QN35: Among students who were less than 18 years of age and who reported current cigarette use, the percentage who usually got their own cigarettes by buying them in a store or gas station during the past 30 days	14.9 (10.4-20.8)	197	14.6 (6.4-30.0)	46			●
QN39: Percentage of students who used chewing tobacco, snuff, or dip on one or more of the past 30 days	6.4 (5.3-7.8)	1,809	13.5 (8.4-21.0)	172		●	
QN40: Percentage of students who smoked cigars, cigarillos, or little cigars on one or more of the past 30 days	11.7 (10.0-13.8)	1,802	22.6 (16.3-30.3)	172		●	
QNANYTOB: Percentage of students who smoked cigarettes or cigars or used chewing tobacco, snuff, or dip on one or more of the past 30 days	17.4 (15.2-19.9)	1,718	33.7 (25.2-43.5)	151		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN41: Percentage of students who had at least one drink of alcohol on one or more days during their life	61.5 (58.5-64.5)	1,758	78.3 (68.8-85.5)	164		●	
QN42: Percentage of students who had their first drink of alcohol other than a few sips before age 13 years	17.2 (15.1-19.6)	1,791	32.4 (24.2-42.0)	170		●	
QN43: Percentage of students who had at least one drink of alcohol on one or more of the past 30 days	31.9 (29.2-34.8)	1,742	52.6 (43.0-62.0)	154		●	
QN44: Percentage of students who had five or more drinks of alcohol in a row, that is, within a couple of hours, on one or more of the past 30 days	18.0 (15.9-20.2)	1,766	26.3 (19.2-35.0)	159		●	
QN47: Percentage of students who used marijuana one or more times during their life	29.0 (26.0-32.1)	1,775	48.7 (39.4-58.0)	160		●	
QN48: Percentage of students who tried marijuana for the first time before age 13 years	5.9 (4.6-7.5)	1,782	13.3 (8.1-21.1)	161		●	
QN49: Percentage of students who used marijuana one or more times during the past 30 days	14.3 (12.4-16.4)	1,778	28.5 (21.1-37.4)	159		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN50: Percentage of students who used any form of cocaine, including powder, crack, or freebase one or more times during their life	4.5 (3.4-6.0)	1,802	15.9 (9.9-24.6)	173		●	
QN51: Percentage of students who sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times during their life	6.6 (5.5-8.1)	1,802	30.6 (22.9-39.7)	173		●	
QN52: Percentage of students who used heroin one or more times during their life	2.4 (1.6-3.4)	1,794	13.5 (8.4-21.1)	174		●	
QN53: Percentage of students who used methamphetamines one or more times during their life	3.1 (2.3-4.3)	1,803	14.6 (9.8-21.2)	175		●	
QN54: Percentage of students who used ecstasy one or more times during their life	5.5 (4.3-7.0)	1,805	23.8 (16.4-33.2)	172		●	
QN55: Percentage of students who took steroid pills or shots without a doctor's prescription one or more times during their life	2.7 (1.9-4.0)	1,800	14.5 (8.4-23.7)	170		●	
QN56: Percentage of students who have taken a prescription drug (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax) without a doctor's prescription one or more times during their life	13.4 (11.5-15.4)	1,807	33.6 (25.9-42.2)	174		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey
Risk Behaviors and Sexual Identity Report

Alcohol/Other Drug Use

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN57: Percentage of students who used a needle to inject any illegal drug into their body one or more times during their life	2.2 (1.6-3.2)	1,800	13.9 (8.3-22.3)	172		●	
QN58: Percentage of students who were offered, sold, or given an illegal drug by someone on school property during the past 12 months	22.9 (21.0-25.0)	1,803	33.7 (26.3-42.0)	174		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN59: Percentage of students who ever had sexual intercourse	38.3 (35.0-41.7)	1,636	57.6 (47.9-66.6)	141		●	
QN60: Percentage of students who had sexual intercourse for the first time before age 13 years	3.9 (2.9-5.2)	1,634	9.9 (5.8-16.4)	141		●	
QN61: Percentage of students who had sexual intercourse with four or more people during their life	8.9 (7.2-10.8)	1,635	16.9 (11.2-24.8)	142		●	
QN62: Percentage of students who had sexual intercourse with one or more people during the past three months	25.4 (22.7-28.3)	1,641	36.9 (28.4-46.2)	144		●	
QN63: Among students who had sexual intercourse during the past three months, the percentage who drank alcohol or used drugs before last sexual intercourse	21.7 (17.9-26.1)	391	29.4 (17.4-45.3)	55			●
QN64: Among students who had sexual intercourse during the past three months, the percentage who did not use a condom during last sexual intercourse	40.9 (36.0-46.1)	388	50.1 (36.3-63.9)	55			●
QN65: Among students who had sexual intercourse during the past three months, the percentage who did not use birth control pills to prevent pregnancy before last sexual intercourse	76.2 (69.8-81.5)	375	71.3 (56.4-82.6)	55			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey Risk Behaviors and Sexual Identity Report

Sexual Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNOTHH: Among students who had sexual intercourse during the past three months, the percentage who did not use an IUD or implant or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	93.2 (90.0-95.5)	375	92.1 (80.5-97.1)	55			●
QNOTHHPL: Among students who had sexual intercourse during the past three months, the percentage who did not use birth control pills, an IUD or implant, or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	69.4 (63.1-75.0)	375	63.4 (48.1-76.3)	55			●
QNDUALBC: Among students who had sexual intercourse during the past three months, the percentage who did not use both a condom during last sexual intercourse and birth control pills, an IUD or implant, or a shot, patch, or birth control ring to prevent pregnancy before last sexual intercourse	89.4 (84.7-92.8)	375	87.7 (75.8-94.2)	54			●
QNBCNONE: Among students who had sexual intercourse during the past three months, the percentage who used no method of birth control to prevent pregnancy before last sexual intercourse	9.4 (6.6-13.4)	375	19.4 (9.8-34.7)	55			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QNOWT: Percentage of students who were overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex) [§]	13.6 (12.0-15.5)	1,691	16.8 (11.7-23.5)	155			●
QNOBESE: Percentage of students who were obese (i.e., at or above the 95th percentile for body mass index, by age and sex) [§]	9.2 (7.7-11.0)	1,691	10.1 (6.0-16.6)	155			●
QN66: Percentage of students who described themselves as slightly or very overweight	27.4 (25.4-29.6)	1,799	38.1 (29.9-47.1)	173		●	
QN71: Percentage of students who did not drink 100% fruit juices during the past seven days	25.3 (23.2-27.7)	1,773	21.7 (15.8-29.2)	169			●
QN72: Percentage of students who did not eat fruit during the past seven days	13.5 (11.4-15.9)	1,790	22.7 (17.1-29.6)	171		●	
QN73: Percentage of students who did not eat green salad during the past seven days	37.4 (34.8-40.0)	1,788	34.8 (27.2-43.2)	171			●
QN74: Percentage of students who did not eat potatoes during the past seven days	30.3 (28.2-32.5)	1,782	35.3 (27.7-43.8)	172			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

[§]Based on reference data from the 2000 CDC Growth Charts.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QN75: Percentage of students who did not eat carrots during the past seven days	55.3 (52.6-58.0)	1,788	52.8 (44.6-60.8)	171			●
QN76: Percentage of students who did not eat other vegetables during the past seven days	15.5 (13.5-17.7)	1,782	20.7 (14.9-27.9)	170			●
QNFRVG: Percentage of students who ate fruits or drank 100% fruit juices or ate vegetables less than five times per day during the past seven days	81.6 (79.1-83.8)	1,744	80.0 (72.4-85.8)	167			●
QNFRUIT: Percentage of students who ate fruits or drank 100% fruit juices less than two times per day during the past seven days	69.1 (66.6-71.4)	1,765	71.5 (64.0-77.9)	168			●
QNFR0: Percentage of students who ate fruits or drank 100% fruit juices 0 times per day during the past seven days	6.9 (5.6-8.4)	1,765	7.4 (4.0-13.2)	168			●
QNFR1: Percentage of students who ate fruits or drank 100% fruit juices less than one time per day during the past seven days	42.3 (39.9-44.9)	1,765	48.1 (40.7-55.5)	168			●
QNFR3: Percentage of students who ate fruits or drank 100% fruit juices less than three times per day during the past seven days	82.2 (80.0-84.1)	1,765	82.2 (74.8-87.7)	168			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey
Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNVEG: Percentage of students who ate vegetables less than three times per day during the past seven days	87.6 (85.6-89.4)	1,770	81.6 (73.9-87.5)	170			●
QNVEG0: Percentage of students who ate vegetables 0 times per day during the past seven days	5.6 (4.5-7.0)	1,770	7.4 (4.5-12.1)	170			●
QNVEG1: Percentage of students who ate vegetables less than one time per day during the past seven days	36.3 (33.5-39.2)	1,770	36.0 (28.5-44.3)	170			●
QNVEG2: Percentage of students who ate vegetables less than two times per day during the past seven days	73.5 (70.8-76.0)	1,770	62.5 (53.8-70.4)	170	●		
QNFRVG2: Percentage of students who ate fruits or drank 100% fruit juices less than two times per day and ate vegetables less than three times per day during the past seven days	91.1 (89.4-92.6)	1,744	89.7 (83.4-93.8)	167			●
QN77: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day during the past seven days	19.5 (17.4-21.8)	1,760	25.0 (19.0-32.1)	167			●
QNSODA0: Percentage of students who drank a can, bottle, or glass of soda or pop during the past seven days	71.2 (68.2-74.0)	1,760	74.9 (67.2-81.4)	167			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey
Risk Behaviors and Sexual Identity Report

Dietary Behaviors

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day during the past seven days	12.0 (10.3-13.9)	1,760	13.3 (8.8-19.7)	167			●
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day during the past seven days	6.6 (5.3-8.3)	1,760	6.9 (3.7-12.6)	167			●
QN79: Percentage of students who did not eat breakfast on all of the past seven days	63.6 (60.6-66.5)	1,782	71.1 (63.8-77.4)	170		●	
QNNOBKFT: Percentage of students who ate breakfast on none of the past seven days	14.0 (12.4-15.8)	1,782	19.3 (13.1-27.6)	170			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey
Risk Behaviors and Sexual Identity Report

Physical Activity

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN80: Percentage of students who were physically active for a total of at least 60 minutes per day less than five of the past seven days	57.1 (54.1-60.0)	1,771	73.1 (64.7-80.0)	171		●	
QNPA0DAY: Percentage of students who were physically active for a total of at least 60 minutes per day on 0 of the past seven days	17.4 (15.2-19.9)	1,771	22.7 (16.8-30.0)	171			●
QNPA7DAY: Percentage of students who were physically active for a total of at least 60 minutes per day on less than seven of the past seven days	76.0 (73.4-78.4)	1,771	85.7 (78.5-90.7)	171		●	
QN81: Percentage of students who watched three or more hours per day of TV on an average school day	31.1 (28.3-34.1)	1,761	30.7 (23.3-39.2)	169			●
QN82: Percentage of students who played video or computer games or used a computer for something that was not school work three or more hours per day on an average school day	32.1 (29.4-34.9)	1,772	41.5 (33.7-49.8)	169		●	
QN83: Percentage of students who did not attend physical education (PE) classes in an average week when they were in school	68.3 (64.1-72.3)	1,758	62.3 (53.2-70.6)	167			●
QNDLYPE: Percentage of students who attended physical education (PE) classes less than daily in an average week when they were in school	72.6 (68.5-76.4)	1,758	74.4 (65.4-81.7)	167			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey
Risk Behaviors and Sexual Identity Report

Other

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN85: Percentage of students who were not sure if they had ever been taught or who had not ever been taught in school about AIDS or HIV infection	12.7 (10.9-14.7)	1,753	18.5 (12.6-26.3)	170			●
QN86: Percentage of students who had ever been told by a doctor or nurse that they had asthma	25.1 (23.1-27.2)	1,756	28.6 (22.0-36.3)	167			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QN87: Percentage of students who tried or used tobacco products (for example, cigarettes, cigars, cigarillos, Black and Milds, little cigars, or smokeless tobacco) for the first time during the past 12 months	19.8 (17.5-22.3)	1,752	38.5 (30.2-47.5)	164		●	
QN89: Percentage of students who used some flavored tobacco product during the past 30 days	13.0 (11.2-15.2)	1,717	21.0 (15.0-28.7)	159		●	
QN90: Percentage of students who used only flavored cigars, cigarillos, little cigars, or only flavored smokeless tobacco products, or both flavored cigar products and flavored smokeless tobacco products during the past 30 days	13.0 (11.1-15.1)	1,717	19.8 (14.1-27.1)	158		●	
QN91: Percentage of students who have smoked a 100 or more cigarettes (5 or more packs) during their life	6.6 (5.2-8.4)	1,778	10.8 (6.6-17.1)	171			●
QN92: Among students who bought or tried to buy cigarettes in a store during past 30 days, the percentage who were not asked to show proof of age	49.8 (39.1-60.6)	107	-	29			
QN94: Percentage of students who usually smoked menthol cigarettes during the past 30 days	7.6 (6.1-9.4)	1,745	16.1 (10.5-24.0)	161		●	
QN95: Percentage of students who usually smoked cigarillos or small cigars, blunts, little cigars that look like brown cigarettes, or some other type of cigar during the past 30 days	10.9 (9.1-13.1)	1,781	20.6 (13.9-29.4)	167		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

- = Fewer than 30 students in this subgroup.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [‡]
	% 95% CI*	N	% 95% CI	N			
QN96: Percentage of students who live with someone who now smokes cigarettes or cigars	42.1 (39.1-45.1)	1,802	52.1 (43.6-60.4)	168		●	
QN97: Percentage of students who describe the rule about smoking inside the home where they live is that it is always allowed inside their home or allowed only at some times or in some places	17.5 (15.4-19.7)	1,800	27.5 (20.2-36.3)	169		●	
QN98: Percentage of students who were in the same room with someone who was smoking cigarettes on one or more of the past 7 days	36.7 (34.0-39.4)	1,805	49.7 (41.6-57.9)	172		●	
QN99: Among students who drank alcohol during the past 30 days and who have a usual type of alcohol that they drink, the percentage who reported liquor, such as vodka, rum, scotch, bourbon, or whisky, as their type of alcohol they usually drank during the past 30 days	50.0 (45.7-54.3)	556	53.1 (41.1-64.8)	85			●
QN100: Percentage of students who took a prescription drug without a doctor's prescription one or more times during the past 30 days	6.2 (5.1-7.5)	1,803	26.7 (20.0-34.8)	173		●	
QN102: Percentage of students who spend less than 20 minutes actually exercising or playing sports during an average physical (PE) class or who do not take PE	67.9 (63.7-71.9)	1,763	68.9 (59.5-77.0)	165			●
QN103: Percentage of students who described their health in general as good, fair, or poor	55.2 (52.5-57.9)	1,759	74.4 (67.5-80.3)	167		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN104: Percentage of students who have had asthma symptoms during the past 12 months	15.0 (13.2-16.9)	1,763	18.2 (12.6-25.5)	166			●
QN105: Percentage of students who used an indoor tanning device such as a sunlamp, sunbed, or tanning booth one or more times during the past 12 months	7.7 (6.3-9.4)	1,763	19.4 (13.6-27.1)	170		●	
QN106: Percentage of students who say definitely yes or probably yes that young people who smoke have more friends	31.4 (28.7-34.3)	1,741	47.5 (39.3-55.8)	165		●	
QN107: Percentage of students who say definitely yes or probably yes that smoking makes young people look cool or fit in	19.2 (17.1-21.4)	1,748	31.8 (25.6-38.8)	167		●	
QN108: Percentage of students who say definitely not or probably not that smokers have shorter lives than non-smokers	8.1 (6.7-9.8)	1,752	16.2 (10.1-25.1)	167		●	
QN109: Percentage of students who were not taught or who were not sure if they were taught in any of their classes about the dangers of tobacco use during the last school year	26.3 (22.5-30.6)	1,742	24.4 (17.5-33.0)	165			●
QN110: Percentage of students who responded that their parents would feel it would be a little wrong, not at all wrong, or not sure for them to drink beer, wine, or hard liquor (such as vodka, whiskey, or gin) at least once or twice a month	33.4 (30.6-36.2)	1,751	32.5 (25.6-40.2)	167			●

*Confidence interval.

[†]Based on t-test analyses, p<0.05.

2013 YOUTH RISK BEHAVIOR SURVEY RESULTS

Saint Mary's County, MD High School Survey Risk Behaviors and Sexual Identity Report

Site-Added

Health Risk Behavior	Heterosexual		Gay, Lesbian, or Bisexual		Heterosexual More Likely Than Gay, Lesbian, or Bisexual [†]	Gay, Lesbian, or Bisexual More Likely Than Heterosexual [†]	No Difference [†]
	% 95% CI*	N	% 95% CI	N			
QN111: Percentage of students who think people are at no risk or slight risk of harming themselves (physically or in other ways) if they have one or two drinks of alcohol (beer, wine, or liquor) nearly every day	38.9 (36.5-41.4)	1,743	39.3 (31.6-47.5)	166			●
QN112: Percentage of students who think people are at no risk or slight risk of harming themselves (physically or in other ways) if they have five or more drinks of alcohol (beer, wine, or liquor) once or twice a week	25.0 (22.4-27.9)	1,737	28.0 (20.5-37.0)	166			●
QN113: Percentage of students who do not have an adult outside of school they can talk to about things that are important to them	12.9 (11.4-14.6)	1,718	28.5 (21.4-36.8)	159		●	
QN114: Percentage of students who would feel comfortable seeking help from 0 adults besides their parents if they had an important question affecting their life	19.2 (17.3-21.4)	1,742	26.6 (20.5-33.7)	168		●	
QN116: Percentage of students who strongly disagree, disagree, or are not sure that their teachers really care about them and give them a lot of encouragement	44.1 (41.5-46.8)	1,745	60.6 (52.0-68.5)	167		●	
QN117: Percentage of students who do not participate in any extracurricular activities at school such as sports, band, drama, clubs, or student government	36.8 (33.5-40.2)	1,725	45.9 (37.8-54.2)	159		●	

*Confidence interval.

[†]Based on t-test analyses, p<0.05.