

Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to MDH for the week ending October 21, 2017

Prepared by the Infectious Disease Epidemiology and Outbreak Response Bureau Prevention and Health Promotion Administration Maryland Department of Health

The data presented in this document are provisional and subject to change as additional reports are received.

SUMMARY


During the week ending October 21, 2017, influenza-like illness (ILI) intensity in Maryland was **MINIMAL** and there was **SPORADIC** geographic activity. The proportion of outpatient visits for ILI reported by Sentinel Providers and Maryland Emergency Departments was low. The proportion of MRITS respondents reporting ILI was also low. Clinical laboratories reported a low number of specimens testing positive for influenza. There were 7 specimens that tested positive for influenza at the MDH lab. There were 5 influenza-associated hospitalizations and 1 respiratory outbreak was reported to MDH.

ILI Intensity Levels	Influenza Geographic Activity	
🖌 Minimal	No Activity	
Low	✔ Sporadic	
Moderate	Local	
High	Regional	
	Widespread	

Click here to visit our influenza surveillance web page

ILINet Sentinel Providers


31 sentinel providers reported a total of 5,647 visits this week. Of those, 69 (1.2%) were visits for ILI. This is below the Maryland baseline of 2.0%.



ILI Visits To Sentinel Providers By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	21 (30%)	19 (23%)	58 (26%)
Age 5-24	21 (30%)	30 (36%)	79 (35%)
Age 25-49	24 (35%)	21 (25%)	56 (25%)
Age 50-64	2 (3%)	10 (12%)	24 (11%)
Age ≥ 65	1 (1%)	4 (5%)	7 (3%)
Total	69 (100%)	84 (100%)	224 (100%)

Visits to Emergency Departments for ILI

Emergency Departments in Maryland reported a total of 39,976 visits this week through the ESSENCE surveillance system. Of those, 511 (1.3%) were visits for ILL.



ILI Visits To Emergency Departments By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4	162 (32%)	174 (29%)	498 (30%)
Age 5-24	129 (25%)	163 (27%)	431 (26%)
Age 25-49	124 (24%)	142 (24%)	427 (26%)
Age 50-64	54 (11%)	68 (11%)	185 (11%)
Age ≥ 65	42 (8%)	46 (8%)	129 (8%)
Total	511 (100%)	593 (100%)	1670 (100%)

 Neighboring states' influenza information:

 Delaware
 http://dhs.delaware.gov/dph/epi/influenzahome.html

 District of Columbia
 http://doh.dc.gov/service/influenza

 Pennsylvania
 http://www.health.pa.gov/My%20Health/Diseases%20and%20Conditions/I-L/Pages/Influenza.aspx#.V-LtaPkrJD8

 Virginia
 http://www.vdh.virginia.gov/epidemiology/influenza-flu-in-virginia/influenza-surveillance/


 West Virginia
 http://dhr.wv.gov/oeps/disease/flu/Pages/fluSurveillance.aspx

Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to MDH for the week ending October 21, 2017

Community-based Influenza Surveillance (MRITS)


MRITS is the Maryland Resident Influenza Tracking System, a weekly survey for influenza-like illness (ILI). A total of 543 residents responded to the <u>MRITS survey</u> this week. Of those, 2 (0.4%) reported having ILI and missing a cumulative 3 days of regular daily activities.




MRITS Respondents Reporting ILI By Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4			1 (14%)
Age 5-24	1 (50%)	1 (50%)	2 (29%)
Age 25-49			
Age 50-64		1 (50%)	2 (29%)
Age ≥ 65	1 (50%)		2 (29%)
Total	2 (100%)	2 (100%)	7 (100%)

Clinical Laboratory Influenza Testing

40 clinical laboratories reported performing 741 influenza diagnostic tests, mostly rapid influenza diagnostic tests (RIDTs). Of those, 18 (2.4%) were positive for influenza. Of those testing positive, 8 (44%) were influenza Type A and 10 (56%) were influenza Type B. The <u>reliability of RIDTs</u> depends largely on the conditions under which they are used. False-positive (and true-negative) results are more likely to occur when the disease prevalence in the community is low, which is generally at the beginning and end of the influenza season and during the summer.




Positive Rapid Flu Tests by Type	This Week Number (%)	Last Week Number (%)	Season Number (%)
Туре А	8 (44%)	13 (54%)	28 (49%)
Туре В	10 (56%)	11 (46%)	29 (51%)
Total	18 (100%)	24 (100%)	57 (100%)



State Laboratories Administration Influenza Testing

The MDH Laboratories Administration performed a total of 78 PCR tests for influenza and 7 (9.0%) specimens tested positive. Six of the specimens were Type A (H3) and 1 was Type B (Yamagata). PCR testing is more reliable than RIDT. The MDH testing identifies subtypes of influenza A and lineages of influenza B, information that is not available from the RIDT results. The table below summarizes results by type, subtype, and lineage.



Positive PCR Tests by Type (Subtype)	This Week Number (%)	Last Week Number (%)	Season Number (%)
Type A (H1)			
Туре А (НЗ)	6 (86%)	3 (100%)	12 (71%)
Type B (Victoria)			
Type B (Yamagata)	1 (14%)		2 (12%)
Type A (H3N2v)			3 (18%)
Total	7 (100%)	3 (100%)	17 (100%)

Where to get an influenza vaccination


Interested in getting a flu vaccine for the 2017-18 influenza season? Go to <u>https://phpa.health.maryland.gov/influenza/Pages/getvaccinated.aspx</u> and click on your county/city of residence. You will be redirected to your local health department website for local information on where to get your flu vaccine.

Maryland Weekly Influenza Surveillance Activity Report

A summary of influenza surveillance indicators reported to MDH for the week ending October 21, 2017

Influenza-associated Hospitalizations


A total of 5 influenza-associated hospitalizations were reported this week. (A person with an overnight hospital stay along with a positive influenza test of any kind, e.g., RIDT or PCR, is considered an "influenza-associated hospitalization" for purposes of influenza surveillance.)



Influenza- Associated Hospitalizations by Age Group	This Week Number (%)	Last Week Number (%)	Season Number (%)
Age 0-4			1 (5%)
Age 5-17			
Age 18-24		1 (17%)	1 (5%)
Age 25-49	2 (40%)		3 (14%)
Age 50-64		2 (33%)	5 (24%)
Age ≥ 65	3 (60%)	3 (50%)	11 (52%)
Total	5 (100%)	6 (100%)	21 (100%)

Outbreaks of Respiratory Disease

There was 1 respiratory outbreak reported to MDH this week. (Disease outbreaks of any kind are reportable in Maryland. Respiratory outbreaks may be reclassified once a causative agent is detected, e.g., from ILI to influenza.)



Respiratory Outbreaks by Type	This Week Number (%)	Last Week Number (%)	Season Number (%)
Influenza		1 (100%)	1 (20%)
Influenza-like Illness	1 (100%)		2 (40%)
Pneumonia			2 (40%)
Other Respiratory			
Total	1 (100%)	1 (100%)	5 (100%)

National Influenza Surveillance (CDC)

During week 42 (October 15-21, 2017), influenza activity was low in the United States.

- <u>Viral Surveillance:</u> The most frequently identified influenza virus type reported by public health laboratories during week 42 was influenza A. The percentage of respiratory specimens testing positive for influenza in clinical laboratories is low.
- O <u>Novel Influenza A Virus:</u> One human infection with a novel influenza A virus was reported.
- Pneumonia and Influenza Mortality: The proportion of deaths attributed to pneumonia and influenza (P&I) was below the system-specific epidemic threshold in the National Center for Health Statistics (NCHS) Mortality Surveillance System.
- o Influenza-associated Pediatric Deaths: One influenza-associated pediatric death was reported.
- Outpatient Illness Surveillance: The proportion of outpatient visits for influenza-like illness (ILI) was 1.3%, which is below the national baseline of 2.2%. All 10 regions reported ILI below region-specific baseline levels. Three states experienced low ILI activity and New York City, the District of Columbia, Puerto Rico and 47 states experienced minimal ILI activity.
- Geographic Spread of Influenza: The geographic spread of influenza in Guam was reported as regional; Puerto Rico and 12 states reported local activity; the District of Columbia and 33 states reported sporadic activity; five states reported no activity; and the U.S. Virgin Islands did not report.

